
From the Pythagorean Dream to the Fermatian Obstruction: Symbolic Representation of $h = \sqrt[p]{a^p + b^p}$ via an Identity Derived from Nicomachus' Cumulative Sum

[Ibar Federico Anderson](#)*

Posted Date: 27 February 2026

doi: 10.20944/preprints202602.0367.v3

Keywords: Universal Anderson–Faulhaber–Bernoulli Identity; finite difference of power; internal structure; complexity $\mathcal{O}(p)$; symbolic representation; arithmetic obstruction; Fermat's Last Theorem; discrete calculus; Bernoulli numbers; combinatorial uniqueness; structural gradation; pedagogical value



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

From the Pythagorean Dream to the Fermatian Obstruction: Symbolic Representation of $h = \sqrt[p]{a^p + b^p}$ via an Identity Derived from Nicomachus' Cumulative Sum

Ibar Federico Anderson

Laboratory of the Department of Industrial Design (LIDDI), Universidad Nacional de La Plata, Argentina; ianderson@empleados.fba.unlp.edu.ar

Abstract

This paper presents, in a unified form and with explicit logical justification, a chain of original results on the internal structure of perfect powers and their connection with Fermat's Last Theorem (FLT). Starting from the historical formula of Nicomachus of Gerasa (c. 100 AD) for the cumulative sum of cubes, $S_3(n) = T_n^2$, and applying the backward finite-difference operator ∇ —formalised by Taylor (1715) and systematised by Boole (1860)—the Anderson Identity (2026) is derived:

$$n^3 = \frac{n^2}{4} [(n+1)^2 - (n-1)^2] = T_n^2 - T_{n-1}^2.$$

This identity, valid exclusively for cubes, is extended to every $p \geq 1$ through the Universal Anderson–Faulhaber–Bernoulli Identity (2026):

$$n^p = \frac{1}{p+1} \sum_{j=0}^p \binom{p+1}{j} B_j^+ \delta_{p+1-j}(n), \quad \delta_m(n) := n^m - (n-1)^m, \text{ (UI)}$$

derived by applying the operator ∇ to the classical Faulhaber–Bernoulli formula for cumulative sums of powers. The quantity $\delta_m(n) = n^m - (n-1)^m$ —the finite difference of the individual m -th power—constitutes the original internal perspective of this work: it reorients the historical cumulative-sum formula toward individual powers, revealing that the internal algebraic complexity of n^p grows as $C(p) = \lfloor p/2 \rfloor + 1$, with $p = 3$ as the unique point of optimal compactness (pure monomial). The second contribution is the Universal Symbolic Representation (2026):

$$h = \sqrt[p]{I_p(a) + I_p(b)}, \quad h \notin \mathbb{Z} \forall p \geq 3, \text{ (SR)}$$

which expresses $h = \sqrt[p]{a^p + b^p}$ through purely integer operations in the radicand and establishes that its irrationality for $p \geq 3$ is a structurally inevitable consequence, not an accidental one. Full step-by-step derivations, explicit expansions for $p = 2, \dots, 8$, 50-digit-precision numerical verifications for 10,000 pairs (a, b) with $1 \leq a \leq b \leq 100$, and the conceptual gradation of the Fermatian obstruction in three regimes—quadratic, cubic, and Bernoulli—are presented. The Structural Stratification Theorem is proved: $C(p) = \lfloor p/2 \rfloor + 1$, with $p = 3$ as the unique point of optimal compactness. Complete chronological historical contextualisation from the Pythagoreans to Wiles, analysis of the originality of the present perspective, generalised symmetry breaking, and genuine pedagogical value are included.

Keywords: Universal Anderson–Faulhaber–Bernoulli Identity; finite difference of power; internal structure; complexity $C(p)$; symbolic representation; arithmetic obstruction; Fermat’s Last Theorem; discrete calculus; Bernoulli numbers; combinatorial uniqueness; structural gradation; pedagogical value

MSC 2020: 11D41; 11B83; 39A70; 11B68; 11A07

1. Introduction

1.1. The Pythagorean Legacy and Quadratic Harmony

The Pythagorean school, founded in Croton (Magna Graecia) around the 6th century BC, established one of the first philosophical–mathematical syntheses in Western history: the belief that “all is number” ($\pi\acute{\alpha}\nu\tau\alpha \acute{\alpha}\rho\iota\theta\mu\acute{o}\varsigma$). For the Pythagoreans, numbers were not mere instruments of calculation but ontological principles that structured cosmic reality. This worldview found its purest expression in the figurate numbers: geometric representations of integers through regular point configurations [11]. The result known today as the Pythagorean theorem—although probably known to Mesopotamian and Egyptian cultures earlier—crystallised this harmony between geometry and arithmetic:

$$h^2 = a^2 + b^2, (1)$$

where h denotes the hypotenuse of a right triangle and a , b its legs. Euclid, in his *Elements* (Book X, Proposition 29, Lemma 1, c. 300 BC), provided the complete parametrisation of primitive Pythagorean triples:

$$a = m^2 - n^2, \quad b = 2mn, \quad h = m^2 + n^2, \quad m > n > 0, \quad \gcd(m, n) = 1, \quad m \not\equiv n \pmod{2}. (2)$$

The abundance of integer solutions in (1) is not accidental: it reflects the fact that the internal structure of n^2 is $\nabla S_2(n) = 2n - 1$, an arithmetic progression flexible enough to admit infinitely many integer combinations.

The Pythagoreans also discovered the figurate numbers—triangular, square, pentagonal— and systematised the study of the integer triples satisfying (1). The crisis of incommensurables—the discovery of $\sqrt{2}$ as irrational, attributed to Hippasus of Metapontum (5th century BC)—fractured the faith in the universality of integers and led to the theory of proportions of Eudoxus (c. 370 BC), a precursor of the modern real number [11].

1.2. The Question That Launched Centuries of Inquiry

This quadratic harmony—the existence of infinitely many integer solutions for a nonlinear Diophantine equation—has captivated generations of mathematicians and naturally suggested an apparently innocent question: if the squares of the legs sum to the square of the hypotenuse in Euclidean plane geometry, could the volumes of “cubic legs” sum to the volume of a “cubic hypotenuse” in three-dimensional space? Formally:

$$h^p = a^p + b^p, \quad h, a, b \in \mathbb{Z}^+, \quad p \geq 3 ? (3)$$

This question, formulated in full generality by Pierre de Fermat in 1637, unleashed more than three centuries of investigation. The answer—always negative—was proved partially by Euler ($p = 3$, c. 1770) and in full generality by Andrew Wiles (1994). The present work does not claim to prove the FLT, but rather to explore its structural geography from elementary discrete calculus.

1.3. Anderson’s Proposal: A Discrete Lens

The starting point is the observation that the historical formula of Nicomachus of Gerasa (c. 100 AD):

$$S_3(n) = \sum_{k=1}^n k^3 = \left(\frac{n(n+1)}{2} \right)^2 = T_n^2, (4)$$

acts as a discrete antiderivative of the sequence $f(n) = n^3$. Applying the backward finite-difference operator $\nabla f(n) := f(n) - f(n-1)$ —formalised by Taylor [16] and systematised by Boole [5]—one obtains the Anderson Identity:

$$n^3 = \frac{n^2}{4} [(n+1)^2 - (n-1)^2] = T_n^2 - T_{n-1}^2. \quad (5)$$

It is an imperative historiographic clarification that Nicomachus neither formulated nor hinted at identity (5) in terms of symmetric differences $(n+1)^2 - (n-1)^2$. His contribution was limited exclusively to the cumulative-sum formula (4). Expression (5) is an algebraic consequence derivable through the finite-difference operator ∇ , a concept developed in the 17th–19th centuries with the advent of discrete calculus [5, 7]. To attribute (5) directly to Nicomachus would constitute an inadmissible historiographic anachronism.

The questions that organise the present work are: (1) Does an identity analogous to (5) exist for $p \geq 4$? (2) What does such a generalisation tell us about the Fermatian obstruction? (3) What is the genuine originality of Anderson’s perspective?

1.4. Scope and Limitations of This Work

Explicit declaration. This work does **not** claim to prove Fermat’s Last Theorem, established definitively by Wiles in 1994 [17]. Its goal is to chart the structural geography of the Fermatian obstruction from elementary discrete calculus, offering an accessible structural map that complements—without competing with—Wiles’ deep proof based on algebraic geometry and modular forms.

2. Theoretical and Historical Framework

2.1. From Pythagoras to Nicomachus

Historical Note 2.1 (Pythagoras of Samos (c. 570–495 BC)). Born on the island of Samos during the Greek archaic period, Pythagoras founded in Croton (southern Italy) a philosophical–religious community with strict ascetic rules. The Pythagoreans believed that numbers were the ultimate essence of reality: “numbers constitute the essence of all things” (Aristotle, *Metaphysics* A, 5). This numerical worldview was manifested in fundamental mathematical discoveries: the Pythagorean theorem, the classification of even and odd numbers, perfect numbers, and figurate numbers (triangular, square, pentagonal) [11].

Historical Note 2.2 (Nicomachus of Gerasa (c. 60–120 AD)). A neo-Pythagorean philosopher born in Gerasa (present-day Jerash, Jordan), Nicomachus systematised in his *Introductio Arithmetica* the classical Greek arithmetical knowledge [14]. His work became the standard arithmetic text in the Greco-Roman world and was translated into Latin by Boethius (c. 500 AD). Unlike Euclid, Nicomachus prioritised intuitive and philosophical understanding over axiomatic rigour, emphasising the mystical and cosmological properties of numbers. Nicomachus systematised the theory of figurate numbers inherited from the Pythagoreans: triangular $T_n = n(n+1)/2$, square $Q_n = n^2$, pentagonal $P_n = n(3n-1)/2$, and hexagonal $H_n = n(2n-1)$.

Theorem 2.3 (Cumulative sum of cubes — Nicomachus, c. 100 AD) [14]. For every $n \in \mathbb{N}$:

$$S_3(n) = \sum_{k=1}^n k^3 = \left(\frac{n(n+1)}{2} \right)^2 = T_n^2 = \frac{n^2(n+1)^2}{4}. \quad (6)$$

Proof. By mathematical induction. Base case ($n = 1$): $S_3(1) = 1^3 = 1 = \frac{1^2 \cdot 1^2}{4} = 1$. *Inductive hypothesis:* valid for n . Inductive step:

$$\begin{aligned} S(n+1) &= S(n) + (n+1)^3 = \frac{n^2(n+1)^2}{4} + (n+1)^3 = (n+1)^2 + \left[\frac{n^2}{4} + (n+1) \right] \\ &= (n+1)^2 + \left[\frac{n^2 + 4n + 4}{4} \right] = \frac{(n+1)^2 \cdot (n+2)^2}{4} = T_n^2 + 1 \end{aligned}$$

Remark 2.4 (Geometric interpretation). Identity (6) possesses an elegant geometric interpretation: the sum of the first n cubes forms a perfect square whose side is the n -th triangular number $T_n = n(n+1)/2$. This dimensional transformation (sum of volumes \rightarrow square area) is unique in figurate arithmetic and anticipates deep structures of modern discrete analysis [6].

2.2. Power Sums: From Faulhaber to Bernoulli

Historical Note 2.5 (Johann Faulhaber (1580–1635)). A German mathematician born in Ulm, recognised by Kepler as one of the finest arithmeticians of his time. In his *Academia Algebrae* (1631, [8]) he computed $S_p(n) = \sum_{k=1}^n k^p$ for p up to at least 17, discovering the polynomial nature of these sums. He observed that $S_3(n) = T_n^2$ is the only case with perfect quadratic factorisation.

Historical Note 2.6 (Jakob Bernoulli (1654–1705)). Bernoulli published posthumously in *Ars Conjectandi* (1713, [4]) the general formula for $S_p(n)$ in terms of the coefficients that today bear his name. The rigorous proof for odd indices was completed by Carl Jacobi in 1834. The Bernoulli numbers B_j vanish for every odd j with $j \geq 3$; a fact fundamental to the Universal Identity.

Theorem 2.7 (Faulhaber–Bernoulli formula – Bernoulli, 1713 [4]). For every $p \geq 1$ and $n \in \mathbb{N}$:

$$S_p(n) = \sum_{k=1}^n k^p = \frac{1}{p+1} \sum_{j=0}^p \binom{p+1}{j} B_j^+ n^{p+1-j}, \quad (7)$$

where $B_0 = 1$, $B_1^+ = 1/2$, $B_2 = 1/6$, $B_3 = 0$, $B_4 = -1/30$, $B_5 = 0$, $B_6 = 1/42$, and $B_j = 0$ for every odd j with $j \geq 3$.

Relevant canonical cases:

$$S_1(n) = \frac{n(n+1)}{2} = T_n, \quad (8)$$

$$S_2(n) = \frac{n(n+1)(2n+1)}{6}, \quad (9)$$

$$S_3(n) = \left(\frac{n(n+1)}{2} \right)^2 = T_n^2. \quad (10)$$

The exceptionality of (10) is fundamental: $S_3(n)$ is the only polynomial of the family $\{S_p\}_{p \geq 1}$ that is a polynomial perfect square [6].

2.3. The Fundamental Theorem of Discrete Calculus

Historical Note 2.8 (Brook Taylor (1685–1731) and George Boole (1815–1864)). Taylor formalised the calculus of finite differences in his *Methodus Incrementorum Directa et Inversa* (1715, [16]). Boole systematised it in *A Treatise on the Calculus of Finite Differences* (1860, [5]), establishing the Fundamental Theorem of Discrete Calculus. The operator ∇ and its inverse Σ are the discrete counterparts of continuous differentiation and integration.

Definition 2.9 (Fundamental discrete operators). Let $f: \mathbb{N} \rightarrow \mathbb{R}$. Define:

$$\nabla f(n) := f(n) - f(n-1), \quad n \geq 2 \quad (\text{backward difference}), \quad (11)$$

$$\Delta f(n) := f(n+1) - f(n) \quad (\text{forward difference}), \quad (12)$$

$$\Sigma f(n) := \sum_{k=1}^n f(k) \quad (\text{cumulative sum}). \quad (13)$$

Theorem 2.10 (Fundamental Theorem of Discrete Calculus — Boole, 1860). If $S(n) = \sum_{k=1}^n f(k)$, then $\nabla S(n) = f(n)$ for every $n \geq 2$.

Proof. $S(n) - S(n-1) = [f(1) + \dots + f(n)] - [f(1) + \dots + f(n-1)] = f(n)$. \square

Remark 2.11. This result is the discrete analogue of the continuous Fundamental Theorem of Calculus $\frac{d}{dx} \int_a^x f(t) dt = f(x)$. The cumulative sum Σ acts as a “discrete integral” and ∇ as a “discrete derivative”; they are mutually inverse [9].

2.4. Symmetric Difference and Uniqueness of the Pure Monomial

Definition 2.12 (Symmetric difference of order k). For $k, n \in \mathbb{N}$:

$$D_k(n) := (n+1)^k - (n-1)^k. \quad (14)$$

Proposition 2.13 (Binomial expansion of $D_k(n)$).

$$D_k(n) = 2 \sum_{\substack{j=1 \\ j \text{ odd}}}^k \binom{k}{j} n^{k-j}. \quad (15)$$

Proof. By the binomial theorem, $(n \pm 1)^k = \sum_{j=0}^k \binom{k}{j} n^{k-j} (\pm 1)^j$. Upon subtraction, terms with even j cancel ($1 - (-1)^j = 0$) and terms with odd j double ($1 - (-1)^j = 2$). \square

Theorem 2.14 (Uniqueness of the pure monomial). $D_k(n)$ is a pure monomial—a single non-constant term—if and only if $k = 2$.

Proof. By the previous proposition the number of non-zero terms in $D_k(n)$ equals the number of odd integers $j \in [1, k]$:

- $k = 1$: only $j = 1 \Rightarrow D_1(n) = 2$ (constant, no variable).
- $k = 2$: only $j = 1 \Rightarrow D_2(n) = 2 \binom{2}{1} n = 4n$ (pure monomial in n).
- $k = 3$: $j = 1, 3$ (two odd values) $\Rightarrow D_3(n) = 6n^2 + 2$ (binomial).
- $k \geq 4$: the number of odd integers in $[1, k]$ is $\lceil k/2 \rceil \geq 2 \Rightarrow$ at least two terms.

Therefore, $k = 2$ is the only value yielding a non-constant pure monomial.

2.5. Historical Context: Fermat, Euler, Germain, Kummer, and Wiles

Historical Note 2.15 (Pierre de Fermat (1607–1665)). A French jurist and amateur mathematician. In 1637, in the margin of his copy of Diophantus’ *Arithmetica* (Bachet edition, 1621), he wrote in Latin: “*Cubum autem in duos cubos, aut quadratoquadratum in duos quadratoquadratos, et generaliter nullam in infinitum ultra quadratum potestatem in duos eiusdem nominis fas est dividere cuius rei demonstrationem mirabilem sane detexi. Hanc marginis exiguitas non caperet.*” Most historians consider that his “marvellous proof” was incomplete or erroneous, possibly based on the method of infinite descent that he did correctly apply to $x^4 + y^4 = z^2$ [15].

Historical Note 2.16 (Leonhard Euler (1707–1783)). He provided the first proof of the FLT for $p = 3$ (c. 1770) by infinite descent in the ring $\mathbb{Z}[\omega]$ of Eisenstein integers, where $\omega = e^{2\pi i/3}$. The key factorisation $x^3 + y^3 = (x+y)(x+\omega y)(x+\omega^2 y) = z^3$ requires unique factorisation in $\mathbb{Z}[\omega]$, a property that Euler assumed tacitly and that was rigorously established by Gauss and Kummer [7].

Historical Note 2.17 (Sophie Germain (1776–1831)). A self-taught French mathematician who faced insurmountable barriers to participate in the formal mathematical community due to her

gender. She adopted the male pseudonym “M. Le Blanc” to correspond with Lagrange and Gauss. After revealing her identity, Gauss wrote in 1807 [13]:

“When a person of the sex which, according to our customs and prejudices, finds infinitely more obstacles than men in familiarising herself with these thorny investigations, succeeds nonetheless in overcoming these obstacles and penetrating the most obscure parts of them, then without doubt she possesses the noblest courage, extraordinary talents, and superior genius.”

Germain developed the first systematic framework for attacking the FLT through modular congruences, verifying conditions for all primes $p < 100$.

Historical Note 2.18 (Ernst Kummer (1810–1893)). He discovered that many rings of algebraic integers lack unique factorisation. For example, in $\mathbb{Z}[\sqrt{-5}]$: $6 = 2 \cdot 3 = (1 + \sqrt{-5})(1 - \sqrt{-5})$, two distinct factorisations into irreducibles. He introduced the revolutionary concept of ideal numbers (predecessors of modern ideals) and proved the FLT for all regular primes [7].

Historical Note 2.19 (Andrew Wiles (1953–) and Richard Taylor (1962–)). Wiles proved in 1994 (published 1995,) the modularity theorem (Taniyama–Shimura conjecture). Gerhard Frey observed in 1984 that a hypothetical solution $a^p + b^p = c^p$ would generate the “non-modular” elliptic curve $y^2 = x(x - a^p)(x + b^p)$. Ken Ribet proved in 1986 that such a curve would violate Taniyama–Shimura. Wiles’ proof, spanning more than 100 pages and with a crucial correction in collaboration with Richard Taylor, unified apparently disjoint mathematical fields: algebraic geometry, representation theory, complex analysis, and number theory [17, 15].

3. Anderson’s Hypotheses

The two central hypotheses of this work are:

Hypothesis 20 (H1 – Universal Identity). *There exists an expression for n^p in terms of finite differences of individual powers $\delta_m(n) = n^m - (n - 1)^m$, deducible via ∇ from the Faulhaber–Bernoulli formula, that exhibits a complexity $C(p) = \lfloor p/2 \rfloor + 1$ increasing with p and that reveals $p = 3$ as the unique point of algebraic compactness.*

Hypothesis 21 (H2 – Symbolic Representation). *The quantity $h = \sqrt[p]{a^p + b^p}$ can be represented exactly through integer operations in the radicand using the identity of H1, and its irrationality for $p \geq 3$ is a structurally inevitable consequence of the gradation of internal complexity, not a property imposed from outside.*

Logical Chain of Deduction

The deductive sequence supporting both hypotheses can be visualised in two steps.

Step 1 – Historical root for $p = 3$:

$$S_3(n) = T_n^2 \xrightarrow{\nabla} n^3 = n^2/4 [(n+1)^2 - (n-1)^2] \xrightarrow{\text{sum}} h = \sqrt[3]{I_3(a) + I_3(b)} \xrightarrow{\text{FLT}} h \notin \mathbb{Z}.$$

Nicomachus, c. 100
particular case of H1
H2 for $p=3$

Step 2 – Generalisation to every $p \geq 1$:

$$S_p(n) = \frac{1}{p+1} \sum_j \binom{p+1}{j} B_j^+ n^{p+1-j} \xrightarrow{\nabla} n^p = I_p(n) \xrightarrow{\text{FLT}} \sqrt[p]{I_p(a) + I_p(b)} \notin \mathbb{Z}.$$

Faulhaber–Bernoulli
H1
H2

The link between Step 1 and Step 2 is algebraic: the case $p = 3$ is the only one in which the action of ∇ on $S_p(n)$ produces a pure monomial, because $S_3(n) = T_n^2$ is the only perfect square in the family $\{S_p\}$. For $p \geq 4$, $\nabla S_p(n)$ inevitably produces a polynomial with $C(p) \geq 3$ terms (Theorem 4.6).

4. Methodology

4.1. The Operator ∇ and the Fundamental Theorem

The operators ∇ , Δ , and Σ have been defined in Definition 2.4, and Theorem 2.5 establishes the essential duality $\nabla \circ \Sigma = \text{id}$.

4.2. The internal perspective: $\delta_m(n)$ versus n^{p+1-j}

The original conceptual key is the following. The Faulhaber–Bernoulli formula (7) expresses $S_p(n)$ as a polynomial in n . Applying ∇ term by term produces differences of the form $n^{p+1-j} - (n-1)^{p+1-j}$. Instead of expanding them immediately, they are named:

$$\delta_m(n) := n^m - (n-1)^m, \quad m \geq 1. \quad (16)$$

These are the finite differences of the individual m -th power. They are not differences of cumulative sums: they measure how much n^m grows when passing from $n-1$ to n . This change of perspective—from global sums to individual powers—is the original conceptual contribution.

The first values are:

$$\begin{aligned} \delta_1(n) &= 1, \\ \delta_2(n) &= 2n - 1, \\ \delta_3(n) &= 3n^2 - 3n + 1, \\ \delta_4(n) &= 4n^3 - 6n^2 + 4n - 1, \\ \delta_5(n) &= 5n^4 - 10n^3 + 10n^2 - 5n + 1. \end{aligned}$$

4.3. Derivation of the Universal Identity (H1)

Theorem 4.1 (Universal Anderson–Faulhaber–Bernoulli Identity, 2026). For every $p \geq 1$ and $n \in \mathbb{N}$:

$$n^p = \frac{1}{p+1} \sum_{j=0}^p \binom{p+1}{j} B_j^+ \delta_{p+1-j}(n), \quad \delta_m(n) = n^m - (n-1)^m. \quad (17)$$

Proof. We apply Theorem 2.5 to formula (7):

$$\begin{aligned} n^p &= S_p(n) - S_p(n-1) \\ &= \frac{1}{p+1} \sum_{j=0}^p \binom{p+1}{j} B_j^+ [n^{p+1-j} - (n-1)^{p+1-j}], \\ &= \frac{1}{p+1} \sum_{j=0}^p \binom{p+1}{j} B_j^+ \delta_{p+1-j}(n). \end{aligned}$$

□

Remark 4.2 (What changes with respect to Faulhaber–Bernoulli). Formula (7) says something about $S_p(n)$: cumulative sums of powers. Identity (20) says something about n^p itself: the individual power. The move is to apply ∇ to extract the internal structure of the individual power from its cumulative sum. This does not modify any known mathematical result: it reformulates the same algebra from a new and more informative perspective.

4.4. Special Formulation: Anderson Identity for $p = 3$

Theorem 4.3 (Anderson Identity – original derivation (2026) from Nicomachus). For every $n \in \mathbb{N}$:

$$n^3 = \frac{n^2}{4} [(n+1)^2 - (n-1)^2] = T_n^2 - T_{n-1}^2. \quad (18)$$

Proof. Applying Theorem 2.5 to (6):

$$n^3 = \nabla S_3(n) = \frac{n^2(n+1)^2}{4} - \frac{(n-1)^2n^2}{4} = \frac{n^2}{4} [(n+1)^2 - (n-1)^2] = \frac{n^2}{4} (4n) = n^3$$

Remark 4.4 (What Anderson contributes beyond Nicomachus). The Nicomachus formula is $S_3(n) = T_n^2$: a property of the cumulative sum. The Anderson Identity (21) is a property of the individual power n^3 : it expresses each cube as the difference of two consecutive triangular squares, exhibiting the adjacent symmetry $(n-1, n, n+1)$ inherent in the structure of n^3 . This change of perspective—from cumulative sums to individual powers—is the original conceptual move of this work. The key factor is $D_2(n) = (n+1)^2 - (n-1)^2 = 4n$: a pure monomial (Table 1), which allows exact cancellation and the compactness of (21). For no other power n^p with $p \neq 3$ does an analogously compact representation exist, because $D_k(n)$ is a pure monomial only for $k = 2$ (Theorem 2.9).

Table 1. Structure of $D_k(n) = (n+1)^k - (n-1)^k$ for $k = 1, \dots, 6$.

k	$D_k(n)$	Type
1	2	Constant
2	$4n$	Pure monomial (unique)
3	$6n^2 + 2$	Binomial
4	$8n^3 + 8n$	Binomial
5	$10n^4 + 20n^2 + 2$	Trinomial
6	$12n^5 + 40n^3 + 12n$	Trinomial

4.5. Structural Stratification Theorem (H1, Continued)

Definition 4.5 (Internal structural complexity). $C(p)$ is the number of indices $j \in \{0, 1, \dots, p\}$ that are active in (20), i.e. those with $B_j^+ \neq 0$.

The Bernoulli numbers satisfy $B_j = 0$ for every odd j with $j \geq 3$. The active indices are $j \in \{0, 1, 2, 4, 6, \dots, 2\lfloor p/2 \rfloor\}$: exactly $\lfloor p/2 \rfloor + 1$ values.

Theorem 4.6 (Structural Stratification – Anderson, 2026).

$$C(p) = \left\lfloor \frac{p}{2} \right\rfloor + 1. \quad (19)$$

$C(p)$ grows monotonically. $p = 3$ is the unique $p \geq 2$ for which (20) reduces to a pure monomial.

Proof. The count of active indices gives (22). The reduction to a pure monomial for $p = 3$ is a consequence of the fact that $S_3(n) = T_n^2$: upon computing ∇T_n^2 , all terms of degree $\neq 3$ cancel exactly, producing pure n^3 . No other $S_p(n)$ is a polynomial perfect square [6], so no other p possesses this cancellative property.

Corollary 4.7 (Absolute uniqueness). The identity $n^3 = \frac{n^2}{4} [(n+1)^2 - (n-1)^2]$ is the only case of (20) that reduces to a pure monomial in n .

4.6. Explicit Expansion Of (20) for $p = 2, \dots, 8$

We use $B_0 = 1$, $B_1^+ = 1/2$, $B_2 = 1/6$, $B_3 = 0$, $B_4 = -1/30$, $B_5 = 0$, $B_6 = 1/42$.
 $p = 2$ — **quadratic regime**, $C(2) = 2$:

$$n^2 = \frac{1}{3} \left(\delta_3(n) + \frac{3}{2} \delta_2(n) \right) = \frac{3n^2 - 3n + 1}{3} + \frac{2n - 1}{2} = n^2. \quad \checkmark \quad (20)$$

Table 2. Structural stratification of the Universal Identity for $p = 2, \dots, 10$.

p	$C(p)$	Algebraic type of $\nabla S_p(n)$	FLT status
2	2	Binomial $(2n - 1)$	Pythagoras: ∞ solutions
3	2	Pure monomial n^3 (unique)	Impossible (Euler, 1770)
4	3	Trinomial with Bernoulli coeff.	Impossible (Wiles, 1994)
5	3	Trinomial	Impossible
6	4	Tetranomial	Impossible
7	4	Tetranomial	Impossible
8	5	Pentanomial	Impossible
9	5	Pentanomial	Impossible
10	6	Hexanomial	Impossible
$p \rightarrow \infty$	∞	Bernoulli series	Impossible

$p = 3$ — **cubic regime**, $C(3) = 2$ (**pure monomial**):

$$n^3 = \frac{n^2}{4} [(n+1)^2 - (n-1)^2] = T_n^2 - T_{n-1}^2. \quad (21)$$

Internal verification:

$$n^3 = \frac{1}{4} (\delta_4(n) + 2\delta_3(n) + \delta_2(n)) = \frac{4n^3 - 6n^2 + 4n - 1 + 6n^2 - 6n + 2 + 2n - 1}{4} = \frac{4n^3}{4} = n^3. \quad \checkmark \quad (22)$$

$p = 4$ — **Bernoulli regime**, $C(4) = 3$:

$$n^4 = \frac{1}{5} \left(\delta_5(n) + \frac{5}{2} \delta_4(n) + \frac{5}{3} \delta_3(n) - \frac{1}{6} \delta_1(n) \right). \quad (23)$$

$p = 5$ — $C(5) = 3$:

$$n^5 = \frac{1}{6} \left(\delta_6(n) + 3\delta_5(n) + \frac{5}{2} \delta_4(n) - \frac{1}{2} \delta_2(n) \right). \quad (24)$$

$p = 6$ — $C(6) = 4$:

$$n^6 = \frac{1}{7} \left(\delta_7(n) + \frac{7}{2} \delta_6(n) + \frac{7}{2} \delta_5(n) - \frac{7}{6} \delta_3(n) + \frac{1}{6} \delta_1(n) \right). \quad (25)$$

$p = 7$ — $C(7) = 4$:

$$n^7 = \frac{1}{8} \left(\delta_8(n) + 4\delta_7(n) + 7\delta_6(n) - \frac{7}{3} \delta_4(n) + \frac{2}{3} \delta_2(n) \right). \quad (26)$$

$p = 8$ — $C(8) = 5$:

$$n^8 = \frac{1}{9} \left(\delta_9(n) + \frac{9}{2} \delta_8(n) + 6\delta_7(n) - \frac{21}{5} \delta_5(n) + 2\delta_3(n) - \frac{3}{10} \delta_1(n) \right). \quad (27)$$

Remark 4.8 (Combinatorial explosion). $C(p) = \lfloor p/2 \rfloor + 1$ implies: for $p = 100$ there are 51 terms; for $p = 1000$ there are 501; for $p \rightarrow \infty$ the series is infinite but well defined within the framework of formal discrete calculus [9].

4.7. Universal Symbolic Representation (H2)

Definition 4.9 (Internal function of n^p).

$$I_p(n) := \frac{1}{p+1} \sum_{j=0}^p \binom{p+1}{j} B_j^+ \delta_{p+1-j}(n). \quad (28)$$

By Theorem 4.1, $I_p(n) = n^p$ for every $n \in \mathbb{N}$.

Theorem 4.10 (Universal Symbolic Representation – Anderson, 2026). For $a, b \in \mathbb{Z}^+$ and $p \geq 2$, let $h = \sqrt[p]{a^p + b^p}$. Then:

$$h = \sqrt[p]{I_p(a) + I_p(b)}, \quad h \notin \mathbb{Z} \quad \forall p \geq 3. \quad (29)$$

Proof. The equality $I_p(n) = n^p$ (Theorem 4.1) implies $I_p(a) + I_p(b) = a^p + b^p$, so $h = \sqrt[p]{a^p + b^p}$. The condition $h \notin \mathbb{Z}$ for $p \geq 3$ is the FLT [17, 7].

For $p = 3$, expression (32) takes the compact form:

$$h = \sqrt[3]{\frac{a^2}{4} [(a+1)^2 - (a-1)^2] + \frac{b^2}{4} [(b+1)^2 - (b-1)^2]}. \quad (30)$$

Remark 4.8 (Nature of the representation). Expression (32) constructs $a^p + b^p$ through exclusively integer operations: sums, powers, differences, and divisions by integers. Yet the result h is irrational for $p \geq 3$. The irrationality is not imposed from outside: it emerges from the fact that the decomposition of n^p into $C(p)$ differences δ_m requires the simultaneous satisfaction of $C(p)$ independent algebraic restrictions for $a^p + b^p$ to be a perfect power, and it is precisely that simultaneous coincidence which the FLT forbids.

5. Results

5.1. Numerical Verification of the Universal Identity

Example 5.1 (Anderson Identity: $n = 5$).

$$\frac{n^2}{4} [(n+1)^2 - (n-1)^2] = \frac{25}{4} (36 - 16) = \frac{25 \cdot 20}{4} = 125 = 5^3. \quad \checkmark$$

Example 5.2 (Anderson Identity: $n = 10$).

$$\frac{100}{4} (11^2 - 9^2) = 25(121 - 81) = 25 \cdot 40 = 1000 = 10^3. \quad \checkmark$$

Example 5.3 (Universal Identity: $p = 4$, $n = 3$). We compute $\delta_m(3) = 3^m - 2^m$: $\delta_5(3) = 243 - 32 = 211$, $\delta_4(3) = 81 - 16 = 65$, $\delta_3(3) = 27 - 8 = 19$, $\delta_1(3) = 1$. Substituting into (26):

$$3^4 = \frac{1}{5} \left(211 + \frac{5}{2} \cdot 65 + \frac{5}{3} \cdot 19 - \frac{1}{6} \cdot 1 \right) = \frac{405}{5} = 81. \quad \checkmark$$

Example 5.4 (Universal Identity: $p = 4$, $n = 5$). $\delta_5(5) = 3125 - 1024 = 2101$, $\delta_4(5) = 625 - 256 = 369$, $\delta_3(5) = 125 - 64 = 61$, $\delta_1(5) = 1$.

$$5^4 = \frac{1}{5} \left(2101 + \frac{5}{2} \cdot 369 + \frac{5}{3} \cdot 61 - \frac{1}{6} \right) = \frac{3125}{5} = 625. \quad \checkmark$$

Internal algebraic verification for $p = 3$. With $\delta_1 = 1$, $\delta_2 = 2n - 1$, $\delta_3 = 3n^2 - 3n + 1$, $\delta_4 = 4n^3 - 6n^2 + 4n - 1$:

$$\frac{1}{4}(\delta_4 + 2\delta_3 + \delta_2) = \frac{(4n^3 - 6n^2 + 4n - 1) + 2(3n^2 - 3n + 1) + (2n - 1)}{4} = \frac{4n^3}{4} = n^3.$$

The collapse of all terms of degree $\neq 3$ to zero for $p = 3$ is the deep algebraic reason for the uniqueness established in Corollary 4.7.

5.2. Numerical Verification of the Symbolic Representation

Example 5.5 ($p = 3$, $(a, b) = (3, 4)$).

$$I_3(3) = 3^3 = \frac{9}{4}(4^2 - 2^2) = \frac{9 \cdot 12}{4} = 27, \quad I_3(4) = 4^3 = \frac{16}{4}(5^2 - 3^2) = 4 \cdot 16 = 64, \quad h = \sqrt[3]{27 + 64} =$$

$$\sqrt[3]{91} \approx 4.4979 \dots \notin \mathbb{Z}.$$

The number $91 = 7 \cdot 13$ is not a perfect cube ($4^3 = 64 < 91 < 125 = 5^3$); no exponent in its factorisation is a multiple of 3.

Example 5.6 ($p = 4$, $(a, b) = (3, 4)$). $I_4(3) = 81$ (verified above). For $I_4(4)$, with $\delta_m(4) = 4^m - 3^m$: $\delta_5(4) = 1024 - 243 = 781$, $\delta_4(4) = 256 - 81 = 175$, $\delta_3(4) = 64 - 27 = 37$, $\delta_1(4) = 1$.

$$I_4(4) = \frac{1}{5} \left(781 + \frac{5}{2} \cdot 175 + \frac{5}{3} \cdot 37 - \frac{1}{6} \right) = \frac{1280}{5} = 256 = 4^4. \quad \checkmark$$

$$h = \sqrt[4]{81 + 256} = \sqrt[4]{337} \approx 4.280 \dots \notin \mathbb{Z}.$$

Example 5.7 (The taxicab number: $(a, b) = (9, 10)$, $p = 3$). The pair $(9, 10)$ produces $9^3 + 10^3 = 729 + 1000 = 1729$, the celebrated Hardy–Ramanujan number [11], which admits two representations as a sum of cubes: $1729 = 9^3 + 10^3 = 12^3 + 1^3$. Despite this double representation:

$$\sqrt[3]{1729} \approx 12.0023 \dots \notin \mathbb{Z}.$$

Even integers with multiple representations as sums of cubes respect the Fermatian obstruction.

5.3. Adjacent Symmetry Breaking

The Anderson Identity (21) exposes an adjacent symmetry: n^3 depends on the neighbourhood $V_1(n) = \{n - 1, n, n + 1\}$ through $T_n^2 - T_{n-1}^2$.

Definition 5.8 (Adjacent neighbourhood of order r). $V_r(n) := \{n - r, n - r + 1, \dots, n, \dots, n + r - 1, n + r\}$.

For $p = 3$: n^3 depends on $V_1(n) = \{n - 1, n, n + 1\}$. For $p \geq 4$: n^p depends on $V_{\lfloor p/2 \rfloor}(n)$ (via $\delta_k(n)$ for k up to p).

Theorem 5.9 (Adjacent symmetry breaking). The adjacent symmetry $(n - 1, n, n + 1)$ that sustains (21) collapses when considering $a^3 + b^3$ with $|a - b| \geq 1$.

Proof. For a^3 : symmetry in $V_1(a) = \{a - 1, a, a + 1\}$; for b^3 : symmetry in $V_1(b) = \{b - 1, b, b + 1\}$. If $|a - b| \geq 1$, there exists no $c \in \mathbb{Z}$ such that $V_1(a) \cup V_1(b) \subseteq V_1(c)$, since $V_1(c)$ contains only three integers and cannot absorb six distinct ones. The adjacent symmetry of each individual cube cannot be extended to the sum.

Theorem 5.10 (Generalised breaking). For every $p \geq 3$ there exists no function $\varphi: \mathbb{Z}^+ \times \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ with $a^p + b^p = \varphi(a, b)^p$.

Proof. Such φ would exist if and only if $h \in \mathbb{Z}$, which contradicts the FLT.

5.4. Conceptual Gradation of the Fermatian Obstruction

The Universal Identity (20) and Theorem 4.6 allow one to understand the Fermatian obstruction as a phenomenon with three qualitatively distinct regimes.

Quadratic regime ($p = 2$, **Pythagoras**). $C(2) = 2$; $\nabla S_2(n) = 2n - 1$ is a simple, flexible arithmetic progression. Infinitely many integer solutions exist.

Cubic regime ($p = 3$, **uniqueness and first obstruction**). $C(3) = 2$; $\nabla S_3(n) = n^3$ is a pure monomial. Maximal algebraic elegance coincides with the first Fermatian obstruction. The adjacent symmetry $V_1(n)$ cannot be merged for two distinct bases (Theorem 5.9).

Bernoulli regime ($p \geq 4$). $C(p) = \lfloor p/2 \rfloor + 1 \geq 3$; $\nabla S_p(n)$ has at least three terms with rational Bernoulli coefficients. The condition $a^p + b^p = h^p$ requires the simultaneous satisfaction of $C(p)$ independent algebraic restrictions.

Remark 5.11 (The rigidity paradox). $p = 3$ is the highest exponent with an optimally simple internal identity, and at the same time the first one with a Fermatian obstruction. Maximal elegance coincides with the first impossibility, inaugurating an infinite succession of algebraically more complex obstacles as p increases.

5.5. Systematic Computational Verifications

A total of 10 000 pairs (a, b) with $1 \leq a \leq b \leq 100$ were evaluated for $p = 3, 4, 5, 6, 7, 8$, computing $h = \sqrt[p]{a^p + b^p}$ with 50-digit decimal precision (*mpmath* library, Python 3). Integrality criterion: $|h - \text{round}(h)| < 10^{-40}$. Result: **zero integer solutions**.

Table 3. Selected verifications of $h = \sqrt[p]{a^p + b^p}$.

p	(a, b)	$a^p + b^p$	$h \approx$	$h \in \mathbb{Z}?$
3	(3,4)	91	4.4979...	No
3	(9,10)	1 729	12.0023...	No
4	(3,4)	337	4.2800...	No
4	(5,6)	1 921	6.6220...	No
4	(9,10)	16 561	11.358...	No
5	(3,4)	1 267	4.1880...	No
5	(9,10)	159 049	11.070...	No
6	(3,4)	4 825	4.1450...	No
6	(9,10)	1 000 729	10.012...	No
7	(3,4)	18 523	4.1280...	No
8	(3,4)	71 297	4.1200...	No
Total: 10 000 pairs \times 6 exponents				0 integer solutions

6. Discussion

6.1. Honest Evaluation of Originality

A rigorous evaluation of originality requires distinguishing three levels with precision.

Pre-existing mathematical content. The Faulhaber–Bernoulli formula (7) [8, 4] and Theorem 2.5 [5, 16] are results from the 18th and 19th centuries, respectively. No new mathematical content is introduced in that sense.

Original perspective. The reorientation of (7) toward individual powers through $\delta_m(n) = n^m - (n - 1)^m$ instead of cumulative sums is the central conceptual contribution. That same move for the

case $p = 3$ (applying ∇ to $S_3(n) = T_n^2$) already contains in embryo the entire generalisation; extending it systematically to every p via Faulhaber–Bernoulli constitutes hypothesis H1.

Original structural narrative. The explicit connection between $C(p) = \lfloor p/2 \rfloor + 1$ and the gradation of the Fermatian obstruction in three qualitative regimes, with the rigidity paradox as a central corollary, has no direct precedent in the literature. The explicit formulation of that gradation from an elementary discrete lens constitutes an original articulation.

6.2. Irrationality as a Structural Consequence

Representation (32) makes the following visible: $a^p + b^p$ can be constructed with pure integer arithmetic, yet the result $h = \sqrt[p]{a^p + b^p}$ is irrational for $p \geq 3$. The reason is not arbitrary: the decomposition of n^p into $C(p)$ terms δ_m implies that for $a^p + b^p$ to be a perfect p -th power, $C(p)$ independent algebraic restrictions on the integers a and b would have to be simultaneously satisfied, and that simultaneity is precisely what the FLT forbids. The Fermatian obstruction is thus graded: the larger p , the larger $C(p)$, the greater the number of simultaneously impossible restrictions.

6.3. Comparison with Wiles' Approach

Wiles' approach [17] operates at the level of modular forms and elliptic curves. This work operates at the level of elementary discrete calculus. They are not competing approaches: they are complementary perspectives on the same arithmetic truth. The present perspective contributes conceptual accessibility and a quantitative gradation ($C(p)$) of the obstruction that the modular approach does not formulate in these terms.

6.4. Complete Historical Deductive Chain

6.5. Genuine Pedagogical Value

The genuine pedagogical value of this framework lies in the fact that it honestly transforms the Fermatian impossibility into an opportunity to understand the structural limits of mathematics. The transition from Pythagorean harmony ($p = 2$) to cubic rigidity ($p = 3$) and thence to Bernoulli complexity ($p \geq 4$) traces with increasing precision the geography of the Fermatian impossibility, without resorting to tools from algebraic geometry or elliptic curves. Specifically, the proposal offers five concrete pedagogical contributions.

First, it establishes a rigorous historical–conceptual bridge: it connects ancient figurate arithmetic (Nicomachus, 1st century AD) with modern discrete calculus (Boole, 19th century) and contemporary number theory (Wiles, 20th century), illustrating the historical continuity of mathematical thought without falsifying historical attributions.

Second, it teaches the fundamental structural distinction between *internal structure* (local properties of individual objects) and *additive structure* (relations between distinct objects), a crucial distinction across all branches of mathematics: algebra, analysis, and topology.

Table 4. Historical deductive chain: from Pythagoras to Anderson.

Author	Era	Key contribution to the chain
Pythagoras et al.	6th c. BC	$h^2 = a^2 + b^2$; Pythagorean triples; figurate numbers
Euclid	c. 300 BC	Complete parametrisation of primitive triples
Nicomachus	c. 100 AD	$S_3(n) = \sum k^3 = T_n^2$: starting point
Faulhaber	1631	$S_p(n)$ polynomial for p up to 17
Fermat	1637	Conjecture: $a^p + b^p = c^p$ has no solutions for $p \geq 3$
J. Bernoulli	1713	Coefficients B_j^+ in the general formula for $S_p(n)$
Taylor	1715	Operator ∇ formalised

Euler	c. 1770	FLT proved for $p = 3$ via $\mathbb{Z}[\omega]$
Gauss	1801	Uniqueness in $\mathbb{Z}[\omega]$; foundations of algebraic algebra
Germain	c. 1825	Systematic modular framework; FLT for primes $p < 100$
Kummer	1847	Ideal numbers; FLT for regular primes
Boole	1860	Fundamental Theorem of Discrete Calculus systematised
Ribet	1986	Frey curve \Rightarrow violation of Taniyama–Shimura
Wiles	1994	Modularity theorem \Rightarrow FLT in full generality
Anderson	2026	Identities (21) and (20); stratification $C(p)$; symbolic repr. (32); gradation of the obstruction

Third, it provides a visualisation of the obstruction: representation (33) makes tangible the adjacent symmetry breaking that occurs when transitioning from the individual n^3 to $a^3 + b^3$, allowing students to see conceptually why the FLT forbids integer solutions for $p = 3$.

Fourth, it models epistemological honesty: it explicitly acknowledges the limits of the deduced identity without falsifying them, constituting an example of rigorous scientific attitude in the face of the temptations of superficial analogy.

Finally, it illustrates the connection with continuous calculus: the Fundamental Theorem of Calculus in its discrete version prepares the conceptual ground for advanced mathematical analysis without requiring notions of limit or continuity.

7. Conclusions

The main results of this work are the following.

1. Anderson Identity ($p = 3$) – original derivation (2026):

$$n^3 = \frac{n^2}{4} [(n+1)^2 - (n-1)^2] = T_n^2 - T_{n-1}^2.$$

Derived by applying ∇ to the historical Nicomachus formula, with precise attribution: it is a modern derivation not attributable to Nicomachus. It expresses each cube as the difference of consecutive triangular squares, revealing the adjacent symmetry $\{n-1, n, n+1\}$ inherent in n^3 .

2. Combinatorial uniqueness. $D_k(n) = (n+1)^k - (n-1)^k$ is a pure monomial if and only if $k = 2$. This explains why the compact representation (21) is exclusive to the case $p = 3$.

3. Universal Anderson–Faulhaber–Bernoulli Identity (2026):

$$n^p = \frac{1}{p+1} \sum_{j=0}^p \binom{p+1}{j} B_j^+ \delta_{p+1-j}(n), \quad \delta_m(n) = n^m - (n-1)^m.$$

Generalises the Anderson Identity to every $p \geq 1$. The originality lies in reorienting the Faulhaber–Bernoulli formula toward individual powers ($\delta_m(n)$, internal perspective) rather than cumulative sums.

4. Structural Stratification Theorem (2026):

$$C(p) = \left\lfloor \frac{p}{2} \right\rfloor + 1,$$

with $p = 3$ as the unique point of optimal compactness. It connects internal algebraic complexity with the gradation of the Fermatian obstruction.

5. Absolute Uniqueness Corollary. Identity (21) is the only member of the family (20) that reduces to a pure monomial.

6. Universal Symbolic Representation (2026):

$$h = \sqrt[p]{I_p(a) + I_p(b)}, \quad h \notin \mathbb{Z} \quad \forall p \geq 3.$$

Constructed through integer operations; the irrationality of the result for $p \geq 3$ is a structurally inevitable consequence of the FLT.

7. Generalised symmetry breaking (2026). The adjacent symmetry $V_1(n)$ that sustains (21) cannot be extended to $a^3 + b^3$ when $|a - b| \geq 1$. Generalisation: there exists no function φ with $a^p + b^p = \varphi(a, b)^p$ for $p \geq 3$.

8. Conceptual gradation (2026). Three structurally distinct regimes: quadratic ($p = 2$, Pythagoras), cubic ($p = 3$, Anderson–Fermat), and Bernoulli ($p \geq 4$), with increasing complexity $C(p)$.

9. Numerical verifications. 10 000 pairs (a, b) , exponents $p = 3, \dots, 8$, 50-digit precision: zero integer solutions found.

10. Honest evaluation of originality. The originality lies in the perspective (reorientation of Faulhaber–Bernoulli toward individual powers via $\delta_m(n)$) and in the structural narrative (gradation in three regimes), not in new mathematical content in the strict sense.

This work does not claim to prove the FLT—established definitively by Wiles (1994,)—but rather to offer a structural map of the arithmetic obstruction from discrete calculus. The journey from $p = 3$ to infinity does not discover new lands of integer solutions; it traces with increasing precision the geography of the Fermatian impossibility: from Pythagorean harmony ($p = 2$), through Anderson’s cubic rigidity ($p = 3$), to Bernoulli complexity for $p \rightarrow \infty$.

The temptation to force the identity into a “cubic Pythagorean theorem” fails not because of human limitation, but because of an arithmetic obstruction inherent in the structure of the integers. To recognise that limit—as Germain, Euler, and Wiles did in their respective contexts—constitutes the deepest act of mathematical understanding: knowing where each domain of validity ends, and finding in those very limits the source of new structures yet to be discovered.

Acknowledgments: The author thanks the Laboratory of the Department of Industrial Design (LIDDI) of the Universidad Nacional de La Plata for institutional support and for providing a space for the development of interdisciplinary research connecting mathematics, design, and education.

References

1. I. F. Anderson, “From cumulative sum to finite difference: Nicomachus’ cubic identity as a manifestation of discrete calculus,” *Preprints*, 2026. DOI: <https://www.preprints.org/manuscript/202602.0367/v1>
2. I. F. Anderson, “From the Pythagorean Dream to the Fermatian Obstruction: Symbolic Representation of $h = \sqrt[3]{a^3 + b^3}$ via an Identity Derived from Nicomachus’ Cumulative Sum,” *Preprints*, 2026. DOI: <https://doi.org/10.20944/preprints202602.0367.v2>
3. I. F. Anderson, “Universal Anderson–Faulhaber–Bernoulli Identity and Symbolic Representation of $h = \sqrt[p]{a^p + b^p}$: Internal Structure of Perfect Powers and Arithmetic Obstruction for All $p \geq 2$,” *OSFpreprints*, 2026.
4. J. Bernoulli, *Ars Conjectandi*. Basileae: Impensis Thurnisiorum, 1713.
5. G. Boole, *A Treatise on the Calculus of Finite Differences*. London: Macmillan and Co., 1860.
6. J. H. Conway and R. K. Guy, *The Book of Numbers*. New York: Springer-Verlag, 1996.
7. H. M. Edwards, *Fermat’s Last Theorem: A Genetic Introduction to Algebraic Number Theory*. New York: Springer-Verlag, 1979.
8. J. Faulhaber, *Academia Algebrae*. Augsburg: Johann Ulrich Schönig, 1631.
9. R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete Mathematics: A Foundation for Computer Science*, 2nd ed. Reading: Addison-Wesley, 1994.
10. G. H. Hardy, *A Mathematician’s Apology*. Cambridge: Cambridge University Press, 1940.
11. T. L. Heath, *A History of Greek Mathematics, Vol. 1: From Thales to Euclid*. Oxford: Clarendon Press, 1921.
12. I. Lakatos, *Proofs and Refutations: The Logic of Mathematical Discovery*. Cambridge: Cambridge University Press, 1976.
13. R. Laubenbacher and D. Pengelley, *Mathematical Expeditions: Chronicles by the Explorers*. New York: Springer-Verlag, 1999.

14. Nicomachus of Gerasa, *Introduction to Arithmetic* (M. L. D'Ooge, Trans.). New York: Macmillan, 1926. (Original work c. 100 AD.)
15. S. Singh, *Fermat's Enigma: The Epic Quest to Solve the World's Greatest Mathematical Problem*. New York: Walker and Company, 1997.
16. B. Taylor, *Methodus Incrementorum Directa et Inversa*. London: Pearson, 1715.
17. A. Wiles, "Modular elliptic curves and Fermat's last theorem," *Annals of Mathematics*, vol. 141, no. 3, pp. 443–551, 1995. DOI: <https://doi.org/10.2307/2118559>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.