

Article

Not peer-reviewed version

Downregulation of Gene Expression by Alpha Satellite Transcripts

Sven Ljubić , [Maja Matulic](#) , [Damir Đermić](#) , Maria Chiara Feliciello , Alfredo Procino , [Francesco Passaro](#) , [Đurđica Ugarković](#) ^{*} , [Isidoro Feliciello](#) ^{*}

Posted Date: 11 September 2025

doi: 10.20944/preprints202509.0957.v1

Keywords: satellite DNA; transcription; gene expression; RNA-DNA hybrid; heterochromatin; histone modifications; alpha satellite DNA



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Article

Downregulation of Gene Expression by Alpha Satellite Transcripts

Sven Ljubić ¹, Maja Matulić ², Damir Đermić ¹, Maria Chiara Feliciello ³, Alfredo Procino ⁴,
Francesco Passaro ⁵, Đurđica Ugarković ^{1,*} and Isidoro Feliciello ^{1,4,*}

¹ Division of Molecular Biology, Ruder Bošković Institute, Bijenička 54, HR-10000 Zagreb, Croatia

² Department of Biology, Faculty of Science, University of Zagreb, HR-10000 Zagreb, Croatia

³ Department of Statistical Science, Alma Mater Studiorum, University of Bologna, 40126 Bologna, Italy

⁴ Department of Clinical Medicine and Surgery, University of Naples Federico II, I-80131 Naples, Italy

⁵ Division of Urology, Istituto Nazionale Tumori, Fondazione G. Pascale, Naples, Italy

* Correspondence: djurdjica.ugarkovic@irb.hr (D.U.); isidoro.feliciello@unina.it (I.F.);

Tel.: +39 0817464317 (I.F.);

Abstract

Satellite DNAs are highly abundant sequences which build functional centromere and pericentromeric heterochromatin in many eukaryotes. Apart from this structural role, their involvement in gene expression modulation has been demonstrated, although the detailed molecular mechanism is still missing. Here, using major human alpha satellite as a model system, we investigate the role of satellite transcripts in gene expression regulation. We generated cell lines with forced, exogenous overexpression of alpha satellite RNA and followed the expression levels of genes containing alpha satellite repeats within introns. Our results reveal a positive correlation between exogenous alpha satellite expression and the downregulation of alpha-associated genes, strongly suggesting that alpha satellite RNA affects their transcription. Notably, the elevated levels of exogenous alpha satellite RNA did not affect histone modifications characteristic of pericentromeric heterochromatin (e.g. H3K9me3 or H3K18Ac) or euchromatin (e.g. H3K4me2) at intronic alpha satellite loci. We propose that alpha satellite RNA directly interacts with homologous DNA at dispersed intronic satellite loci by forming RNA-DNA hybrid structures, which may affect chromatin structure and transcriptional activity. The results demonstrate that alpha satellite RNA is not only involved in centromere and heterochromatin assembly but, as shown here for the first time, also plays a role in modulating the expression of alpha-associated genes.

Keywords: satellite DNA; transcription; gene expression; RNA-DNA hybrid; heterochromatin; histone modifications; alpha satellite DNA

1. Introduction

Alpha satellite DNA represents a major human satellite which is composed of tandemly arranged, diverged 171 bp long monomers, often organized in complex higher order repeats [1]. The satellite comprises up to 10% of the human genome and is located in the centromeric and pericentromeric regions of all chromosomes in the form of long, Mb size arrays [2]. In addition to the (peri)centromeric location, a bioinformatic search of the human genome revealed the presence of short arrays of alpha satellite repeats within the euchromatic regions of the genome, including introns and areas near genes [3]. As previously described, one possible origin of these euchromatic, dispersed form of satellite repeats is due to the molecular mechanism of satellite DNA evolution based on the rolling circle amplification [3–5]. This non-canonical occurrence of dispersed satellite repeats has been observed in various species to date and is proposed to be a characteristic feature of all satellite DNAs [6–9]. In the case of beetle *Tribolium castaneum*, it was demonstrated that dispersed satellite repeats

can influence the expression of neighbouring genes upon specific conditions, such as heat stress [10]. In general, heat stress as well other kinds of stress like antibiotic treatment [11] induce transcription of satellite DNA. These transcripts are proposed to guide the deposition of repressive histone marks at homologous satellite sequences, affecting both heterochromatic and euchromatic regions [10,12].

In humans, RNA polymerase II (Pol II) transcribes alpha satellite DNA repeats into long non-coding RNA and the predominant factors controlling transcription seem to be the presence of centromere–nucleolar contacts [13] and Topoisomerase I (TopI) [14]. In addition, a role of N6-methyladenosine (m6A) modification in the regulation of alpha satellite transcription was proposed [15]. Alpha satellite RNA levels fluctuate throughout the cell cycle, peaking in the G2/M phase and the transcripts are not exported to the cytoplasm [13]. Transcripts of alpha satellite DNA contribute to essential chromosomal functions such as centromere assembly and kinetochore formation [2] and are necessary for proper heterochromatin formation in humans [16,17]. Transcription of alpha satellite DNA is also induced upon heat stress as shown by studies on different cell lines [18,19]. The increased levels of alpha satellite transcripts after heat stress correlates with the downregulation of genes containing alpha satellite repeats within introns or in the gene vicinity, indicating possible influence of alpha satellite transcripts on gene expression modulation [19].

To analyse the possible gene modulatory role of alpha satellite transcripts more deeply and to exclude the effect of heat stress itself or of other stressors on gene expression, we have now developed cell lines with forced, exogenous overexpression of alpha satellite RNA. In such modified cell lines we monitored the level of expression of genes containing alpha satellite repeats within introns. The results reveal a positive correlation between exogenous alpha satellite expression and downregulation of alpha-associated genes, strongly suggesting an effect of alpha satellite transcripts on gene expression. Furthermore, we performed an analysis of different histone marks on intronic alpha satellite repeats in cells with exogenous expression of alpha satellite RNA, and propose a possible molecular mechanism by which alpha satellite transcripts modulate gene expression.

2. Results

2.1. Alpha Satellite Transcription After Transfection

To investigate possible effect of alpha satellite RNA on gene expression we constructed vectors expressing alpha satellite monomers in both orientations (171Fw and 171Rev) and monitored their transcription dynamics in transfected human MJ90-hTERT cell line by RT-qPCR. The exogenous alpha satellite RNA was quantified 24, 48, 72 and 96 hours after transfection and compared to its endogenous variant. Primers used for transcriptional analysis of endogenous alpha satellite RNA were able to amplify only tandemly arranged repeats (Figure S1) and since in human pericentromeric heterochromatin alpha satellite DNA is organized in tandemly arranged monomers [2], it is expected that the primers preferentially recognize transcripts deriving from pericentromeric regions.

The results revealed a significant level of exogenous alpha satellite expression from both 171Fw and 171Rev vectors, the highest being achieved 24 hours after transfection and rapidly decreasing afterwards (Figure 1), primarily as a consequence of cell division related plasmid loss and lack of selective pressure. Additionally, both vector variants demonstrated mutually similar levels of expression during the experiments. Endogenous alpha satellite expression retained its basal levels throughout all investigated time points, indicating standard physiological conditions and lack of stress or toxicity for the cells post treatment.




Figure 1. Levels of exogenous alpha satellite RNA expressed from pCMV6-A-GFP vectors 24, 48, 72 and 96 hours after transfection. 171Fw indicates the 171 bp alpha satellite monomer insert in forward orientation and 171Rev its inverted form. No value is expressed in arbitrary fluorescence units and is calculated by taking into account PCR efficiency and baseline fluorescence. Columns show averages of two independent experiments and error bars indicate standard deviations.

2.2. Expression Analysis of Alpha-Associated Genes After Transfection

The expression profiles of genes containing dispersed alpha satellite repeats within their intronic regions, described in [19], were explored over the course of four consecutive days, after transfection of MJ90-hTERT cell line with satellite-expressing vectors compared to controls transfected in the same way with unmodified vectors. These included: *SLC30A6* (solute carrier family 30 member 6, ID: 55676), *STAM* (signal transducing adapter molecule 1, ID: 8027), *MYO1E* (myosin IE, ID: 4643), *MAP7* (MAP7 domain containing 2, ID: 256714), *ZNF675* (zinc finger protein 675, ID: 171392), *VAV1* (vav guanine nucleotide exchange factor 1, ID: 7409), *PRIM2* (DNA primase subunit 2, ID: 5558) and *DLG2* (discs large homolog 2, ID: 1740). Glucuronidase beta gene (*GUSB*, ID: 2990) was used as endogenous control for expression normalization as well as a negative control against alpha repeat-associated genes.

24 hours post transfection, a significant downregulation of gene expression was detected for six tested genes relative to controls. These included *SLC30A6*, *STAM*, *MYO1E*, *MAP7*, *ZNF675* and *PRIM2*. Out of those, first five genes demonstrated highest level of downregulation between transfected samples and controls ($P<10^{-3}$) and *PRIM2*, although to a lesser extent, was also significant ($P=0.02$) (Figure 2). Also, there were no observed differences in vectors' downregulation efficiency between 171Fw and 171Rev transfected cells in all cases, indicating comparable levels of effectiveness regardless of alpha satellite insert orientation. Other investigated time points (48, 72 and 96 hours post transfection) showed no significant differences in gene expression of candidate genes between transfected samples and controls (Figures S2, S3, S4). This is likely due to vector expression dynamics, signifying the most impactful effect of satellite transcription 24 hours post transfection, and subsequent rapid decline (Figure 1). The genes *VAV1* and *DLG2* showed no detectable expression in any of the samples at any of the analyzed time points, indicating they are tissue-specific and, therefore, not expressed in the MJ90-hTERT cell line. *GUSB* gene was stably expressed at all times

during the experiments, with no significant variability between samples and controls (Figure S5) confirming its role as a dependable normalization gene for relative RT-qPCR quantification.




Figure 2. Expression profiles of genes containing alpha satellite repeats within intronic regions, in MJ90hTERT cell lines transfected with satellite expressing vectors and controls, 24 hours after treatment. 171Fw denotes the vector with satellite insert in forward orientation and 171Rev its inverted counterpart. Control refers to unaltered pCMV6-A-GFP vector. Gene downregulation rates of transfected samples compared to controls are shown above error bars which represent standard deviations. N₀ represents normalized average N₀ value expressed in arbitrary fluorescence units.

Additionally, it should be noted that our observed results are likely to be understated to a certain degree. MJ90-hTERT human cell line is quite difficult to transfect effectively without using viral vectors (such as lentivirus) which is often accompanied by higher levels of toxicity and may induce adverse effects for the cells. To avoid these problems, using lipofection method, we managed to achieve approximately 30-35% transfection efficiency in our experiments by monitoring constitutively expressed GFP inside the cells, an inherent additional trait of our vectors. With higher transfection efficiency it is likely that the observed downregulatory effects would be more pronounced or even become significant in later time points (48, 72 and 96 hours after transfection).

2.3. H3K9me3, H3K18ac and H3K4me2 Levels at Alpha Repeats Dispersed Within Genes After Transfection

We analysed the distribution of silent histone mark H3K9me3, H3K18ac mark characteristic for transcriptional activation of heterochromatin and H3K4me2, typical for open euchromatin, on alpha repeats dispersed within introns of six genes previously tested for expression. Histone marks were analysed in MJ90hTERT cells, 24 hours after transfection with 171Fw, 171Rev and unmodified control expression vectors. In order to investigate whether the previously observed downregulation of gene expression could be related to above-mentioned epigenetic changes, we performed chromatin immunoprecipitation (ChIP) coupled with quantitative real-time PCR, using specific primers for histone modification level analyses of alpha repeats associated with genes (Table S2). ChIP assay was performed on chromatin isolated from MJ90hTERT cells. The levels of tested histone modifications were measured immediately after 24 hours transfection period and compared to the level of control transfected in the same way with unmodified vector, using the unpaired t-test. In addition, we followed the IgG binding dynamics to investigated loci and the amount of bound IgG was very low, resulting in a signal below the qPCR threshold.

The transfection of MJ90hTERT cells with 171Fw alpha satellite expression vector resulted in no significant differences with regard to tested histone modifications at alpha repeats in six genes of interest compared to control samples, 24 hours after treatment (Figure 3). Similarly, no differences were observed after repeated experiments with 171Rev vector containing inverted alpha satellite insert (Figure S6). Both vectors showed similar results compared to controls, mirroring their close performance from gene expression analyses (Figure 2). On one hand, it is possible that this outcome is a consequence of previously mentioned limited transfection potential of MJ90hTERT cells or lower sensitivity of ChIP experiments. However, it might also be an indicator of other potential mechanism(s) modulating gene expression of alpha satellite repeat-associated genes.




Figure 3. Levels of H3K9me3, H3K18ac and H3K4me2 histone modifications at alpha satellite repeats associated with six genes after 24 hour transfection with 171Fw and unmodified control vectors. Levels of histone modifications were measured by ChIP coupled with quantitative real-time PCR on MJ90hTERT chromatin immediately after each treatment. No values were normalized using No values of input fractions and represent the levels of histone modifications. Columns show averages of two independent experiments and error bars indicate standard deviations.

2.4. Alpha Satellite RNA Level Analysis After RNase H digestion of RNA:DNA Hybrids

To determine whether alpha satellite transcripts actively form RNA:DNA hybrids under standard physiological conditions, RNase H digestion assay was performed. RNase H specifically degrades the RNA strand of an RNA-DNA hybrid. We detected a modest (around 25%) but significant (Student's t-test, $P < 0.05$) reduction of alpha transcripts in samples treated with RNase H compared to untreated controls (Figure 4a). Alpha satellite RNA was quantified by RT-qPCR using the same primer pair as in previous experiments (Fw 5'-CACTCTTTTGTAGAATCTGC-3' ; Rev 5'-AATGCACATATCACTATGTAC-3'). *GUSB* gene was used as endogenous housekeeper for normalization as well as a negative control, being stably and uniformly expressed, showing no significant variation between samples before and after RNase H treatment (Figure 4b).




Figure 4. (a) Alpha satellite RNA levels in samples treated with RNase H and untreated controls. N₀ represents normalized average N₀ values expressed in arbitrary fluorescence units. Columns show averages of two different RT-qPCR experiments performed in triplicates and error bars represent standard deviations. Statistical significance between controls and treated samples was calculated using the Student's t-test (P<0.05). (b) Expression profile of housekeeping gene glucuronidase beta (*GUSB*) in MJ90hTERT cell line after treatment with RNase H enzyme and in untreated controls. Three independent RT-qPCR experiments were performed. Error bars represent standard deviations and averaged N₀ values are expressed in arbitrary fluorescence units. No significant differences in gene expression were observed between treated samples and controls in all cases (Student's t-test, P>0.1).

3. Discussion

While transcription of satellite DNAs is tightly regulated under physiological conditions, upon specific conditions such as heat stress it is significantly changed and satellite transcripts were proposed to be involved in gene expression modulation [20]. Human satellite III transcription is particularly strongly activated upon heat stress [21] and the transcripts mediate the recruitment of a number of RNA binding proteins involved in pre-mRNA processing participating in the control of gene expression at the level of splicing regulation [22,23]. In the case of the major TCAST1 satellite DNA of beetle *Tribolium castaneum*, increased levels of H3K9me2/3 are detected after heat stress at regions of (peri)centromeric heterochromatin as well as at dispersed satellite repeats and their flanking regions up to 2 kb from the insertion site, indicating that satellite transcripts can act in *trans*, targeting homologous regions in euchromatin. Increased levels of H3K9me2/3 at euchromatic satellite repeats correlates with transient suppression of neighbouring genes and indicates a role for TCAST1 satellite siRNAs in the modulation of gene expression [10,12].

In the present study we transiently expressed MJ90-hTERT fibroblasts with the plasmid expressing alpha satellite DNA monomer and followed expression of alpha satellite as well as alpha-associated genes at 24h, 48h, 72 hr and 96hr post transfection. Maximal exogenous expression of alpha satellite exceeding endogenous transcription for more than 10x was obtained 24 hr after transfection, coinciding with the downregulation of all alpha-associated genes. The endogenous transcription of alpha satellite was not only significantly lower than the exogenous expression, but also remained constant before and after transfection under all tested conditions, indicating that exogenous alpha satellite expression did not affect the endogenous gene regulation. Our present results showed that gene silencing by satellite RNA is not solely mediated by histone modifications but may also involve other forms of transcriptional interference. In humans, ChIP analyses did not reveal significant changes in histone marks typically associated with pericentromeric heterochromatin, such as H3K9me3 and H3K18Ac, nor in euchromatin-associated marks like H3K4me2 at the level of dispersed intronic alpha satellite repeats. This does not imply a fundamentally different mechanism between *Tribolium* and humans; rather, it suggests an additional silencing pathway alongside the one previously characterized in *Tribolium*. Although ChIP-seq is a powerful technique, conventional protocols often underperform when dealing with low-input material, as in the case of our experiments, where only about 30–35% of cells were efficiently transfected. To conclusively determine whether histone marks also contribute to gene silencing in humans, further experiments using higher-resolution methods are necessary. In particular, single-cell chromatin immunocleavage sequencing (scChIC-seq) may allow the detection of histone modification changes that remain undetectable using conventional bulk ChIP methods [24,25].

Our findings further add a layer of complexity to the mechanisms underlying gene expression downregulation by satellite DNA transcripts. Specifically, the transient expression of exogenous alpha satellite DNAs, produced by transcription from both DNA strands, exerts the same silencing effect on alpha-associated genes, independently of the strand of origin. Based on this observation and on susceptibility of alpha satellite RNA to RNase H treatment, we propose that alpha satellite RNA interacts directly with homologous DNA at dispersed intronic satellite loci by forming hybrid structures in the form of triple helices (RNA: DNA:DNA) or R-loops, thereby affecting the transcription of neighbouring genes (Figure 5). A mechanism of gene expression modulation based on direct interaction between non-coding RNA and DNA was demonstrated [26]. Unlike classical Watson-Crick base pairing between the strands of the DNA double helix, RNA-DNA hybrids are characterised by Hoogsteen hydrogen bonding between nucleic acid bases. This type of interaction is characterised by weaker and more flexible association between DNA and RNA molecules, supporting the notion of a mechanism of transient interference based on alpha satellite sequence homology. This could explain also the observed differences in the level of gene suppression among the investigated alpha satellite repeat-associated genes (Figure 2). Given that alpha satellite DNA is polymorphic in nature, with potential monomer sequence variability of up to 45%, and the vector expressed satellite RNA being a single clone sequence, it is plausible that the observed variability differences in gene repression can be attributed to different degrees of sequence homology. Additionally, the gene-modulatory capabilities of alpha satellite DNA do not depend on gene polarity since the similar levels of gene downregulation were observed using either vector, regardless of the direction of transcription (Figure 2). This would imply that the formation of this hybrid genomic structure plays a key role, either directly or by guiding RNA-associated regulatory proteins to specific genomic locations. The formation of triple helices and R-loops seems to be widespread and essential for regulatory activity of many non-coding RNAs [27,28]. Although many R-loop forming long non-coding RNAs act in *cis*, R-loops can also form in *trans*, influencing the expression of protein coding genes [29]. Triplex formation by long non-coding RNAs was also shown to be able to regulate gene expression in *trans* [28,30]. In mice, pericentromeric satellite DNA transcripts have been shown to form RNA:DNA hybrids which enable the retention of heterochromatin protein 1 (HP1) and the histone methyltransferases SUV39h1 and SUV39h2, which are necessary for heterochromatin formation [17,31].

In conclusion, our results demonstrate that alpha satellite RNA is not only involved in centromere and heterochromatin assembly, but, for the first time reveal its role in the modulation of gene expression.




Figure 5. Proposed model of alpha satellite RNA gene-modulatory potential. Depicted is a part of an intronic region of an alpha satellite repeat-associated gene. Termination of transcriptional activity occurs at the boundary between transcriptional machinery and RNA-DNA triplex genomic structure. Pol II signifies eukaryotic RNA Polymerase II enzyme, mRNA is the nascent messenger RNA molecule and alpha satellite RNA sequence is highlighted in red.

4. Materials and Methods

4.1. Human Cell Line

Human diploid fibroblast strain MJ90hTERT (HCA2hTERT) was kindly provided by Dr Olivia M. Pereira-Smith (University of Texas, Health Science Center, San Antonio, TX, USA) [32–34]. Cells were cultured in appropriate medium (DMEM) supplemented with 10% FBS and 5% CO₂ at 37°C.

4.2. Construction of Vectors

The alpha satellite 171 bp long monomer was amplified by PCR from MJ90-hTERT genomic DNA and cloned into the pCMV6-A-GFP plasmid vector (OriGene). Two separate constructs were created; the first containing the satellite in forward orientation (171Fw) and the second with the insert inverted (171Rev). Inserts were cloned into the vector's multiple cloning site using modified alpha satellite primers specific for consensus sequence of 171 bp alpha satellite monomer [35] with inbuilt restriction site sequences. The forward-oriented insert was generated using forward primer (5'-TGCATTGGATCCCATTCTCAGAACTTCTTTGTG-3') and reverse primer (5'-TGCATTCTCGAGCTTCTGTCTAGTTTTTATGTGAAG-3') while the inverted insert was obtained using forward primer (5'-GCATATCTCGAGCATTCTCAGAACTTCTTTGTG-3') and reverse primer (5'-TGGCTGGGATCCCTTCTGTCTAGTTTTTATGTGAAG-3'). The unmodified pCMV6-A-GFP vector was used as a negative control. Amplicon fidelity was tested by agarose gel electrophoresis before restriction and ligation into vectors. The JM109 bacterial strain (Promega) was subsequently transformed with above mentioned recombinant vectors; clones were selected on

ampicillin-containing plates, screened by colony PCR and plasmid vectors were isolated using GenElute HP Plasmid Midiprep Kit (Sigma-Aldrich). The sequences of the cloned fragments were validated by Sanger DNA sequencing.

4.3. MJ90hTERT Cell Line Transfection

MJ90hTERT (immortalized human skin fibroblasts) cells were transfected with 171Fw, 171Rev and unmodified pCMV6-A-GFP control vector using Lipofectamine 3000 Transfection Reagent (Thermo Fisher Scientific), according to manufacturer's instructions. Afterwards, cells were incubated for 24, 48, 72 and 96 hrs, respectively, at 37°C in complete medium.

4.4. RNA Isolation and Reverse Transcription

For RNA isolation from cell cultures, RNeasy Plus Mini Kit (Qiagen) was used, at each specified time point. Lysis buffer was added directly after the PBS washing step, avoiding trypsin treatment. RNA was quantified with the Quant-IT RNA assay kit using a Qubit fluorometer (Invitrogen). Integrity of RNA was checked by gel electrophoresis. Approximately 1 µg of RNA was reverse transcribed using the PrimeScript RT reagent Kit with gDNA Eraser (perfect Real Time, Takara) in 20 µl reaction using RT Primer Mix for gene expression analyses and the specifically modified primer for alpha satellite (Rev 5'-AATGCACATATCACTATGTAC-3'), designed to produce cDNA molecules that differ from genomic DNA in order to avoid DNA contamination [36]. For all samples, negative controls without reverse transcriptase were used.

4.5. Quantitative Real-Time PCR (qPCR) Analyses

qPCR analyses were performed 24, 48, 72 and 96 hrs post transfection, according to the previously published protocol [10]. Primers used for transcriptional analysis of endogenous alpha satellite RNA were constructed based on consensus sequence derived from cloned alpha satellite monomers of wide-ranging chromosomal origins ([35], and the same modified primer used previously in reverse transcription (Rev 5'-AATGCACATATCACTATGTAC-3') was used in qPCR amplification in order to avoid any potential DNA contamination along with the second primer (Fw 5'-CACTCTTTTGTAGAATCTGC-3'). Exogenous 171 bp alpha satellite RNA derived from vectors in both orientations was quantified using forward (Fw 5'-CATTCTCAGAACTTCTTTGTG-3') or reverse (Rev 5'-CTTCTGTCTAGTTTTATGTGAAG-3') primers in combination with vector-specific T7 promoter forward primer (5'-TAATACGACTCACTATAGGG-3'). This strategy ensures the specific amplification of exogenous alpha RNA. Candidate genes of interest with intronic alpha satellite repeats were identified by bioinformatic analyses, as described in Feliciello et al. 2020a, b, and the primer combinations used for their expression analysis are listed in Table S1. Glucuronidase beta (*GUSB*) was used as an endogenous control for normalization in human samples as well as a negative control for aforementioned genes. *GUSB* gene (Gene ID: 2990) was stably expressed at all time points without any variation among samples after transfection. Sequences of *GUSB* gene primers are (Fw 5'-GAAAATACGTGTTGGAGAGCTCATT-3') and (Rev 5'-CCGAGTGAAGATCCCCTTTTA-3'). The following thermal cycling conditions were used: 50°C 2 min, 95°C 7 min, 95°C 15 s, 60°C 1 min for 40 cycles followed by dissociation stage: 95°C for 15 s, 60°C for 1 min, 95°C for 15 s and 60°C for 15 s. Amplification specificity was confirmed by dissociation curve analysis and specificity of amplified products was tested on agarose gel. Control without template (NTC) was included in each run. Post-run data were analysed using LinRegPCR software v.11.1. [37,38] which enables calculation of the starting concentration of amplicon ("N₀ value"). N₀ value is expressed in arbitrary fluorescence units and is calculated by taking into account PCR efficiency and baseline fluorescence. N₀ value determined for each technical replicate was averaged and the averaged N₀ values were divided by the N₀ values of the endogenous control. Statistical analysis of qPCR data was done using GraphPad v.6.01 and normalized N₀ values were compared using the unpaired t-test which compares the means of two unmatched groups.

4.6. Chromatin Immunoprecipitation

24 hours after transfection, MJ90hTERT cells were grown to subconfluence, washed in PBS, scraped in Nuclear Isolation buffer (10 mM MOPS; 5 mM KCl; 10 mM EDTA; 0.6% Triton X-100) with protease inhibitor cocktail CompleteMini (Roche) and chromatin immunoprecipitation was performed according to the published protocol [10,19], with the exception of sonication step which was performed 30 times for 30 s on ice, high sonicator amplitude. The antibodies used were: Anti-Histone H3 (tri methyl K9, Abcam, ab8898), Anti-Histone H3 (acetyl K18, Abcam, ab1191), Anti-Histone H3 (di methyl K4, Abcam, ab7766) and IgG (Santa Cruz Biotechnology, sc2027). Binding of precipitated target was monitored by qPCR using the SYBR Green PCR Master mix (Bio-Rad). Primers used for H3K9me3, H3K18ac and H3K4me2 level analyses of genes with intronic alpha satellite repeats are listed in Table S2. The N_0 values were normalized using N_0 values of input fractions.

4.7. Alpha Satellite RNA:DNA Hybrid Detection Assay

Approximately 2.5×10^6 of MJ90hTERT cells per sample were washed in PBS, harvested in medium, transferred into tubes and centrifuged at room temperature for 5 min at 400 g. The pellets were resuspended in 600 μ L of Lairds buffer (100 mM Tris pH8.5, 200 mM NaCl, 5 mM EDTA, 0.2% SDS). The samples were sonicated (45 s OFF, 15 s ON, 12 cycles) using a sonicator (high amplitude), centrifuged for 5 min at 12000 rpm (4°C) and supernatants transferred to new tubes. Total nucleic acids were purified with phenol:chloroform:isoamyl alcohol (Roth, A156.3), precipitated with 2.5 volumes of cold ethanol (96%) and 0.1 volume of 3 M sodium acetate (pH 5.2) and washed with 70% ethanol. The air-dried pellets were resuspended in 50 μ L of nuclease-free water. 10 μ g of chromatin-associated, phenol/chloroform-isolated nucleic acids per sample were incubated for 30 min at 37°C with 13 U of RNase H (NEB) in 1 $\times$ buffer (NEB) in a total volume of 50 μ L. The untreated (without RNase H) controls (10 μ g) were also incubated in the same way. RNase H treated and untreated samples were then double DNase I (Rnase-Free Dnase Set, QIAGEN) digested and total RNA was purified using RNeasy Mini Kit (QIAGEN), according to manufacturer instructions. All nucleic acid quantifications were carried out using Qubit fluorometer (Invitrogen).

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org Fig. S1: Consensus sequence of alpha satellite monomer; Fig. S2: Expression profiles of genes containing alpha satellite repeats within intronic regions, in MJ90hTERT cell lines transfected with satellite expressing vectors and controls, 48 hours after treatment; Fig. S3: Expression profiles of genes containing alpha satellite repeats within intronic regions, in MJ90hTERT cell lines transfected with satellite expressing vectors and controls, 72 hours after treatment; Fig. S4: Expression profiles of genes containing alpha satellite repeats within intronic regions, in MJ90hTERT cell lines transfected with satellite expressing vectors and controls, 96 hours after treatment; Fig.S5: Glucuronidase beta gene expression profile in MJ90hTERT cell line 24, 48, 72 and 96 hours after transfection; Fig. S6: Levels of H3K9me3, H3K18ac and H3K4me2 histone modifications at alpha satellite repeats associated with six genes after 24 hour transfection with 171Rev and unmodified control vectors. Table S1: List of primers used for expression analysis of alpha repeat-associated genes; Table S2: List of primers used for histone modification levels analyses of alpha repeats located within introns of genes, in ChIP-qPCR experiments.

Author Contributions: Conceptualization, I.F., Đ.U.; methodology, S.L., M.M.; formal analysis, S.L., M.M., D.Đ., M.C.F., A.P., I.F.; investigation, S.L., M.M., I.F.; writing original draft preparation, I.F. and Đ.U.; writing—review and editing, Đ.U., I.F., D.Đ.; supervision, I.F; All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by Croatian Science Foundation grants: IP-2019-04-6915 to Đ. Ugarković and by the Italian Ministry of Education, University and Research (MIUR), FFABR2017, fund for Investments on Basic Research (FIRB), by the International Staff Mobility Program of University of Naples Federico II to I. Feliciello.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The original contributions presented in the study are included in the article/supplementary material, further inquiries can be directed to the corresponding author/s.

Acknowledgments: We thank Marianna Feliciello for the graphical support.

Conflicts of Interest: The authors declare no conflict of interests.

References

1. Lee, C.; Wevrick, R.; Fisher, R.B.; Ferguson-Smith, M.A.; Lin, C.C. Human centromeric DNAs. *Hum. Genet.* 1997, 100, 291–304.
2. McNulty, S.M.; Sullivan, B.A. Alpha satellite DNA biology: Finding function in the recesses of the genome. *Chromosom. Res.* 2018, 26, 115–138.
3. Feliciello, I.; Pezer, Z.; Kordiš, D.; Bruvo-Madžarić, B.; Ugarković, Đ. Evolutionary history of α satellite DNA repeats dispersed within human genome euchromatin. *Genome Biol. Evol.* 2020a, 12, 2125–2138.
4. Feliciello, I.; Picariello, O.; Chinali, G. The first characterisation of the overall variability of repetitive units in a species reveals unexpected features of satellite DNA. *Gene* 2005, 349, 153–164.
5. Feliciello, I.; Picariello, O.; Chinali, G. Intra-specific variability and unusual organization of the repetitive units in a satellite DNA from *Rana dalmatina*: molecular evidence of a new mechanism of DNA repair acting on satellite DNA. *Gene* 2006, 383, 81–92.
6. Brajković, J.; Feliciello, I.; Bruvo-Madžarić, B.; Ugarković, Đ. Satellite DNA-Like Elements Associated with Genes Within Euchromatin of the Beetle *Tribolium castaneum*. *G3 Genes/Genomes/Genetics* 2012, 2, 931–941.
7. Feliciello, I.; Akrap, I.; Brajković, J.; Zlatar, I.; Ugarković, Đ. Satellite DNA as a driver of population divergence in the red flour beetle *Tribolium castaneum*. *Genome Biol. Evol.* 2015a, 7, 228–239.
8. Kuhn, G.C.; Küttler, H.; Moreira-Filho, O.; Heslop-Harrison, J.S. The 1.688 Repetitive DNA of *Drosophila*: Concerted Evolution at Different Genomic Scales and Association with Genes. *Mol. Biol. Evol.* 2011, 29, 7–11.
9. Ruiz-Ruano, F.J.; López-León, M.D.; Cabrero, J.; Camacho, J.P.M. High-throughput analysis of the satellitome illuminates satellite DNA evolution. *Sci. Rep.* 2016, 6, 28333.
10. Feliciello, I.; Akrap, I.; Ugarković, Đ. Satellite DNA Modulates Gene Expression in the Beetle *Tribolium castaneum* after Heat Stress. *PLoS Genet.* 2015b, 11, e1005466.
11. Ljubić, S.; Matulić, M.; Đermić, D.; Feliciello, I.; Procino, A.; Ugarković, Đ.; Feliciello, I. Antibiotics induce overexpression of alpha satellite DNA accompanied with epigenetic changes at alpha satellite arrays as well as genome-wide. *Epigenetics&Chromatin*, 2025, in press.
12. Sermek, A.; Feliciello, I.; Ugarković, Đ. Distinct Regulation of the Expression of Satellite DNAs in the Beetle *Tribolium castaneum*. *Int. J. Mol. Sci.* 2021, 22, 296.
13. Bury, L.; Moodie, B.; Ly, J.; McKay, L.S.; Miga, K.H.; Cheeseman, I.M. α -satellite RNA transcripts are repressed by centromere-nucleolus associations. *eLife* 2020, 9, e59770.
14. Teng Z, Yang L, Zhang Q, Chen Y, Wang X, Zheng Y, Tian A, Tian D, Lin Z, Deng WM, Liu H. Topoisomerase I is an evolutionarily conserved key regulator for satellite DNA transcription. *Nat Commun.* 2024;17;15(1), 5151.
15. Huang C, Shu X, Zhou S, Mi Y, Bian H, Li T, Li T, Ying X, Cheng C, Liu D, Gao M, Wen Y, Ma Q, Wang F, Cao J, Wang J, Liu J. Nuclear m6A modification regulates satellite transcription and chromosome segregation. *Nat Chem Biol.* 2025, doi: 10.1038/s41589-025-01900-9. Online ahead of print.
16. Johnson, W.L.; Yewdell, W.T.; Bell, J.; McNulty, S.M.; Duda, Z.; O'Neill, R.J.; Sullivan, B.A.; Straight, A. RNA-dependent stabilization of SUV39H1 at constitutive heterochromatin. *eLife* 2017, 6.
17. Velazquez Camacho, O.; Galan, C.; Swist-Rosowska, K.; Ching, R.; Gamalinda, M.; Karabiber, F.; De La Rosa-Velazquez, I.; Engist, B.; Koschorz, B.; Shukeir, N.; et al. Major satellite repeat RNA stabilize

heterochromatin retention of Suv39h enzymes by RNA- nucleosome association and RNA:DNA hybrid formation. *eLife* 2017, 6, 817.

18. Eymery, A.; Horard, B.; el Atifi-Borel, M.; Fourel, G.; Berger, F.; Vitte, A.-L.; Broeck, A.V.D.; Brambilla, E.; Fournier, A.; Callanan, M.; et al. A transcriptomic analysis of human centromeric and pericentric sequences in normal and tumor cells. *Nucleic Acids Res.* 2009, 37, 6340–6354.

19. Feliciello, I.; Sermek, A.; Pezer, Ž.; Matulić, M.; Ugarković, Đ. Heat stress affects H3K9me3 level at human α satellite DNA repeats. *Genes* 2020b, 11, 663.

20. Ugarković Đ, Sermek A, Ljubić S, Feliciello I. Satellite DNAs in Health and Disease. *Genes* 2022; 13(7), 1154.

21. Jolly, C.; Metz, A.; Govin, J.; Vigneron, M.; Turner, B.M.; Khochbin, S.; Vourc, H.C. Stress-induced transcription of satellite III repeats. *J. Cell Biol.* 2003, 164, 25–33.

22. Ninomiya, K.; Adachi, S.; Natsume, T.; Iwakiri, J.; Terai, G.; Asai, K.; Hirose, T. LncRNA-dependent nuclear stress bodies promote intron retention through SR protein phosphorylation. *EMBO J.* 2019, 39, e102729.

23. Ninomiya, K.; Iwakiri, J.; Aly, M.K.; Sakaguchi, Y.; Adachi, S.; Natsume, T.; Terai, G.; Asai, K.; Suzuki, T.; Hirose, T. m⁶A modification of HSATIII lncRNAs regulates temperature-dependent splicing. *EMBO J.* 2021, 40, e107976.

24. Ku WL, Nakamura K, Gao W, Cui K, Hu G, Tang Q, Ni B, Zhao K. Single-cell chromatin immunocleavage sequencing (scChIC-seq) to profile histone modification. *Nat Methods.* 2019, 16(4), 323–325.

25. Ku WL, Pan L, Cao Y, Gao W, Zhao K. Profiling single-cell histone modifications using indexing chromatin immunocleavage sequencing. *Genome Res.* 2021, 31(10),1831-1842.

26. Statello, L., Guo, C.J., Chen, L.L. et al. Gene regulation by long non-coding RNAs and its biological functions. *Nat Rev Mol Cell Biol* 2021, 22, 96–118.

27. Li, Y.; Syed, J.; Sugiyama, H. RNA-DNA Triplex Formation by Long Noncoding RNAs. *Cell Chem Biol.* 2016, 23(11):1325-1333.

28. Warwick, T.; Brandes, R.P.; Leisegang, M.S. Computational Methods to Study DNA:DNA:RNA Triplex Formation by lncRNAs. *Non-Coding RNA* 2023, 9, 10.

29. Ariel F et al. R-Loop Mediated trans Action of the APOLO Long Noncoding RNA. *Mol Cell* 2020, 77(5), 1055-1065.e4

30. Mondal, T.; Subhash, S.; Vaid, R.; Enroth, S.; Uday, S.; Reinius, B.; Mitra, S.; Mohammed, A.; James, A.R.; Hoberg, E.; et al. MEG3 long noncoding RNA regulates the TGF- β pathway genes through formation of RNA-DNA triplex structures. *Nat. Commun.* 2015, 6, 7743.

31. Duda, K.J.; Ching, R.W.; Jerabek, L.; Shukeir, N.; Erikson, G.; Engist, B.; Onishi-Seebacher, M.; Perrera, V.; Richter, F.; Mittler, G.; et al. m⁶A RNA methylation of major satellite repeat transcripts facilitates chromatin association and RNA:DNA hybrid formation in mouse heterochromatin. *Nucleic Acids Res.* 2021, 49, 5568–5587.

32. Cukusic Kalajzic A, Vidacek NS, Huzak M, Ivankovic M, Rubelj I. Telomere Q-PNA-FISH--reliable results from stochastic signals. *PLoS One.* 2014, 9(3), e92559.

33. Gorbunova V, Seluanov A, Pereira-Smith OM. Expression of human telomerase (hTERT) does not prevent stress-induced senescence in normal human fibroblasts but protects the cells from stress-induced apoptosis and necrosis. *J Biol Chem* 2002, 277, 38540–38549.

34. Young JI, Smith JR. DNA Methyltransferase Inhibition in Normal Human Fibroblasts Induces a p21-dependent Cell Cycle Withdrawal. *J Biol Chem* 2001, 276, 19610–19616.

35. Choo KH, Vissel B, Nagy A, Earle E, Kalitsis P. A survey of the genomic distribution of alpha satellite DNA on all the human chromosomes, and derivation of a new consensus sequence. *Nucleic Acids Res.* 1991, 19(6),1179-82

36. Đermić D, Ljubić S, Matulić M, Procino A, Feliciello MC, Ugarković Đ, Feliciello I. Reverse transcription-quantitative PCR (RT-qPCR) without the need for prior removal of DNA. *Sci Rep.* 2023;13(1),11470.

37. Ruijter, J.M.; Ramakers, C.; Hoogaars, W.M.H.; Karlen, Y.; Bakker, O.; Hoff, M.J.B.V.D.; Moorman, A.F.M. Amplification efficiency: Linking baseline and bias in the analysis of quantitative PCR data. *Nucleic Acids Res.* 2009, 37, e45.
38. Ruijter, J.M.; Pfaffl, M.W.; Zhao, S.; Spiess, A.N.; Boggy, G.; Blom, J.; Rutledge, R.G.; Sisti, D.; Lievens, A.; de Preter, K.; et al. Evaluation of qPCR curve analysis methods for reliable biomarker discovery: Bias, resolution, precision, and implications. *Methods* 2013, 59, 32–46.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.