

Article

Not peer-reviewed version

Nutritional Characterization of Annual and Perennial Glassworts from the Apulia Region (Italy)

[Luigi Giuseppe Duri](#) , [Lucia Botticella](#) , [Corrado Lazzizzera](#) , [Enrico Vito Perrino](#) , [Angelica Giancaspro](#) , [Annarita Cammerino](#) , [Anna Bonasia](#) , [Antonio Elia](#) , [Giulia Conversa](#) *

Posted Date: 1 September 2025

doi: 10.20944/preprints202509.0023.v1

Keywords: Halophytes; Mineral nutrients; Antioxidants; Phenols; Iodine; *Salicornia europaea*; *Sarcocornia fruticosa*; *Arthrocaulon macrostachyum*



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Nutritional Characterization of Annual and Perennial Glassworts from the Apulia Region (Italy)

Luigi Giuseppe Duri, Lucia Botticella, Corrado Lazzizzera, Enrico Vito Perrino, Angelica Giancaspro, Annarita Cammerino, Anna Bonasia, Antonio Elia and Giulia Conversa *

Department of Agriculture, Food, Natural Resources and Engineering, University of Foggia, Via Napoli 25, Foggia, 71100, Italy

* Correspondence: giulia.conversa@unifg.it

Abstract

Halophytes are increasingly recognized as sustainable crops that offer a wide range of nutrients. This study provides a nutritional characterization of annual (*Salicornia europaea*) and perennial (*Sarcocornia fruticosa*, *Arthrocaulon macrostachyum*) species of glasswort, collected from different coastal habitats in southern Italy. *S. europaea* was also cultivated under non-saline conditions. Results showed differences in mineral content, and bioactive compounds among genotypes, but they were modulated by environmental conditions, leading to significant site-specific variation. *S. europaea*, regardless of the collecting sites, exhibited the highest concentration of minerals (K, Ca, and Mg), chlorophylls, carotenoids, and phenolic compounds as well as antioxidant activity. *A. macrostachyum* stood out for its high flavonoid and sterol content, exhibiting other nutritional traits comparable to *S. europaea* when collected in a more arid site. *A. macrostachyum* and *S. fruticosa* displayed similar compositional features, showing the highest anthocyanin and iodine (187.8 $\mu\text{g } 100 \text{ g}^{-1} \text{ FW}$, on average) content. Sodium and potassium levels varied, exceeding the recommended Na/K ratio (1) for human consumption, especially in *A. macrostachyum* grown close to the sea. The Na/K closest to 1 was observed in non-saline cultivated *S. europaea*. Overall findings confirm the potential of both annual and perennial glassworts as nutritionally rich, sustainable crops for marginal environments.

Keywords: halophytes; mineral nutrients; antioxidants; phenols; iodine; *Salicornia europaea*; *Sarcocornia fruticosa*; *Arthrocaulon macrostachyum*

1. Introduction

Soil salinization and freshwater scarcity are growing global issues, making the increasing salinization of arable land a critical point of land degradation [1]. This, coupled with the increasing challenges of climate change, is a goal that the Food and Agriculture Organization of the United Nations [2] considers a priority, highlighting the importance of research on sustainable and resilient food sources. Halophytes are a group of plants that thrive in saline environments (coastal and inland areas) and have attracted significant interest as a potentially sustainable option for agriculture [3]. These plants can grow in marginal lands that face multiple constraints such as drought, fluctuating temperatures, high light irradiance, and elevated salinity [4,5], without competing for primary resources like freshwater. Many halophyte species are edible plants possessing unique biochemical properties that make them valuable as novel foods [6]. To complete their life cycle in saline environments, these plants have developed adaptive mechanisms that involve the production of secondary antioxidant metabolites [4,5].

Salicornioideae (Amaranthaceae/Chenopodiaceae *sensu* The Angiosperm Phylogeny Group [7]) is an important halophyte subfamily with many genera, such as *Salicornia* L., *Sarcocornia* A.J. Scott, and *Arthrocaulon* Piirainen & G.Kadereit.

Among these genera, *Salicornia europaea* Auct. (syn.: *Salicornia ramosissima* Auct., *Salicornia patula* Duval-Jouve, *Salicornia perennans* Willd. subsp. *perennans*) [8,9] is an annual taxon with a Western

European geographic distribution colonising periodically inundated muds and sands of marine or interior salt marshes.

Sarcocornia fruticosa (L.) A.J. Scott (homotypic synonyms: *Arthrocnemum fruticosum* (L.) Moq., *Salicornia europaea* var. *fruticosa* L., *Salicornia fruticosa* (L.) A.J. Scott) [8–10] is a perennial species with a Euro-Mediterranean and South African distribution, that colonizes coastal salty marshes periodically flooded by tides with close to marine salinity. It grows in silty-clayey soils, poorly drained, flooded for more or less long periods by salt water.

Arthrocaulon macrostachyum (Moric.) Piirainen & G.Kadereit [homotypic synonyms: *Arthrocnemum fruticosum* var. *macrostachyum* (Moric.) Moq., *Arthrocnemum macrostachyum* (Moric.) K. Koch, *Salicornia fruticosa* var. *pachystachya* W.D.J. Koch, *Salicornia macrostachya* Moric. [8–10] is a Macaronesian Mediterranean distribution taxon which differs ecologically from *S. fruticosa* as it prefers salty coastal places with salinity higher than marine and often with a rocky substrate. It is in salt marshes, in sites not directly affected by tide variations, where they grow on the reliefs between the ponds.

Due to difficult taxonomy, leafless, succulent and fleshy stems of species belonging to *Salicornia* [11,12] and, more recently, those of *Sarcocornia* [13] and *Arthrocaulon* [14] genera are commercialised and/or consumed mixed as a green leafy vegetable, commonly known as glassworts or, particularly in most European countries, as ‘sea asparagus’ [13].

The product is considered a niche offering because this salt-tolerant vegetable has traditionally been harvested from wild plants and sold in local markets. However, the expanding cultivation of this vegetable in both soil [15] and soilless systems [16] has been driven by increasing demand. As a result, its commercial economic importance is on the rise [17].

In the Mediterranean basin, studies evaluating various aspects of the nutritional composition of these plants have been conducted mainly in the Iberian Peninsula and some North African and Middle Eastern countries. The genus *Salicornia* is rich in compounds associated with oxidative stress (drought, salinity), such as vitamin A, fatty acids, ascorbic acid, carotenoids and phenolic compounds, as discussed in the reviews by Cárdenas-Pérez et al. [12], Ekanayake et al. [17], and Sánchez-Gavilán et al. [18]. Similar nutritional aspects were noted in species *Sarcocornia* [19] and *Arthrocaulon* [14,20]. However, considering the effects of both genotype and environmental factors, along with their interactions, there have been reports of changes in the nutritional profile of glasswort products. Higher antioxidant and mineral contents were observed in the wild *Sarcocornia perennis* than in *Salicornia ramosissima* collected in the same salt marshes [19]. Changes in fatty acid and phenolic profiles were detected between *Arthrocnemum* and *Sarcocornia* genera, despite collecting from the same areas in Spain and Portugal [21]. Cultivated *Salicornia* ecotypes exhibited enhanced fatty acid (total and profile) compared to those of the *Sarcocornia* [22].

Besides the genotype, the pedoclimatic conditions present in natural habitats and/or cultivation systems significantly influence the nutritional composition of glassworts.

Wild *Salicornia patula* had a variable phenol profile based on its mainland or coastal origin [23]. Similarly, the fatty acid profiles of *S. ramosissima* differed depending on their geographical origin [24]. When ecotypes of *Sarcocornia* and *Salicornia* were cultivated in a soilless system with increasing seawater salinity, they showed increased mineral, carotenoid [25], and phenolic content [22]. Drought stress in cultivated *S. ramosissima* also led to an increase in total phenols, fatty acids, and oxalates [26]. Overall, wild *Sarcocornia* had higher levels of carbohydrates, lipids, and phenolic compounds [27] but lower mineral content [28] compared to cultivated varieties.

In the Apulian region (Italy), both annual and perennial glassworts are consumed, mainly in the Foggia province, characterised by wide coastal salt marshes, followed by the Brindisi and Taranto areas, through the gathering of wild plants. In Foggia province, the cultivation of the annual *Salicornia* species, which started more than 40 years ago for family consumption, has now evolved into a cash crop covering a few dozen hectares (Conversa, personal communication). Despite their importance for food purposes, to our knowledge, there is a lack of comprehensive research characterizing the nutritional profiles of both cultivated and wild Apulian glassworts. Based on the above, the study

aims to investigate the mineral nutrients, antioxidative compounds and antinutritional traits characterizing species of *Salicornia*, *Sarcocornia* and *Arthrocaulon* genera cultivated or collected along the coasts of the Apulia region.

2. Materials and Methods

2.1. Sampling Area and Glasswort Description

Glasswort samples were collected in Apulian coastal areas (the geographical coordinates are in the WGS84 system):

1) Margherita di Savoia Saltworks (MSS), [Margherita di Savoia Barletta–Andria- Trani province (BT)] (site a: 41° 25' 54.39" N; 16° 0' 31.395" E; site b: 41° 25' 39.512" N; 16° 0' 28.46" E);

2) Laguna del Re Oasis (LRO) [Manfredonia, Foggia province (FG)] (site a: 41° 34' 40.8" N; 15° 53' 5.8" E; site b: 41° 34' 47.86" N; 15° 53' 0.12" E);

3) Varano lake (VL) (Cagnano Varano, FG) (41° 54' 51.5" N; 15° 48' 17.6" E);

4) Monti d'Arena-Bosco Caggione (MA-BC) [Taranto province (TA)] (40° 20' 54.4" N; 17° 22' 31.9" E);

5) 'Spirito Contadino' (SC) commercial farm (Borgo Tressanti, FG) (41° 23' 44.8" N; 15° 49' 25.2" E).

Samples of wild perennial species were collected at 1) MSS site a (*Sarcocornia fruticosa* L.), and site b (*Arthrocaulon macrostachyum* Moric.), 2) VL (*Sarcocornia fruticosa* L.), and 3) MA-BC (*Arthrocaulon macrostachyum* Moric.). At location LRO, we harvested samples of glasswort both from the wild perennial (*Arthrocaulon macrostachyum* Moric., site a) and cultivated annual plants (*Salicornia europaea* Auct. site b); at site SC, conventionally cultivated (no saline conditions) annual plants (*Salicornia europaea* Auct.) were collected.

Locations MSS, LRO, and VL were within coastal marsh wetlands with different soils, from predominantly sandy to clayey, to rarely rocky, whereas MA-BC is a fully rocky coastal site. The SC site falls within the Tavoliere plain, where intensive non-saline agriculture is practised [29].


In the MSS area, *S. fruticosa* (*S. fruticosa*-MSS) plants were located along the edge and at the mouth of the brackish water Aloisa canal (site a), whereas *A. macrostachyum* (*A. macrostachyum*-MSS) was found adjacent to the salt pan area, 0.5 km away from the coast (site b). *A. macrostachyum* was also gathered at the MA-BC (*A. macrostachyum*-MA-BC) site, on the rocky seacoast, 1-2 m high. At the VL site, *S. fruticosa* (*S. fruticosa*-VL) was in a brackish area that is rarely flooded, close to the mouth of the salt Varano lake. The LRO location is characterized by the alternation of flooded areas and dry land for agricultural use, both with highly saline and sodic, often clayey soil [30]. Specifically, *A. macrostachyum* (*A. macrostachyum*-LRO) was along the edges of a brackish water stream (site a), whereas *S. europaea* was cultivated on reclaimed land for agriculture (site b) (*S. europaea*-LRO). The samples collected were summarized in Table 1, and Figure 1 shows the location of the sampling sites.

For cultivation, plantlets of *S. europaea* were provided by the nursery 'Aurora' (San Nicandro Garganico, FG- Italy). The transplant was performed in April (SC) and May (LRO) 2023. At site LRO-b, the soil and irrigation water's electric conductivity (EC) were 1.9 and 21.5 mS cm⁻¹, respectively [30]. In contrast, EC of soil and irrigation water was, respectively, 0.2 and 0.6 mS cm⁻¹ at the SC site.

Edible tips (2 kg) from at least 20 plants of each site were randomly collected during the April-June 2023 period (wild samples) and the August-September period (cultivated samples). For each site, the harvested plants were divided into three subsamples, then washed with distilled water, dried with paper, and subjected to freeze-drying (ScanVac CoolSafe 55-9 Pro; LaboGene ApS, Lyngø, Denmark) for dry mass determination (g·kg⁻¹ of fresh weight, FW) and subsequent analyses, performed with six repetitions for each subsample (n=18).

Table 1. Sampling areas, taxonomic classification and sample identification of the collected glassworts.

Sampling Area	Taxonomic Classification	Sample Identification
Margherita di Savoia Saltworks (MSS), site a	<i>Sarcocornia fruticosa</i> L.	<i>S. fruticosa</i> -MSS
Margherita di Savoia Saltworks (MSS), site b	<i>Arthrocaulon macrostachyum</i> Moric.	<i>A. macrostachyum</i> -MSS
Laguna del Re Oasis (LRO), site a	<i>Arthrocaulon macrostachyum</i> Moric.	<i>A. macrostachyum</i> -LRO
Laguna del Re Oasis (LRO), site b	<i>Salicornia europaea</i> Auct.	<i>S. europaea</i> -LRO
Varano lake (VL)	<i>Sarcocornia fruticosa</i> L.	<i>S. fruticosa</i> -VL
Monti d'Arena-Bosco Caggione (MA-BC)	<i>Arthrocaulon macrostachyum</i> Moric.	<i>A. macrostachyum</i> -MA-BC
Spirito Contadino farm (SC)	<i>Salicornia europaea</i> Auct.	<i>S. europaea</i> -SC

**Figure 1.** Satellite image of sampling sites. Please see Table 1 for sample identification.

2.2. Inorganic Ions and Iodine Determination

2.2.1. Anions and Cations

Anions were extracted from 0.5 g of lyophilized sample with 50 mL of eluent solution in a shaking water bath at room temperature for 30 min and filtered using first Whatman no. 2 paper and then a 0.22 μm Millipore filter, before injection into the ion chromatography system (ICS-3000, Dionex Corp., Milan, Italy). The system was equipped with an isocratic pump (AS-DV autosampler), a self-generating ASR anion suppressor (4 mm), a Dionex Ion-Pac AS23 (Dionex Corp., Milan, Italy) analytical column (4 mm \times 250 mm), and a guard column (4 mm \times 50 mm) maintained at 35 $^{\circ}\text{C}$. The eluent consisted of 3.5 mM sodium carbonate and 1 mM sodium bicarbonate at a flow rate of 1 mL min^{-1} .

The cations were extracted from dried samples (0.3 g), previously ashed (in a muffle furnace at 550 $^{\circ}\text{C}$ for 6 h) and acid digested (20 mL of 1 mol L^{-1} HCl in boiling water for 30 min). The resulting

solution was filtered through a 0.22 μm Millipore filter, diluted and analyzed by ion chromatography (Dionex ICS 3000; Dionex-ThermoFisher Scientific, Waltham, MA, USA). The ion chromatography system was equipped with: an isocratic pump; conductivity detector; a model AS-DV auto-sampler (ThermoFisher Scientific, Waltham, MA, USA) a self-generating DRS-600 suppressor (4 mm), a Dionex IonPack CS12A analytical column (4 \times 250 mm, 5 μm), and an eluent solution (20 mM methanesulfonic acid) at a flow rate of 1 mL min⁻¹.

Anions and cations were identified by retention times in comparison with standards, using Dionex Chromeleon software (version 6.80, Thermo Scientific) for peak areas analysis. The results are expressed as g·kg⁻¹ FW.

2.2.2. Iodine

For the extraction, 0.1 g of the sample was treated with 25 mL of hot distilled water (60 °C), stirring at room temperature for 30 min. When cooled, 25 mL of distilled water was added and centrifuged for 15 min at 6000 rpm, then filtered twice, the first using Whatman filter paper no. 2 and the second involving a syringe membrane filter with 0.2 μm of mesh [31]. Thereafter, 0.5 mL of extracted sample with iodate standard solution was added to 4.5 mL of distilled water, and treated with 1 mL of KSCN 2.37 mM, 2 mL of NH₄Fe(SO₄)₂·12H₂O (7.7 % m/v) in 2.4 M HNO₃ and 2 mL of NaNO₂ (0.02 % m/v). The mixed solution was incubated for 1 h at 60 \pm 2 °C in a water bath. To stop the reaction, the mixture was placed in a water-ice bath. After extraction, iodine determination was performed using a spectrophotometer (Shimadzu UV-1800, Shimadzu Italia S.r.l. Milano, Italia) set at 454 nm. The quantification was determined using a calibration curve (0–12 $\mu\text{g L}^{-1}$; R₂ = 0.996), and the results are expressed as $\mu\text{g 100 g}^{-1}$ FW.

2.3. Pigments and Antioxidant Compounds

2.3.1. Chlorophylls and Anthocyanins

Chlorophyll a, b, and total and carotenoid concentrations were determined by modifying the spectrophotometric method proposed by Sumanta et al. [32]. Briefly, 1.5 mL 80% Ethanol (ethanol:water=80:20) containing 0.1% hydrochloric acid was mixed with 0.05 g of lyophilized glasswort into a 10 mL tube with a screw cap. The mixture was placed for 30 min at room temperature in an ultrasonic bath (DU-32 Digital ultrasonic cleaner, Argo Lab, Carpi, MO, Italy). After sonication, the sample was centrifuged in a refrigerated (4°C) centrifuge (Beckman Coulter Allegra™ 25, Fullerton, CA, USA) for 15 min at 4,000 g, then the supernatant was collected. The steps were repeated twice, combining the supernatants. The extract was filtered at 0.22 μm . To measure the filtrate was employed a spectrophotometer (Evolution 201UV-Visible Spectrophotometers, Thermo Scientific Waltham, MA, USA) set at 664, 649, and 470 nm (for chlorophyll a, b, and carotenoid, respectively).

To quantify chlorophylls and carotenoids, the following equations were used:

$$\text{Chl-a} = 13.36(A_{664}) - 5.19(A_{649})$$

$$\text{Chl-b} = 27.43(A_{649}) - 8.12(A_{664})$$

$$\text{Chl-total} = 17.32(A_{649}) + 7.18(A_{664})$$

$$\text{Total carotenoid} = (1000(A_{470}) - 2.13 \text{ Chl-a} - 97.63 \text{ Chl-b})/209$$

Chlorophyll results were expressed in $\mu\text{g g}^{-1}$ FW, while the carotenoids in mg 100 g⁻¹ FW.

Samples (0.1 g) were mixed with 3 mL 80% ethanol (ethanol:water=80:20) containing 0.1% hydrochloric acid into a 10 mL tube with a screw cap. DU-32 Digital ultrasonic cleaner (Argo Lab, Carpi, MO, Italy) was employed to sonicate the mixture for 30 min at room temperature. After sonication, the samples were centrifuged in a refrigerated centrifuge (Beckman Coulter Allegra™ 25, Fullerton, CA, USA) for 15 min at 4,000 g at 4°C, and the supernatant was collected. The protocol mentioned above was repeated twice, and supernatants were combined. The extract was filtered at 0.22 μm and measured with the spectrophotometer at 650 and 525 nm. The equation (anthocyanin absorbance = A₅₂₉ - (0.288 \times A₆₅₀)) proposed by Sims and Gamon [33] was involved in correcting the interferences due to chlorophylls. Total anthocyanin content was then calculated in cyanidin-3-

glucoside using this corrected absorbance and a molar absorbance coefficient for anthocyanin at 525 nm of $26.900 \text{ L mol}^{-1} \text{ cm}^{-1}$ [34].

2.3.2. Phenols and Flavonoids

Total phenols and flavonoids extraction were carried out on 30 mg of the sample with 1 mL of water:methanol (20:80, v:v) at room temperature in an ultrasonic cleaner bath for 15 min; then the mixture was centrifuged in a refrigerated centrifuge (14,000 rpm for 15 min at 4 °C) and the supernatant was collected. The extraction was repeated two times, and the supernatants were combined. The extracts were stored at -20 °C and measured within 24 h.

Total polyphenol concentration was determined on methanolic extracts according to the method of Singleton and Rossi [35]. Briefly, 100 μL of the extracts were diluted with 3 mL of distilled water, mixed with 0.5 mL of Folin–Ciocalteu reagent, and placed at room temperature for 5 min. Then 1.0 mL of 20% Na_2CO_3 was added to the mixture. After 45 min at 30 °C.

On the other hand, the determination of total flavonoid content is conducted based on the AlCl_3 method [36]. In detail, 200 μL of extract were mixed with 40 μL of 10% AlCl_3 , 40 μL of 1 M potassium acetate, and 1120 μL of distillate water. Samples were incubated for 30 minutes at room temperature. The same was performed with standard quercetin solutions.

The absorbances were read at 750 and 415 nm, respectively, for phenols and flavonoids (Shimadzu UV-1800, Shimadzu Scientific Instruments, North America, USA).

The results are expressed in mg 100 g^{-1} of FW as gallic acid equivalents (GAE) for phenols and in QE (Quercetin equivalent) for flavonoids.

2.3.3. Phytosterols

Total sterols were quantified as reported by Xiang et al. [37] with some modifications. In a screw-cap tube, 100 mg of lyophilized glasswort was mixed with 3 mL of sulfuric acid solution (0.5 mol L^{-1} in 95% ethanol) and vortexed for 10 seconds. The addition of 3 mg of tert-butyl-hydroquinone prevented the thermal decomposition of phytosterols. Acid hydrolysis was performed at 55 °C for 1 h. Alkali saponification (performed at 55 °C for 1 h) was carried out by adding 3 mL of sodium hydroxide solution (2.0 mol L^{-1} in 95% ethanol). Then, 3 mL of sodium chloride solution (1.0 mol L^{-1}) was added to prevent emulsification and facilitate hexane extraction. A double extraction was performed by adding hexane (5 mL). After stirring the mixture (15 minutes), the two supernatants were combined. Subsequently dried using a rotary evaporator (Strike 300, Steroglass, San Martino in Campania, AV, Italy) and then dissolved in absolute ethanol (1.0 mL) creating the crude extract.

The crude extract was purified by solid phase extraction (SPE) column with a 500 mg-C18-cartridge (Strata C18-E, Phenomenex, Castel Maggiore, BO, Italy). The complete SPE procedure includes methanol activation, water equilibrium, sample loading, and elution.

The concentration of total phytosterols was measured according to the spectrophotometry method described by Feng et al. [38], placing 10 g of $\text{FeCl}_3 \cdot (6\text{H}_2\text{O})$ into a volumetric flask (100 mL) and diluting with phosphoric acid (85%) to reach the volume. Then, 1.5 mL of FeCl_3 solution (10%) was transferred to a 100 mL volumetric flask and diluted with concentrated sulfuric acid. Subsequently, the present sulfuric-phosphoric- FeCl_3 reagent (2 mL) was combined with stigmasterol in ethanol solutions (4 mL) at concentrations ranging from 19 to 45.5 $\mu\text{g mL}^{-1}$ and with the sample extract solution (4 mL). Left reacting for 15 minutes, afterwards the absorption was measured using a spectrophotometer at a wavelength of 520 nm. The concentration was calculated based on a standard curve. The results were expressed as mg of stigmasterol equivalent per 100 g^{-1} FW.

2.4. Antioxidant Capacity

We assessed the antioxidant capacity by employing three assays to determine the radical scavenging capacity, differentiating the lipophilic fraction from the hydrophilic one, namely the

ABTS, DPPH, and FRAP methods [39,40], followed described in detail, and the detection of three assays were expressed in $\mu\text{mol eq. trolox g}^{-1}$ FW.

2.4.1. ABTS Assay

A slightly modified version of the TEAC (Trolox Equivalent Antioxidant Capacity) method was involved in this detection, applying the same decolorization assay for both hydrophilic and lipophilic antioxidants, delving into 30 mg of lyophilized sample was extracted with 1 mL of 80% methanol in an ultrasonic bath (DU-32 Digital ultrasonic power, Argo Lab, Carpi, MO, Italy) for 15 min. Then, it was centrifuged at 13,000 rpm for 10 min, collecting the supernatant in the end. This step was repeated two times, and the two supernatants were combined to determine the hydrophilic fraction. The residue was extracted twice with 1 mL of hexane in ultrasonic power for 15 min, combining both into a lipophilic fraction. The extracts were stored at 20°C and measured within 24 h. The measure of absorbance was performed at 734 nm after 6 min standing. Values were determined from a calibration curve prepared with Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid) solutions prepared following the same procedure as for the extracted sample. The Trolox stock solution was 1.100 μM , and the concentration range of the calibration curve was 0-15 μM .

2.4.2. DPPH Assay

The antioxidant activity using the DPPH (2,2- diphenyl-1-picrylhydrazyl) method was assessed only for a hydrophilic fraction (methanolic extracts). The extraction of the lyophilized sample was carried out following the same procedure applied in the ABTS method for the hydrophilic fraction. After the storage at 20°C for 24h DPPH was progressively solubilized in ethanol until a concentration producing an absorbance ranged from 1.12 to 1.08 at 517 nm. Preparing the analysis solution by mixing 1.900 μL of diluted DPPH and 100 μL of extract sample or Trolox standard solution. The Trolox stock solution was 1.000 μM and the concentration range of the calibration curve was 0-800 μM . The absorbance was measured at 517 nm after 30 min, standing in the dark. Values were determined from a calibration curve prepared with Trolox solutions following the same procedure as for the extracted sample.

2.4.3. FRAP Assay

The antioxidant activity was assessed using a slightly modified version of Ferric Reducing Antioxidant Power Assay (FRAP) method. The extraction of lyophilized was carried out like the other assays. The FRAP reagent was prepared by combining 2.5 mL of a TPTZ solution (10 mM) in HCl (40 mM), 2.5 mL of a FeCl_3 solution (20 mM), and 25 mL of NaOAc buffer (300 mM, pH 3.6). Thereafter, 1.8 μL of FRAP reagent was mixed with 40 μL of the sample extracts and 200 μL of deionized water. Keeping the reaction for 4 min at room temperature and measuring the absorbance at 593 nm. Values were determined from a calibration curve obtained with Trolox solutions at fixed concentrations, following the same procedure as for the extracted sample. The Trolox stock solution was 1.000 μM , and the range of the calibration curve was 0-800 μM .

2.5. Statistical Analysis

One-way statistical analysis was performed with the Statistical Analysis System software (SAS 9.1; SAS Institute, Cary, NC, USA) using the General Linear Model (GLM Proc of the SAS Software). The mean comparison was performed using Tukey's HSD test with a significance level of $\alpha = 0.05$.

The principal component analysis (PCA) was performed using the PAST3 Software (<http://folk.uio.no/ohammer/past>) on mean standardized data. The data standardization was performed using the formula: $(x - \text{mean}) / \text{standard deviation}$. The data matrix considered all glasswort collected samples with relative replications.

3. Results

3.1. Dry Mass and Mineral Concentrations

The examined glassworts had similar dry mass (DM) concentrations, except for the highest value for *Arthrocaulon macrostachyum* samples collected in Margherita di Savoia saltworks (MSS) (*A. macrostachyum*-MSS) and the lowest one registered for *A. macrostachyum* sample from Laguna del Re Oasis (*A. macrostachyum*-LRO) (Table 2).

Fluctuation in sodium concentration ranged from 6.9 to 20.9 g·kg⁻¹ FW, with *A. macrostachyum* samples exhibiting a larger variability (Table 2). The highest K, Mg and Ca levels were found for *S. europaea*-SC and -LRO, especially compared to *A. macrostachyum*-LRO samples.

A. macrostachyum-MSS showed a higher chloride concentration, whereas Cl ranged from 8.9 to 21.7 g·kg⁻¹ FW in the other sampled plants. Reduced Cl levels were registered for *Sarcocornia fruticosa* from MSS (*S. fruticosa*-MSS) (Table 2).

Nitrate level was the highest in *A. macrostachyum* collected from Taranto province locality (*A. macrostachyum*-MA-BC) and *Salicornia europaea* cultivated at the commercial farm (*S. europaea*-SC). This anion decreased in other genotypes up to not detectable values in *A. macrostachyum*-LRO (Table 2).

Variability in oxalate levels was observed, with *Sarcocornia fruticosa* from Varano Lake (*S. fruticosa*-VL) showing the highest value, followed by samples belonging to *A. macrostachyum* species.

The maximum level of iodine was observed for genotypes collected from the LRO site (*S. europaea*-LRO and *A. macrostachyum*-LRO) while the lower values were found in samples collected from MSS (both *S. fruticosa* and *A. macrostachyum*) as well as in *S. europaea* from the SC farm (Table 2).

Table 2. Dry mass and mineral contents of annual and perennial glassworts collected from different areas of the Apulia region.

Glasswort sample	Dry mass	Na	K	Mg	Ca	Cl	NO ₃	C ₂ O ₄	I
				g·kg ⁻¹ FW					μg 100 g ⁻¹ FW
<i>S. europaea</i> -LRO	178.3 b ⁽¹⁾	20.9 a	8.7 ab	2.2 ab	6.1 ab	21.9 b	0.1 b	0.2 d	349.4 a
<i>S. europaea</i> -SC	194.2 b	14.8 abc	9.0 a	2.8 a	6.9 a	15.4 c	0.4 a	0.8 cd	96.5 c
<i>S. fruticosa</i> -MSS	187.0 b	13.4bcd	4.0 bc	1.4 bc	4.2 abc	1.9 e	0.1 b	0.0 d	104.8 c
<i>S. fruticosa</i> -VL	189.0 b	14.9 abc	4.1 abc	1.3 c	2.5 bc	12.9 cd	0.1 b	6.6 a	212.0 abc
<i>A. macrostachyum</i> -MA-BC	187.0 b	17.1 ab	3.4 c	1.2 c	3.6 abc	21.7 b	0.4 a	1.5 c	223.7 abc
<i>A. macrostachyum</i> -LRO	104.6 c	6.9 d	1.9 c	0.8 c	1.8 c	8.9 d	0.0 b	0.8 cd	277.4 ab
<i>A. macrostachyum</i> -MSS	242.2 a	9.8 cd	5.4 abc	1.5 bc	5.7 ab	37.5 a	0.1 b	3.0 b	112.1 bc
Significance ⁽²⁾	***	***	***	***	***	***	***	***	***

⁽¹⁾ Within each column, different letters indicate significant differences according to Tukey's HSD test at $\alpha \leq 0.05$;

⁽²⁾ ***: significant at $p < 0.001$. Values are the means of eighteen subsamples (n = 18).

3.2. Bioactive Compound Concentrations and Antioxidant Capacity

Chlorophyll levels (CHLa, CHLb) in *S. europaea* and *A. macrostachyum*-MSS were significantly higher (more than 10 times) than in other collected glassworts (Table 3). Carotenoid concentrations were also elevated in *S. europaea*, particularly in the commercial farm sample (*S. europaea*-SC), followed by *S. fruticosa*-MSS. Total phenol (TP) content in *S. europaea* and *A. macrostachyum*-MSS was approximately 4.5 times higher than that in the other samples. Higher variability was detected for flavonoids (FL) with *A. macrostachyum*-MSS having the greatest amount. FL concentration was halved in *S. europaea* and decreased to about one-fourth in other *A. macrostachyum* genotypes, whereas negligible was the level of FL in *S. fruticosa*-MSS (Table 3). Anthocyanin content fluctuated, with clear differences observed between *S. fruticosa*-MSS (higher values) and *A. macrostachyum*-MSS (Table 3).

Sterol concentration in *A. macrostachyum*-MSS was 3.6-fold higher than in *A. macrostachyum*-MA-BA, *S. fruticosa*-VL and *S. europaea*-SC. The glasswort of Laguna del Re Oasis (LRO) had very low sterol levels (Table 3).

The antioxidant capacity (Table 4), tested through three assays, was greater for *S. europaea*-SC followed by the other annual sample (*S. europaea*-LRO) and *A. macrostachyum*-MSS. Whereas lower values were measured in the other perennial plants belonging to the species *A. macrostachyum* and *S. fruticosa*, indistinctly from the native site.

Table 3. Pigment and antioxidant compound contents of annual and perennial glassworts collected from different areas of the Apulia region. Specifically: chlorophyll a, b, and total (CHL_a, b, t), carotenoids (Car), Anthocyanins (Anth), Total Phenols (TP), Flavonoids (FL), and Sterols.

Glasswort sample	CHL _a	CHL _b	CHL _t	Car	TP	FL	Anth	Sterols
	μg g ⁻¹ FW			mg 100 g ⁻¹ FW	mg g.a.e. ⁽²⁾ 100 g ⁻¹ FW	mg q.e. ⁽²⁾ 100 g ⁻¹ FW	mg c.g. ⁽²⁾ 100 g ⁻¹ FW	mg 100 g ⁻¹ FW
<i>S. europaea</i> -LRO	134.4 a ⁽¹⁾	49.5 a	180.8 a	4.1 b	258.3 a	31.7 bc	0.7 bc	7.4 c
<i>S. europaea</i> -SC	127.0 a	43.0 a	167.7 a	6.2 a	192.9 a	39.3 b	1.0 abc	12.2 bc
<i>S. fruticosa</i> -MSS	11.5 b	8.0 b	18.7 b	3.5 b	19.3 b	0.3 e	1.6 a	n.d. ⁽³⁾
<i>S. fruticosa</i> -VL	7.8 b	6.3 b	13.4 b	1.7 c	26.6 b	26.2 cd	1.1 abc	17.0 b
<i>A. macrostachyum</i> -MA-BC	12.3 b	10.8 b	21.8 b	1.3 c	78.4 b	22.0 cd	1.4 ab	15.3 bc
<i>A. macrostachyum</i> -LRO	6.9 b	4.7 b	11.0 b	1.2 c	83.7 b	18.0 d	0.6 bc	7.3 c
<i>A. macrostachyum</i> -MSS	119.7 a	41.5 a	159.0 a	2.8 bc	256.2 a	73.9 a	0.4 c	53.3 a
Significance ⁽⁴⁾	***	***	***	***	***	***	***	***

⁽¹⁾ Within each column, different letters indicate significant differences according to Tukey's HSD test at $\alpha \leq 0.05$;

⁽²⁾ g.a.e.= gallic acid equivalents; q.e.= quercetin equivalent; c.g.= cyanidin-3-glucoside. ⁽³⁾ n.d.= not detected; ⁽⁴⁾ *** = significant at $p < 0.001$. Values are the mean of eighteen subsamples ($n = 18$).

Table 4. Antioxidant capacity of annual and perennial glassworts collected from different areas of the Apulia region tested through DPPH, ABTS and FRAP assays.

Glasswort sample	DPPH	ABTS			FRAP		
		HA ⁽²⁾	LA ⁽³⁾	Total	HA ⁽²⁾	LA ⁽³⁾	Total
μmol eq trolox g ⁻¹ FW							
<i>S. europaea</i> -LRO	8.2 b ⁽¹⁾	12.4 c	0.6 a	13.0 c	19.5 b	0.00 c	19.5 b
<i>S. europaea</i> -SC	13.0 a	22.9 a	0.7 a	23.7 a	71.6 a	0.00 c	71.6 a
<i>S. fruticosa</i> -MSS	3.6 d	6.9 d	0.0 c	6.9 d	5.8 d	0.26 b	6.2 d
<i>S. fruticosa</i> -VL	5.8 c	7.3 d	0.0 c	7.3 d	8.4 cd	0.43 a	8.8 cd
<i>A. macrostachyum</i> -MA-BC	2.6 d	5.4 d	0.0 c	5.4 d	5.0 d	0.33 ab	5.3 d
<i>A. macrostachyum</i> -LRO	4.1 d	4.4 d	0.0 c	4.4 d	7.8 d	0.40 ab	7.5 d
<i>A. macrostachyum</i> -MSS	9.4 b	17.8 b	0.3 b	18.1 b	14.3 bc	0.10 c	14.4 bc
Significance ⁽⁴⁾	***	***	***	***	***	***	***

⁽¹⁾ Within each column, different letters indicate significant differences according to Tukey's HSD test at $\alpha \leq 0.05$;

⁽²⁾ HA= hydrophilic fraction of antioxidants; ⁽³⁾ LA= lipophilic fraction of antioxidants; ⁽⁴⁾ ***= significant at $p < 0.001$. Values are the mean of eighteen subsamples ($n = 18$).

4. Discussion

Glassworts, belonging to different genera of the Salicornioideae subfamily, can provide valuable functional foods and dietary supplements, raising the interest of the scientific community for their nutraceutical profile [6]. In the Mediterranean region, research has focused on the nutritional properties of these halophytes. In contrast, there is a relative lack of studies on the nutraceutical profile of both spontaneous and cultivated glassworts in Italy. This work explored their potential as a novel food source by examining both their functional properties and antinutritional traits, as well as how these factors change with species and provenance. To the best of our knowledge, this is the first study to report a comprehensive nutritional characterization of Italian glassworts, specifically from the Apulian region. A comparison was made with products from other Mediterranean areas.

4.1. Mineral Nutrients and Antinutrients

The mineral composition of halophytes is governed not only by environmental conditions but also by genotype [27]. In our study, Na and Cl were the dominant elements in the mineral profiles of all samples. Sodium concentrations ranged from 40 to 120 mg g⁻¹ DW, while chloride concentrations ranged from 10 to 150 mg g⁻¹ DW. Notable genotypic differences in sodium accumulation were observed. Among the perennial species, *Arthrocaulon macrostachyum* generally accumulated less sodium than *Sarcocornia fruticosa*. A clear exception was the *A. macrostachyum* sample from Monti d'Arena-Bosco Caggione (MA-BC), a site located immediately adjacent to the sea. In contrast, the annual species *Salicornia europaea* exhibited a pronounced tendency to accumulate sodium. This was highlighted by a significant disparity between samples from Laguna del Re Oasis (LRO), where *S. europaea* (*S. europaea*-LRO) had the highest sodium concentration and *A. macrostachyum* (*A. macrostachyum*-LRO) the lowest, despite similar soil characteristics and water salinity. *A. macrostachyum* showed low Na level even when grown close to a salt pan (*A. macrostachyum* from Margherita di Savoia Saltworks -MSS).

Halophytes employ various physiological mechanisms to manage salinity stress, including sodium exclusion at the root level, excretion via specialized structures (e.g., bladder cells), and inclusion-diluting absorbed sodium within succulent tissues [41]. Our data confirm that *S. europaea* is a sodium-including species, likely using Na for osmoregulation [42,43]. This obligate halophyte's growth is stimulated by sodium availability [44], a trait further evidenced by the sample from the conventional Spirito Contadino (SC) farm, which was richer in Na than other samples from brackish areas. Furthermore, it demonstrates the potential role of this species in the remediation of Na-affected soils. Some studies have reported the introduction of halophytes, including *S. europaea* [30], in intercropping systems for salt soil remediation, demonstrating beneficial effects for companion crops due to a reduction of soil salinity [45].

The generally low sodium accumulation in *A. macrostachyum* was surprising, as this species is often suggested for phytoremediation due to its high biomass and NaCl uptake [14,46]. This discrepancy may be explained by the partitioning of sodium within the plant [47]. It should be considered that in the present work, only fresh stems representing the edible portion were analysed. Ramírez et al. [14] found that in the *Arthrocaulon* genus, sodium accumulates primarily in biologically less active tissues like the xylem, vascular bundles of lignified stems, and epidermis. Therefore, the sodium in our sampled *A. macrostachyum* was likely stored in roots and lignified parts, which were not analyzed.

Reported sodium concentrations in glassworts are highly variable, attributable to differing environments. Our values are consistent with studies on the same species from Spain and Portugal [1,11,48]. However, our measured sodium content in *A. macrostachyum* was lower than the 80–310 mg g⁻¹ DW range reported by Ramírez et al. [14] across Mediterranean countries (sampling in both coastal and inland areas).

Sodium is an element of major concern for human health, due to its implications in vascular dysfunction. Indeed, the World Health Organization (WHO) and the European Food Safety

Authority (EFSA) recommend not exceeding the sodium acceptable daily intake dose of 2.0 g [49,50]. Furthermore, to mitigate vascular dysfunction and the risk of high blood pressure, it is recommended to consume foods with a sodium-to-potassium (Na/K) ratio of less than 1 [51], as potassium is essential for the human body, reducing blood pressure and stroke incidence [52]. The samples of our survey showed a Na/K ratio ranging from 1.6 for *S. europaea*-SC (cultivated in non-saline soil) to 5 for *A. macrostachyum*-MA-BC (grown close to the sea). However, despite their high sodium content, several studies have demonstrated that dried glassworts, when consumed as a salt substitute or 'green salt,' can exhibit antihypertensive effects. This is attributed to their rich composition of bioactive compounds, including trans-ferulic acid, which has been found to reduce hypertension [12,16,53].

In the perennial species, chlorine levels were generally higher in *A. macrostachyum* than in *S. fruticosa*, suggesting differences in chlorine regulation mechanisms [54]. Moreover, the lower chlorine levels observed in *S. fruticosa*-MSS and *A. macrostachyum*-LRO, compared to other samples of the same species, may be due to specific conditions at the sampling sites. The occasional supply of fresh water could decrease the chlorine concentration in the soil, since both collecting sites (MSSa and LROa) were located where salt and fresh water converge, creating a unique ecological environment. As expected, low Cl content characterized the conventionally cultivated *S. europaea* (*S. europaea*-SC) (Table 2). The chlorine concentrations of our samples turn out to be slightly lower compared to literature findings [55,56].

Concentrations of nitrate, potassium, calcium and magnesium ranged from 0.4–2.1, 21–48, 13–35, and 6–15 mg g⁻¹ DW, respectively. As expected, cultivated *S. europaea* accumulated more nutrients due to the application of fertilizers. Specifically, a greater amount of NO₃ was detected in the sample harvested from the SC farm, where an intensive agricultural system is practiced. The elevated NO₃ in wild *A. macrostachyum*-MA-BC was likely due to runoff from upstream cultivated land.

Within perennial species, the mineral content results in line with what is found in the literature for the same genera [11,28,57]. Instead, the mineral content of cultivated *S. europaea* is significantly higher than reported by other authors in cultivated plants of the same species [25,58], likely for differences in soil and crop management.

The nitrate accumulation in green leafy vegetables poses a potential food safety hazard, as the nitrates can be reduced into nitrites and nitrosamines (carcinogenic compounds [59]). All nitrate values for the glasswort samples were well below European Community (EU 2023/915) safety thresholds for leafy vegetables.

Oxalate, an antinutrient that can chelate minerals (e.g. calcium) and can lead to kidney stones [60]. To date, the mechanism regulating the synthesis of oxalates is still complex, as they act as plant mineral regulators, and the Chenopodioidae uses the complexation of free oxalic acid to maintain ionic equilibrium [26]. Some authors reported that the content of oxalates in halophytes increases as substrate salinity rises [61]. On the contrary, in *S. ramosissima* under saline irrigation conditions, it was observed that a rise in temperature and CO₂ lowered oxalate levels [26]. Among our samples, *S. fruticosa* from Varano lake (*S. fruticosa*-VL) and *A. macrostachyum*-MSS had the highest value, likely due to site-specific environmental conditions. Accordingly, on wild plants of *Salicornia patula*, oxalates increased from 1.5 to 16.9 g kg⁻¹ FW when moving from coastal to inland areas [18]. Except for these two, our samples showed lower oxalates than those observed in *S. ramosissima* (3.5 and 4 g kg⁻¹ FW) in the unstressed plants [26]. Comparing our glassworts with vegetables well known for high oxalate contents, it emerged that only *S. fruticosa*-VL is richer in oxalate than spinach (6.0 g kg⁻¹ FW), and in *A. macrostachyum*-MSS it is similar to Swiss chard (3.0 g kg⁻¹ FW).

Iodine is an important micronutrient involved in the synthesis of thyroid hormones, which regulate various physiological processes [62]. There is some evidence of the potential for halophytes to be an interesting source of this oligoelement [17,63]. Iodine content varied significantly with sampling location, with the highest levels found in LRO, VL, and MA-BC sites. Changes in iodine concentration could be ascribed to site-specific variability in soil and/or sea spray contribution in

accumulating this ion. The low iodine content of the product from the Spirito Contadino farm is attributable to a non-saline cultivation system.

Studies analyzing the iodine content in glassworts are limited, but they reported lower concentrations than those found in our samples. Oliveira-Alves et al. [48] in *S. ramosissima* found iodine values below $0.13 \mu\text{g g}^{-1}$ FW, and Duarte et al. [64] found values around $15 \mu\text{g g}^{-1}$ DW on *S. fruticosa*.

The recommended daily intake (RDI) of iodine for an adult is $150 \mu\text{g}$, varying between 90 and $250 \mu\text{g}$ according to age, sex, and other aspects of life (such as pregnancy) [65]. The iodine RDI could be satisfied by consuming portions of glasswort close to 40-150 g, depending on the product's provenance. The consumption of the iodine-rich glasswort (*S. europaea*-LRO, $349 \mu\text{g}$ per 100 g FW) does not pose a risk of toxic effects, as an intake of up to $1500 \mu\text{g}$ per day is accepted as safe [66].

4.2. Visual Quality, Nutritional and Antioxidative Traits

Consumer acceptance of glassworts is influenced by both visual appeals and nutritional quality. From a nutritional standpoint, glasswort is particularly appealing since it is characterized by a rich profile of antioxidant compounds, including pigments, phenolics, and sterols, which contribute to its overall health benefits.

Chlorophylls are primary determinants of the green colour in leafy vegetables, a key visual quality indicator for consumers [11]. Beyond aesthetics, they offer significant health benefits, such as anti-mutagenic, anti-cancer, and anti-inflammatory properties [4]. Early-life dietary intake may also help prevent obesity and reduce the risk of type 1 diabetes [67].

In our study, the annual species *S. europaea* and the perennial *A. macrostachyum* from the Margherita di Savoia Saltworks (MSS) site exhibited total chlorophyll concentrations an order of magnitude higher than other samples (Table 3), making them particularly visually appealing. Conversely, *S. fruticosa*, *A. macrostachyum*-LRO, and *A. macrostachyum*-MA-BC had lower chlorophyll levels, which could suggest a protective response to environmental stresses. These conditions are reported to lead to excessive plant energy exposure, exceeding its capacity to use light effectively in photosynthesis [68]. In these samples, chlorophyll degradation and a reduced Chl a /Chl b ratio (averaging 1.3 vs. 2.8) likely served as a protective mechanism [69], a phenomenon also reported in other halophytes, including *Sarcocornia* [28]. This hypothesis is supported by their consistently higher anthocyanin concentrations (Table 3), known photosystem protectants. Our chlorophyll values are generally consistent with some literature [19], though lower than values reported for *S. ramosissima* from Portugal [11].

Carotenoids are pigments with potent antioxidant and immunological properties [11]. Lutein and β -carotene, for example, support vision, protect against macular degeneration, maintain skin health, and combat free radicals [70]. The European Food Safety Authority [71] guidelines consider carotenoids (like β -carotene) only as a supplement to meet vitamin A requirements.

In plants, they absorb excess light energy, prevent ROS formation, and protect the photosynthetic apparatus from salt stress in a species-dependent manner [68,72]. A rise in carotenoids was observed in *A. macrostachyum* and *S. fruticosa* with the increase of NaCl in the nutrient solution, while *S. europaea* showed the opposite trend [72]. Notably, the *S. europaea* sample from the non-saline Spirito Contadino (SC) farm was richest in carotenoids (Table 3), aligning with findings that this species may reduce carotenoid accumulation under salinity [72]. Our carotenoid values are similar to those reported for Mediterranean glassworts from different areas of Portugal, Spain, and Tunisia [19,28,47] but lower than those found in plants from Portugal's Algarve region [11].

Phenolic compounds, particularly flavonoids and anthocyanins, are strong radical scavengers [59] that confer anti-inflammatory, antiviral, and antibacterial benefits [73,74]. Adequate anthocyanin intake is linked to improved neurological and cardiovascular health [75].

In plants, phenol synthesis is a primary defense mechanism against oxidative stress [76]. Anthocyanins, specifically, filter light to prevent ROS production [63].

In halophytes, phenolic compounds are known to help alleviate the harmful effects of harsh environmental conditions. As soil salinity or air temperature rised, their biosynthesis increases was reported , though the response was genotype specific [64]. In glassworts, when salt stress was applied under a greenhouse-soilless system, moderate salt concentration at the root level led to an improvement of TP in *Salicornia* and *Sarcocornia* species [25]. Moreover, *A. macrostachyum* showed significant increases in phenols under moderate/high salinity, while *S. europaea* levels often remained constant [72]. When comparing wild and soilless cultivated *S. fruticosa*, it was suggested that phenol content was positively affected by stresses encountered in wild conditions, beyond just salinity [28], such as drought and unbalanced nutrient availability . Therefore, strong interaction between genotypes and various abiotic/biotic stress is expected regarding phenol content.

We observed clear differences in total phenol (TP) content. Specifically, *A. macrostachyum* collected near saltworks (MSS) and *S. europaea* cultivated under both saline (LRO) and non-saline (SC) conditions showed significantly more TP compared to other samples. Similarly, a slight improvement in flavonoid content was found in these same plants (Table 3). The likely explanation for this overproduction of TP, as an antioxidative response, is the reduced ability of these plants to prevent oxidative damage by decreasing chlorophylls, a phenomenon observed in other samples. It can be argued that *S. europaea* appears predisposed to synthesizing phenols as a mechanism to mitigate environmental stress without involving photosynthetic pigments. It's noteworthy that the increase in TP was also observed in plants grown on non-saline commercial farms, suggesting growth-stressing conditions even there. In *S. europaea* samples, a further possible explanation for the higher TP levels could be related to the harvest period. Both cultivated glassworts were collected during the hottest months (August-September), while the wild samples were collected during the spring period. This seasonal variation may have influenced the plants' secondary metabolite production, including phenolics. An improvement of Folin-Ciocalteu TPs was observed in *S. ramosissima* harvested in July compared to May in Portugal [19]. The *A. macrostachyum* samples near saltworks (*A. macrostachyum*-MSS) represent a strong example of genotype-environment interaction. This is evident when considering its lower TP and chlorophyll contents compared to the same species collected from other areas (*A. macrostachyum*-MA-BC; *A. macrostachyum*-LRO) (Table 3), suggesting a varied response to the environmental pressure across the sites.

As expected, TP content widely varies among Mediterranean glassworts. Our TP values for *S. europaea* and *S. fruticosa* align with those from cultivated plants [1,25] but are generally lower than those from wild plants in Portugal, Turkey, and Spain [11,19,28,77]. *A. macrostachyum* values were consistent with some studies [21,78] but lower than others [11,77,79]. Flavonoid content was slightly higher across our samples compared to literature values for the *Sarcocornia*, *Salicornia*, and *Arthrocaulon* genera [11,28,77,78]. Anthocyanin levels in *S. europaea* aligned with previous reports, while those in *S. fruticosa* were slightly higher [80]; no studies on anthocyanin content in *A. macrostachyum* were found.

Phytosterols possess anti-tumorigenic properties and contribute to heart health by lowering blood cholesterol levels [81,82]. In plants, they are essential for embryogenesis and for maintaining membrane fluidity and permeability, especially under stress [83]. Their biosynthesis is influenced by soil and climate; for instance, nitrogen availability, higher temperatures, and lower rainfall can increase sterol content [83–85]. Research on sterol changes in halophytes is limited, despite evidence that they are crucial for protecting membrane lipids against salt stress, even at the cost of increased permeability [86,87].

As reported for the other compounds mentioned so far, sterol content was significantly influenced by plant location, with the *A. macrostachyum*-MSS sample significantly exceeding other glassworts.

This finding was likely due to the arid conditions of its specific sampling site, which was marked by a lack of water during certain times of the year. In contrast, the *S. fruticosa* from the same area (MSS) was collected from a flooded spot, resulting in lower values, demonstrating how microenvironments may dictate physiological responses.

Most of our samples had sterol values similar to *S. ramosissima* [88], while *A. macrostachyum*-MSS reached levels comparable to *Salicornia perennis* under moderate stress [86]. Compared to commonly consumed vegetables, our glassworts are similar to Chinese cabbage and carrot [81], with *A. macrostachyum*-MSS overcoming the sterol content of broccoli, Brussels sprouts, and spinach, and matching that of peas and black olives [81,89].

The antioxidant activity (AC) of complex systems (such as plant extracts) cannot be evaluated only with one specific assay, because it involves distinct radicals or antioxidant sources [90]. Therefore, the antioxidant activity was assessed by three complementary and widely used assays, including antioxidant reaction with a Fe (III) complex (FRAP), reaction with organic radical cation (ABTS), and reaction with organic radical (DPPH). Antioxidant activity increases in conjunction with stress conditions and is typically related to the production of secondary metabolism compounds.

The results of the three assays showed a similar trend, as also noted by Rumpf et al. [90] on different vegetable aqueous extracts. The hydrophilic antioxidant capacity (HA: from phenolics, flavonoids, anthocyanins, vitamin C) was the dominant contributor, vastly exceeding the lipophilic capacity (LA: from carotenoids, chlorophylls, sterols).

The two annual glasswort samples had improved antioxidant activity compared with the perennial species. An exception was found in the *A. macrostachyum*-MSS sample, which exhibited antioxidant levels that were comparable to those of *S. europaea*. It is plausible to hypothesize that the observed increase in antioxidant activity was driven by higher phenolics; however, it is important to note that we did not investigate other hydrophilic compounds, such as ascorbic acid, which might also have played a significant role in influencing hydrophilic antioxidant capacity. Regarding the *Arthrocaulon* of MSS, the higher AC confirmed a stronger site-specific stress pressure, leading the plant to exhibit an elevated antioxidant response, even when compared to other samples of the same species.

In *Salicornia ramosissima* and *Sarcocornia perennis*, Hupel et al. [91] and Antunes et al. [19] reported that in response to oxidative stress, the amount of carotenoids and phenols increased, which resulted in a similar magnitude of changes in antioxidant capacity.

4.3. Principal Component Analysis

Principal component analysis (PCA) was employed to visualize the key biochemical traits differentiating glasswort species and their growing environments across Apulia region (Figures 2A and B). The first three principal components (PCs), with eigenvalues >1, collectively explained 88% of the total variance in the dataset. The biplot of the first two components (PC1 and PC2), which accounted for 75.5% of the variance, revealed distinct groupings (Figure 2A). PC1 was positively correlated with essential minerals (K, Mg, Ca, Cl), photosynthetic pigments (chlorophylls, carotenoids), and phenolic compounds. PC2 was positively associated with anthocyanins, sodium, and nitrate, and negatively correlated with chlorine, sterols, and flavonoids. A third component, PC3, was primarily defined by a strong positive correlation with iodine and a weaker one with Na, and a negative correlation with sterols and nitrate (Figure 2B).

The first two PCA components (Figure 2A) clearly differentiated the two annual *S. europaea* samples and the perennial *A. macrostachyum*-MSS from the other perennial species. The *S. europaea* samples, clustering in the upper right quadrant, were characterized by their high mineral and chlorophyll content. This aligns with ANOVA results and suggests a strong influence of nutrient availability, likely from fertilization practices at both the non-saline (SC) and saline (LRO) cultivation sites. Notably, *S. europaea*-LRO's accumulation of essential cations (K, Ca, Mg) despite high soil NaCl competition highlights its efficient ion selectivity.

In contrast, *A. macrostachyum*-MSS, located in the lower right quadrant, was distinguished by its high chlorine content and bioactive profile, rich in sterols and flavonoids but poorer in anthocyanins. This clustering suggests that an excessive soil concentration and plant uptake of chlorine could have provoked a response to the oxidative stress in this sample involving sterols and flavonoids biosynthesis [92].




Figure 2. Principal Component Analysis bi-plot (A – PC1 vs. PC2; B- PC1 vs. PC3) reporting the minerals and biochemicals. Specifically: Nitrate (NO₃), Sodium (Na), Potassium (K), Chlorine (Cl), Calcium (Ca), Magnesium (Mg), Iodine, chlorophyll a, and b (CHLa, b), carotenoids (Car), Anthocyanins (Anth), Total Phenols (TP), Flavonoids (FL), and Sterols. Please see Tables 1 for sample identification.

The remaining perennial *Sarcocornia* and *Arthrocaulon* samples clustered on the left side of PC1, showing a strong positive association with anthocyanins. This contrast with the *S. europaea* and *A. macrostachyum*-MSS clusters confirms that different halophyte species employ distinct biochemical

strategies to cope with environmental stress and that the specific environmental condition effect may also mediate genotype response.

Furthermore, the importance of sodium for glasswort physiology was underscored by its significant loading on both PC2 and PC3. The latter was particularly useful in separating samples based on their iodine content, with *S. europaea*-LRO, *A. macrostachyum*-LRO, and *S. fruticosa*-VL forming a group characterized by higher accumulation. Conversely, the same perennial species collected at the MSS site, along with *S. europaea*-SC, were the poorest in iodine. This pattern can be attributed to the site-specific bioavailability of this ion. In the *S. europaea*-SC sample, it was expected because of its cultivation on a conventional farm. For the wild glassworts, it can be explained by *A. macrostachyum* was found in a location shielded from frequent marine flooding, and *S. fruticosa* was located at the mouth of brackish water. Points may be potentially less rich in iodine, which could have led to less accumulation of this element in plants.

5. Conclusions

This study provides a comprehensive biochemical and nutritional profile of both wild and cultivated glassworts from the Apulia region, offering critical insights for their development as a novel food source. Our results confirm that genotypes (*Salicornia europaea*, *Arthrocaulon macrostachyum*, *Sarcocornia fruticosa*) are the primary driver of mineral and bioactive compound composition, but its expression may be deeply modulated by environmental conditions, leading to significant site-specific variation.

In support of the initial aim, we found that these halophytes possess great nutritional potential. The annual species *S. europaea* consistently demonstrated promising nutraceutical traits, accumulating high levels of essential minerals (K, Ca, Mg), carotenoids, and phenolic compounds, even under conventional non-saline conditions. This confirms its suitability as a nutrient-dense, sustainable crop. Furthermore, all species served as an excellent source of iodine, with a single portion capable of meeting the recommended daily intake without risk of toxicity. However, the elevated levels of Na and oxalates in some samples may raise concerns regarding their dietary suitability, highlighting the importance of assessing product provenance. Implementing appropriate agricultural practices could help manage anti-nutritional compounds, thereby reducing the sodium-to-potassium (Na/K) ratio and limiting oxalate accumulation.

This research reinforces the existing literature on the value of these halophytes but moves the discussion forward by highlighting the critical importance of provenance and genotype selection. Future efforts must focus on elucidating the specific mechanisms behind the observed genotype-environment interactions, particularly the physiological strategies governing ion uptake and regulation, and antioxidant production. This will be essential for standardizing product quality, ensuring food safety, and ultimately leveraging the full potential of glassworts as a sustainable and nutritionally enriched food source for a resilient agricultural system.

Author Contributions: Conceptualization, G.C.; methodology, G.C.; formal analysis, L.G.D., A.B. and G.C.; investigation, L.B., C.L., E.V.P., A.G., A.C. and A.B.; data curation, L.B., C.L. A.B.; resources: L.B., C.L., E.V.P., A.G., A.C.; writing—original draft preparation, L.G.D. and G.C.; writing—review and editing, L.G.D., A.E. and G.C.; visualization, A.E. A.C.; project administration, G.C.; funding acquisition, G.C.. All authors have read and agreed to the published version of the manuscript

Funding: This research was supported by (1) the Agritech National Research Center and received funding from the European Union Next-GenerationEU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR)—MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.4—D.D. 1032 17/06/2022, CN00000022) (This manuscript reflects only the authors' views and opinions, neither the European Union nor the European Commission can be considered responsible for them); (2) Regione Apulia Project 'Biodiversity of Apulian vegetable species (BiodiverSO Veg)', Measure 10, Sub measure 10.2, Operation 1 "Program for the conservation and the valorization of the genetic resources in agri-culture".

Acknowledgments: The authors would like to acknowledge Ruggiero Piazzola for his assistance during field activities.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Agudelo, A.; Carvajal, M.; Martinez-Ballesta, M. del C. Halophytes of the Mediterranean Basin—Underutilized Species with the Potential to Be Nutritious Crops in the Scenario of the Climate Change. *Foods* **2021**, *10*, 119, doi:10.3390/foods10010119.
2. FAO *The State of Food and Agriculture 2020. Overcoming Water Challenges in Agriculture.*; FAO: Rome, 2020; ISBN 978-92-5-133441-6.
3. Navarro-Torre, S.; Garcia-Caparrós, P.; Nogales, A.; Abreu, M.M.; Santos, E.; Cortinhas, A.L.; Caperta, A.D. Sustainable Agricultural Management of Saline Soils in Arid and Semi-Arid Mediterranean Regions through Halophytes, Microbial and Soil-Based Technologies. *Environ. Exp. Bot.* **2023**, *212*, doi:10.1016/j.envexpbot.2023.105397.
4. Hulkko, L.S.S.; Chaturvedi, T.; Thomsen, M.H. Extraction and Quantification of Chlorophylls, Carotenoids, Phenolic Compounds, and Vitamins from Halophyte Biomasses. *Appl. Sci.* **2022**, *12*, 840, doi:10.3390/app12020840.
5. Lopes, A.; Rodrigues, M.J.; Pereira, C.; Oliveira, M.; Barreira, L.; Varela, J.; Trampetti, F.; Custódio, L. Natural Products from Extreme Marine Environments: Searching for Potential Industrial Uses within Extremophile Plants. *Ind. Crops Prod.* **2016**, *94*, 299–307, doi:10.1016/j.indcrop.2016.08.040.
6. Petropoulos, S.A.; Karkanis, A.; Martins, N.; Ferreira, I.C.F.R. Edible Halophytes of the Mediterranean Basin: Potential Candidates for Novel Food Products. *Trends Food Sci. Technol.* **2018**, *74*, 69–84, doi:10.1016/j.tifs.2018.02.006.
7. The Angiosperm Phylogeny Group.; Chase, M.W.; Christenhusz, M.J.M.; Fay, M.F.; Byng, J.W.; Judd, W.S.; Soltis, D.E.; Mabberley, D.J.; Sennikov, A.N.; Soltis, P.S.; et al. An Update of the Angiosperm Phylogeny Group Classification for the Orders and Families of Flowering Plants: APG IV. *Bot. J. Linn. Soc.* **2016**, *181*, 1–20, doi:10.1111/boj.12385.
8. Bartolucci, F.; Peruzzi, L.; Galasso, G.; Alessandrini, A.; Ardenghi, N.M.G.; Bacchetta, G.; Banfi, E.; Barberis, G.; Bernardo, L.; Bouvet, D.; et al. A Second Update to the Checklist of the Vascular Flora Native to Italy. *Plant Biosyst. - An Int. J. Deal. with all Asp. Plant Biol.* **2024**, *158*, 219–296, doi:10.1080/11263504.2024.2320126.
9. Pignatti, S.; Guarino, R.; La Rosa, M. *Flora d'Italia*; Edagricole, Ed.; 2017; ISBN 978-88-506-5242-6.
10. POWO Plants of the World Online Available online: <http://www.plantsoftheworldonline.org/>.
11. Barreira, L.; Resek, E.; Rodrigues, M.J.; Rocha, M.I.; Pereira, H.; Bandarra, N.; da Silva, M.M.; Varela, J.; Custódio, L. Halophytes: Gourmet Food with Nutritional Health Benefits? *J. Food Compos. Anal.* **2017**, *59*, 35–42, doi:10.1016/j.jfca.2017.02.003.
12. Cárdenas-Pérez, S.; Piernik, A.; Chanona-Pérez, J.J.; Grigore, M.N.; Perea-Flores, M.J. An Overview of the Emerging Trends of the *Salicornia* L. Genus as a Sustainable Crop. *Environ. Exp. Bot.* **2021**, *191*, 104606, doi:10.1016/j.envexpbot.2021.104606.
13. Custódio, L.; Rodrigues, M.J.; Pereira, C.G.; Castañeda-Loaiza, V.; Fernandes, E.; Standing, D.; Neori, A.; Shpigel, M.; Sagi, M. A Review on *Sarcocornia* Species: Ethnopharmacology, Nutritional Properties, Phytochemistry, Biological Activities and Propagation. *Foods* **2021**, *10*, 2778, doi:10.3390/foods10112778.
14. Ramírez, E.; Rodríguez, N.; de la Fuente, V. *Arthrocnemum* Moq.: Unlocking Opportunities for Biosaline Agriculture and Improved Human Nutrition. *Plants* **2024**, *13*, 496, doi:10.3390/plants13040496.
15. Loconsole, D.; Cristiano, G.; De Lucia, B. Glassworts: From Wild Salt Marsh Species to Sustainable Edible Crops. *Agriculture* **2019**, *9*, 14, doi:10.3390/agriculture9010014.
16. Oliveira-Alves, S.C.; Andrade, F.; Sousa, J.; Bento-Silva, A.; Duarte, B.; Caçador, I.; Salazar, M.; Mecha, E.; Serra, A.T.; Bronze, M.R. Soilless Cultivated Halophyte Plants: Volatile, Nutritional, Phytochemical, and Biological Differences. *Antioxidants* **2023**, *12*, 1161, doi:10.3390/antiox12061161.
17. Ekanayake, S.; Egodawatta, C.; Attanayake, R.N.; Perera, D. *From Salt Pan to Saucepan: Salicornia, a Halophytic Vegetable with an Array of Potential Health Benefits*; 2023; Vol. 4; ISBN 2518695931.

18. Sánchez Gavilán, I.; Velázquez Ybarzabal, D.; de la Fuente, V.; Cámara, R.M.; Sánchez-Mata, M.C.; Cámara, M. Valorization of *Salicornia Patula Duval-Jouve* Young Shoots in Healthy and Sustainable Diets. *Nutrients* **2024**, *16*, 358, doi:10.3390/nu16030358.
19. Antunes, M.D.; Gago, C.; Guerreiro, A.; Sousa, A.R.; Julião, M.; Miguel, M.G.; Faleiro, M.L.; Panagopoulos, T. Nutritional Characterization and Storage Ability of *Salicornia Ramosissima* and *Sarcocornia Perennis* for Fresh Vegetable Salads. *Horticulturae* **2021**, *7*, 1–12, doi:10.3390/horticulturae7010006.
20. Lopes, M.; Sanches-Silva, A.; Castilho, M.; Cavaleiro, C.; Ramos, F. Halophytes as Source of Bioactive Phenolic Compounds and Their Potential Applications. *Crit. Rev. Food Sci. Nutr.* **2023**, *63*, 1078–1101, doi:10.1080/10408398.2021.1959295.
21. Sánchez-Gavilán, I.; Chueca, E.R.; García, V. de la F. Bioactive Compounds in *Sarcocornia* and *Arthrocnemum*, Two Wild Halophilic Genera from the Iberian Peninsula. *Plants* **2021**, *10*, doi:10.3390/plants10102218.
22. Ventura, Y.; Wuddineh, W.A.; Myrzabayeva, M.; Alikulov, Z.; Khozin-Goldberg, I.; Shpigel, M.; Samocha, T.M.; Sagi, M. Effect of Seawater Concentration on the Productivity and Nutritional Value of Annual *Salicornia* and Perennial *Sarcocornia* Halophytes as Leafy Vegetable Crops. *Sci. Hortic. (Amsterdam)*. **2011**, *128*, 189–196, doi:10.1016/j.scienta.2011.02.001.
23. Sánchez-Gavilán, I.; Rufo, L.; Rodríguez, N.; de la Fuente, V. On the Elemental Composition of the Mediterranean Euhalophyte *Salicornia Patula Duval-Jouve* (Chenopodiaceae) from Saline Habitats in Spain (Huelva, Toledo and Zamora). *Environ. Sci. Pollut. Res.* **2021**, *28*, 2719–2727, doi:10.1007/s11356-020-10663-w.
24. Ricardo, F.; Veríssimo, A.C.; Maciel, E.; Domingues, M.R.; Calado, R. Fatty Acid Profiling as a Tool for Fostering the Traceability of the Halophyte Plant *Salicornia Ramosissima* and Contributing to Its Nutritional Valorization. *Plants* **2024**, *13*, doi:10.3390/plants13040545.
25. Lima, A.R.; Castañeda-Loaiza, V.; Salazar, M.; Nunes, C.; Quintas, C.; Gama, F.; Pestana, M.; Correia, P.J.; Santos, T.; Varela, J.; et al. Influence of Cultivation Salinity in the Nutritional Composition, Antioxidant Capacity and Microbial Quality of *Salicornia Ramosissima* Commercially Produced in Soilless Systems. *Food Chem.* **2020**, *333*, 127525, doi:10.1016/j.foodchem.2020.127525.
26. Mesa-Marín, J.; Mateos-Naranjo, E.; Carreiras, J.; Feijão, E.; Duarte, B.; Matos, A.R.; Betti, M.; Del Rio, C.; Romero-Bernal, M.; Montaner, J.; et al. Interactive Temperature and CO₂ Rise, Salinity, Drought, and Bacterial Inoculation Alter the Content of Fatty Acids, Total Phenols, and Oxalates in the Edible Halophyte *Salicornia Ramosissima*. *Plants* **2023**, *12*, 1395, doi:10.3390/plants12061395.
27. Bertin, R.L.; Gonzaga, L.V.; Borges, G. da S.C.; Azevedo, M.S.; Maltez, H.F.; Heller, M.; Micke, G.A.; Tavares, L.B.B.; Fett, R. Nutrient Composition and, Identification/Quantification of Major Phenolic Compounds in *Sarcocornia Ambigua* (Amaranthaceae) Using HPLC–ESI-MS/MS. *Food Res. Int.* **2014**, *55*, 404–411, doi:10.1016/j.foodres.2013.11.036.
28. Castañeda-Loaiza, V.; Oliveira, M.; Santos, T.; Schüler, L.; Lima, A.R.; Gama, F.; Salazar, M.; Neng, N.R.; Nogueira, J.M.F.; Varela, J.; et al. Wild vs Cultivated Halophytes: Nutritional and Functional Differences. *Food Chem.* **2020**, *333*, 127536, doi:10.1016/j.foodchem.2020.127536.
29. Conversa, G.; Lazzizzera, C.; Bonasia, A.; Cifarelli, S.; Losavio, F.; Sonnante, G.; Elia, A. Exploring On-Farm Agro-Biodiversity: A Study Case of Vegetable Landraces from Puglia Region (Italy). *Biodivers. Conserv.* **2020**, *29*, 747–770, doi:10.1007/s10531-019-01908-3.
30. Cammerino, A.R.B.; Ingaramo, M.; Rizzi, V.; Gioiosa, M.; Monteleone, M. Glasswort as a Strategic Crop in Coastal Wetlands: Intercropping Results with Swiss Chard. *Agronomy* **2025**, *15*, 158, doi:10.3390/agronomy15010158.
31. Somma, A.; Palmitessa, O.D.; Conversa, G.; Serio, F.; Santamaria, P. Localized Foliar Application of Iodine on Tomato: An Effective Approach for Targeted Agronomic Biofortification. *Sci. Hortic. (Amsterdam)*. **2024**, *327*, 112807, doi:10.1016/j.scienta.2023.112807.
32. Sumanta, N.; Haque, C.I.; Nishika, J.; Suprakash, R. Spectrophotometric Analysis of Chlorophylls and Carotenoids from Commonly Grown Fern Species by Using Various Extracting Solvents. *Res. J. Chem. Sci.* **2014**, *4*, 63–69.

33. Sims, D.A.; Gamon, J.A. Relationships between Leaf Pigment Content and Spectral Reflectance across a Wide Range of Species, Leaf Structures and Developmental Stages. *Remote Sens. Environ.* **2002**, *81*, 337–354, doi:10.1016/S0034-4257(02)00010-X.
34. Murray, J.R.; Hackett, W.P. Dihydroflavonol Reductase Activity in Relation to Differential Anthocyanin Accumulation in Juvenile and Mature Phase Hedera Helix L. *Plant Physiol.* **1991**, *97*, 343–351, doi:10.1104/pp.97.1.343.
35. Singleton, V.L.; Rossi, J.A. Colorimetry of Total Phenolics with Phosphomolybdic-Phosphotungstic Acid Reagents. *Am. J. Enol. Vitic.* **1965**, *16*, 144–158, doi:10.5344/ajev.1965.16.3.144.
36. Chandra, S.; Khan, S.; Avula, B.; Lata, H.; Yang, M.H.; ElSohly, M.A.; Khan, I.A. Assessment of Total Phenolic and Flavonoid Content, Antioxidant Properties, and Yield of Aeroponically and Conventionally Grown Leafy Vegetables and Fruit Crops: A Comparative Study. *Evidence-Based Complement. Altern. Med.* **2014**, *2014*, doi:10.1155/2014/253875.
37. Xiang, L.; Liu, Y.; Li, H.; Lin, J. Simultaneous Extraction and Determination of Free and Conjugated Phytosterols in Tobacco. *J. Sep. Sci.* **2016**, *39*, 2466–2473, doi:10.1002/jssc.201600247.
38. Feng, S.; Wang, L.; Belwal, T.; Li, L.; Luo, Z. Phytosterols Extraction from Hickory (*Carya Cathayensis* Sarg.) Husk with a Green Direct Citric Acid Hydrolysis Extraction Method. *Food Chem.* **2020**, *315*, 126217, doi:10.1016/j.foodchem.2020.126217.
39. Re, R.; Pellegrini, N.; Proteggente, A.; Pannala, A.; Yang, M.; Rice-Evans, C. Antioxidant Activity Applying an Improved ABTS Radical Cation Decolorization Assay. *Free Radic. Biol. Med.* **1999**, *26*, 1231–1237, doi:10.1016/S0891-5849(98)00315-3.
40. Lisanti, A.; Formica, V.; Ianni, F.; Albertini, B.; Marinozzi, M.; Sardella, R.; Natalini, B. Antioxidant Activity of Phenolic Extracts from Different Cultivars of Italian Onion (*Allium Cepa*) and Relative Human Immune Cell Proliferative Induction. *Pharm. Biol.* **2016**, *54*, 799–806, doi:10.3109/13880209.2015.1080733.
41. Mann, A.; Lata, C.; Kumar, N.; Kumar, A.; Kumar, A.; Sheoran, P. Halophytes as New Model Plant Species for Salt Tolerance Strategies. *Front. Plant Sci.* **2023**, *14*, doi:10.3389/fpls.2023.1137211.
42. Mendis, C.L.; Padmathilake, R.E.; Attanayake, R.N.; Perera, D. Learning from *Salicornia*: Physiological, Biochemical, and Molecular Mechanisms of Salinity Tolerance. *Int. J. Mol. Sci.* **2025**, *26*, 5936, doi:10.3390/ijms26135936.
43. Homayouni, H.; Razi, H.; Izadi, M.; Alemzadeh, A.; Kazemeini, S.A.; Niazi, A.; Vicente, O. Temporal Changes in Biochemical Responses to Salt Stress in Three *Salicornia* Species. *Plants* **2024**, *13*, doi:10.3390/plants13070979.
44. Puccinelli, M.; Marchioni, I.; Botrini, L.; Carmassi, G.; Pardossi, A.; Pistelli, L. Growing *Salicornia Europaea* L. with Saline Hydroponic or Aquaculture Wastewater. *Horticulturae* **2024**, *10*, 196, doi:10.3390/horticulturae10020196.
45. Castagna, A.; Mariottini, G.; Gabriele, M.; Longo, V.; Souid, A.; Dauvergne, X.; Magné, C.; Foggi, G.; Conte, G.; Santin, M.; et al. Nutritional Composition and Bioactivity of *Salicornia Europaea* L. Plants Grown in Monoculture or Intercropped with Tomato Plants in Salt-Affected Soils. *Horticulturae* **2022**, *8*, 828, doi:10.3390/horticulturae8090828.
46. Yasseen, B.T.; Al-Thani, R.F. Endophytes and Halophytes to Remediate Industrial Wastewater and Saline Soils: Perspectives from Qatar. *Plants* **2022**, *11*, 1497, doi:10.3390/plants11111497.
47. Rahmani, R.; Arbi, K. El; Aydi, S.S.; Hzami, A.; Tlahig, S.; Najjar, R.; Aydi, S.; Debouba, M. Biochemical Composition and Biological Activities of *Salicornia Europaea* L. from Southern Tunisia. *J. Food Meas. Charact.* **2022**, *16*, 4833–4846, doi:10.1007/s11694-022-01574-0.
48. Oliveira-Alves, S.C.; Andrade, F.; Prazeres, I.; Silva, A.B.; Capelo, J.; Duarte, B.; Caçador, I.; Coelho, J.; Serra, A.T.; Bronze, M.R. Impact of Drying Processes on the Nutritional Composition, Volatile Profile, Phytochemical Content and Bioactivity of *Salicornia Ramosissima* J. Woods. *Antioxidants* **2021**, *10*, 1312, doi:10.3390/antiox10081312.
49. WHO *Guideline: Sodium Intake for Adults and Children*; World Health Organization, Ed.; Geneva, 2012; ISBN 978-92-4-150483-6.

50. Turck, D.; Castenmiller, J.; de Henauw, S.; Hirsch-Ernst, K.; Kearney, J.; Knutsen, H.K.; Maciuk, A.; Mangelsdorf, I.; McArdle, H.J.; Pelaez, C.; et al. Dietary Reference Values for Sodium. *EFSA J.* **2019**, *17*, doi:10.2903/j.efsa.2019.5778.
51. Park, J.; Kwock, C.; Yang, Y. The Effect of the Sodium to Potassium Ratio on Hypertension Prevalence: A Propensity Score Matching Approach. *Nutrients* **2016**, *8*, 482, doi:10.3390/nu8080482.
52. Bonasia, A.; Conversa, G.; Lazzizzera, C.; Gambacorta, G.; Elia, A. Morpho-Biometrical, Nutritional and Phytochemical Characterization of Carrot Landraces from Puglia Region (Southern Italy). *Sustainability* **2021**, *13*, 3940, doi:10.3390/su13073940.
53. Alfheaid, H.A.; Raheem, D.; Ahmed, F.; Alhodieb, F.S.; Alsharari, Z.D.; Alhaji, J.H.; BinMowyna, M.N.; Saraiva, A.; Raposo, A. *Salicornia Bigelovii*, *S. Brachiata* and *S. Herbacea*: Their Nutritional Characteristics and an Evaluation of Their Potential as Salt Substitutes. *Foods* **2022**, *11*, 3402, doi:10.3390/foods11213402.
54. Bazihizina, N.; Colmer, T.D.; Cuin, T.A.; Mancuso, S.; Shabala, S. Friend or Foe? Chloride Patterning in Halophytes. *Trends Plant Sci.* **2019**, *24*, 142–151, doi:10.1016/j.tplants.2018.11.003.
55. Ralph, Y.; Manley, S.L. Spatial and Temporal Variation in Tissue Halide Levels of *Salicornia Virginica*. *Wetlands* **2006**, *26*, 97–106, doi:https://doi.org/10.1672/0277-5212(2006)26[97:SATVIT]2.0.CO;2.
56. Lv, S.; Nie, L.; Fan, P.; Wang, X.; Jiang, D.; Chen, X.; Li, Y. Sodium Plays a More Important Role than Potassium and Chloride in Growth of *Salicornia Europaea*. *Acta Physiol. Plant.* **2012**, *34*, 503–513, doi:10.1007/s11738-011-0847-0.
57. Zahran, M.A.; El-Amier, Y.A. Non-Traditional Fodders from the Halophytic Vegetation of the Deltaic Mediterranean Coastal Desert, Egypt. *J. Biol. Sci.* **2013**, *13*, 226–233, doi:10.3923/jbs.2013.226.233.
58. Lopes, M.; Silva, A.S.; Séndon, R.; Barbosa-Pereira, L.; Cavaleiro, C.; Ramos, F. Towards the Sustainable Exploitation of Salt-Tolerant Plants: Nutritional Characterisation, Phenolics Composition, and Potential Contaminants Analysis of *Salicornia Ramosissima* and *Sarcocornia Perennis* Alpini. *Molecules* **2023**, *28*, doi:10.3390/molecules28062726.
59. Conversa, G.; Bonasia, A.; Lazzizzera, C.; La Rotonda, P.; Elia, A. Reduction of Nitrate Content in Baby-Leaf Lettuce and *Cichorium Endivia* through the Soilless Cultivation System, Electrical Conductivity and Management of Nutrient Solution. *Front. Plant Sci.* **2021**, *12*, doi:10.3389/fpls.2021.645671.
60. Santamaria, P.; Elia, A.; Serio, F.; Todaro, E. A Survey of Nitrate and Oxalate Content in Fresh Vegetables. *J. Sci. Food Agric.* **1999**, *79*, 1882–1888, doi:10.1002/(SICI)1097-0010(199910)79:13<1882::AID-JSFA450>3.0.CO;2-D.
61. Shi, R.; Liang, L.; Liu, W.; Zeb, A. *Kochia Scoparia* L., a Newfound Candidate Halophyte, for Phytoremediation of Cadmium-Contaminated Saline Soils. *Environ. Sci. Pollut. Res.* **2022**, *29*, 44759–44768, doi:10.1007/s11356-022-18895-8.
62. Haldimann, M.; Alt, A.; Blanc, A.; Blondeau, K. Iodine Content of Food Groups. *J. Food Compos. Anal.* **2005**, *18*, 461–471, doi:10.1016/j.jfca.2004.06.003.
63. Conversa, G.; Botticella, L.; Lazzizzera, C.; Bonasia, A.; Elia, A. Ecophysiological and Nutritional Characterisation of Two Morphotypes of *Cakile Maritima* Subsp. *Maritima* Scop. from Puglia Region, Southern Italy. *Front. Plant Sci.* **2024**, *15*, doi:10.3389/fpls.2024.1397852.
64. Duarte, B.; Feijão, E.; Pinto, M.V.; Matos, A.R.; Silva, A.; Figueiredo, A.; Fonseca, V.F.; Reis-Santos, P.; Caçador, I. Nutritional Valuation and Food Safety of Endemic Mediterranean Halophytes Species Cultivated in Abandoned Salt Pans under a Natural Irrigation Scheme. *Estuar. Coast. Shelf Sci.* **2022**, *265*, doi:10.1016/j.ecss.2021.107733.
65. EFSA NDA Panel (EFSA Panel on Panel on Dietetic Products Nutrition and Allergies) Scientific Opinion on Dietary Reference Values for Iodine. *EFSA J.* **2014**, *12*, 1–57, doi:10.2903/j.efsa.2014.3660.
66. Backer, H.; Hollowell, J. Use of Iodine for Water Disinfection: Iodine Toxicity and Maximum Recommended Dose. *Environ. Health Perspect.* **2000**, *108*, 679–684, doi:10.1289/ehp.00108679.
67. Ebrahimi, P.; Shokramraji, Z.; Tavakkoli, S.; Mihaylova, D.; Lante, A. Chlorophylls as Natural Bioactive Compounds Existing in Food By-Products: A Critical Review. *Plants* **2023**, *12*, 1533, doi:10.3390/plants12071533.
68. Ashikhmin, A.; Bolshakov, M.; Pashkovskiy, P.; Vereshchagin, M.; Khudyakova, A.; Shirshikova, G.; Kozhevnikova, A.; Kosobryukhov, A.; Kreslavski, V.; Kuznetsov, V.; et al. The Adaptive Role of Carotenoids

- and Anthocyanins in Solanum Lycopersicum Pigment Mutants under High Irradiance. *Cells* **2023**, *12*, 2569, doi:10.3390/cells12212569.
69. Tanaka, R.; Tanaka, A. Chlorophyll Cycle Regulates the Construction and Destruction of the Light-Harvesting Complexes. *Biochim. Biophys. Acta - Bioenerg.* **2011**, *1807*, 968–976, doi:10.1016/j.bbabi.2011.01.002.
 70. D'ambrosio, C.; Stigliani, A.L.; Giorio, G. Food from Genetically Engineered Plants: Tomato with Increased β -Carotene, Lutein, and Xanthophylls Contents. In *Genetically Modified Organisms in Food: Production, Safety, Regulation and Public Health*; Elsevier Inc., 2015; pp. 361–380 ISBN 978-012802530-7.
 71. Turck, D.; Bohn, T.; Castenmiller, J.; de Henauw, S.; Hirsch-Ernst, K.; Knutsen, H.K.; Maciuk, A.; Mangelsdorf, I.; McArdle, H.J.; Pentieva, K.; et al. Scientific Opinion on the Tolerable Upper Intake Level for Preformed Vitamin A and B-carotene. *EFSA J.* **2024**, *22*, doi:10.2903/j.efsa.2024.8814.
 72. Ghanem, A.-M.F.M.; Mohamed, E.; Kasem, A.M.M.A.; El-Ghamery, A.A. Differential Salt Tolerance Strategies in Three Halophytes from the Same Ecological Habitat: Augmentation of Antioxidant Enzymes and Compounds. *Plants* **2021**, *10*, 1100, doi:10.3390/plants10061100.
 73. Gargouri, M.; Magné, C.; Dauvergne, X.; Ksouri, R.; El Feki, A.; Metges, M.-A.G.; Talarmin, H. Cytoprotective and Antioxidant Effects of the Edible Halophyte *Sarcocornia Perennis* L. (Swampfire) against Lead-Induced Toxicity in Renal Cells. *Ecotoxicol. Environ. Saf.* **2013**, *95*, 44–51, doi:10.1016/j.ecoenv.2013.05.011.
 74. Gonzali, S.; Perata, P. Anthocyanins from Purple Tomatoes as Novel Antioxidants to Promote Human Health. *Antioxidants* **2020**, *9*, 1017, doi:10.3390/antiox9101017.
 75. Khoo, H.E.; Azlan, A.; Tang, S.T.; Lim, S.M. Anthocyanidins and Anthocyanins: Colored Pigments as Food, Pharmaceutical Ingredients, and the Potential Health Benefits. *Food Nutr. Res.* **2017**, *61*, 1361779, doi:10.1080/16546628.2017.1361779.
 76. Shao, H.-B.; Chu, L.-Y.; Jaleel, C.A.; Zhao, C.-X. Water-Deficit Stress-Induced Anatomical Changes in Higher Plants. *Comptes Rendus. Biol.* **2008**, *331*, 215–225, doi:10.1016/j.crv.2008.01.002.
 77. Zengin, G.; Aumeeruddy-Elalfi, Z.; Mollica, A.; Yilmaz, M.A.; Mahomoodally, M.F. In Vitro and in Silico Perspectives on Biological and Phytochemical Profile of Three Halophyte Species—A Source of Innovative Phytopharmaceuticals from Nature. *Phytomedicine* **2018**, *38*, 35–44, doi:10.1016/j.phymed.2017.10.017.
 78. Chekroun-Bechlaghem, N.; Belyagoubi-Benhammou, N.; Belyagoubi, L.; Gismondi, A.; Nanni, V.; Di Marco, G.; Canuti, L.; Canini, A.; El Haci, I.A.; Atik Bekkara, F. Phytochemical Analysis and Antioxidant Activity of *Tamarix Africana*, *Arthrocnemum Macrostachyum* and *Suaeda Fruticosa*, Three Halophyte Species from Algeria. *Plant Biosyst. - An Int. J. Deal. with all Asp. Plant Biol.* **2019**, *153*, 843–852, doi:10.1080/11263504.2018.1555191.
 79. Rodrigues, M.; Gangadhar, K.; Vizetto-Duarte, C.; Wubshet, S.; Nyberg, N.; Barreira, L.; Varela, J.; Custódio, L. Maritime Halophyte Species from Southern Portugal as Sources of Bioactive Molecules. *Mar. Drugs* **2014**, *12*, 2228–2244, doi:10.3390/md12042228.
 80. Stanković, M.; Stojanović-Radić, Z.; Jakovljević, D.; Zlatić, N.; Luković, M.; Dajić-Stevanović, Z. Coastal Halophytes: Potent Source of Bioactive Molecules from Saline Environment. *Plants* **2023**, *12*, 1857, doi:10.3390/plants12091857.
 81. Normén, L.; Johnsson, M.; Andersson, H.; van Gameren, Y.; Dutta, P. Plant Sterols in Vegetables and Fruits Commonly Consumed in Sweden. *Eur. J. Nutr.* **1999**, *38*, 84–89, doi:10.1007/s003940050048.
 82. Laitinen, K.; Gylling, H. Dose-Dependent LDL-Cholesterol Lowering Effect by Plant Stanol Ester Consumption: Clinical Evidence. *Lipids Health Dis.* **2012**, *11*, 140, doi:10.1186/1476-511X-11-140.
 83. Roche, J.; Alignan, M.; Bouniols, A.; Cerny, M.; Mouloungui, Z.; Vear, F.; Merah, O. Sterol Content in Sunflower Seeds (*Helianthus Annuus* L.) as Affected by Genotypes and Environmental Conditions. *Food Chem.* **2010**, *121*, 990–995, doi:10.1016/j.foodchem.2010.01.036.
 84. Pavlík, M.; Pavlíková, D.; Balík, J.; Neuberger, M. The Contents of Amino Acids and Sterols in Maize Plants Growing under Different Nitrogen Conditions. *Plant, Soil Environ.* **2010**, *56*, 125–132, doi:10.17221/214/2009-PSE.
 85. Roche, J.; Mouloungui, Z.; Cerny, M.; Merah, O. Effect of Sowing Dates on Fatty Acids and Phytosterols Patterns of *Carthamus Tinctorius* L. *Appl. Sci.* **2019**, *9*, 2839, doi:10.3390/app9142839.

86. Magni, N.N.; Veríssimo, A.C.S.; Silva, H.; Pinto, D.C.G.A. Metabolomic Profile of *Salicornia Perennis* Plant's Organs under Diverse in Situ Stress: The Ria de Aveiro Salt Marshes Case. *Metabolites* **2023**, *13*, 280, doi:10.3390/metabo13020280.
87. Rozentsvet, O.A.; Nesterov, V.N.; Bogdanova, E.S. Lipids of Halophyte Species Growing in Lake Elton Region (South East of the European Part of Russia). In *Handbook of Halophytes*; Grigore, M.N., Ed.; Springer Cham., 2021; pp. 2013–2037 ISBN 978-3-030-57634-9.
88. Isca, V.M.S.; Seca, A.M.L.; Pinto, D.C.G.A.; Silva, H.; Silva, A.M.S. Lipophilic Profile of the Edible Halophyte *Salicornia Ramosissima*. *Food Chem.* **2014**, *165*, 330–336, doi:10.1016/j.foodchem.2014.05.117.
89. Han, J.-H.; Yang, Y.-X.; Feng, M.-Y. Contents of Phytosterols in Vegetables and Fruits Commonly Consumed in China. *Biomed. Environ. Sci.* **2008**, *21*, 449–453, doi:10.1016/S0895-3988(09)60001-5.
90. Rumpf, J.; Burger, R.; Schulze, M. Statistical Evaluation of DPPH, ABTS, FRAP, and Folin-Ciocalteu Assays to Assess the Antioxidant Capacity of Lignins. *Int. J. Biol. Macromol.* **2023**, *233*, 123470, doi:10.1016/j.ijbiomac.2023.123470.
91. Hupel, M.; Lecointre, C.; Meudec, A.; Poupart, N.; Gall, E.A. Comparison of Photoprotective Responses to UV Radiation in the Brown Seaweed *Pelvetia Canaliculata* and the Marine Angiosperm *Salicornia Ramosissima*. *J. Exp. Mar. Bio. Ecol.* **2011**, *401*, 36–47, doi:10.1016/j.jembe.2011.03.004.
92. Kumari, N.; Rani, B.; Manne, H.; Jattan, M.; Sushil; Avtar, R.; Kumari, A.; Duhan, J.; Kodidhala, V. Antioxidative Response Mechanisms in Halophytes: Their Role in Stress Defence. In *Halophytes vis-à-vis Saline Agriculture*; Springer Nature Singapore: Singapore, 2024; pp. 329–350 ISBN 978-981-97-3156-5.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.