

Article

Not peer-reviewed version

On General Covariance

[Shoude Li](#) *

Posted Date: 27 October 2025

doi: 10.20944/preprints202505.0745.v6

Keywords: general covariance; general relativity; gravitational redshift; gravitational acceleration; differential geometry; covariant differential; Christoffel symbols; Lagrangian; energy momentum conservativeness; light angular momentum; geodesic line; kinematics; planet ring model; relativistic release; relativistic emission; relativistic absorption; relativistic redshift; broad line region; narrow line region



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

On General Covariance

Li Shoude

Hohai University, Nanjing 210098, China; lishoude@hhu.edu.cn

Abstract

If the invariance of light speed is absolutely universal in covariant space rather than contra variant space, the following researches and conclusions on general covariance would be so far as to catch up with very high probabilities and reliabilities. As has been verified, it is. Basically, general covariance is tested controversial after the investigations on gravitational redshift and acceleration. Further inspections on differential geometry indicate the opportunities of inequality of mixed derivatives of bases for the transformations between Riemannian spaces that will then lead to failure of the classical equations of Christoffel symbols, which is one of the reasons that causes controversies on general covariance. In fact, Christoffel symbols and base derivatives both are valid methodologies for analysis in a non-Euclidean space. Measurable experiments on gravitational redshifts and accelerations have been carried out to support the theoretical results. Conclusions have been drawn that light speed keeps general covariance in gravitational fields, but light energy momentum would not, may as well the motions of massive matters in gravitational fields do not perform general covariance thoroughly. Consequently, inferences on kinematics and relativistic release were put into researches, which have got surprising verifications in applications. In the researches on applicable problems of kinematics and relativistic release, the conclusions and inferences have been forcefully verified furtherly. With the ingenious proposition of the concept of gravitational metrics, the so-called geodesic equations have been falsified to be matter's kinematic equations anymore. The conservativeness of light angular momentum has been discovered in a most surprising form that has help to find out mistakes in traditional methodologies both on the issues of light deflect and radar echo delay. Renovated solutions for light rays as well as massive matters have been carried out that fabulously impact the traditional methodologies on kinematics of trajectory and time delay, and comprehensively interpret the realities. Newtonian ballistic method was put forward to make numerical analyses on close-to-light-speed motions, and the invalidity of that method on light propagation could be seen as another support to the conclusions and inferences. Especially, the forceful falsifications on energy momentum conservativeness carried out in the discussions on the traditional treatment of time delay of close-to-light-speed particles, have thoroughly exposed the essential mistakes in the way of methodologies. Exponent expression of dynamic mass has been carried out to create dynamic equations that prompt more generalized expressions of kinematics, for light ray's propagation, for massive matters traveling, especially for that thereby, massive matters moving in strong fields, the problem of planet perihelion precession drawn to argue about the only item that the renovated calculations shows controversial to the declared observations. Dynamic models of fluid planet rings were founded to interpret the evolutions of accretions of quasars and active galactic nuclei and the mechanism of relativistic release. The equation of relativistic release was developed after the conclusions and inferences. It is predicted that the peak release of an inflow is at 1.5 of gravitational radius and the peak luminosity of an accretion may locate at the position of 1.33 of gravitational radius. Relativistic frequency shift interprets the mechanism of giant redshift that predict the probability of observational redshift might be up to more than 20.0. The width equations of emission and absorption lines indicate the mechanism and the positions of broad line regions and narrow line regions. It could be imagined that the relativistic emissions and relativistic absorptions with relativistic redshifts would have involved with fantastic mystery of intrinsic structures of matters that we know less.

Keywords: general covariance; general relativity; gravitational redshift; gravitational acceleration; differential geometry; covariant differential; Christoffel symbols; Lagrangian; energy momentum

conservativeness; light angular momentum; geodesic line; kinematics; planet ring model; relativistic release; relativistic emission; relativistic absorption; relativistic redshift; broad line region; narrow line region

1. Preface

Einstein carried out the equivalence principle after discussing the equivalence of gravitational mass and inertial mass, and then he generalized the equivalent principle to create general relativity [1], which predicts same physics in curve space of gravity geometrization as that in no gravity space, that could be called general covariance. Theoretically, general covariance should include but not be limited in the performances of motion inertia, energy and momentum conservations as well as equilibrium states in complicated systems. However, we will find that quite amount of observations and evidences perform against general covariance.

It is believed that Riemannian geometry has been employed in general relativity for gravity geometrization [2]. But in fact, a transformation of a space does not really determine physics, what on earth the realities do. It is said that it is not the geometry but only the general covariance that matters. In fact, we could find out plenty of contradictions in classical theory of general relativity. It could be verified that even the motions of matter’s freefalling on the Earth cannot be well interpreted in the frames of the classical theory. So that it is time to sponsor a series of inspections and perceptions to insight into the topics on general covariance.

The gravitational redshift and gravitational acceleration are of the two typical effects that gravity acts on the light rays and massive matters respectively. Hence, researches on these topics would be greatly forceful to probe into the investigations and realizations on general covariance.

To discover realities is more important than to carry out new equations and theorems.

2. Contradictions on Gravitational Redshift and Acceleration

2.1. Newtonian Gravitational Redshift

The equation of gravitational redshift can be drawn via Doppler redshift in a thought experiment that a light ray be emitted from the ceiling to the bottom or that is reversely performed from bottom to the ceiling, in a freely falling elevator cabin in a center source field [3]. It could be verified that any observer in the cabin would detect no frequency shift whatever the ways the light emitted. Suppose another observer outside the cabin keeping rest so that to have a relative velocity against the cabin, who will then detect a frequency shift other than the freefalling observer. What I want to say is that whoever freefalling will detect no frequency shift, no matter inside or outside the cabin. It is the relative motion that eliminates the gravitational frequency shift. Once the relative velocities catch up to a relativistic velocity, the gravitational frequency shift will then cannot be eliminated anymore.

As a light ray is down ward emitted from a cabin ceiling and at the same time the cabin is released to freely fall, the light front should spend a very short time interval to reach the bottom that

$$\Delta t = \frac{\Delta H}{c} \tag{1}$$

where, ΔH is the distance that the light ray travels from the start to the end which may approximately equal to the height of the cabin, c is light speed.

Thus, the cabin velocity increase is

$$\Delta v = a\Delta t \tag{2}$$

where a is gravitational acceleration, which approximately equals to gravity g case velocity does not reach relativistic level. We know it exactly has a minus value as its direction pointing to center source.

Doppler redshift is frequency shift between two observers that one has a relative motion to another, to detect frequency. Here the rest observer outside would have a relative velocity $-\Delta v$ or $+\Delta v$ to the cabin so that when they receive the light ray at the same time with the inner, the Doppler redshift could be calculated as

$$z_D = \frac{\Delta v}{c} = \frac{g\Delta H}{c^2} = \frac{\Delta\phi}{mc^2} \quad (3)$$

With the calculation of Doppler redshift, we know that the gravitational redshift has happened in the same value. Notwithstanding, I prefer to suggest a new methodology to get an equation for gravitational redshift, in which a proposal should be given that every photon at any position in a center source field could be assumed to have experienced a travel from a farthest point to the current position. This attempt may bring about more physical significances and comprehensive understandings.

A photon in one source field at position r with frequency $\nu_{0(r)}$ is set to have an imaginary primary frequency $\nu_{0(\infty)}$ at a farthest point

$$E_\infty = h\nu_{0(\infty)} \quad (4)$$

where h is Planck's constant. NB, we are not talking about quantum character of photons so that photonic energy momentum mentioned refers to statistic quantities.

The corresponding dynamic mass comes from the mass-energy equation is

$$m = E_\infty c^{-2} \quad (5)$$

where c is light speed.

Then the gravitational potential at position r , especially as is in weak field with $r \gg r^*$, could be written as

$$\phi_r = \int_\infty^r \frac{GMm_r}{r^2} dr = -\frac{GM\bar{m}}{r} \approx -\frac{GMm}{r} = -\frac{r^*}{2r} mc^2 \quad (6)$$

where G is gravitational constant, M is the mass of the center source with $M \gg m_r$, m_r is photonic mass at position r , $\bar{m}$ is the mean mass for the integral and it could be approximately replaced by m case in weak field, and r^* is Schwarzschild radius written as $r^* = 2GM/c^2$.

The current dynamic energy is the summation of primary dynamic energy and released potential

$$E_r = h\nu_{0(r)} = E_\infty + [\phi_\infty - \phi_r] \approx mc^2 + \frac{r^*}{2r} mc^2 \approx h\nu_{0(\infty)} \left(1 + \frac{r^*}{2r}\right) \quad (7)$$

Case a light ray travels from positions r_1 to r_2 , as is shown in Figure 1, gravitational redshift happens.




Figure 1. Center source field.

Gravitational redshift could be defined traditionally as

$$z_{g\lambda} = \frac{\lambda^1_{(2)} - \lambda^1_{(1)}}{\lambda^1_{(1)}} = \frac{\nu_{0(1)} - \nu_{0(2)}}{\nu_{0(2)}} \approx \frac{r^*}{r_{(1)}} \cdot \frac{r_{(2)} - r_{(1)}}{2r_{(2)} + r^*} \quad (8)$$

Considering weak field effects, the gravitational redshift is

$$z_{g\lambda} \approx r^* \cdot \frac{r_{(2)} - r_{(1)}}{2r_{(1)}r_{(2)}} = \frac{r^*}{2r_{(1)}} - \frac{r^*}{2r_{(2)}} = \frac{\phi_{(2)} - \phi_{(1)}}{mc^2} = \frac{\Delta\phi}{mc^2} \quad (9)$$

They are the forms of frequency shift of wave length based expression in weak fields, called redshift, where λ^1 and ν_0 are wave length and frequency tensors in contra variant space. Of course, a frequency shift could also be defined based on frequency. But we have been used to the forms previous in tradition. In the wave length based equation, redshift may go up to more than 1.0, while blueshift must have been limited in -1 to 0. Case in the form of frequency based equations, one could get blueshift greater than 1.0 but redshift limited in -1 to 0.

We could also carry out new forms of frequency shift for conveniences in special discussions, that a differential of wave length based redshift could be defined as

$$dz_\lambda = \frac{d\lambda^1(r)}{\lambda^1(r)} = d\ln\lambda^1(r) \quad (10)$$

So that the integral form for $r_{(1)}$ to $r_{(2)}$

$$z_\lambda = \ln\lambda^1(r) \Big|_{r_{(1)}}^{r_{(2)}} = \ln\frac{\lambda^1(2)}{\lambda^1(1)} = \ln\frac{\nu_{0(1)}}{\nu_{0(2)}} \quad (11)$$

Or a differential form of frequency based

$$dz_\nu = \frac{d\nu_{0(r)}}{\nu_{0(r)}} = d\ln\nu_{0(r)} \quad (12)$$

So that

$$z_\nu = \ln\nu_{0(r)} \Big|_{r_{(1)}}^{r_{(2)}} = \ln\frac{\nu_{0(2)}}{\nu_{0(1)}} = \ln\frac{\lambda^1(1)}{\lambda^1(2)} = -\ln\frac{\lambda^1(2)}{\lambda^1(1)} \quad (13)$$

Considering the energy expression of the Equation (467) in to the Equation (11), one can gain the redshift result as same as the Equation (9). But they are based on different definitions of frequency shift so that they would perform physical implications a little different.

It is said that after the definitions of Equation (11) and Equation (13) there is

$$z_\lambda = -z_\nu \quad (14)$$

We have seen that they have shown difference from traditional equations. But for small frequency shift, both the two integral equations could be use instead of traditional equation.

Moreover, with Equation (7), the gravitational frequency differential

$$d\nu_{0(r)} \approx d[\nu_{0(\infty)} \left(1 + \frac{r^*}{2r}\right)] = -\frac{r^*}{2r^2} \nu_{0(\infty)} dr \quad (15)$$

Then the differential frequency shift goes

$$dz_{g\nu} = \frac{d\nu_{0(r)}}{\nu_{0(r)}} \approx \frac{\frac{r^*}{2r^2}}{1 + \frac{r^*}{2r}} dr = -\frac{r^* dr}{r(2r + r^*)} \quad (16)$$

This equation will be also taken into further discussions in the next sections. In following sections, the subscripts of frequency shift symbols will be neglected for convenience and in most cases we use the concept redshift to present frequency shifts.

2.2. Errors in the Equation of the So-Called Revisit Gravitational Redshift

A thought experiment has ever been employed to present the concept of the so-called revisit gravitational redshift [3,4], in which a pulse of light ray is supposed to be emitted from position 1, lasting for a time interval $\Delta t_{(1)}$, and be received at position 2 within a time interval $\Delta t_{(2)}$. The world lines of the photons are shown in Figure 2. We then know that the two drawn lines are literally the world lines of the first photon and the final photon of the light pulse.




Figure 2. World lines for a pulse of photons.

Believing that the two world lines besides the $\Delta t_{(1)}$ and $\Delta t_{(2)}$ are parallel, it is known that the time interval $\Delta t_{(1)}$ equals to that of $\Delta t_{(2)}$. As the light frequency being inversely proportional to proper time interval $\Delta \tau$, the proper forms of frequency ratio were written in some textbooks as [3,4]

$$\frac{\nu_{(1)}}{\nu_{(2)}} = \frac{\Delta \tau_{(2)}}{\Delta \tau_{(1)}} = \frac{\sqrt{g_{00(2)} \Delta t_{(2)}}}{\sqrt{g_{00(1)} \Delta t_{(1)}}} = \frac{\sqrt{g_{00(2)}}}{\sqrt{g_{00(1)}}} \approx 1 + \frac{\phi_{(2)} - \phi_{(1)}}{mc^2} \quad (17)$$

where $\nu_{(1)}$ and $\nu_{(2)}$ are proper frequencies corresponding to position 1 and position 2, g_{00} is time metric of Schwarzschild. The right hand side of this equation is an approximate result with Schwarzschild solution of metric g_{00} in condition that $r_2 > r_1 \gg r^*$.

Thus, the so-called revisit redshift is

$$z_{revisit} = \frac{\nu_{(1)}}{\nu_{(2)}} - 1 \approx \frac{\Delta \phi}{mc^2} \quad (18)$$

It is seemingly that the revisit form of equation for gravitational redshift was worked out.

But there are quite many errors in above equations. (i) The two world lines belong to the first photon and the final photon respectively, thus they might be controlled by emitter, so that the time intervals between the two lines do nothing with any light frequencies. (ii) Any time intervals between neighboring photons could be randomly assigned, so that these intervals also do nothing with any light frequencies. (iii) Frequency of a photon is the reciprocal of its photonic period and is the intrinsic property of a photon, so that it is independent to its positions relating to other photons and any variation of the frequency will not change its position in the pulse of photons. (iv) Detections of frequencies must involve with wave numbers and time intervals together, while this equation has made a mistake by comparing time intervals only. (v) It is said that the time intervals $\Delta t_{(1)}$ and $\Delta t_{(2)}$ do nothing with the physical events of frequency shift, they are just the records of the physical event of emission and receiving.

2.3. Further Investigation into the Revisit Gravitational Redshift

Following the rules in classical frames, the tensor of light wave period is a tensor with upper index

$$T^0 = \frac{dt}{dn} \quad (19)$$

where, T^0 is contra variant light period that is described by coordinate time, and dt is coordinate time lasting in which the number of dn waves may have traveled across a specific position. And the proper form of wave period

$$T = \frac{d\tau}{dn} \quad (20)$$

where, $d\tau$ is proper time lasting for dn number of waves to cross the position. So that there is

$$T = e_0 T^0 \text{ or } T^0 = e^0 T \quad (21)$$

where, e_0 is the first component of covariant time base, and it should be noted that the base e_0 is a vector but the component e_0 is not vector even though it is still a tensor. One can get more understandings for these expressions I would sponsor here and in followings.

In this way, we know that different values of the contra variant period and proper period correspond to a same physical issue of wave counting. Then it leads to a frequency tensor

$$\nu_0 = \frac{1}{T_0} = e_0 \nu = \frac{dn}{dt} \quad (22)$$

We have seen that, ν_0 is called covariant tensor traditionally, and ν is a proper tensor. This may have brought about confusions in that ν is actually covariant and yet ν_0 has been named the name. I am not going to change the naming methodology thoroughly right now because that may cause more difficulties and sounds more trivial.

Generally, some pure one-order tensors seem to be infinite small quantities such as dr , but for ν_0 , it is dn divided by dt , so that it is not an infinite small quantity. As for velocity tensors, they are really mixed tensors.

Theoretically, the covariant derivative of a frequency in a falling process into a center source can be written as

$$\frac{D\nu_0}{dr} = \frac{\partial \nu_0}{\partial r} - \Gamma_{10}^0 \nu_0 \quad (23)$$

where, $\Gamma_{\nu\mu}^\lambda$ is Christoffel symbols.

We know that, as has been presented in Equation (15), the contra variant derivative is

$$\frac{\partial \nu_0}{\partial r} = -\frac{r^*}{2r^2} \nu_{0(\infty)} \quad (24)$$

Let us calculate the Christoffel symbols in Equation (23) that

$$\Gamma_{10}^0 = \frac{1}{2} g^{0\lambda} \left(\frac{\partial g_{1\lambda}}{\partial x^0} + \frac{\partial g_{0\lambda}}{\partial x^1} - \frac{\partial g_{10}}{\partial x^\lambda} \right) \quad (25)$$

The Einstein summation convention has been and will be adopt unless additional declarations.

It is found that only in the condition of $\lambda = 0$ there is a nonvanishing item in the bracket of right hand side of Equation (25), so that with Schwarzschild metrics it turns to be

$$\Gamma_{10}^0 = \frac{1}{2} g^{00} \frac{\partial g_{00}}{\partial x^1} = \frac{1}{2} \left(-1 + \frac{r^*}{r} \right)^{-1} \left(-1 + \frac{r^*}{r} \right)' = \frac{r^*}{2r^2} \left(1 + \frac{r^*}{r-r^*} \right) \quad (26)$$

The covariant derivative will be calculated to be

$$\begin{aligned} \frac{D\nu_0}{dr} &= -\frac{r^*}{2r^2} \nu_{0(\infty)} - \frac{r^*}{2r^2} \left(1 + \frac{r^*}{r-r^*} \right) \nu_0 \\ &= -\frac{r^*}{2r^2} \nu_{0(\infty)} - \frac{r^*}{2r^2} \left(1 + \frac{r^*}{r-r^*} \right) \left(1 + \frac{r^*}{2r} \right) \nu_{0(\infty)} \end{aligned} \quad (27)$$

We could get an approximate solution for weak field that

$$\frac{D\nu_0}{dr} \approx 2 \frac{\partial \nu_0}{\partial r} \quad (28)$$

As we have seen, it shows that the values of covariant redshift doubles that of the contra variant redshift.

2.4. Additional Discussions on the Thought Experiment of Freefalling Elevator Cabin

The thought experiment that observer in freefalling elevator cabin will detect no frequency shift is usually employed to discuss the equivalent principle for the support of general covariance. But in fact, that issue does nothing with covariance. It is just because that the gravitational redshift happens to be offset by Doppler redshift. In fact, this experiment is a comprehensive event that relates both to gravitational redshift and gravitational acceleration.

We know that in a freefalling cabin, freefalling observer will not detect any frequency shift no matter the light ray emitter is on the bottom or on the top to emit up or down to the receiver.

Nevertheless, even if the light ray is not vertical, freefalling observer would also observe no frequency shift in the cabin.

For a case that a light ray is emitted to the top with an angle θ to the vertical line

The time interval for light ray from bottom to the top is

$$\Delta t = \frac{1}{c} \frac{\Delta H}{\cos \theta} \quad (29)$$

And the velocity increase of freefall cabin is

$$\Delta v = a \Delta t = \frac{g}{c} \frac{\Delta H}{\cos \theta} \quad (30)$$

Then the velocity increase component at the direction of light ray

$$\Delta v_l = \Delta v \cos \theta = g \frac{\Delta H}{c} \quad (31)$$

So that we get the Doppler redshift again as

$$z_D = \frac{\Delta v_l}{c} = \frac{g \Delta H}{c^2} = \frac{\Delta \phi}{mc^2} \quad (32)$$

In fact, the Doppler redshift for the detector in freefalling cabin could eliminate the gravitational frequency shift, even if the emitter is either outside of the cabin or with any low initial velocity. The real reason for non-detectable frequency shift in freefalling cabin is that the relative motion formed Doppler redshift just has a minus approximate value of gravitational frequency shift.

I prefer to put forward the case that the gravitational frequency shift cannot be eliminated by Doppler redshift. In the case that cabin has a relativistic initial velocity, the geometrical acceleration is not the total gravitational acceleration again as that in Equation (297) as

$$a = \eta g, \eta < 1 \quad (33)$$

Thus, for light rays passing across the cabin there is the Doppler velocity of detector

$$\Delta v_l = a \Delta t = \eta g \frac{\Delta H}{c} \quad (34)$$

so that

$$z_D = \frac{\Delta v_l}{c} = \eta \frac{g \Delta H}{c^2} = \eta \frac{\Delta \phi}{mc^2} \quad (35)$$

While the gravitational frequency is still the form

$$z_g = \frac{g \Delta H}{c^2} = \frac{\Delta \phi}{mc^2} \quad (36)$$

so that in this case

$$z_D \neq z_g \quad (37)$$

We will see that in some situations in freefalling cabin one could detect frequency shift again, so that the thought experiment cannot support general covariance thoroughly. Of course, one can continue to argue that relativistic motion may bring about more sophisticated conditions on frequency shift.

2.5. An Investigation into Gravitational Acceleration

For a freefalling massive matter in gravitational field, the component of the velocity at the direction of radius of center source is

$$V^1 = \frac{dx^1}{d\tau} \quad (38)$$

where, dx^1 is number one component of vector of moving distance, in this case it is dr .

There is the covariant derivative at one of coordinate direction

$$\frac{DV^1}{d\lambda} = \frac{\partial V^1}{\partial \lambda} + \Gamma_{\lambda\mu}^1 V^\mu \quad (39)$$

Because $dx^0 = cdt$, and $dt = e^0 d\tau$, where e^0 is the nonvanishing component of the base e^0 . Thus, the covariant derivative by τ is

$$\frac{DV^1}{d\tau} = \frac{dV^1}{d\tau} + \frac{dx^0}{d\tau} \Gamma_{0\mu}^1 V^\mu = a^1 + ce^0 \Gamma_{0\mu}^1 V^\mu \quad (40)$$

where, $a^1 = \frac{dV^1}{d\tau}$ is the contra variant acceleration of the matter, c is light speed.

NB, accelerations we have and will discuss refer to geometrical accelerations, which will be something different from gravity g , in that the latter sometimes may also be called gravitational accelerations in some writings but in fact matters may not experience accelerating as that.

With the equation of Christoffel symbols

$$\Gamma_{0\lambda}^1 = \frac{1}{2} g^{1\rho} \left(\frac{\partial g_{0\rho}}{\partial x^\lambda} + \frac{\partial g_{\lambda\rho}}{\partial x^0} - \frac{\partial g_{0\lambda}}{\partial x^\rho} \right) \quad (41)$$

case $\lambda = 1$ in this equation, there is $\Gamma_{01}^1 = 0$. Then considering the condition of $\lambda = 0$, it is

$$\Gamma_{00}^1 = \frac{1}{2} g^{1\rho} \left(\frac{\partial g_{0\rho}}{\partial x^0} + \frac{\partial g_{0\rho}}{\partial x^0} - \frac{\partial g_{00}}{\partial x^\rho} \right) \quad (42)$$

In this condition, only in the case of $\rho = 1$ there is the nonvanishing item in the bracket of right hand side, so that

$$\Gamma_{00}^1 = \frac{1}{2} g^{11} \left(-\frac{\partial g_{00}}{\partial x^1} \right) = \frac{1}{2} \left(1 - \frac{r^*}{r} \right) \left(1 - \frac{r^*}{r} \right)' = \frac{1}{2} \left(1 - \frac{r^*}{r} \right) \frac{r^*}{r^2} \quad (43)$$

And with $V^0 = \frac{dx^0}{d\tau} = \frac{cdt}{d\tau} = e^0 c$, Equation (40) turns to be

$$\frac{DV^1}{d\tau} = a^1 + ce^0 \Gamma_{00}^1 V^0 = a^1 + e^0 e^0 c^2 \left(1 - \frac{r^*}{r} \right) \frac{r^*}{2r^2} \quad (44)$$

As a matter freefalls on to the earth with irrelativistic velocity, its acceleration could be calculated to be

$$a^1 = -e^0 e^0 \frac{GM}{r^2} = -e^0 e^0 c^2 \frac{r^*}{2r^2} \quad (45)$$

Thus, there is the weak field solution

$$\frac{DV^1}{d\tau} \approx 0 \quad (46)$$

2.6. Discussions and Controversies

Now we have got more complex results that the revisit gravitational redshift calculated doubles that of contra variant value while covariant acceleration of freefalling goes to zero. The latter seems to fit with general covariance but the previous does not. And there is still a problem that revisit gravitational redshift solved in the way of covariant derivation goes contradictory to classical solution. Moreover, the minus form of g_{00} also involves more matters that have caused the result of zero acceleration, deserving of further discussions in next sections.

It is extraordinary that the covariant derivative of light frequency goes nonvanishing in the view of general covariance. Even more, it is investigated that covariant derivative analysis shows that the classical solution of revisit redshift may be false in that the frequency has been wrong defined. Someone may argue that the frequency is not a tensor, but that makes no sense because light frequency is the reciprocal of its wave period which involves with time coordinate.

What I urgently want to say is that these discussions are not enough. The most significant problem is that, the item in original covariant differential in Equation (44) that matter's velocities multiplied with the base's differential, has been calculated to do nothing with the realistic velocity. We should know that the multiplied item in Equation (39) originally indicates a base variation ratio multiplied with the very tensors, but the final equation has given up the effects of initial velocity that does lead to contradictions, and the time speed V^0 is a virtual velocity which might have been abused. These contradictions really bothered me until it is occasionally gone through one day, that the real problem is deeply hidden in the equations of Christoffel symbols.

Notwithstanding, a differential of velocity is the differential of that on the trajectory of matter's motion. Thus, the covariant time derivatives cannot be treated directly as ordinary derivatives anymore as in Equation (40). We are going to carry out detailed discussions on trajectory derivatives in next sections so that to interpret covariant time derivatives correctly.

3. Theoretical Investigations on Christoffel Symbols

3.1. Classical Equations of Christoffel Symbols

Christoffel symbols have been defined in the equation

$$\frac{\partial e^\mu}{\partial x^\nu} = -\Gamma_{\nu\lambda}^\mu e^\lambda \quad (47)$$

There is nothing wrong with the definition in that the derivative of a base must have a direction and so that to be written as a linear combination of the total bases. The key problem is what the Christoffel symbols are.

In this and following sections, all symbols of vectors and matrix would be boldly written while their components and that of other quantities may be simply written, no matter they are tensors or not.

In most conditions, Christoffel symbols could be discussed by the derivation of metrics as

$$\frac{\partial g_{\mu\nu}}{\partial x^\lambda} = \frac{\partial}{\partial x^\lambda} (e_\mu \cdot e_\nu) = \frac{\partial e_\mu}{\partial x^\lambda} \cdot e_\nu + e_\mu \cdot \frac{\partial e_\nu}{\partial x^\lambda} = \Gamma_{\lambda\mu}^\rho e_\rho \cdot e_\nu + \Gamma_{\lambda\nu}^\rho e_\rho \cdot e_\mu \quad (48)$$

For the derivative forms with alternative indexes mathematically, there will be

$$\frac{\partial g_{\mu\nu}}{\partial x^\lambda} - \Gamma_{\lambda\mu}^\rho g_{\rho\nu} - \Gamma_{\lambda\nu}^\rho g_{\rho\mu} = 0 \quad (49)$$

$$\frac{\partial g_{\lambda\nu}}{\partial x^\mu} - \Gamma_{\mu\lambda}^\rho g_{\rho\nu} - \Gamma_{\mu\nu}^\rho g_{\rho\lambda} = 0 \quad (50)$$

$$\frac{\partial g_{\mu\lambda}}{\partial x^\nu} - \Gamma_{\nu\mu}^\rho g_{\rho\lambda} - \Gamma_{\nu\lambda}^\rho g_{\rho\mu} = 0 \quad (51)$$

In the case the so-called torsions $S_{\mu\nu}^\rho = \Gamma_{\mu\nu}^\rho - \Gamma_{\nu\mu}^\rho$ are set zero, the summation of the previous two equations minus the last one that

$$\frac{\partial g_{\mu\nu}}{\partial x^\lambda} + \frac{\partial g_{\lambda\nu}}{\partial x^\mu} - \frac{\partial g_{\mu\lambda}}{\partial x^\nu} - 2\Gamma_{\lambda\mu}^\rho g_{\rho\nu} = 0 \quad (52)$$

Thereafter the equations of Christoffel symbols will be solved as [2,3,5]

$$\Gamma_{\lambda\mu}^\rho = \frac{1}{2} g^{\rho\nu} \left(\frac{\partial g_{\mu\nu}}{\partial x^\lambda} + \frac{\partial g_{\lambda\nu}}{\partial x^\mu} - \frac{\partial g_{\mu\lambda}}{\partial x^\nu} \right) \quad (53)$$

Generally, Equation (49) to Equation (51) could also be rewritten as the original forms

$$\frac{\partial g_{\mu\nu}}{\partial x^\lambda} - \frac{\partial e_\mu}{\partial x^\lambda} \cdot e_\nu - \frac{\partial e_\nu}{\partial x^\lambda} \cdot e_\mu = 0 \quad (54)$$

$$\frac{\partial g_{\lambda\nu}}{\partial x^\mu} - \frac{\partial e_\lambda}{\partial x^\mu} \cdot e_\nu - \frac{\partial e_\nu}{\partial x^\mu} \cdot e_\lambda = 0 \quad (55)$$

$$\frac{\partial g_{\mu\lambda}}{\partial x^\nu} - \frac{\partial e_\mu}{\partial x^\nu} \cdot e_\lambda - \frac{\partial e_\lambda}{\partial x^\nu} \cdot e_\mu = 0 \quad (56)$$

We will find that in some conditions the torsions do not always equal to zero. It is said that the mixed derivatives of bases $\frac{\partial e_\mu}{\partial x^\nu}$ and $\frac{\partial e_\nu}{\partial x^\mu}$ do not always equal, and then the Christoffel symbols with mixed subscripts $\Gamma_{\mu\nu}^\rho$ and $\Gamma_{\nu\mu}^\rho$ do not always equal, so that the Equation (53) might be invalid in those conditions.

3.2. The Inequality of Christoffel Symbols of Mixed Subscripts

In Riemannian differential geometry, the equality of Christoffel symbols of mixed subscripts is usually adopted. But no forceful researches could provide reliable supports. The truth is that the problem of mixed derivatives of bases in a Riemannian space are far different from the problem of normal mixed derivatives of a 3-dimensional surface in Euclidean geometry.

We could find out the truth that in the deduction of Γ_{00}^1 in Equation (43), $\frac{\partial g_{00}}{\partial x^1}$ has been used instead of $\frac{\partial g_{11}}{\partial x^0}$ so that to gain $\Gamma_{00}^1 = \frac{1}{2}g^{11}(-\frac{\partial g_{00}}{\partial x^1})$. But we can easily calculate that $\frac{\partial e_0}{\partial x^1}$ and $\frac{\partial e_1}{\partial x^0}$ are not equal. It is said that the Equation (43) could lead to nonvanishing value of Γ_{00}^1 , which really relates to time derivatives of bases that must be determined to be zero in rest field. We might have found out the problems.

3.2.1. Coordinate Transformations and Bases

Any points in a Riemannian space of Riemannian manifold of dimension n has a neighborhood homeomorphic to a subset of Euclidean space of dimension n , so that there must be probable maps between the neighborhoods and the corresponding subsets. It is just to say that the coordinates of any points in Riemannian space could be expressed with the coordinates of corresponding points of Euclidean space, and reversely. If a part or the entire of a Riemannian space are continuous and differentiable, Euclidean coordinate lines could be drawn in the part or the entire of the Riemannian space. On the other side, coordinate lines of Riemannian space could also be drawn in the corresponding Euclidean space. For convenience, the Riemannian space could be called covariant space, and the corresponding Euclidean space could be called contra variant space. A contra variant space is curved in the view of its covariant space, and the covariant space is also curved in the view of contra variant space.

It is obvious that transformations of spaces are actual coordinate transformations. These transformations could happen between covariant space and contra variant space, as well as they could happen among homeomorphic Riemannian spaces. Coordinate transformations may perform in the way with unequal metrics as well as the way with equal metrics.

The more effective method for coordinate transformation is to define bases and distances for spaces. Two examples would be presented firstly for definitions and for following discussions.

Example 1: Bases of Riemannian manifold of super surface

The derivative vectors of 3-dimensional surfaces were usually employed to form bases in classical differential geometry. The curve space has an extra dimension than a plane space that could be called the super surface. The Riemannian manifold of the super surface in the 3-dimensional space (u, v, w) would have a homeomorphic Euclidean space (x, y) in the space (x, y, z) . The coordinate lines x and y in contra variant space could be transformed to be $\xi(x)$ and $\eta(y)$ in covariant space, while u , v and w in covariant space be transformed to be $\alpha(u)$, $\beta(v)$ and $\gamma(w)$ in the space (x, y, z) as shown in Figure 3 and Fig.4.




Figure 3. A super surface as covariant space in 3-dimensional space .




Figure 4. A coordinate plane (x, y) as contra variant space in 3-dimensional space.

The super surface could be determined by a vector function $\boldsymbol{\rho}$

$$\boldsymbol{\rho} = \boldsymbol{\rho}(u, v, w)|_{2 \text{ of variables independent}} \tag{57}$$

And the function could also be written as

$$\boldsymbol{\rho} = \boldsymbol{\rho}(\xi, \eta) \tag{58}$$

or

$$\boldsymbol{\rho} = \boldsymbol{\rho}(x, y) \tag{59}$$

At the same time, the contra variant space could be defined by $\boldsymbol{r}$

$$\boldsymbol{r} = \boldsymbol{r}(x, y, z)|_{z=const.} = \boldsymbol{r}(\alpha, \beta, \gamma)|_{2 \text{ of variables independent}} = \boldsymbol{r}(x, y) = \boldsymbol{r}(\xi, \eta) \tag{60}$$

There will be varieties of available ways to develop the expressions of bases and distances. I prefer to put forward the followings might as well.

The way in super space:

In super space, the differential $d\boldsymbol{\rho}$ has 3 components

$$d\boldsymbol{\rho} = \begin{pmatrix} du \\ dv \\ dw \end{pmatrix} \tag{61}$$

That of differential $d\boldsymbol{r}$ could be simplified to be 2 dimensional because it just locates in the space (x, y)

$$d\boldsymbol{r} = \begin{pmatrix} dx \\ dy \end{pmatrix} \tag{62}$$

To define a set of covariant bases for a position in covariant space by

$$\boldsymbol{e}_x = \frac{\partial \boldsymbol{\rho}}{\partial x}, \boldsymbol{e}_y = \frac{\partial \boldsymbol{\rho}}{\partial y} \tag{63}$$

It should be pointed out that, in some publications, coordinate and vector symbols have been used reversely like $d\boldsymbol{\rho}$ and $d\boldsymbol{r}$, which would have brought about confusions.

In covariant space, the differential $d\boldsymbol{\rho}$ expressed by $d\boldsymbol{r}$ with covariant bases

$$d\boldsymbol{\rho} = dx\boldsymbol{e}_x + dy\boldsymbol{e}_y \tag{64}$$

Differential distance could be defined as

$$ds^2 = du^2 + dv^2 + dw^2 = d\boldsymbol{\rho} \cdot d\boldsymbol{\rho} = (dx\boldsymbol{e}_x + dy\boldsymbol{e}_y) \cdot (dx\boldsymbol{e}_x + dy\boldsymbol{e}_y) \tag{65}$$

If the bases are orthogonal, there is

$$ds^2 = \boldsymbol{e}_x \cdot \boldsymbol{e}_x dx^2 + \boldsymbol{e}_y \cdot \boldsymbol{e}_y dy^2 \tag{66}$$

We have seen that the covariant bases are defined in covariant space to help contra variant coordinates to form covariant distances.

There are more complexities for a transformation between a super surface in 3-dimensional space and R^2 that the 2 covariant bases would have 3 components

$$\boldsymbol{e}_x = \begin{pmatrix} \frac{\partial u}{\partial x} \\ \frac{\partial v}{\partial x} \\ \frac{\partial w}{\partial x} \end{pmatrix}, \boldsymbol{e}_y = \begin{pmatrix} \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial y} \\ \frac{\partial w}{\partial y} \end{pmatrix} \tag{67}$$

Define the contra variant bases for a point in the plane that

$$\boldsymbol{e}^u = \frac{\partial \boldsymbol{r}}{\partial u}, \boldsymbol{e}^v = \frac{\partial \boldsymbol{r}}{\partial v}, \boldsymbol{e}^w = \frac{\partial \boldsymbol{r}}{\partial w} \tag{68}$$

The 3 contra variant bases all have 2 components as

$$\mathbf{e}^u = \begin{pmatrix} \frac{\partial x}{\partial u} \\ \frac{\partial y}{\partial u} \end{pmatrix}, \mathbf{e}^v = \begin{pmatrix} \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial v} \end{pmatrix}, \mathbf{e}^w = \begin{pmatrix} \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial w} \end{pmatrix} \quad (69)$$

The differential $d\mathbf{r}$ expressed by $d\boldsymbol{\rho}$ in contra variant space

$$d\mathbf{r} = du\mathbf{e}^u + dv\mathbf{e}^v + dw\mathbf{e}^w \quad (70)$$

Of course, one can create transformation matrix to perform the relationship between $d\mathbf{r}$ and $d\boldsymbol{\rho}$ directly.

The distance could be defined as

$$d\zeta^2 = dx^2 + dy^2 = d\mathbf{r} \cdot d\mathbf{r} = (du\mathbf{e}^u + dv\mathbf{e}^v + dw\mathbf{e}^w) \cdot (du\mathbf{e}^u + dv\mathbf{e}^v + dw\mathbf{e}^w) \quad (71)$$

If the bases are orthogonal

$$d\zeta^2 = \mathbf{e}^u \cdot \mathbf{e}^u du^2 + \mathbf{e}^v \cdot \mathbf{e}^v dv^2 + \mathbf{e}^w \cdot \mathbf{e}^w dw^2 \quad (72)$$

One could imagine that this condition you cannot give the relationship of metrics that g_{ii} equals to $1/g^{ii}$, in that the covariant bases have 3 components and contra variant bases have 2.

The way in tangent space:

Consequently, the issues could be simplified in tangent spaces. At a position $\boldsymbol{\rho}$ in the covariant space, there is a neighborhood which will be labeled with coordinate lines of $\xi(x)$ and $\eta(y)$, at the same time at the position $\mathbf{r}$, there is a corresponding neighborhood in contra variant space labeled with coordinate lines of x and y , as shown in Figure 5 and Figure 6. Generally, coordinate lines could be set orthogonal. In most of publications, $\xi(x)$ and $\eta(y)$ were seen as x and y , but one should realize that the difference really matters.




Figure 5. Covariant tangent space.




Figure 6. Contra variant tangent space.

One could define the differential vector in covariant space

$$d\boldsymbol{\rho} = \begin{pmatrix} d\xi \\ d\eta \end{pmatrix} \quad (73)$$

As a result, the bases

$$\mathbf{e}_x = \frac{\partial \boldsymbol{\rho}}{\partial x} = \begin{pmatrix} \frac{\partial \xi}{\partial x} \\ \frac{\partial \eta}{\partial x} \end{pmatrix}, \mathbf{e}_y = \frac{\partial \boldsymbol{\rho}}{\partial y} = \begin{pmatrix} \frac{\partial \xi}{\partial y} \\ \frac{\partial \eta}{\partial y} \end{pmatrix} \quad (74)$$

Again, there is the distance

$$ds^2 = d\xi^2 + d\eta^2 = d\boldsymbol{\rho} \cdot d\boldsymbol{\rho} = \mathbf{e}_x \cdot \mathbf{e}_x dx^2 + \mathbf{e}_y \cdot \mathbf{e}_y dy^2 \quad (75)$$

The differential $d\mathbf{r}$ keep the form as Equation (62), so that the contra variant bases could be defined as

$$\mathbf{e}^\xi = \frac{\partial \mathbf{r}}{\partial \xi} = \begin{pmatrix} \frac{\partial x}{\partial \xi} \\ \frac{\partial y}{\partial \xi} \end{pmatrix}, \mathbf{e}^\eta = \frac{\partial \mathbf{r}}{\partial \eta} = \begin{pmatrix} \frac{\partial x}{\partial \eta} \\ \frac{\partial y}{\partial \eta} \end{pmatrix} \quad (76)$$

Also, there is

$$d\zeta^2 = dx^2 + dy^2 = d\mathbf{r} \cdot d\mathbf{r} = \mathbf{e}^\xi \cdot \mathbf{e}^\xi d\xi^2 + \mathbf{e}^\eta \cdot \mathbf{e}^\eta d\eta^2 \quad (77)$$

In this case, the relationship of metrics go harmonized that g_{ii} equals to $1/g^{ii}$. It should be pointed out that the ways of expressions of bases are all equivalent except that the substitutions of coordinate lines ξ and η might have hidden some information of super surface, so that I would like to make analysis within super space in most cases.

Example 2: Bases of Riemannian manifold of equal dimension

As a Riemannian manifold has equal dimension with its contra variant space, it could be called equal dimension manifold. A plane space (u, v) maps to a plane space (x, y) could be taken for granted, as shown in Figure 7 and Figure 8.




Figure 7. Covariant space.




Figure 8. Contra variant space.

A differential vector in covariant space is

$$d\mathbf{r} = \begin{pmatrix} dx \\ dy \end{pmatrix} \quad (78)$$

The differential vector in contra variant space is

$$d\boldsymbol{\rho} = \begin{pmatrix} du \\ dv \end{pmatrix} \quad (79)$$

Thus, the definition of contra variant bases could be

$$\mathbf{e}^u = \frac{\partial \mathbf{r}}{\partial u}, \mathbf{e}^v = \frac{\partial \mathbf{r}}{\partial v} \quad (80)$$

To express $d\mathbf{r}$ with $d\boldsymbol{\rho}$

$$d\mathbf{r} = du\mathbf{e}^u + dv\mathbf{e}^v \quad (81)$$

Also, there is the definition of covariant bases

$$\mathbf{e}_x = \frac{\partial \boldsymbol{\rho}}{\partial x}, \mathbf{e}_y = \frac{\partial \boldsymbol{\rho}}{\partial y} \quad (82)$$

So that the expression of $d\boldsymbol{\rho}$ by $d\mathbf{r}$ should be

$$d\boldsymbol{\rho} = dx\mathbf{e}_x + dy\mathbf{e}_y \quad (83)$$

Something different is that a covariant base has 2 components

$$\mathbf{e}_x = \begin{pmatrix} \frac{\partial u}{\partial x} \\ \frac{\partial v}{\partial x} \end{pmatrix}, \mathbf{e}_y = \begin{pmatrix} \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial y} \end{pmatrix} \quad (84)$$

And a contra variant base also has 2 components

$$\mathbf{e}^u = \begin{pmatrix} \frac{\partial x}{\partial u} \\ \frac{\partial y}{\partial u} \end{pmatrix}, \mathbf{e}^v = \begin{pmatrix} \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial v} \end{pmatrix} \quad (85)$$

In the case that bases are orthogonal, the distance

$$ds^2 = du^2 + dv^2 = d\boldsymbol{\rho} \cdot d\boldsymbol{\rho} = \mathbf{e}_x \cdot \mathbf{e}_x dx^2 + \mathbf{e}_y \cdot \mathbf{e}_y dy^2 \quad (86)$$

and

$$d\zeta^2 = dx^2 + dy^2 = d\mathbf{r} \cdot d\mathbf{r} = \mathbf{e}^u \cdot \mathbf{e}^u du^2 + \mathbf{e}^v \cdot \mathbf{e}^v dv^2 \quad (87)$$

3.2.2. Inequalities of Mixed Derivatives of Bases

Now it is the time to carry out the first discussion on the inequality of mixed derivatives of bases. The mixed derivatives of bases are just special defined for bases alternative derivations. As transformation from contra variant space to covariant space is concerned, the covariant bases could be considered to be derivated by the coordinate lines in chain rule

$$\mathbf{e}_x = \frac{\partial \boldsymbol{\rho}}{\partial x} = \frac{\partial \boldsymbol{\rho}}{\partial \xi} \frac{\partial \xi}{\partial x}, \mathbf{e}_y = \frac{\partial \boldsymbol{\rho}}{\partial y} = \frac{\partial \boldsymbol{\rho}}{\partial \eta} \frac{\partial \eta}{\partial y} \quad (88)$$

where, $\frac{\partial \boldsymbol{\rho}}{\partial \xi}$ and $\frac{\partial \boldsymbol{\rho}}{\partial \eta}$ are the direction derivatives along the coordinate lines $\xi(x)$ and $\eta(y)$ in covariant space, and $d\xi$ and $d\eta$ are their differential lengths in covariant space, which could be called the covariant lengths. And there will be a setting that Einstein summation convention does not act on double $d\xi$ and double $d\eta$.

It should be pointed out that, in most mathematics and physics, mixed derivatives being confirmed to be equal is because in the Equation (88) dx and dy is incorrectly understood to be the differential length in covariant space (u, v, w) , but they are really the lengths in contra variant space (x, y, z) . That is the reason we have carried out the concept of covariant length $d\xi$ and $d\eta$.

Thus, the mixed derivatives will be

$$\frac{\partial \mathbf{e}_x}{\partial y} = \frac{\partial^2 \boldsymbol{\rho}}{\partial \xi \partial \eta} \frac{\partial \xi}{\partial x} \frac{\partial \eta}{\partial y} + \frac{\partial \boldsymbol{\rho}}{\partial \xi} \frac{\partial^2 \xi}{\partial x \partial y} \quad (89)$$

and

$$\frac{\partial \mathbf{e}_y}{\partial x} = \frac{\partial^2 \boldsymbol{\rho}}{\partial \eta \partial \xi} \frac{\partial \eta}{\partial y} \frac{\partial \xi}{\partial x} + \frac{\partial \boldsymbol{\rho}}{\partial \eta} \frac{\partial^2 \eta}{\partial y \partial x} \quad (90)$$

Consequently 3 conditions could be focused on:

Condition 1:

If there is an equality between the first items of the two equations that

$$\frac{\partial^2 \boldsymbol{\rho}}{\partial \xi \partial \eta} = \frac{\partial^2 \boldsymbol{\rho}}{\partial \eta \partial \xi} \quad (91)$$

For example, in the super surface, the mixed derivatives of course have the equality just as the equality of normal mixed derivatives of a 3-dimensional surface in a Euclidean space.

In this case and if there is another equality for the last items of the two equations that

$$\frac{\partial \boldsymbol{\rho}}{\partial \xi} \frac{\partial^2 \xi}{\partial x \partial y} = \frac{\partial \boldsymbol{\rho}}{\partial \eta} \frac{\partial^2 \eta}{\partial y \partial x} \quad (92)$$

Then that must come to the conclusion

$$\frac{\partial \mathbf{e}_x}{\partial x} = \frac{\partial \mathbf{e}_y}{\partial y} \quad (93)$$

Otherwise, that depends.

It should be pointed out that $\frac{\partial \boldsymbol{\rho}}{\partial \xi}$ and $\frac{\partial \boldsymbol{\rho}}{\partial \eta}$ do not equal in general conditions because they have different directions and, in most cases, they are usually set orthogonal, so that if that equality of Equation (92) happens, it asks for

$$\frac{\partial^2 \xi}{\partial x \partial y} = \frac{\partial^2 \eta}{\partial y \partial x} = 0 \quad (94)$$

We will see that in some cases it is really well satisfied.

Condition 2:

Most special if

$$\frac{\partial^2 \rho}{\partial \xi \partial \eta} \neq \frac{\partial^2 \rho}{\partial \eta \partial \xi} \quad (95)$$

that indicate the first items of the two equations are not equal, but at the same time if the total equations are still equal that

$$\frac{\partial^2 \rho}{\partial \xi \partial \eta} \frac{\partial \xi}{\partial x} \frac{\partial \eta}{\partial y} + \frac{\partial \rho}{\partial \xi} \frac{\partial^2 \xi}{\partial x \partial y} = \frac{\partial^2 \rho}{\partial \eta \partial \xi} \frac{\partial \eta}{\partial y} \frac{\partial \xi}{\partial x} + \frac{\partial \rho}{\partial \eta} \frac{\partial^2 \eta}{\partial y \partial x} \quad (96)$$

We will still obtain the equality that

$$\frac{\partial e_x}{\partial x} = \frac{\partial e_y}{\partial y} \quad (97)$$

Condition 3:

This is the condition after the previous two conditions and else to them. Generally if

$$\frac{\partial^2 \rho}{\partial \xi \partial \eta} \frac{\partial \xi}{\partial x} \frac{\partial \eta}{\partial y} + \frac{\partial \rho}{\partial \xi} \frac{\partial^2 \xi}{\partial x \partial y} \neq \frac{\partial^2 \rho}{\partial \eta \partial \xi} \frac{\partial \eta}{\partial y} \frac{\partial \xi}{\partial x} + \frac{\partial \rho}{\partial \eta} \frac{\partial^2 \eta}{\partial y \partial x} \quad (98)$$

No matter the first items of the two equations are equal or not, the mixed derivatives will perform inequality

$$\frac{\partial e_x}{\partial x} \neq \frac{\partial e_y}{\partial y} \quad (99)$$

Then, turn to the issue of geometrical influence that the inequality of mixed derivatives will cause closure errors [6,7]. I prefer to give a brief presentation. Consider a differential in a curve line coordinate system expressed by bases along different coordinate paths as shown in Figure 9 that




Figure 9. Bases vary in different paths.

in path 1,

$$(d\rho)_1 = \int_x^{x+dx} e_x dx + \int_y^{y+dy} (e_y + \frac{\partial e_y}{\partial x} dx) dy \quad (100)$$

The irregular expressions of same symbols of integral variables and integral range could be adopted in special cases.

By Taylor's approximation, it could be written as

$$(d\rho)_1 \approx e_x dx + \frac{1}{2} \frac{\partial e_x}{\partial x} dx^2 + e_y dy + \frac{\partial e_y}{\partial x} dx dy + \frac{1}{2} (\frac{\partial e_y}{\partial y} + \frac{\partial e_y}{\partial x \partial y} dx) dy^2 \quad (101)$$

We could also get the differential in path 2,

$$(d\rho)_2 \approx e_y dy + \frac{1}{2} \frac{\partial e_y}{\partial y} dy^2 + e_x dx + \frac{\partial e_x}{\partial y} dy dx + \frac{1}{2} (\frac{\partial e_x}{\partial x} + \frac{\partial e_x}{\partial y \partial x} dy) dx^2 \quad (102)$$

Trimming off the 3-order infinite small quantities, the difference of $(d\rho)_1$ and $(d\rho)_2$ is

$$(d\rho)_1 - (d\rho)_2 \approx (\frac{\partial e_y}{\partial x} - \frac{\partial e_x}{\partial y}) dy dx \quad (103)$$

There will be a closure error in close path if the mixed derivatives of bases do not equal.

3.2.3. Verifications and Discussions

Example 1: Polar coordinate system

Polar coordinate system that we are familiar with is a transformation from its contra variant space (r, θ) , as shown in Figure 10 and Figure 11.




Figure 10. Covariant space.




Figure 11. Contra variant space.

A position in contra variant space could be expressed by vector

$$\mathbf{r} = \mathbf{r}(r, \theta) \quad (104)$$

and the differential is

$$d\mathbf{r} = \begin{pmatrix} dr \\ d\theta \end{pmatrix} \quad (105)$$

Corresponding position in covariant space, will be expressed by

$$\boldsymbol{\rho} = \boldsymbol{\rho}(u, v) \quad (106)$$

and the differential is

$$d\boldsymbol{\rho} = \begin{pmatrix} du \\ dv \end{pmatrix} \quad (107)$$

In the contra variant space, the differential distance between two positions could be defined as

$$d\zeta^2 = d\mathbf{r} \cdot d\mathbf{r} = dr^2 + d\theta^2 \quad (108)$$

The system we have used to is the one that has experienced transformation from space (r, θ) with

$$u = r \cos \theta, v = r \sin \theta \quad (109)$$

The bases

$$\mathbf{e}_r = \frac{\partial \boldsymbol{\rho}}{\partial r} = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \quad (110)$$

$$\mathbf{e}_\theta = \frac{\partial \boldsymbol{\rho}}{\partial \theta} = \begin{pmatrix} -r \sin \theta \\ r \cos \theta \end{pmatrix} \quad (111)$$

so that

$$d\boldsymbol{\rho} = \mathbf{e}_r dr + \mathbf{e}_\theta d\theta \quad (112)$$

Thus, there is the covariant distance

$$ds^2 = du^2 + dv^2 = d\boldsymbol{\rho} \cdot d\boldsymbol{\rho} = \mathbf{e}_r \cdot \mathbf{e}_r dr^2 + \mathbf{e}_\theta \cdot \mathbf{e}_\theta d\theta^2 = dr^2 + r^2 d\theta^2 \quad (113)$$

The mixed derivatives of bases

$$\frac{\partial \mathbf{e}_r}{\partial \theta} = \frac{\partial}{\partial \theta} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \quad (114)$$

$$\frac{\partial \mathbf{e}_\theta}{\partial r} = \frac{\partial}{\partial r} \begin{pmatrix} -r \sin \theta \\ r \cos \theta \end{pmatrix} = \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \quad (115)$$

We have seen the mixed derivatives got equal

$$\frac{\partial \mathbf{e}_r}{\partial \theta} = \frac{\partial \mathbf{e}_\theta}{\partial r} \quad (116)$$

It could also be verified in Equation (89) and Equation (90) that

$$\frac{\partial \mathbf{e}_r}{\partial \theta} = \frac{\partial^2 \boldsymbol{\rho}}{\partial \xi_{(r)} \partial \eta_{(\theta)}} \frac{\partial \xi_{(r)}}{\partial r} \frac{\partial \eta_{(\theta)}}{\partial \theta} + \frac{\partial \boldsymbol{\rho}}{\partial \xi_{(r)}} \frac{\partial^2 \xi_{(r)}}{\partial r \partial \theta} \quad (117)$$

and

$$\frac{\partial \mathbf{e}_\theta}{\partial r} = \frac{\partial^2 \boldsymbol{\rho}}{\partial \eta_{(\theta)} \partial \xi_{(r)}} \frac{\partial \eta_{(\theta)}}{\partial \theta} \frac{\partial \xi_{(r)}}{\partial r} + \frac{\partial \boldsymbol{\rho}}{\partial \eta_{(\theta)}} \frac{\partial^2 \eta_{(\theta)}}{\partial \theta \partial r} \quad (118)$$

The vector $\boldsymbol{\rho}$ is

$$\boldsymbol{\rho} = \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} r \cos \theta \\ r \sin \theta \end{pmatrix} \quad (119)$$

Because $d\xi_{(r)}$ is radius length dr , and $d\eta_{(\theta)}$ is arc length $r d\theta$, then

$$\frac{\partial \boldsymbol{\rho}}{\partial \xi_{(r)}} = \frac{\partial}{\partial r} \begin{pmatrix} r \cos \theta \\ r \sin \theta \end{pmatrix} = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \quad (120)$$

$$\frac{\partial \boldsymbol{\rho}}{\partial \eta_{(\theta)}} = \frac{\partial}{\partial \theta} \begin{pmatrix} r \cos \theta \\ r \sin \theta \end{pmatrix} = \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \quad (121)$$

Thus, the first item of Equation (117) is

$$\frac{\partial^2 \boldsymbol{\rho}}{\partial \xi_{(r)} \partial \eta_{(\theta)}} \frac{\partial \xi_{(r)}}{\partial r} \frac{\partial \eta_{(\theta)}}{\partial \theta} = \frac{\partial}{\partial \theta} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \frac{\partial r}{\partial r} \frac{\partial \theta}{\partial \theta} = \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \quad (122)$$

The second item is

$$\frac{\partial \boldsymbol{\rho}}{\partial \xi_{(r)}} \frac{\partial^2 \xi_{(r)}}{\partial r \partial \theta} = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \frac{\partial}{\partial \theta} \frac{\partial r}{\partial r} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (123)$$

And the first item of Equation (118)

$$\frac{\partial^2 \boldsymbol{\rho}}{\partial \eta_{(\theta)} \partial \xi_{(r)}} \frac{\partial \eta_{(\theta)}}{\partial \theta} \frac{\partial \xi_{(r)}}{\partial r} = \frac{\partial}{\partial r} \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \frac{\partial \theta}{\partial \theta} \frac{\partial r}{\partial r} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (124)$$

The second item

$$\frac{\partial \boldsymbol{\rho}}{\partial \eta_{(\theta)}} \frac{\partial^2 \eta_{(\theta)}}{\partial \theta \partial r} = \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \frac{\partial}{\partial r} \frac{\partial \theta}{\partial \theta} = \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \quad (125)$$

so that

$$\frac{\partial \mathbf{e}_r}{\partial \theta} = \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \end{pmatrix} = \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \quad (126)$$

and

$$\frac{\partial \mathbf{e}_\theta}{\partial r} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} = \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \quad (127)$$

Obviously there is

$$\frac{\partial^2 \boldsymbol{\rho}}{\partial \xi_{(r)} \partial \eta_{(\theta)}} \neq \frac{\partial^2 \boldsymbol{\rho}}{\partial \eta_{(\theta)} \partial \xi_{(r)}} \quad (128)$$

and

$$\frac{\partial \boldsymbol{\rho}}{\partial \xi_{(r)}} \frac{\partial^2 \xi_{(r)}}{\partial r \partial \theta} \neq \frac{\partial \boldsymbol{\rho}}{\partial \eta_{(\theta)}} \frac{\partial^2 \eta_{(\theta)}}{\partial \theta \partial r} \quad (129)$$

but the mixed derivatives are equal totally that

$$\frac{\partial e_r}{\partial \theta} = \frac{\partial e_\theta}{\partial r} \quad (130)$$

Again, we have got the equality of mixed derivatives. But to our surprise is that this solution really subject to condition 2. It is said that the first items of Equation (117) and Equation (118) do not equal. One of the reasons in this case, is that there is no super surface.

Example 2: Spherical surface coordinate system

A spherical surface coordinate system is also a transformation of the corresponding contra variant space as in Figure 12 and Figure 13, in which




Figure 12. Covariant space.




Figure 13. Contra variant space.

$$\mathbf{r} = \mathbf{r}(r, \theta, \varphi)|_{r=R} = \begin{pmatrix} \theta \\ \varphi \end{pmatrix} \Big|_{r=R} \quad (131)$$

Differential distance is

$$d\zeta^2 = d\mathbf{r} \cdot d\mathbf{r} = d\theta^2 + d\varphi^2 \quad (132)$$

And the coordinates of covariant space will be expressed with

$$\boldsymbol{\rho} = \boldsymbol{\rho}(u, v, w) = \begin{pmatrix} u \\ v \\ w \end{pmatrix} \quad (133)$$

The spherical coordinates could be transformed to Cartesian coordinates,

$$u = R \sin \theta \cos \varphi, v = R \sin \theta \sin \varphi, w = R \cos \theta \quad (134)$$

The bases could be defined as

$$\mathbf{e}_\theta = \frac{\partial \boldsymbol{\rho}}{\partial \theta} = \begin{pmatrix} R \cos \theta \cos \varphi \\ R \cos \theta \sin \varphi \\ -R \sin \theta \end{pmatrix}, \mathbf{e}_\varphi = \frac{\partial \boldsymbol{\rho}}{\partial \varphi} = \begin{pmatrix} -R \sin \theta \sin \varphi \\ R \sin \theta \cos \varphi \\ 0 \end{pmatrix} \quad (135)$$

and

$$d\boldsymbol{\rho} = \mathbf{e}_\theta d\theta + \mathbf{e}_\varphi d\varphi \quad (136)$$

Thus, there is the covariant distance

$$ds^2 = d\boldsymbol{\rho} \cdot d\boldsymbol{\rho} = \mathbf{e}_\theta \cdot \mathbf{e}_\theta dr^2 + \mathbf{e}_\varphi \cdot \mathbf{e}_\varphi dr^2 = R^2 d\theta^2 + R^2 \sin^2 \theta d\varphi^2 \quad (137)$$

The derivatives

$$\frac{\partial \mathbf{e}_\theta}{\partial \varphi} = \begin{pmatrix} -R \cos \theta \sin \varphi \\ R \cos \theta \cos \varphi \\ 0 \end{pmatrix}, \frac{\partial \mathbf{e}_\varphi}{\partial \theta} = \begin{pmatrix} -R \cos \theta \sin \varphi \\ R \cos \theta \cos \varphi \\ 0 \end{pmatrix} \quad (138)$$

so that

$$\frac{\partial \mathbf{e}_\theta}{\partial \varphi} = \frac{\partial \mathbf{e}_\varphi}{\partial \theta} \quad (139)$$

It could also be verified in Equation (89) and Equation (90) that

$$\frac{\partial \mathbf{e}_\theta}{\partial \varphi} = \frac{\partial^2 \boldsymbol{\rho}}{\partial \eta_{(\theta)} \partial \zeta_{(\varphi)}} \frac{\partial \eta_{(\theta)}}{\partial \theta} \frac{\partial \zeta_{(\varphi)}}{\partial \varphi} + \frac{\partial \boldsymbol{\rho}}{\partial \eta_{(\theta)}} \frac{\partial^2 \eta_{(\theta)}}{\partial \theta \partial \varphi} \quad (140)$$

and

$$\frac{\partial \mathbf{e}_\varphi}{\partial \theta} = \frac{\partial^2 \boldsymbol{\rho}}{\partial \zeta_{(\varphi)} \partial \eta_{(\theta)}} \frac{\partial \zeta_{(\varphi)}}{\partial \varphi} \frac{\partial \eta_{(\theta)}}{\partial \theta} + \frac{\partial \boldsymbol{\rho}}{\partial \zeta_{(\varphi)}} \frac{\partial^2 \zeta_{(\varphi)}}{\partial \varphi \partial \theta} \quad (141)$$

The vector $\boldsymbol{\rho}$ is

$$\boldsymbol{\rho} = \begin{pmatrix} u \\ v \\ w \end{pmatrix} = \begin{pmatrix} R \sin \theta \cos \varphi \\ R \sin \theta \sin \varphi \\ R \cos \theta \end{pmatrix} \quad (142)$$

Because $d\eta_{(\theta)}$ is arc length $Rd\theta$, and $d\zeta_{(\varphi)}$ is arc length $Rd\varphi$, then

$$\frac{\partial \boldsymbol{\rho}}{\partial \eta_{(\theta)}} = \frac{\partial}{R \partial \theta} \begin{pmatrix} R \sin \theta \cos \varphi \\ R \sin \theta \sin \varphi \\ R \cos \theta \end{pmatrix} = \begin{pmatrix} \cos \theta \cos \varphi \\ \cos \theta \sin \varphi \\ -\sin \theta \end{pmatrix} \quad (143)$$

$$\frac{\partial \boldsymbol{\rho}}{\partial \zeta_{(\varphi)}} = \frac{\partial}{R \partial \varphi} \begin{pmatrix} R \sin \theta \cos \varphi \\ R \sin \theta \sin \varphi \\ R \cos \theta \end{pmatrix} = \begin{pmatrix} -\sin \theta \sin \varphi \\ \sin \theta \cos \varphi \\ 0 \end{pmatrix} \quad (144)$$

Thus, the first item of Equation (140) is

$$\frac{\partial^2 \boldsymbol{\rho}}{\partial \eta_{(\theta)} \partial \zeta_{(\varphi)}} \frac{\partial \eta_{(\theta)}}{\partial \theta} \frac{\partial \zeta_{(\varphi)}}{\partial \varphi} = \frac{\partial}{R \partial \theta} \begin{pmatrix} \cos \theta \cos \varphi \\ \cos \theta \sin \varphi \\ -\sin \theta \end{pmatrix} \frac{R \partial \theta}{\partial \theta} \frac{R \partial \varphi}{\partial \varphi} = \begin{pmatrix} -R \cos \theta \sin \varphi \\ R \cos \theta \cos \varphi \\ 0 \end{pmatrix} \quad (145)$$

The second item is

$$\frac{\partial \boldsymbol{\rho}}{\partial \eta_{(\theta)}} \frac{\partial^2 \eta_{(\theta)}}{\partial \theta \partial \varphi} = \begin{pmatrix} \cos \theta \cos \varphi \\ \cos \theta \sin \varphi \\ -\sin \theta \end{pmatrix} \frac{\partial}{\partial \varphi} \frac{R \partial \theta}{\partial \theta} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \quad (146)$$

And the first item of Equation (141)

$$\frac{\partial^2 \boldsymbol{\rho}}{\partial \zeta_{(\varphi)} \partial \eta_{(\theta)}} \frac{\partial \zeta_{(\varphi)}}{\partial \varphi} \frac{\partial \eta_{(\theta)}}{\partial \theta} = \frac{\partial}{R \partial \theta} \begin{pmatrix} -\sin \theta \sin \varphi \\ \sin \theta \cos \varphi \\ 0 \end{pmatrix} \frac{R \partial \varphi}{\partial \varphi} \frac{R \partial \theta}{\partial \theta} = \begin{pmatrix} -R \cos \theta \sin \varphi \\ R \cos \theta \cos \varphi \\ 0 \end{pmatrix} \quad (147)$$

The second item

$$\frac{\partial \boldsymbol{\rho}}{\partial \zeta_{(\varphi)}} \frac{\partial^2 \zeta_{(\varphi)}}{\partial \varphi \partial \theta} = \begin{pmatrix} -\sin \theta \sin \varphi \\ \sin \theta \cos \varphi \\ 0 \end{pmatrix} \frac{\partial}{\partial \theta} \frac{R \partial \varphi}{\partial \varphi} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \quad (148)$$

With Equation (145) to Equation (148) we found that

$$\frac{\partial^2 \boldsymbol{\rho}}{\partial \eta_{(\theta)} \partial \zeta_{(\varphi)}} = \frac{\partial^2 \boldsymbol{\rho}}{\partial \zeta_{(\varphi)} \partial \eta_{(\theta)}} \quad (149)$$

and

$$\frac{\partial \boldsymbol{\rho}}{\partial \eta_{(\theta)}} \frac{\partial^2 \eta_{(\theta)}}{\partial \theta \partial \varphi} = \frac{\partial \boldsymbol{\rho}}{\partial \zeta_{(\varphi)}} \frac{\partial^2 \zeta_{(\varphi)}}{\partial \varphi \partial \theta} \quad (150)$$

so that there is

$$\frac{\partial \mathbf{e}_\theta}{\partial \varphi} = \frac{\partial \mathbf{e}_\varphi}{\partial \theta} \quad (151)$$

One can see that this is of condition 1.

Example 3: Spherical coordinate system

A spherical coordinate system is also a transformation of the corresponding contra variant space as in Figure 14 and Figure 15, in which




Figure 14. Covariant space.




Figure 15. Contra variant space.

$$\mathbf{r} = \mathbf{r}(r, \theta, \varphi) = \begin{pmatrix} r \\ \theta \\ \varphi \end{pmatrix} \quad (152)$$

Differential distance is

$$d\zeta^2 = d\mathbf{r} \cdot d\mathbf{r} = dr^2 + d\theta^2 + d\varphi^2 \quad (153)$$

And the coordinates of covariant space will be expressed with

$$\boldsymbol{\rho} = \boldsymbol{\rho}(u, v, w) = \begin{pmatrix} u \\ v \\ w \end{pmatrix} \quad (154)$$

The spherical coordinates could be transformed to Cartesian coordinates,

$$u = r \sin \theta \cos \varphi, v = r \sin \theta \sin \varphi, w = r \cos \theta \quad (155)$$

The bases could be defined as

$$\mathbf{e}_r = \frac{\partial \boldsymbol{\rho}}{\partial r} = \begin{pmatrix} \sin \theta \cos \varphi \\ \sin \theta \sin \varphi \\ \cos \theta \end{pmatrix}, \mathbf{e}_\theta = \frac{\partial \boldsymbol{\rho}}{\partial \theta} = \begin{pmatrix} r \cos \theta \cos \varphi \\ r \cos \theta \sin \varphi \\ -r \sin \theta \end{pmatrix}, \mathbf{e}_\varphi = \frac{\partial \boldsymbol{\rho}}{\partial \varphi} = \begin{pmatrix} -r \sin \theta \sin \varphi \\ r \sin \theta \cos \varphi \\ 0 \end{pmatrix} \quad (156)$$

and

$$d\boldsymbol{\rho} = \mathbf{e}_r dr + \mathbf{e}_\theta d\theta + \mathbf{e}_\varphi d\varphi \quad (157)$$

Thus, there is the covariant distance

$$ds^2 = d\boldsymbol{\rho} \cdot d\boldsymbol{\rho} = \mathbf{e}_r \cdot \mathbf{e}_r dr^2 + \mathbf{e}_\theta \cdot \mathbf{e}_\theta d\theta^2 + \mathbf{e}_\varphi \cdot \mathbf{e}_\varphi d\varphi^2 = dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2 \quad (158)$$

The derivatives

$$\begin{aligned} \frac{\partial \mathbf{e}_r}{\partial \theta} &= \begin{pmatrix} \cos \theta \cos \varphi \\ \cos \theta \sin \varphi \\ -\sin \theta \end{pmatrix}, \frac{\partial \mathbf{e}_\theta}{\partial r} = \begin{pmatrix} \cos \theta \cos \varphi \\ \cos \theta \sin \varphi \\ -\sin \theta \end{pmatrix}, \frac{\partial \mathbf{e}_\theta}{\partial \varphi} = \begin{pmatrix} -r \cos \theta \sin \varphi \\ r \cos \theta \cos \varphi \\ 0 \end{pmatrix} \\ \frac{\partial \mathbf{e}_\varphi}{\partial \theta} &= \begin{pmatrix} -r \cos \theta \sin \varphi \\ r \cos \theta \cos \varphi \\ 0 \end{pmatrix}, \frac{\partial \mathbf{e}_\varphi}{\partial r} = \begin{pmatrix} -\sin \theta \sin \varphi \\ \sin \theta \cos \varphi \\ 0 \end{pmatrix}, \frac{\partial \mathbf{e}_r}{\partial \varphi} = \begin{pmatrix} -\sin \theta \sin \varphi \\ \sin \theta \cos \varphi \\ 0 \end{pmatrix} \end{aligned} \quad (159)$$

so that

$$\frac{\partial \mathbf{e}_r}{\partial \theta} = \frac{\partial \mathbf{e}_\theta}{\partial r}, \frac{\partial \mathbf{e}_\theta}{\partial \varphi} = \frac{\partial \mathbf{e}_\varphi}{\partial \theta}, \frac{\partial \mathbf{e}_\varphi}{\partial r} = \frac{\partial \mathbf{e}_r}{\partial \varphi} \quad (160)$$

It could also be verified in Equation (89) and Equation (90) that

$$\frac{\partial \mathbf{e}_r}{\partial \theta} = \frac{\partial^2 \boldsymbol{\rho}}{\partial \xi_{(r)} \partial \eta_{(\theta)}} \frac{\partial \xi_{(r)}}{\partial r} \frac{\partial \eta_{(\theta)}}{\partial \theta} + \frac{\partial \boldsymbol{\rho}}{\partial \xi_{(r)}} \frac{\partial^2 \xi_{(r)}}{\partial r \partial \theta} \quad (161)$$

and

$$\frac{\partial \mathbf{e}_\theta}{\partial r} = \frac{\partial^2 \boldsymbol{\rho}}{\partial \eta_{(\theta)} \partial \xi_{(r)}} \frac{\partial \eta_{(\theta)}}{\partial \theta} \frac{\partial \xi_{(r)}}{\partial r} + \frac{\partial \boldsymbol{\rho}}{\partial \eta_{(\theta)}} \frac{\partial^2 \eta_{(\theta)}}{\partial \theta \partial r} \quad (162)$$

The vector $\boldsymbol{\rho}$ is

$$\boldsymbol{\rho} = \begin{pmatrix} u \\ v \\ w \end{pmatrix} = \begin{pmatrix} r \sin \theta \cos \varphi \\ r \sin \theta \sin \varphi \\ r \cos \theta \end{pmatrix} \quad (163)$$

Because $d\xi_{(r)}$ is radius length dr , $d\eta_{(\theta)}$ is arc length $r d\theta$, and $d\zeta_{(\varphi)}$ is arc length $r d\varphi$, then

$$\frac{\partial \boldsymbol{\rho}}{\partial \xi_{(r)}} = \frac{\partial}{\partial r} \begin{pmatrix} r \sin \theta \cos \varphi \\ r \sin \theta \sin \varphi \\ r \cos \theta \end{pmatrix} = \begin{pmatrix} \sin \theta \cos \varphi \\ \sin \theta \sin \varphi \\ \cos \theta \end{pmatrix} \quad (164)$$

$$\frac{\partial \boldsymbol{\rho}}{\partial \eta_{(\theta)}} = \frac{\partial}{r \partial \theta} \begin{pmatrix} r \sin \theta \cos \varphi \\ r \sin \theta \sin \varphi \\ r \cos \theta \end{pmatrix} = \begin{pmatrix} \cos \theta \cos \varphi \\ \cos \theta \sin \varphi \\ -\sin \theta \end{pmatrix} \quad (165)$$

$$\frac{\partial \boldsymbol{\rho}}{\partial \zeta_{(\varphi)}} = \frac{\partial}{r \partial \varphi} \begin{pmatrix} r \sin \theta \cos \varphi \\ r \sin \theta \sin \varphi \\ r \cos \theta \end{pmatrix} = \begin{pmatrix} -\sin \theta \sin \varphi \\ \sin \theta \cos \varphi \\ 0 \end{pmatrix} \quad (166)$$

Thus, the first item of Equation (161) is

$$\frac{\partial^2 \boldsymbol{\rho}}{\partial \xi_{(r)} \partial \eta_{(\theta)}} \frac{\partial \xi_{(r)}}{\partial r} \frac{\partial \eta_{(\theta)}}{\partial \theta} = \frac{\partial}{r \partial \theta} \begin{pmatrix} \sin \theta \cos \varphi \\ \sin \theta \sin \varphi \\ \cos \theta \end{pmatrix} \frac{\partial r}{\partial r} \frac{\partial \theta}{\partial \theta} = \begin{pmatrix} \cos \theta \cos \varphi \\ \cos \theta \sin \varphi \\ -\sin \theta \end{pmatrix} \quad (167)$$

The second item is

$$\frac{\partial \boldsymbol{\rho}}{\partial \xi_{(r)}} \frac{\partial^2 \xi_{(r)}}{\partial r \partial \theta} = \begin{pmatrix} \sin \theta \cos \varphi \\ \sin \theta \sin \varphi \\ \cos \theta \end{pmatrix} \frac{\partial}{\partial \theta} \frac{\partial r}{\partial r} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \quad (168)$$

And the first item of Equation (162)

$$\frac{\partial^2 \boldsymbol{\rho}}{\partial \eta_{(\theta)} \partial \xi_{(r)}} \frac{\partial \eta_{(\theta)}}{\partial \theta} \frac{\partial \xi_{(r)}}{\partial r} = \frac{\partial}{\partial r} \begin{pmatrix} \cos \theta \cos \varphi \\ \cos \theta \sin \varphi \\ -\sin \theta \end{pmatrix} \frac{r \partial \theta}{\partial \theta} \frac{\partial r}{\partial r} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \quad (169)$$

The second item

$$\frac{\partial \boldsymbol{\rho}}{\partial \eta_{(\theta)}} \frac{\partial^2 \eta_{(\theta)}}{\partial \theta \partial r} = \begin{pmatrix} \cos \theta \cos \varphi \\ \cos \theta \sin \varphi \\ -\sin \theta \end{pmatrix} \frac{\partial}{\partial r} \frac{r \partial \theta}{\partial \theta} = \begin{pmatrix} \cos \theta \cos \varphi \\ \cos \theta \sin \varphi \\ -\sin \theta \end{pmatrix} \quad (170)$$

With Equation (167) and Equation (170) we found that

$$\frac{\partial^2 \boldsymbol{\rho}}{\partial \xi_{(r)} \partial \eta_{(\theta)}} \neq \frac{\partial^2 \boldsymbol{\rho}}{\partial \eta_{(\theta)} \partial \xi_{(r)}} \quad (171)$$

but there is

$$\frac{\partial \mathbf{e}_r}{\partial \theta} = \frac{\partial \mathbf{e}_\theta}{\partial r} \quad (172)$$

One can also calculate that

$$\frac{\partial^2 \boldsymbol{\rho}}{\partial \eta_{(\theta)} \partial \zeta_{(\varphi)}} = \frac{\partial^2 \boldsymbol{\rho}}{\partial \zeta_{(\varphi)} \partial \eta_{(\theta)}}, \quad \frac{\partial^2 \boldsymbol{\rho}}{\partial \zeta_{(\varphi)} \partial \xi_{(r)}} \neq \frac{\partial^2 \boldsymbol{\rho}}{\partial \xi_{(r)} \partial \zeta_{(\varphi)}} \quad (173)$$

At the end we can obtain

$$\frac{\partial \mathbf{e}_\theta}{\partial \varphi} = \frac{\partial \mathbf{e}_\varphi}{\partial \theta}, \quad \frac{\partial \mathbf{e}_\varphi}{\partial r} = \frac{\partial \mathbf{e}_r}{\partial \varphi} \quad (174)$$

One can see that one of them is of condition 1 and the others of them are of condition 2.

Additional discussion: deformed bases of example 3

If one of the bases in example 3 be set deformed as

$$\mathbf{e}_r = f(r) \begin{pmatrix} \sin\theta\cos\varphi \\ \sin\theta\sin\varphi \\ \cos\theta \end{pmatrix} \quad (175)$$

where, $f(r) \neq \text{const.}$ is a function of coordinate r .

One will find that the derivatives

$$\frac{\partial \mathbf{e}_r}{\partial \theta} = f(r) \begin{pmatrix} \cos\theta\cos\varphi \\ \cos\theta\sin\varphi \\ -\sin\theta \end{pmatrix} + \frac{\partial f(r)}{\partial \theta} \begin{pmatrix} \sin\theta\cos\varphi \\ \sin\theta\sin\varphi \\ \cos\theta \end{pmatrix} = f(r) \begin{pmatrix} \cos\theta\cos\varphi \\ \cos\theta\sin\varphi \\ -\sin\theta \end{pmatrix} \quad (176)$$

and

$$\frac{\partial \mathbf{e}_r}{\partial \varphi} = f(r) \begin{pmatrix} -\sin\theta\sin\varphi \\ \sin\theta\cos\varphi \\ 0 \end{pmatrix} + \frac{\partial f(r)}{\partial \varphi} \begin{pmatrix} \sin\theta\cos\varphi \\ \sin\theta\sin\varphi \\ \cos\theta \end{pmatrix} = f(r) \begin{pmatrix} -\sin\theta\sin\varphi \\ \sin\theta\cos\varphi \\ 0 \end{pmatrix} \quad (177)$$

while $\frac{\partial \mathbf{e}_\theta}{\partial r}$ and $\frac{\partial \mathbf{e}_\varphi}{\partial r}$ will still keep the results as in Equation (159), that will cause

$$\frac{\partial \mathbf{e}_r}{\partial \theta} \neq \frac{\partial \mathbf{e}_\theta}{\partial r}, \frac{\partial \mathbf{e}_\varphi}{\partial r} \neq \frac{\partial \mathbf{e}_r}{\partial \varphi} \quad (178)$$

This result reminds us that the same performance would have happened in gravitational space time that will be put into discussions in next section.

4. Metrics and Covariant Derivatives in Space Time

4.1. Metrics in Pseudo Riemannian Space

Pseudo Riemannian space is raised for the description of space time for general relativity, after Minkowski space for special relativity [8,9]. Invariant distance for flat space time could be written as

$$ds^2 = -c^2 dt^2 + dx^2 + dy^2 + dz^2 \quad (179)$$

We know that there is a minus signal in the equation. The minus signal does not come from transformations of spaces or coordinates. It is a kind of mathematical and physical setting. Former researchers have made efforts on this topic, for example, the concept of plural employed to reform the base $\mathbf{e}_0$ [6]. But plural bases for relativity is not a good idea. Another treatment is to define $x^0 = ict$, which looks like more reasonable [7]. The more important is that the metrics for this condition should be carefully treated so that the metric g_{00} will not be minus.

In general relativity, spherical coordinates are usually suggested for one source problem. So that there will be contra variant space $(cdt, dr, d\theta, d\varphi)$ and covariant space $(cd\tau, d\rho, d\theta, d\varphi)$. The invariant distance of Schwarzschild solution is

$$ds^2 = d\boldsymbol{\rho} \cdot d\boldsymbol{\rho} = -(1 - \frac{r^*}{r})(cdt)^2 + (1 - \frac{r^*}{r})^{-1}dr^2 + r^2 d\theta^2 + r^2 \sin^2\theta d\varphi^2 \quad (180)$$

Most of publications have set the metric [3,4,10–12] to be

$$g_{00} = -(1 - \frac{r^*}{r}) \quad (181)$$

So that there could be a brief expression of invariant distance

$$ds^2 = g_{ij} dx^i dx^j \quad (182)$$

As we have discussed, that will cause plural bases. Anyway, minus metrics is improper. In fact, it is one of the reasons that cause the wrong result of acceleration calculation in Equation (44).

Considering the above discussions, I prefer to give the invariant distance of one source field as

$$ds^2 = -g_{00}(cdt)^2 + g_{11}dr^2 + g_{22}d\theta^2 + g_{33}d\varphi^2 \quad (183)$$

Thus, we will prevent from plural items. That doesn't hurt the expressions of relativity. Metrics and bases defined are just employed for the transformation from Minkowski space time to pseudo Riemannian space time.

It should be highlighted that the bases still would have sophisticated forms that

$$\mathbf{e}_0 = \begin{pmatrix} 0 \\ (1 - \frac{r^*}{r})^{1/2} \\ 0 \\ 0 \end{pmatrix}, \mathbf{e}_1 = \begin{pmatrix} 0 \\ \sin\theta\cos\varphi(1 - \frac{r^*}{r})^{-1/2} \\ \sin\theta\sin\varphi(1 - \frac{r^*}{r})^{-1/2} \\ \cos\theta(1 - \frac{r^*}{r})^{-1/2} \end{pmatrix}, \mathbf{e}_2 = \begin{pmatrix} 0 \\ r\cos\theta\cos\varphi \\ r\cos\theta\sin\varphi \\ -r\sin\theta \end{pmatrix}, \mathbf{e}_3 = \begin{pmatrix} 0 \\ -r\sin\theta\sin\varphi \\ r\sin\theta\cos\varphi \\ 0 \end{pmatrix} \quad (184)$$

That is because the spherical space we have discussed has already experienced coordinates transformations. The transformed coordinates are not real spherical coordinates, they are as the Cartesian coordinates (u, v, w) expressed by parameters of r, θ, φ , which could be called parameterized Cartesian coordinates. One can learn from previous section for the reasons.

It could be calculated that

$$\begin{aligned} \frac{\partial \mathbf{e}_1}{\partial t} &= \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \frac{\partial \mathbf{e}_1}{\partial \theta} = \begin{pmatrix} 0 \\ \cos\theta\cos\varphi(1 - \frac{r^*}{r})^{-1/2} \\ \cos\theta\sin\varphi(1 - \frac{r^*}{r})^{-1/2} \\ -\sin\theta(1 - \frac{r^*}{r})^{-1/2} \end{pmatrix}, \frac{\partial \mathbf{e}_1}{\partial \varphi} = \begin{pmatrix} 0 \\ -\sin\theta\sin\varphi(1 - \frac{r^*}{r})^{-1/2} \\ \sin\theta\cos\varphi(1 - \frac{r^*}{r})^{-1/2} \\ 0 \end{pmatrix} \\ \frac{\partial \mathbf{e}_0}{\partial r} &= \begin{pmatrix} \frac{r^*}{2r^2}(1 - \frac{r^*}{r})^{-1/2} \\ 0 \\ 0 \\ 0 \end{pmatrix}, \frac{\partial \mathbf{e}_2}{\partial r} = \begin{pmatrix} 0 \\ \cos\theta\cos\varphi \\ \cos\theta\sin\varphi \\ -\sin\theta \end{pmatrix}, \frac{\partial \mathbf{e}_3}{\partial r} = \begin{pmatrix} 0 \\ -\sin\theta\sin\varphi \\ \sin\theta\cos\varphi \\ 0 \end{pmatrix} \end{aligned} \quad (185)$$

there are

$$\frac{\partial \mathbf{e}_1}{\partial t} \neq \frac{\partial \mathbf{e}_0}{\partial r}, \frac{\partial \mathbf{e}_1}{\partial \theta} \neq \frac{\partial \mathbf{e}_2}{\partial r}, \frac{\partial \mathbf{e}_1}{\partial \varphi} \neq \frac{\partial \mathbf{e}_3}{\partial r} \quad (186)$$

In general relativity, it could be suggested to simplify the presentation that we only focus on the gravitational transformation. It is a transformation from contra variant tangent space $(cdt, dr, rd\theta, r\sin\theta d\varphi)$ to covariant tangent space $(cd\tau, d\rho, rd\theta, r\sin\theta d\varphi)$. So that the contra variant space is no longer of real spherical coordinates, which could be set as parameterized Cartesian coordinates, and the covariant space could also be the form of parameterized Cartesian. Thus, the invariant distance could be expressed as

$$ds^2 = -g_{00}(cdt)^2 + g_{11}dr^2 + g_{22}r^2d\theta^2 + g_{33}r^2\sin^2\theta d\varphi^2 \quad (187)$$

In which, $g_{22} = g_{33} = 1$.

These metrics could be called the gravitational metrics, while the metrics previous could be called total metrics and the metrics before gravitational transformation could be called original metrics or pseudo spherical transformation metrics.

For gravitational metrics of Schwarzschild solution, we have the bases

$$\mathbf{e}_0 = \begin{pmatrix} 0 \\ (1 - \frac{r^*}{r})^{1/2} \\ 0 \\ 0 \end{pmatrix}, \mathbf{e}_1 = \begin{pmatrix} 0 \\ (1 - \frac{r^*}{r})^{-1/2} \\ 0 \\ 0 \end{pmatrix}, \mathbf{e}_2 = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \mathbf{e}_3 = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} \quad (188)$$

It could be calculated that

$$\frac{\partial e_0}{\partial r} = \begin{pmatrix} \frac{r^*}{2r^2} (1 - \frac{r^*}{r})^{-1/2} \\ 0 \\ 0 \\ 0 \end{pmatrix}, \frac{\partial e_1}{\partial t} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad (189)$$

so that

$$\frac{\partial e_0}{\partial r} \neq \frac{\partial e_1}{\partial t} \quad (190)$$

This is of the condition 3 that we have discussed in Section 3.2.2. One can also calculate the inequality of mixed derivatives of bases of total metrics.

That reminds us that the inequality of mixed derivatives of bases will cause closure errors in space time, which would be left for more discussions elsewhere.

4.2. Discussions on Bases, Tensors, and Their Derivatives

Tensors could be recognized as the quantities relating to coordinates in space time. Case a tensor varies in space time, the variation ratio could be inspected by derivation. The simplest tensor is position vector $\rho(x_0, x_1, x_2, x_3,)$. You have seen that we are going to use middle subscriptions to express coordinates and tensors in covariant space time, though they are rarely mentioned in most of references. To study its variation in space time, one could define the distance variation ratio to form the bases

$$e_i = \frac{\partial \rho}{\partial x^i} \quad (191)$$

For any tensors involved with bases, such as a proper tensor

$$A = A^i e_i \quad (192)$$

where, A^i is the number i component of the total contra variant tensor.

The derivatives of the tensor are

$$\frac{\partial A}{\partial x^j} = e_i \frac{\partial A^i}{\partial x^j} + A^i \frac{\partial e_i}{\partial x^j} \quad (193)$$

The differential of a proper tensor dA could be defined to be covariant differential labeled as DA , and then the derivative $\frac{\partial A}{\partial x^j}$ to be covariant derivative $\frac{DA}{\partial x^j}$

$$\frac{DA}{\partial x^j} = \frac{\partial A}{\partial x^j} = e_i \frac{\partial A^i}{\partial x^j} + A^i \frac{\partial e_i}{\partial x^j} \quad (194)$$

In these equations, the middle subscriptions of proper tensors maybe neglected conventionally so that it is expressed as A . And the tensor A could be called proper tensor because A_i has already been named covariant tensor conventionally. Because A^i or A_i is just a component, we could imagine that there must be the total quantity. That will be expressed to be $A^\dagger = (A^0, A^1, A^2, A^3)$ or $A_\downarrow = (A_0, A_1, A_2, A_3)$ for convenience. Sometimes, row forms of expressions of vectors are employed to interpret components which are equivalent to the column forms.

Bases sometimes look like one-order tensors since they have a single index in expressions. But in fact, they really are two-order mixed tensors. For example, a component of contra variant base of collinear transformation, i.e. coordinate lines of contra variant space and covariant space coinciding, could be written as

$$e^\mu \nu = \frac{\partial x^\mu}{\partial x^\nu} \quad (195)$$

where the contra variant base and proper coordinate differential are all labeled with middle index ν .

So that the base really is

$$\mathbf{e}^\mu = \begin{pmatrix} e^\mu_0 \\ e^\mu_1 \\ e^\mu_2 \\ e^\mu_3 \end{pmatrix} \quad (196)$$

If $e^\mu_\nu=0$ case $\mu \neq \nu$, we could use e^μ instead of e^μ_μ for convenience, as it is the only nonvanishing component.

For any one order tensors there is a transformation

$$A^\mu = \mathbf{e}^\mu \cdot \mathbf{A} \text{ or } A_\mu = \mathbf{e}_\mu \cdot \mathbf{A} \quad (197)$$

But for two order tensors, there are some things different. For example, a component of contra variant velocity could be transformed from proper velocity

$$V^\mu = \frac{dx^\mu}{d\tau} = \mathbf{e}^\mu \cdot \frac{d\mathbf{p}}{d\tau} = \mathbf{e}^\mu \cdot \mathbf{V} \quad (198)$$

We know that it may be seen as one-order tensor in practice, but it is really two order tensor.

As for whole contra variant velocity as $V^1_0 = \frac{dx^1}{dt}$, that would not be worked out by direct vector product. In fact, it could be composed independently by dx^1 and dt .

$$V^1_0 = \frac{dx^1}{dt} = \frac{e^1 \cdot d\mathbf{p}}{e^0 \cdot d\mathbf{p}} = \frac{e^1 dx^1}{e^0 d\tau} = e^1 e_0 \frac{dx^1}{d\tau} \quad (199)$$

It is impossible to get $e^1 e_0$ from $\mathbf{e}^1 \cdot \mathbf{e}_0$ and the latter is zero. Notwithstanding, a velocity is a derivative on matter's trajectory rather than a direct derivative. That will be further discussed in next sections.

4.3. Derivation via Christoffel Symbols

Christoffel symbols were put forward to perform geometrical relationship that takes similar effects with that of bases. They are defined in the equations [3,13]

$$\frac{\partial \mathbf{e}^\mu}{\partial x^\lambda} = -\Gamma_{\lambda\nu}^\mu \mathbf{e}^\nu \text{ or } \frac{\partial \mathbf{e}_\mu}{\partial x^\lambda} = \Gamma_{\lambda\mu}^\nu \mathbf{e}_\nu \quad (200)$$

Taking the first one for example, the purpose of the equation is to consider the derivatives to be a function of bases, so that the right hand item is really a kind of trivial types. In the summation items, $\mathbf{e}^\nu$ just act as direction indicators that would give out whole basic vectors of entire dimensions. And then, $\Gamma_{\lambda\nu}^\mu$ provide the coefficients of all directions. It is said, this definition has just provided an error-free frame for the functions of derivatives. It means there may be redundant designs for the coefficients.

Since there is the probability of inequality of mixed derivatives of bases, we should define a specific sequence for subscripts of Christoffel symbols. For the traditional reasons, $\Gamma_{\lambda\nu}^\mu$ will be defined as the coefficient of a derivative of $\mathbf{e}^\mu$ that is derivated by x^λ , on a direction of $\mathbf{e}^\nu$, that requires unexchangeable subscripts of $\Gamma_{\lambda\nu}^\mu$.

We know that a covariant differential is exactly a differential of a proper tensor

$$\frac{D\mathbf{A}}{dx^\lambda} = \frac{\partial \mathbf{A}}{\partial x^\lambda} \quad (201)$$

This highlightable concept is essentially carried out to perform general covariance.

The contra variant form also performs the same covariance as that

$$\frac{DA^\mu}{\partial x^\lambda} = \mathbf{e}^\mu \cdot \frac{D\mathbf{A}}{\partial x^\lambda} \quad (202)$$

You might have found that the tensor component that has been expressed by a total tensor is exactly partial expression rather than a whole expression. It is just of traditional operations. One can of course carry out whole form expression of $\mathbf{A}^\dagger$ expressed by $\mathbf{A}$ with base matrix $[\mathbf{e}^\dagger]$. But too more renovations in a performance will bring about more reading difficulties. So I prefer to present equations in traditional forms as far as possible.

A component of contra variant tensor transformed from covariant one

$$A^\mu = \mathbf{e}^\mu \cdot \mathbf{A} \quad (203)$$

Its derivatives is

$$\frac{\partial A^\mu}{\partial x^\nu} = \frac{\partial}{\partial x^\nu} (\mathbf{e}^\mu \cdot \mathbf{A}) = \frac{\partial \mathbf{e}^\mu}{\partial x^\nu} \cdot \mathbf{A} + \mathbf{e}^\mu \cdot \frac{\partial \mathbf{A}}{\partial x^\nu} = \frac{\partial \mathbf{e}^\mu}{\partial x^\nu} \cdot \mathbf{A} + \frac{DA^\mu}{\partial x^\nu} \quad (204)$$

so that

$$\frac{DA^\mu}{\partial x^\nu} = \frac{\partial A^\mu}{\partial x^\nu} - \frac{\partial \mathbf{e}^\mu}{\partial x^\nu} \cdot \mathbf{A} = \frac{\partial A^\mu}{\partial x^\nu} + \Gamma_{\nu\lambda}^\mu A^\lambda \quad (205)$$

It is easy to study those covariant derivatives for covariant tensors

$$\frac{DA_\mu}{\partial x^\nu} = \frac{\partial A_\mu}{\partial x^\nu} - \frac{\partial \mathbf{e}_\mu}{\partial x^\nu} \cdot \mathbf{A} = \frac{\partial A_\mu}{\partial x^\nu} - \Gamma_{\nu\mu}^\lambda A_\lambda \quad (206)$$

We have seen that the methodologies of Christoffel symbols and the derivation directly from bases are actually equivalent treatments that present the covariant derivatives. That of course may be use to inspect the problems of equations of Christoffel symbols. Since we have known that part of Christoffel symbols with mixed subscripts do not equal in space time, it is necessary to do more discussions.

It is convenient to discuss the case that a space is defined by orthogonal bases. In practice, metrics are usually taken in to Christoffel connections analysis. For a series of bases $\mathbf{e}_0, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$, there is the metric

$$g_{ij} = \mathbf{e}_i \cdot \mathbf{e}_j \quad (207)$$

For whole orthogonal coordinate spaces,

$$\begin{cases} g_{ij} = 0, & i \neq j \\ g_{ij} = \mathbf{e}_i \cdot \mathbf{e}_i \neq 0, & i = j \end{cases} \quad (208)$$

The nonvanishing derivatives

$$\frac{\partial g_{ij}}{\partial x^\lambda} = \frac{\partial \mathbf{e}_i}{\partial x^\lambda} \cdot \mathbf{e}_j + \frac{\partial \mathbf{e}_j}{\partial x^\lambda} \cdot \mathbf{e}_i = 2 \frac{\partial \mathbf{e}_i}{\partial x^\lambda} \cdot \mathbf{e}_j, \quad i = j \quad (209)$$

Then the equation could have nonvanishing value, that

$$\frac{\partial g_{ij}}{\partial x^\lambda} = 2\Gamma_{\lambda i}^k \mathbf{e}_k \cdot \mathbf{e}_j = 2\Gamma_{\lambda i}^k g_{kj}, \quad i = j \quad (210)$$

Then

$$\Gamma_{\lambda i}^k = \frac{1}{2} g^{kj} \frac{\partial g_{ij}}{\partial x^\lambda}, \quad i = j \quad (211)$$

This equation could be called revised equation for Christoffel symbols in general relativity. In the equation, it also need $k = i = j$ for the $\Gamma_{\lambda i}^k$ to get nonvanishing.

For a result, this equation could be verified in a covariant derivative directly as that in Equation (206) that

$$\Gamma_{\lambda i}^k A_k = \frac{1}{2} g^{kj} \frac{\partial g_{ij}}{\partial x^\lambda} A_k = \frac{1}{2} (\mathbf{e}^k \cdot \mathbf{e}^j) (2 \frac{\partial \mathbf{e}_i}{\partial x^\lambda} \cdot \mathbf{e}_j) A_k = A_k \mathbf{e}^k \cdot \frac{\partial \mathbf{e}_i}{\partial x^\lambda} = \frac{\partial \mathbf{e}_i}{\partial x^\lambda} \cdot \mathbf{A} \quad (212)$$

It should be pointed out that the Christoffel symbols are not necessary because that the issue only started from derivatives of bases, as consequences they surely might be taken the place by the operations of bases.

4.4. Derivatives on Matter's Trajectory

The calculation of time derivatives may cause to another mathematical abuse in classical theory. For a one source field, it could be seen as rest field in general. Thus, a time derivative of a field quantity should be zero. Case a matter moves in a space, it is the issue that the matter changes its position in a time interval and forms a motion trajectory. In this condition, to learn the acceleration is to study the position variation rather than the field variation. It is one of the reasons that make errors

in covariant derivative calculation in Equation (43), in that the direct derivative has been used instead of trajectory derivative.

It is valuable to reclassify tensors to be field tensors and motion tensors, thus field tensors may vary with field while motion tensors should vary both with field and matter's motion. For example, the bases only depend on gravitational field, while velocity of a matter may vary due to positions changed. For example, case in one source field, a space derivative of a base may be nonvanishing, but a time derivative of a base must be zero, nevertheless, the base relating to the matter's moving in space time would vary because the coordinates varied on trajectory.

A trajectory of motions of a matter should be a directed curve line in a space, from the start to the end. It is said that the trajectory must be single parametrical curve line. Theoretically, the parameter maybe natural i.e. the line it pass through, and as same it could be time that the motion experienced. What worthy of highlight is that these parameters are simultaneous. That is said a record of the parameter corresponds to a sole record of another. The parameter indicates the sequences.

It is no harm to discuss the trajectory vector λ as a curve line in contra variant space, in that the trajectory is a function of single variable. There is

$$\lambda(x^0, x^1, x^2, x^3) = \lambda(\lambda) = \lambda(t) \quad (213)$$

where λ is the length of trajectory.

It is said that the trajectory is of

$$x^0 = x^0(t), x^1 = x^1(t), x^2 = x^2(t), x^3 = x^3(t) \quad (214)$$

A tensor variation ratio during a time interval on trajectory could be defined to be trajectory derivative that

$$\left(\frac{DA}{dx^\mu}\right)_{tr} = \frac{DA}{d\lambda} \frac{d\lambda}{dx^\mu} \quad (215)$$

For example

$$\left(\frac{DA}{dt}\right)_{tr} = \frac{DA}{d\lambda} \frac{d\lambda}{dt} \quad (216)$$

where, Einstein summation convention does not act on double λ because trajectory is just a single line. The differential length $d\lambda$ is the differential of matter's trajectory, so that $\frac{DA}{d\lambda} d\lambda$ is the covariant differential of tensor A between two neighbourhood positions on the trajectory. Thus, the so-called trajectory derivative is really a kind of line derivative that is derivated by a parameter of trajectory.

It should be noted that a trajectory of matter's motion indicates time sequences corresponding to positions on the trajectory. There is the substantial difference between trajectory derivatives and primary derivatives. A differential on trajectory is the distance interval that a matter has past across, so that the velocity is a trajectory derivative

$$V_0^\dagger = \left(\frac{d\lambda}{dt}\right)_{tr} = \frac{d\lambda}{dt} = \left(\frac{dx^0(t)}{dt}, \frac{dx^1(t)}{dt}, \frac{dx^2(t)}{dt}, \frac{dx^3(t)}{dt}\right) \quad (217)$$

As mentioned above, bases are defined by direct derivatives such as

$$e^\mu = \frac{\partial r}{\partial x^\mu} = \left(\frac{\partial x^0}{\partial x^\mu}, \frac{\partial x^1}{\partial x^\mu}, \frac{\partial x^2}{\partial x^\mu}, \frac{\partial x^3}{\partial x^\mu}\right) \quad (218)$$

where, dx^μ is a proper coordinate differential, so that it is middle labeled.

For the velocity, the differential dx^μ is exactly defined on a trajectory of a matter, so that there is the probability that

$$V_0^\dagger \neq 0 \quad (219)$$

It is said that velocity tensor itself is literally trajectory derivatives.

Trajectory derivatives of general tensor should be labeled for discrepancy.

For example, the bases along a trajectory in rest field

$$\left(\frac{de^i}{dt}\right)_{tr} = \frac{de^i}{d\lambda} \frac{d\lambda}{dt} \neq 0 \quad (220)$$

while direct derivatives with $i \neq 0$

$$\frac{\partial e^i}{\partial t} = 0 \quad (221)$$

Equation (216) could be calculated as

$$\left(\frac{DA}{dt}\right)_{tr} = \frac{D(A^i e_i)}{d\lambda} \frac{d\lambda}{dt} = e_i \frac{dA^i}{d\lambda} \frac{d\lambda}{dt} + A^i \frac{de_i}{d\lambda} \frac{d\lambda}{dt} \quad (222)$$

and $\frac{dA^i}{d\lambda} \frac{d\lambda}{dt}$ is also time derivative, and there is

$$\left(\frac{dA^i}{dt}\right)_{tr} = \frac{dA^i}{d\lambda} \frac{d\lambda}{dt} \quad (223)$$

where, Einstein summation convention does not act on double λ .

Thus, we have seen the difference between a trajectory derivative and a primary derivative.

On the other hand, the value of matter's velocity

$$V_0^\lambda = \frac{d\lambda}{dt} \quad (224)$$

so that

$$\begin{aligned} \left(\frac{DA}{dt}\right)_{tr} &= e_i \frac{dA^i}{d\lambda} \frac{d\lambda}{dt} + A^i \frac{de_i}{d\lambda} \frac{d\lambda}{dt} \\ &= e_i \left(\frac{dA^i}{dt}\right)_{tr} + A^i \frac{de_i}{d\lambda} V_0^\lambda \\ &= e_i \left(\frac{dA^i}{dt}\right)_{tr} + A^i \frac{\partial e_i}{\partial x^\mu} \frac{\partial x^\mu}{\partial x^\lambda} V_0^\lambda \\ &= e_i \left(\frac{dA^i}{dt}\right)_{tr} + A^i \frac{\partial e_i}{\partial x^\mu} \frac{\partial x^\mu}{\partial x^\lambda} \cdot V_0^\lambda \\ &= e_i \left(\frac{dA^i}{dt}\right)_{tr} + A^i \frac{\partial e_i}{\partial x^\mu} \alpha_\lambda^\mu \cdot V_0^\lambda \\ &= e_i \left(\frac{dA^i}{dt}\right)_{tr} + A^i \frac{\partial e_i}{\partial x^\mu} V_0^\mu \end{aligned} \quad (225)$$

where, Einstein summation convention does not act on double λ , and $\alpha_\lambda^\mu = \frac{\partial x^\mu}{\partial x^\lambda}$ is the direction cosine on the direction μ of vector line λ , so that $V_0^\mu = \alpha_\lambda^\mu \cdot V_0^\lambda$ is of component of V_0^λ on that direction and V_0^λ is vector form of V_0^λ so that it could also be written as V_0^λ equivalently in this equation specially for matter's motion on the trajectory.

It could be expanded to be

$$\left(\frac{DA}{dt}\right)_{tr} = e_i \left(\frac{dA^i}{dt}\right)_{tr} + A^i \left(\frac{\partial e_i}{\partial x^0} V_0^0 + \frac{\partial e_i}{\partial x^1} V_0^1 + \frac{\partial e_i}{\partial x^2} V_0^2 + \frac{\partial e_i}{\partial x^3} V_0^3 \right) \quad (226)$$

The expression in component forms could also be worked out as

$$\left(\frac{DA^\mu}{dt}\right)_{tr} = \left(\frac{dA^\mu}{dt}\right)_{tr} + A^i \left(\frac{\partial e_i}{\partial x^0} \cdot e^\mu V_0^0 + \frac{\partial e_i}{\partial x^1} \cdot e^\mu V_0^1 + \frac{\partial e_i}{\partial x^2} \cdot e^\mu V_0^2 + \frac{\partial e_i}{\partial x^3} \cdot e^\mu V_0^3 \right) \quad (227)$$

The expression in the way of Christoffel symbols as

$$\left(\frac{DA^\mu}{dt}\right)_{tr} = \left(\frac{dA^\mu}{dt}\right)_{tr} + \Gamma_{0i}^\mu A^i V_0^0 + \Gamma_{1i}^\mu A^i V_0^1 + \Gamma_{2i}^\mu A^i V_0^2 + \Gamma_{3i}^\mu A^i V_0^3 \quad (228)$$

In rest fields, there will be $\frac{\partial e_i}{\partial x^0} = 0$, so that $\Gamma_{0i}^\mu = 0$. It is said that for one source field, a trajectory derivative could be based on a real trajectory that

$$\left(\frac{DA}{dt}\right)_{tr} = e_i \left(\frac{dA^i}{dt}\right)_{tr} + A^i \left(\frac{\partial e_i}{\partial x^1} V_0^1 + \frac{\partial e_i}{\partial x^2} V_0^2 + \frac{\partial e_i}{\partial x^3} V_0^3 \right) \quad (229)$$

and

$$\left(\frac{DA^\mu}{dt}\right)_{tr} = \left(\frac{dA^\mu}{dt}\right)_{tr} + \Gamma_{1i}^\mu A^i V_0^1 + \Gamma_{2i}^\mu A^i V_0^2 + \Gamma_{3i}^\mu A^i V_0^3 \quad (230)$$

It should be highlighted that the trajectory derivatives could also be defined in distance derivatives as

$$\left(\frac{DA^\mu}{d\lambda}\right)_{tr} = \frac{DA^\mu}{d\lambda} \quad (231)$$

which just performs a special appearance of trajectory derivatives.

For free falling trajectory, it is

$$\left(\frac{DA^\mu}{dr}\right)_{tr} = \frac{DA^\mu}{dr} \quad (232)$$

Trajectory derivative is derivative on a curve line in space time. Sometimes it is presented with time because that the time variable is used in the parametric equation. Christoffel symbols in the equations owe to derivatives of space differentials rather than trajectory differentials. We will see that the concept of trajectory derivative help to describe frequency shift and acceleration, as well as to falsify the concept of geodesic line in section 8.2.

5. Theoretical Verifications on Gravitational Redshifts and Accelerations

Because of the inequality of Christoffel symbols of mixed subscripts, the classical Christoffel symbol equations could not be used any more in the theory of general relativity. The covariant derivatives in gravitational field should be considered in their correct forms.

5.1. On Gravitational Redshifts

Taking light propagation at vertical direction for granted, a distance derivative of contra variant frequency is exactly trajectory derivative

$$\left(\frac{Dv_0}{dr}\right)_{tr} = \left(\frac{dv_0}{dr}\right)_{tr} - \Gamma_{10}^0 v_0 \quad (233)$$

where, the first item of right side of the equation is contra variant derivative that has been drawn in Equation (15).

It is sure to consider the tensor of frequency and its derivative to be vectors, but in traditions it is not of a rare necessity. It has been mentioned that Christoffel symbol Γ_{10}^0 were employed correctly with the form $\Gamma_{10}^0 = \frac{1}{2} g^{00} \frac{\partial g_{00}}{\partial x^1}$ as has been shown in Equation (26). As a result, it will lead to a real answer

$$\Gamma_{10}^0 = \frac{1}{2} g^{00} \frac{\partial g_{00}}{\partial r} = \frac{1}{2} (1 - \frac{r^*}{r})^{-1} (1 - \frac{r^*}{r})' = \frac{r^*}{2r^2} (1 + \frac{r^*}{r-r^*}) \quad (234)$$

The covariant derivative will be calculated to be

$$\left(\frac{Dv_0}{dr}\right)_{tr} = -\frac{r^*}{2r^2} v_{0(\infty)} - \frac{r^*}{2r^2} (1 + \frac{r^*}{r-r^*}) (1 + \frac{r^*}{2r}) v_{0(\infty)} \quad (235)$$

The approximate solution for weak field is that

$$\left(\frac{Dv_0}{dr}\right)_{tr} \approx 2 \frac{dv_0}{dr} \quad (236)$$

It is said that the solution Equation (27) and Equation (28) are confirmed again, and they have revealed that the covariant redshift is approximately double of the contra variant one in weak field.

Notwithstanding, one can also make another derivative as

$$\left(\frac{Dv_0}{dt}\right)_{tr} = \frac{dv_0}{dr} \frac{dr}{dt} - \Gamma_{10}^0 v_0 \frac{dr}{dt} \quad (237)$$

For light rays at that trajectory, $\frac{dr}{dt} = c_0^1$. It becomes

$$(\frac{Dv_0}{dt})_{tr} = (\frac{dv_0}{dt})_{tr} - c_0^1 \Gamma_{10}^0 v_0 \quad (238)$$

The first item of right hand side can also be transformed to be

$$(\frac{dv_0}{dt})_{tr} = \frac{dv_0}{dr} \frac{dr}{dt} = c_0^1 \frac{dv_0}{dr} \quad (239)$$

so that

$$(\frac{Dv_0}{dt})_{tr} = c_0^1 (\frac{dv_0}{dr} - \Gamma_{10}^0 v_0) = c_0^1 (\frac{Dv_0}{dr})_{tr} \approx 2c_0^1 \frac{dv_0}{dr} \quad (240)$$

5.2. On Accelerations

One can take matter's freefalling for example to study the acceleration in gravitational fields. The whole contra variant acceleration is the derivative of whole contra variant velocity as that

$$a_{00}^1 = (\frac{dV_0^1}{dt})_{tr} = \frac{d^2 r}{dt^2} \quad (241)$$

And whole covariant acceleration is a pure covariant derivative

$$a_{1/00} = (\frac{DV_{1/0}^1}{d\tau})_{tr} = \frac{d^2 \rho}{d\tau^2} \quad (242)$$

One may find that some tensors have been labeled with detailed middle index hence they may help to provide explicit expressions, in which the symbol / is employed to divide the middle upper and middle lower indexes.

With the relationship between covariant derivatives, it could be drawn that

$$e^1 e_0 a_{1/00} = (\frac{DV_0^1}{dt})_{tr} \quad (243)$$

To study the covariant derivative expression with Christoffel symbols, there is

$$(\frac{DV_0^1}{dt})_{tr} = (\frac{dV_0^1}{dt})_{tr} + \Gamma_{11}^1 \frac{dr}{dt} V_0^1 - \Gamma_{10}^0 \frac{dr}{dt} V_0^1 \quad (244)$$

With $\frac{dr}{dt} = V_0^1$, it becomes

$$(\frac{DV_0^1}{dt})_{tr} = (\frac{dV_0^1}{dt})_{tr} + \Gamma_{11}^1 (V_0^1)^2 - \Gamma_{10}^0 (V_0^1)^2 \quad (245)$$

With $\Gamma_{11}^1 = e^1 \frac{\partial e_1}{\partial r}$, $\Gamma_{10}^0 = e^0 \frac{\partial e_0}{\partial r}$, $e^0 \partial e_0 = -e_0 \partial e^0$ and $(\frac{dV_0^1}{dt})_{tr} = a_{00}^1$, it is

$$\begin{aligned} (\frac{DV_0^1}{dt})_{tr} &= a_{00}^1 + e^1 \frac{\partial e_1}{\partial r} (V_0^1)^2 - e^0 \frac{\partial e_0}{\partial r} (V_0^1)^2 \\ &= a_{00}^1 + e^1 \frac{\partial e_1}{\partial r} (V_0^1)^2 + e_0 \frac{\partial e^0}{\partial r} (V_0^1)^2 \\ &= a_{00}^1 + e^1 e_0 \frac{\partial}{\partial r} (e_1 e^0) (V_0^1)^2 \end{aligned} \quad (246)$$

With Equation (243), we obtain

$$a_{1/00} = e^0 e^0 e_1 (\frac{DV_0^1}{dt})_{tr} = e^0 e^0 e_1 a_{00}^1 + e^0 \frac{\partial}{\partial r} (e_1 e^0) (V_0^1)^2 \quad (247)$$

One can find that we have use Γ_{11}^1 and Γ_{10}^0 in the calculation of $a_{1/00}$, rather than Γ_{00}^1 that has been used in Equation (44). As we have discussed, the value of Γ_{00}^1 is really of zero.

As has been said that covariant derivatives could also be developed in a direct way without Christoffel symbols

$$a_{1/00} = (\frac{DV_{1/0}^1}{d\tau})_{tr} = (\frac{d^2 \rho}{d\tau^2})_{tr} = [\frac{d}{d\tau} (\frac{e_1 dr}{e_0 dt})]_{tr} = [\frac{d}{e_0 dt} (\frac{e_1 dr}{e_0 dt})]_{tr}$$

$$\begin{aligned}
&= e^0 \frac{e_1}{e_0} \left[\frac{d}{dt} \left(\frac{dr}{dt} \right) \right]_{tr} + e^0 \frac{dr}{dt} \left[\frac{d}{dt} \left(\frac{e_1}{e_0} \right) \right]_{tr} \\
&= e^0 e^0 e_1 a_{00}^1 + e^0 V_0^1 \left[\frac{d}{dt} (e^0 e_1) \right]_{tr} \\
&= e^0 e^0 e_1 a_{00}^1 + e^0 V_0^1 \frac{\partial}{\partial r} (e^0 e_1) \frac{dr}{dt} \\
&= e^0 e^0 e_1 a_{00}^1 + e^0 \frac{\partial}{\partial r} (e^0 e_1) (V_0^1)^2
\end{aligned} \tag{248}$$

It could be found that this equation has been far different from the Equation (44), because errors in Christoffel symbol equation have been eliminated and at the same time the concept of trajectory derivate help to calculate an acceleration in right way. These discussions have presented further verifications for the revised equation of Christoffel symbols of Equation (211).

By the way, it is interesting to take some discussions on some trivial concepts such as a_{00}^0 and the covariant form $a_0/00$ of massive matters. Since V_0^0 and $V_0/0$ are the velocities of contra variant time and proper time but not the real velocity of light

$$V_0^0 = \frac{dx^0}{dt} = \frac{d(ct)}{dt} = c \tag{249}$$

and

$$V_0/0 = \frac{e_0 dx^0}{d\tau} = \frac{e_0 d(ct)}{d\tau} = c \tag{250}$$

Then their derivatives are just the accelerations of time coordinates that

$$a_{00}^0 = \left(\frac{dV_0^0}{dt} \right)_{tr} = 0 \tag{251}$$

while

$$\left(\frac{dV_0^0}{dt} \right)_{tr} = \left(\frac{dV_0^0}{dt} \right)_{tr} + \Gamma_{10}^0 V_0^0 V_0^1 - \Gamma_{10}^0 V_0^0 V_0^1 = 0 \tag{252}$$

so that

$$a_0/00 = 0 \tag{253}$$

As light propagation at a direction of a radius is concerned, we know that light speed c keeps invariant in covariant space, so that there is

$$\frac{Dc}{d\tau} = 0 \tag{254}$$

Case discussing the performance of contra variant light speed, with invariant distance, there is

$$ds^2 = -g_{00}c^2 dt^2 + g_{11}dr^2 = 0 \tag{255}$$

and then the light speed in contra variant space will be

$$c_0^1 = \frac{dr}{dt} = \sqrt{\frac{g_{00}}{g_{11}}} c = e_0 e^1 c \tag{256}$$

where, positive g_{00} is set instead of a minus g_{00} as has suggested previous.

Then the acceleration

$$a_{00}^1 = \left(\frac{dc_0^1}{dt} \right)_{tr} = \left[\frac{d}{dt} (e_0 e^1 c) \right]_{tr} = c \frac{\partial}{\partial r} (e_0 e^1) \frac{\partial r}{\partial t} = c c_0^1 \frac{\partial}{\partial r} (e_0 e^1) = c^2 e_0 e^1 \frac{\partial}{\partial r} (e_0 e^1) \tag{257}$$

With Equation (246) the covariant derivative is

$$\begin{aligned}
\left(\frac{Dc_0^1}{dt} \right)_{tr} &= \left(\frac{dc_0^1}{dt} \right)_{tr} - e_1 e^0 \frac{\partial}{\partial r} (e^1 e_0) (c_0^1)^2 \\
&= a_{00}^1 - e_1 e^0 \frac{\partial}{\partial r} (e^1 e_0) (c_0^1)^2
\end{aligned}$$

$$= c^2 e_0 e^1 \frac{\partial}{\partial r} (e_0 e^1) - c^2 e_0 e^1 \frac{\partial}{\partial r} (e^1 e_0) = 0 \tag{258}$$

Of course, with Equation (246) and Equation (243), we could obtain the result only by a judgment that

$$\left(\frac{Dc_0^1}{dt}\right)_{tr} = e^1 e_0 e_0 \frac{Dc}{d\tau} = 0 \tag{259}$$

6. Experimental Verifications on Gravitational Redshifts and Accelerations

Every tensor involved with measurable quantities could have probabilities to be performed in practice with measured quantities to verify their theoretical expressions. In space time, space intervals and time intervals are all measurable quantities so that they surely could be employed to perform the space and time dependent tensors.

The methodology of the so-called revisit gravitational redshift encourages me to sponsor a realistic analysis method to further verify the general covariance, which will present solutions all based on physical events of realities. Physical events always have substantial existence so that they can help to create irrefutable conclusions. We know that physical events may be record both in contra variant space and covariant space that might provide different values for physical quantities, but both of them actually represent the same physical realities.

6.1. On Measurable Experiments

Measurable quantities could be used to describe physical events, which may be coordinate independent or not. Coordinate independent quantities of course show invariance in physical events in different spaces, such as wave numbers, which could be record as images or texts at specific times and positions. However, coordinate dependent quantities measured in site may really depend. For examples, distance measurements not only depend on in-site space coordinates but also depend on the in-site rulers, so as well, time measurements also depend on both in-site time coordinates and the in-site clocks. We may imagine that the space rulers and clocks their selves maybe also vary. Logically, records of these quantities are recognizable even if they are in farthest distance to the bystanders.

Case a measurement equipment varies with time space, whether the measurement quantity measured is in contra variant space quantity or covariant space quantity? With general covariance, it has been believed that rulers will shrink when they go closer to the center source corresponding to the space interval to become shorter. And also, it has been expected that clocks will go variant corresponding to their dynamic conditions.

However, after those inspections in previous sections, we know that general covariance does not work in some circumstances. Energy and momentum of a matter may not keep covariant in covariant space, while light speed may keep covariant spectacularly. On another side, our discussions may have led to a theoretical inference that matters may experience relativistic emission when they go to a center source and then shrink because of the variation of energy structure. That could be called covariant deformations.

Once we measure space and time intervals at a position, maybe they are not committed to be contra variant quantities or covariant quantities, because our rulers may vary uncommitted. But we could do make measurements anyway. In another word, we could indeed measure something so that they will correspond to any others. Thus, it is not harmful to suppose one of the series of measurable quantities could be measured in following discussions, for example, the contra variant distances or contra variant time intervals. And then they will be valid to be transformed from one to another. That will help us to do more analysis for comparisons and discussions.

6.2. Measurable Verifications for Gravitational Redshift

For the issue of redshift, we are going to sponsor the physical events of wave number counting. It is known that light frequency investigation should be accomplished by indirect techniques and sometimes it may come out with deviations. But it is supposed here that the wave number of the light is countable, or it is believed that light wave could be seen and record. This assumption actually may not do harm to our understanding to the realities, because that indeed will not change the realities and the events of wave counting in that the measurements themselves are also physical processes.

The event of wave number counting could be specified as the record of a number of waves to past a position in a time interval, and it could also be simplified to be one wave corresponding to a time lasting of the light ray propagating a wave length distance. On another side indirectly, one can get wave number by measuring wave length, based on the assumption of invariant light speed. But the apparent light speed might be variable so that the indirect method is not a good idea.

If there is a photon propagating from position 1 to position 2 in a one source field as shown in Figure 16, which correspond to coordinates $r_{(1)}$ and $r_{(2)}$,




Figure 16. A photon traveling in center source field.

The wave number counting events should be carried out at the time that the photon passes the position 1 and position 2. In very short time $\Delta t_{(1)}$ and $\Delta t_{(2)}$, we will count the corresponding wave number $\Delta n_{(1)}$ and $\Delta n_{(2)}$. It should be pointed out that the time intervals here do nothing with the time intervals in Figure 2 in Section 2.2, in that they belong to different physical events that the ones we are talking are of the records of wave counting events and the ones in Figure 2 are of the records of pulse emission and receiving events.

Since the frequencies should be calculated as

$$\nu_{0(1)} = \frac{\Delta n_{(1)}}{\Delta t_{(1)}} \text{ and } \nu_{0(2)} = \frac{\Delta n_{(2)}}{\Delta t_{(2)}} \quad (260)$$

Redshift has been defined as

$$z_{\text{contra}} = \frac{\nu_{0(1)} - \nu_{0(2)}}{\nu_{0(2)}} \quad (261)$$

With the measurable records, it could be rewritten as

$$z_{\text{contra}} = \frac{\Delta n_{(1)} \Delta t_{(2)}}{\Delta n_{(2)} \Delta t_{(1)}} - 1 \quad (262)$$

In the event of wave counting, wave numbers are invariant in both contra variant space and covariant space, but the time intervals $\Delta t_{(1)}$ and $\Delta t_{(2)}$ will vary to be $\Delta \tau_{(1)}$ and $\Delta \tau_{(2)}$. In fact, every physical event keeps the only one event, whereas the different describing metrics lead to different results in the different spaces.

Naturally, gravitational redshift in covariant space is

$$z_{\text{revisit}} = \frac{\nu_{(1)} - \nu_{(2)}}{\nu_{(2)}} = \frac{\Delta n_{(1)} \Delta \tau_{(2)}}{\Delta n_{(2)} \Delta \tau_{(1)}} - 1 \quad (263)$$

where, this redshift symbol labeled with revisit is because it corresponds to that one named in classical equations.

As we know,

$$\Delta \tau = e_0 \Delta t \quad (264)$$

It turns to be

$$z_{\text{revisit}} = \frac{e_{0(2)} \Delta n_{(1)} \Delta t_{(2)}}{e_{0(1)} \Delta n_{(2)} \Delta t_{(1)}} - 1 \quad (265)$$

As in a field of a center source, the metrics take the forms of Schwarzschild solution, the equation will be drawn as

$$z_{\text{revisit}} = \left(1 + \frac{\phi_2 - \phi_1}{c^2} + o\right) \frac{\Delta n_{(1)} \Delta t_{(2)}}{\Delta n_{(2)} \Delta t_{(1)}} - 1 \quad (266)$$

We know that the z_{contra} in Equation (262) could have been measured in the physical event of wave number counting that of course equals to that in Equation (9), so that

$$\frac{\Delta n_{(1)} \Delta t_{(2)}}{\Delta n_{(2)} \Delta t_{(1)}} = \left(1 + \frac{\phi_2 - \phi_1}{c^2} + o\right) \quad (267)$$

Thus, the covariant redshift in weak field is obtained

$$z_{\text{revisit}} = \left(1 + \frac{\phi_2 - \phi_1}{c^2} + o\right)^2 - 1 \approx 2 \frac{\Delta \phi}{mc^2} \approx 2(z)_{\text{contra}} \quad (268)$$

It is said that, the revisit gravitational redshift is double of that of contra variant one.

As the equation of contra variant redshift is concerned, we know that it could be of course drawn by counting two wave numbers in two equivalently specified time intervals. For example, set $\Delta t_{(1)} = \Delta t_{(2)} = \Delta t$ which are measured at positions of $r_{(1)}$ and $r_{(2)}$, then $\Delta n_{(1)}$ and $\Delta n_{(2)}$ should represent the difference of frequency without time intervals. So that

$$z_{\text{contra}} = \frac{\Delta n_{(1)} \Delta t}{\Delta n_{(2)} \Delta t} - 1 = \frac{\Delta n_{(1)}}{\Delta n_{(2)}} - 1 \quad (269)$$

We know that $\Delta n_{(1)}$ and $\Delta n_{(2)}$ present the wave numbers with respect to $\Delta t_{(1)} = \Delta t_{(2)} = \Delta t$.

As for revisit redshift, one will still get different covariant time intervals because the metrics go varied. Thus, it is always doubled of the previous.

$$z_{\text{revisit}} = \frac{\Delta n_{(1)} e_{0(2)} \Delta t}{\Delta n_{(2)} e_{0(1)} \Delta t} - 1 \approx 2(z)_{\text{contra}} \quad (270)$$

It should be pointed out that in some experiments on gravitational redshift, only one timing clock was designed for time interval measurement. In this case, a wrong setting of proper time intervals may be taken into consideration, for example some sole clock timing experiments, so that the experimental redshift may be presented as

$$z_{\text{experimental}} = \frac{\Delta n_{(1)} \Delta \tau_{(2)}}{\Delta n_{(2)} \Delta \tau_{(1)}} - 1 = \frac{\Delta n_{(2)} e_{0(x)} \Delta t_{(2)}}{\Delta n_{(1)} e_{0(x)} \Delta t_{(1)}} - 1 = (z)_{\text{contra}} \quad (271)$$

where, $e_{0(x)}$ is base component at clock position of $r_{(1)}$ or $r_{(2)}$ or any position others to them.

We can find out those completed experiments observations [14–16] on gravitational redshift will be easy to be verified to have only worked out the results of contra variant frequency shift.

Of course, one can calculate the real proper time intervals by time interval transformation between sole timing position and frequency shift positions. That will finally help to work out revisit gravitational redshift as have discussed.

6.3. Measurable Verifications for Acceleration

6.3.1. Measurable Quantities and Measurable Acceleration

Firstly, I prefer to rise a controversy of a free falling on the Earth that if a matter freefalls from rest with velocity $V_{0(1)}^1 = 0$ as well as $V1/0_{(1)} = 0$ by nature, we do know that it will move quite faster with velocity $V_{0(2)}^1$ after traveling a distance in a time interval because of gravity. In traditional theory, we know that the proper velocity $V1/0_{(2)} = e_0 e^1 V_{0(2)}^1$. Considering the weak field effect, there is $e_0 e^1 \approx 1$. Hence comes the controversy that the covariant acceleration must be great than zero because the matter has started from rest to a quite apparent motion. That is really contradicted with the principle of general covariance with a requirement of zero covariant acceleration.

Nevertheless, considering that $V_{0(2)}^1$ and $V_{1(2)}^1$ are still non-relativistic velocities, it is easy to estimated that the accelerations are also approximately equal that $a_{1/00} \approx a_{00}^1$. The following works of so-called realistic verifications in this section are exactly to be sponsored to solve these controversies thoroughly.

A freefalling test with initial velocity is going to be put forward, in which a matter freely falls to source center from a position $r_{(1)}$ to a position $r_{(2)}$ as shown in Figure 17. Once the velocities at the two positions are measured, the average values could be estimated with the velocities difference and the interval distance. Considering the condition on the surface of the Earth, a freefalling with a rarely big velocity and a rarely small traveling would be performed. For example, a velocity of more than 10000 m/s, could be seen as a constant accelerating motion even in covariant space, in that a covariant derivative is expected to be linear with velocity.




Figure 17. Free falling measurement.

It is supposed that contra variant distances and time intervals are measurable. Based on the measurements, velocity at position 1 could be written as

$$V_{0(1)}^1 = \frac{\Delta r_{(1)}}{\Delta t_{(1)}} \quad (272)$$

where the $\Delta r_{(1)}$ and $\Delta t_{(1)}$ are measured distance and time intervals when the matter goes by the position $r_{(1)}$, so that they are both tensors of contra variant space. For reasons of convenience their bracketed sub indexes here are only employed to represent positions.

As well as that at position 2

$$V_{0(2)}^1 = \frac{\Delta r_{(2)}}{\Delta t_{(2)}} \quad (273)$$

Because the velocity at position 2 is the result of acceleration, it could be written in integral form

$$\begin{aligned} V_{0(2)}^1 &= V_{0(1)}^1 + \int_{t_{(1)}}^{t_{(2)}} a_{00}^1 dt \\ &= V_{0(1)}^1 + \overline{a_{00}^1} [t_{(2)} - t_{(1)}] \end{aligned} \quad (274)$$

where, the mean acceleration $\overline{a_{00}^1}$ is the integral point value, and $t_{(2)} - t_{(1)}$ is time interval for the matter traveling from $r_{(1)}$ to $r_{(2)}$.

So that the mean acceleration is

$$\overline{a_{00}^1} = \frac{V_{0(2)}^1 - V_{0(1)}^1}{t_{(2)} - t_{(1)}} \quad (275)$$

On the other hand, with covariant bases there are the relationships of contra variant quantities and proper ones

$$\Delta \rho_{(1)} = e_{1(1)} \Delta r_{(1)}, \Delta \tau_{(1)} = e_{0(1)} \Delta t_{(1)}, \Delta \rho_{(2)} = e_{1(2)} \Delta r_{(2)}, \Delta \tau_{(2)} = e_{0(2)} \Delta t_{(2)} \quad (276)$$

One can use the mean metric to calculate the proper time intervals from position 1 to position 2

$$\tau_{(2)} - \tau_{(1)} = \int_{t_{(1)}}^{t_{(2)}} e_0 dt = \overline{e_0} (t_{(2)} - t_{(1)}) \quad (277)$$

where the $\overline{e_0}$ is the value at integral point, and it is suggested to be evaluated approximately as following in weak field

$$\bar{e}_0 \approx 0.5(e_{0(1)} + e_{0(2)}) \quad (278)$$

The proper velocities

$$V1/0_{(1)} = \frac{\Delta\rho_{(1)}}{\Delta\tau_{(1)}} \quad (279)$$

$$V1/0_{(2)} = \frac{\Delta\rho_{(2)}}{\Delta\tau_{(2)}} \quad (280)$$

And the integral relationship in covariant space that

$$\begin{aligned} V1/0_{(2)} &= V1/0_{(1)} + \int_{\tau_{(1)}}^{\tau_{(2)}} a1/00 d\tau \\ &= V1/0_{(1)} + \overline{a1/00}(\tau_{(2)} - \tau_{(1)}) \end{aligned} \quad (281)$$

And also, we get the mean covariant acceleration with Lagrangian mean value theorem of integration that

$$\overline{a1/00} = \frac{V1/0_{(2)} - V1/0_{(1)}}{\tau_{(2)} - \tau_{(1)}} \quad (282)$$

It could be rewritten as

$$\overline{a1/00} = \frac{\frac{e_{1(2)}\Delta r_{(2)}}{e_{0(2)}\Delta t_{(2)}} - \frac{e_{1(1)}\Delta r_{(1)}}{e_{0(1)}\Delta t_{(1)}}}{\bar{e}_0(t_{(2)} - t_{(1)})} = \frac{\frac{e_{1(2)}V_{0(2)}^1}{e_{0(2)}} - \frac{e_{1(1)}V_{0(1)}^1}{e_{0(1)}}}{\bar{e}_0(t_{(2)} - t_{(1)})} \quad (283)$$

It is of course the measuring forms of covariant acceleration of a freefalling. And then it could be compared to that of contra variant one.

We would like substitute the equation of contra variant velocity 2 of Equation (274) into this equation. That is

$$\begin{aligned} \overline{a1/00} &= \bar{e}^0 \frac{\frac{e_{1(2)}}{e_{0(2)}} [(t_{(2)} - t_{(1)})\overline{a_{00}^1} + V_{0(1)}^1] - \frac{e_{1(1)}}{e_{0(1)}} V_{0(1)}^1}{t_{(2)} - t_{(1)}} \\ &= \bar{e}^0 \frac{e_{1(2)}}{e_{0(2)}} \overline{a_{00}^1} + \bar{e}^0 \frac{\frac{e_{1(2)}}{e_{0(2)}} - \frac{e_{1(1)}}{e_{0(1)}}}{t_{(2)} - t_{(1)}} V_{0(1)}^1 \\ &= \bar{e}^0 e_{1(2)} e_{0(2)}^0 \overline{a_{00}^1} + \bar{e}^0 \frac{e_{1(2)}e_{0(2)}^0 - e_{1(1)}e_{0(1)}^0}{t_{(2)} - t_{(1)}} V_{0(1)}^1 \end{aligned} \quad (284)$$

It could be transformed to be

$$\begin{aligned} \overline{a1/00} &= \bar{e}^0 e_{1(2)} e_{0(2)}^0 \overline{a_{00}^1} + \bar{e}^0 \frac{e_{1(2)}e_{0(2)}^0 - e_{1(1)}e_{0(1)}^0}{t_{(2)} - t_{(1)}} V_{0(1)}^1 \frac{r_{(2)} - r_{(1)}}{r_{(2)} - r_{(1)}} \\ &= \bar{e}^0 e_{1(2)} e_{0(2)}^0 \overline{a_{00}^1} + \bar{e}^0 \frac{e_{1(2)}e_{0(2)}^0 - e_{1(1)}e_{0(1)}^0}{r_{(2)} - r_{(1)}} V_{0(1)}^1 \overline{V_0^1} \end{aligned} \quad (285)$$

Here we have got the transformed form of covariant acceleration of freefalling.

Nevertheless, with Lagrangian differential mean value theorem, we can write down the differential form as

$$\overline{a1/00} = \bar{e}^0 e_{1(2)} e_{0(2)}^0 \overline{a_{00}^1} + \bar{e}^0 \overline{\frac{\partial}{\partial r} (e_1 e_0^0)} V_{0(1)}^1 \overline{V_0^1} \quad (286)$$

Or the form of reverse bases

$$\overline{a1/00} = \bar{e}^0 e_{1(2)} e_{0(2)}^0 \overline{a_{00}^1} - \bar{e}^0 e_{1(2)} e_{1(1)} e_{0(2)}^0 e_{0(1)}^0 \overline{\frac{\partial}{\partial r} (e^1 e_0)} V_{0(1)}^1 \overline{V_0^1} \quad (287)$$

Thus, by the way, another kind of proof of differential analysis of the Equation (247) or Equation (248) has been completed, in the way of measurable experiment.

6.3.2. Examples

Some terrestrial experiments are going to be put forward, that matters with initial velocity freefall in vacuum circumstance with in 1000m height to the ground. Both at the start point position 1 and end point position 2, the matter’s velocities will be measured. And of course, the space and time intervals between position 1 and 2 that depend on the so-called geodesic line will be measured together so that to calculate the mean accelerations.

Some basic data of the Earth have already been tested certainly, so that we can take the standard value for our experiments, such as the total mass of the Earth $M = 5.97237 \times 10^{24}kg$, and the position on the ground could be assigned to have a radial coordinate $R_{\oplus} = 6.371393 \times 10^6m$. We could also take the gravitational constant $G = 6.67259 \times 10^{-11}Nm^2/kg^2$, with the light speed $c = 299792458\text{ m/s}$ thus the gravitational radius will be calculated as

$$r^* = \frac{2GM}{c^2} = 8.8680827 \times 10^{-3}m \tag{288}$$

With Newtonian equation and Schwarzschild’s solution, some positional data could be listed in Table 1.

Table 1. Base components, their derivatives and gravity at experimental positions.

r	e_1	e_0	GM/r^2	$\partial(e_1e^0)/\partial r$
6372.393	1.00000000069582045	0.99999999930417955	9.81377376	$2.183859184 \times 10^{-16}$
6371.493	1.00000000069591874	0.99999999930408126	9.81654643	$2.184476187 \times 10^{-16}$
6371.393	1.00000000069592966	0.99999999930407034	9.81685457	$2.184544758 \times 10^{-16}$

So far as we have discussed, the accelerations a^1_{00} and $a_{1/00}$ are really geological quantities, and now it is necessary to make an extending study. We know that all kinds of interactions could be seen as momentum exchanges between matters, as that

$$dP = d(mv) = vdm + m dv \tag{289}$$

For the convenience, some quantities discussed in this section will not be marked with tensor index anymore.

In conditions of low velocity motions, the theory of special relativity indicates small mass variations, thus

$$dP \approx m dv \tag{290}$$

For the cases of high velocity motions, one should take a total analysis. Now the total acceleration could be defined

$$\Lambda = \frac{d(mv)}{mdt} = \frac{vdm}{mdt} + \frac{dv}{dt} \tag{291}$$

It is said, the total acceleration includes mass variant acceleration and velocity variant acceleration, and the latter also could be called geometrical acceleration.

With a momentum variation, kinetic energy will vary a difference

$$dE_k = v dP = v^2 dm + m v dv \tag{292}$$

At the same time, the mass energy equation of differential form is

$$dE_k = c^2 dm \tag{293}$$

Thus, there will be

$$c^2 dm = v^2 dm + m v dv \tag{294}$$

To be divided by time differential, that will lead to the expression of acceleration

$$a^1_{00} = \frac{dv}{dt} = \frac{c^2 - v^2}{mv} \frac{dm}{dt} \tag{295}$$

Now one can define a coefficient of geometrical acceleration

$$\eta = \frac{\frac{dv}{dt}}{\Lambda} = \frac{\frac{c^2 - v^2}{mv} \frac{dm}{dt}}{\frac{v dm}{mdt} + \frac{c^2 - v^2}{mv} \frac{dm}{dt}} = 1 - \frac{v^2}{c^2} \quad (296)$$

In one source field, single acting gravitational geometrical acceleration is

$$a_{00}^1 = \eta \frac{GM}{r^2} = \eta g \quad (297)$$

where g is the total acceleration of gravity.

We will see that geometrical acceleration declines as velocity goes up to a relativistic level, and it goes to zero as velocity closely catches up to light speed.

If a matter is accelerated from rest, the total energy includes rest part and kinetic part

$$mc^2 = m_0 c^2 + \xi m v^2 \quad (298)$$

where m is relativistic mass and m_0 is rest mass.

We know that in special relativity there is

$$m = \frac{m_0}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (299)$$

Back to Equation (298), there is

$$\xi = (1 - \sqrt{1 - \frac{v^2}{c^2}}) \frac{c^2}{v^2} \quad (300)$$

Then the total kinetic energy

$$E_k = \xi m v^2 = (1 - \sqrt{1 - \frac{v^2}{c^2}}) \frac{c^2}{v^2} m v^2 \quad (301)$$

Case v/c is a small value, ξ would be close to 0.5, so that

$$E_k \approx \frac{1}{2} m v^2 \quad (302)$$

On the occasion of freefalling the variation of kinetic energy

$$\xi_2 m_2 v^2 - \xi_1 m_1 v_1^2 = \int_{r(1)}^{r(2)} m g ds = \bar{m} \bar{g} s \quad (303)$$

Again with Equation (294) the energy difference in an experiment

$$\Delta m c^2 \approx \Delta m v^2 + m_1 v \Delta v \quad (304)$$

Thus, there is

$$\Delta m = m_1 \frac{v \Delta v}{c^2 - v^2} \quad (305)$$

Considering $\eta_1 \approx \eta_2$ in experiments and v_1 is quite big, the difference of kinetic energy could be written as

$$\begin{aligned} \Delta E_k &= \xi_2 m_2 v^2 - \xi_1 m_1 v_1^2 = \xi_2 (m_1 + \Delta m) (v_1 + \Delta v)^2 - \xi_1 m_1 v_1^2 \\ &\approx \xi_1 (m_1 + \Delta m) (v_1^2 + 2v_1 \Delta v) - \xi_1 m_1 v_1^2 \\ &= \xi_1 (\Delta m v_1^2 + 2m_1 v_1 \Delta v) \\ &= \xi_1 m_1 \left(\frac{v_1 \Delta v}{c^2 - v_1^2} v_1^2 + 2v_1 \Delta v \right) \\ &\approx m_1 g s \end{aligned} \quad (306)$$

so that

$$\Delta v \approx \frac{gs}{\xi_1 v_1 \left(\frac{v_1^2}{c^2 - v_1^2} + 2 \right)} \quad (307)$$

Thus,

$$t_{(2)} - t_{(1)} = \Delta t = \Delta v / a_{00}^1 \quad (308)$$

Unfortunately, this solution cannot come up with a higher accuracy than that

$$\Delta t = \frac{s}{v_1 + 0.5\Delta v}$$

(309)

After then, we are going to sponsor series of freefalling experiments. Contra variant accelerations and covariant accelerations for every position are easy to calculate. While measurable covariant acceleration $\overline{a_{1/00}}$ could be obtained with measured distances and time intervals via Equation (283). But it is convenient to calculate with Equation (284), in that the latter is just a transformation of the previous. And in this equation, space intervals would be gained with Newtonian equations and time intervals $t_{(2)} - t_{(1)}$ with Equation (309) for convenience. One may argue that the measured quantities might come from calculation. That doesn't matter, because the equation has been verified for hundreds of years, therefore it is sure that the calculated quantities have equal value with that by measuring. And then the mean covariant acceleration will be taken to compare with the contra variant acceleration a_{00}^1 calculated with Equation (297) and covariant acceleration $a_{1/00}$ calculated with Equation (245) or Equation (248). Calculation results have been listed in Table 2 and Table 3.

Table 2. Terrestrial freefalling experiments from position 6371.493 to 6371.393.

Analysis method	Theoretical analysis				Measurable analysis	
Positions	$r_{(1)}=6371.493$		$r_{(2)}=6371.393$		From $r_{(1)}$ to $r_{(2)}$	
Initial velocity	Contra variant acceleration	Covariant acceleration	Contra variant acceleration	Covariant acceleration	measured time intervals	Covariant acceleration
$V_{0(1)}^1$ (m/s)	$a_{00(1)}^1$ (m/s ²)	$a_{1/00(1)}$ (m/s ²)	$a_{00(2)}^1$ (m/s ²)	$a_{1/00(2)}$ (m/s ²)	$t_{(2)} - t_{(1)}$ (s)	$\overline{a_{1/00}}$ (m/s ²)
0	9.8165464	9.8165464	9.8168546	9.8168546	4.5137	9.8165
100000	9.8165453	9.8165453	9.8168535	9.8168535	1.0×10^{-3}	9.8165
10000000	9.8056240	9.8058425	9.8059318	9.8061503	1.0×10^{-5}	9.8275
100000000	8.7243083	10.908784	8.7245822	10.909127	1.0×10^{-6}	10.908
200000000	5.4475940	14.185499	5.4477651	14.185944	5.0×10^{-7}	14.184
250000000	2.9900583	16.643034	2.9901522	16.643557	4.0×10^{-7}	16.640
299792458	0	19.633093	0	19.633709	3.3356×10^{-7}	19.629

Table 3. Terrestrial freefalling experiments from position 6372.393 to 6371.393.

Analysis method	Theoretical analysis				Measurable analysis	
Positions	$r_{(1)}=6372.393$		$r_{(2)}=6371.393$		From $r_{(1)}$ to $r_{(2)}$	
Initial velocity	Contra variant acceleration	Covariant acceleration	Contra variant acceleration	Covariant acceleration	measured time intervals	Covariant acceleration
$V_{0(1)}^1$ (m/s)	$a_{00(1)}^1$ (m/s ²)	$a_{1/00(1)}$ (m/s ²)	$a_{00(2)}^1$ (m/s ²)	$a_{1/00(2)}$ (m/s ²)	Δt (s)	$\overline{a_{1/00}}$ (m/s ²)
0	9.8137738	9.8137738	9.8168546	9.8168546	14.276	9.8138
100000	9.8137730	9.8137751	9.8168535	9.8168535	1.0×10^{-2}	9.8138
10000000	9.8028548	9.8246934	9.8059318	9.8061503	1.0×10^{-4}	9.8247
100000000	8.7218446	10.905704	8.7245822	10.909127	1.0×10^{-5}	10.906
200000000	5.4460559	14.181493	5.4477651	14.185944	5.0×10^{-6}	14.182
250000000	2.9892144	16.383343	2.9901522	16.643557	4.0×10^{-6}	16.639
299792458	0	19.627547	0	19.633709	3.3356×10^{-6}	19.629

7. Conclusions and Inferences and Their Applications

7.1. Conclusions

Previous discussions will lead to two conclusions for matters' motions in gravitational fields:
1) For light: Light speed keeps general covariance in covariant space, but light frequency keeps conservation in contra variant space.

2) For massive matters: Massive matter's velocity does not perform general covariance, but it will keep conservation in contra variant space.

7.2. Inferences

The conclusions are really different from classical theory of general relativity and they will then lead to natural inferences. I prefer to focus on the inferences on kinematics and relativistic release:

1) For kinematics: General covariance goes break by a large range. During the motions in gravitational field, all matters, including light rays, will keep energy and momentum conservation in contra variant space rather than that in covariant space. Only for light rays they may keep velocity invariant in covariant space, but their energy and momentum will still keep conservation in contra variant space. Energy momentum conservations are the conservations under the condition of gravitational potential conversions. It is said that there is only one exception in realities, the light speed invariance, which will lead to the validity Lagrangian of light ray propagation. While for massive matters, Lagrangian goes invalid. At any positions in gravity fields, massive matters always have opportunities to be accelerated up to and keep velocities close to absolute-light-speed.

2) For relativistic release: Since apparent light speed may vary in gravitational field, that will bring changes to interaction efficiency between and within particles of massive matters so as to influence fine structures. For electromagnet forces, there will be of variations of momentum exchanges. It is also reasonable to predict that the speed of gluons relating to the strong interactions is general covariant like that of photons. On the other side, these interactions keep energy momentum conservations at the same time. Therefore, case massive matters inflow enough distance in gravitational fields, they might get to excited state and release energy, which could be called relativistic release, just as excited electrons in atoms might do. The difference is that relativistic releases might be releases experiencing thoroughly exciting of the matters both in intrinsic structures and exterior structures, including exciting of electrons. Matters may also experience covariant deformation after relativistic release because of new equivalent state. Moreover, relativistic emissions and absorptions may reveal the mystery of intrinsic structures just as the exciting electrons in atoms might do.

7.3. Applications

Detailed discussions on some applications will be sponsored consequently that will greatly support the conclusions and inferences.

1) On kinematics: General covariance and physical quantity conservativeness are of the handles to rectify the classical equations, especially the principle of moment and mass energy conservations. It would be seen that those efforts to employ the geodesic equation or covariant derivatives to build kinetic equations have already gone failed, in that covariant derivatives might be nonvanishing. Renovated solutions on light ray propagations and massive matter's motion as applications might provide forceful verifications on conclusions and inferences.

2) On relativistic release: The concept of equivalent state would be carried out to estimate the energy exceeding for inflow matters so that to discuss energy release, which will then lead to relativistic redshift of emission and absorption. Equivalent state also relates to relativistic deformation that might perform another kind of covariance. Researches on relativistic release will bring about fantastic interpretations on tremendous observations on quasars and active galactic nuclei.

7.4. On Conservativeness

If a physical quantity being conservative in covariant space or contra variant space is to say the quantity keep invariant at any positions. Consequently, the forms of conservative quantities must be expressed in subsidiary conservatives. It should be highlighted that a position in gravitational fields refers to a definite physical reality no matter how does it be described in covariant space or contra

variant space. Thus, a quantity at a position always refers to a definite physical quantity whatever described in any coordinate space. A physical quantity in a gravitational field could be conservative or not in one of those descriptions. For example, light speed is a conservative quantity at any positions in covariant space but it is variable in contra variant space. If a physics expressed by conservative quantities, one could give their values directly, else if not, the variations should be performed by coordinates of space time. In a word, the physical quantities defined should perform physical realities.

Momentum and energy keep conservations in contra variant space is something special that they are keeping conservations after considerations of the momentum and energy exchanges with gravitational fields.

A quantity conservative in a space may be formed with subsidiary quantities in another space or not. We will see in next section that the angular moment of light rays keep conservation in contra variant space but the expression would be written in form of covariant light speed.

8. Kinematics and Dynamics

8.1. The Most Important

The second Newtonian law interprets the mechanism of accelerative motions of massive matters so that to form the dynamics. Case in the conditions that matters have relativistic velocities, forces acting on matters will cause not only the variations of velocities but also the variations of matter's mass. It should be highlighted that a force really is of a statistic quantity rather than an essential physical quantity. In fact, a force is just a performance of exchange of momentum, as well as mass energy at the mean time. Thus, that physics could be called the relativistic dynamics.

But for light propagations, the second Newtonian law will not take effects anymore. Even in the case that a force is vertical to a light ray, we will see that the second Newtonian law remains invalid. That is the reason we suggest the concept of kinematics that others to the concept of dynamics. If we persistently employ the concept of dynamics, it should be a new one.

No matter the space time been determined by what kind of metrics and labeled by what kind of coordinates, it is just a methodology for descriptions for physical events. None of them would have priorities. Physics is on earth depends on its nature rather on spaces defined. The most important is the conservative principles in realities.

It is easy to imagine that geodesic line could be employed for the solution of kinematic trajectories of matters, because general relativity expects conservations in curve space. But we will find out that geodesic equation or covariant derivatives have not really taken effect in the solving of the kinematics in the past century. We know the reason is that covariant derivatives may be nonvanishing so that those imposed settings of vanishing covariant derivatives might cause discrepancies with respect to realities.

Most of methodologies for kinematic trajectories published were based on the so-called Lagrangian. Besides these conditions, contra variant angular momentum conservation has been used in all those solutions. One can imagine that this condition is apparently contradicted with general covariance. In fact, it is always the greatest reason for me to persist in this issue with more efforts.

Finally, the Lorentz covariance is also a kind of constraint condition, since it has been involved in the settings of Minkowski space and pseudo Riemannian space.

8.2. Falsification of the Employment of Geodesic Line for Kinematic Equation

We have drawn the conclusion that covariant accelerations of matters in gravitational fields may be nonvanishing so that the geodesic lines which ask for zero value of the covariant derivatives cannot be employed to be the kinematic equation for matter's motions. We will find great many faults in those equations with which they declare the kinematic equations coming from geodesic lines.

The geodesic line equations presented by Weinberg [17] with metrics given by

$$ds^2 = B(r)dt^2 - A(r)dr^2 - r^2d\theta^2 - r^2\sin^2\theta d\varphi^2 \quad (310)$$

is that

$$\frac{d^2 x^\mu}{d\tau^2} + \Gamma_{\nu\lambda}^\mu \frac{dx^\nu}{d\tau} \frac{dx^\lambda}{d\tau} = 0 \quad (311)$$

That has been calculated to perform as components as [17]

$$0 = \frac{d^2 r}{d\tau^2} + \frac{A'(r)}{2A(r)} \left(\frac{dr}{d\tau}\right)^2 - \frac{r}{A(r)} \left(\frac{d\theta}{d\tau}\right)^2 - r \frac{\sin^2 \theta}{A(r)} \left(\frac{d\varphi}{d\tau}\right)^2 + \frac{B'(r)}{2A(r)} \left(\frac{dt}{d\tau}\right)^2 \quad (312)$$

$$0 = \frac{d^2 \theta}{d\tau^2} + \frac{2}{r} \frac{d\theta}{d\tau} \frac{dr}{d\tau} - \sin\theta \cos\theta \left(\frac{d\varphi}{d\tau}\right)^2 \quad (313)$$

$$0 = \frac{d^2 \varphi}{d\tau^2} + \frac{2}{r} \frac{d\varphi}{d\tau} \frac{dr}{d\tau} + 2 \cot\theta \frac{d\varphi}{d\tau} \frac{d\theta}{d\tau} \quad (314)$$

$$0 = \frac{d^2 t}{d\tau^2} + \frac{B'(r)}{B(r)} \frac{dt}{d\tau} \frac{dr}{d\tau} \quad (315)$$

And then, with $\theta = \pi/2$, the so-called kinematic equation, were finally drawn as

$$A(r) \left(\frac{dr}{d\tau}\right)^2 + \frac{L^2}{r^2} - \frac{1}{B(r)} = -E \quad (316)$$

where, $L = r^2 \frac{d\varphi}{d\tau}$ and E are set constants.

It seems that the kinematic equation has been created by covariant derivatives. But it should be pointed out that Equation (311) to Equation (316) have gone wrong. Because we have discussed that the covariant derivatives could be nonvanishing in some occasions, so that there must be something wrong involved.

We are going to sponsor investigations in two ways for a comparison. Firstly, a transformation from the tangent space defined as $(cdt, dr, d\theta, d\varphi)$ to that defined as $(cd\tau, dp, rd\theta, r\sin\theta d\varphi)$, just as the same as previous discussions, will be put into considerations. For convenience, the invariant distance should be rewritten as

$$\begin{aligned} ds^2 &= -B(r)dt^2 + A(r)dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2 \\ &= -g_{00}dt^2 + g_{11}dr^2 + g_{22}d\theta^2 + g_{33}d\varphi^2 \end{aligned} \quad (317)$$

where, signs in the equation have been modified to traditional forms and the metrics would be defined as $g_{00} = B(r)$, $g_{11} = A(r)$, $g_{22} = r^2$, $g_{33} = r^2 \sin^2 \theta$, in which metrics should be positive as has been discussed.

The Equation (228) could be employed for the solution, so that the derivative components could be performed with and then

$$\begin{aligned} \left(\frac{DV^1}{\partial\tau}\right)_{tr} &= \left(\frac{D}{\partial\tau} \frac{dr}{d\tau}\right)_{tr} = \left(\frac{dV^1}{d\tau}\right)_{tr} + \Gamma_{0i}^1 V^i V^0 + \Gamma_{1i}^1 V^i V^1 + \Gamma_{2i}^1 V^i V^2 + \Gamma_{3i}^1 V^i V^3 \\ &= \left(\frac{dV^1}{d\tau}\right)_{tr} + \Gamma_{01}^1 V^1 V^0 + \Gamma_{11}^1 V^1 V^1 + \Gamma_{21}^1 V^1 V^2 + \Gamma_{31}^1 V^1 V^3 \\ &= \left(\frac{dV^1}{d\tau}\right)_{tr} + \Gamma_{11}^1 V^1 V^1 \\ &= \frac{d^2 r}{d\tau^2} + \frac{A'(r)}{2A(r)} \left(\frac{dr}{d\tau}\right)^2 \end{aligned} \quad (318)$$

$$\begin{aligned} \left(\frac{DV^2}{\partial\tau}\right)_{tr} &= \left(\frac{D}{\partial\tau} \frac{d\theta}{d\tau}\right)_{tr} = \left(\frac{dV^2}{d\tau}\right)_{tr} + \Gamma_{0i}^2 V^i V^0 + \Gamma_{1i}^2 V^i V^1 + \Gamma_{2i}^2 V^i V^2 + \Gamma_{3i}^2 V^i V^3 \\ &= \left(\frac{dV^2}{d\tau}\right)_{tr} + \Gamma_{02}^2 V^2 V^0 + \Gamma_{12}^2 V^2 V^1 + \Gamma_{22}^2 V^2 V^2 + \Gamma_{32}^2 V^2 V^3 \\ &= \left(\frac{dV^2}{d\tau}\right)_{tr} + \Gamma_{12}^2 V^2 V^1 \\ &= \frac{d^2 \theta}{d\tau^2} + \frac{1}{r} \frac{d\theta}{d\tau} \frac{dr}{d\tau} \end{aligned} \quad (319)$$

$$\begin{aligned}
\left(\frac{DV^3}{\partial\tau}\right)_{tr} &= \left(\frac{D}{d\tau}\frac{d\varphi}{d\tau}\right)_{tr} = \left(\frac{dV^3}{d\tau}\right)_{tr} + \Gamma_{0i}^3 V^i V^0 + \Gamma_{1i}^3 V^i V^1 + \Gamma_{2i}^3 V^i V^2 + \Gamma_{3i}^3 V^i V^3 \\
&= \left(\frac{dV^3}{d\tau}\right)_{tr} + \Gamma_{03}^3 V^3 V^0 + \Gamma_{13}^3 V^3 V^1 + \Gamma_{23}^3 V^3 V^2 + \Gamma_{33}^3 V^3 V^3 \\
&= \left(\frac{dV^3}{d\tau}\right)_{tr} + \Gamma_{13}^3 V^3 V^1 + \Gamma_{23}^3 V^3 V^2 \\
&= \frac{d^2\varphi}{d\tau^2} + \frac{1}{r} \frac{d\varphi}{d\tau} \frac{dr}{d\tau} + \cot\theta \frac{d\varphi}{d\tau} \frac{d\theta}{d\tau}
\end{aligned} \tag{320}$$

$$\begin{aligned}
\left(\frac{DV^0}{\partial\tau}\right)_{tr} &= \left(\frac{D}{d\tau}\frac{dt}{d\tau}\right)_{tr} = \left(\frac{dV^0}{d\tau}\right)_{tr} + \Gamma_{0i}^0 V^i V^0 + \Gamma_{1i}^0 V^i V^1 + \Gamma_{2i}^0 V^i V^2 + \Gamma_{3i}^0 V^i V^3 \\
&= \left(\frac{dV^0}{d\tau}\right)_{tr} + \Gamma_{00}^0 V^0 V^0 + \Gamma_{10}^0 V^0 V^1 + \Gamma_{20}^0 V^0 V^2 + \Gamma_{30}^0 V^0 V^3 \\
&= \left(\frac{dV^0}{d\tau}\right)_{tr} + \Gamma_{10}^0 V^0 V^1 \\
&= \frac{d^2t}{d\tau^2} + \frac{B'(r)}{2B(r)} \frac{dt}{d\tau} \frac{dr}{d\tau}
\end{aligned} \tag{321}$$

where, $V^\mu = \frac{dx^\mu}{d\tau}$ is number μ component of contra variant velocity.

We have seen that some errors in the Equation (314) to Equation (316) have been rectified. In fact, it is easy to find out calculation errors. If any $i \neq j$ the Christoffel symbol of $\Gamma_{\mu i}^j=0$, in that the bases we discussed are orthogonal.

Secondly, we are going to study another condition that the tangent spaces from $(cdt, dr, rd\theta, r\sin\theta d\varphi)$ to $(cd\tau, d\rho, rd\theta, r\sin\theta d\varphi)$. The invariant distance could be written as

$$\begin{aligned}
ds^2 &= -B(r)(dt)^2 + A(r)dr^2 + (rd\theta)^2 + (r\sin\theta d\varphi)^2 \\
&= -g_{00}dt^2 + g_{11}dr^2 + g_{22}(rd\theta)^2 + g_{33}(r\sin\theta d\varphi)^2
\end{aligned} \tag{322}$$

Thus, $g_{00} = B(r)$, $g_{11} = A(r)$, $g_{22} = 1$, $g_{33} = 1$. We can see that only gravitational metrics has been considered in the expression, as has been mentioned. This kind of settings would easily help us to verify the calculation previous. One may argue that this transformation has overcome Riemannian manifold definition because the contra variant space is not a R^4 . But on earth in mathematics, that doesn't matter because we know that the space could map to a R^4 at all. The derivation regulars are still available. Then the derivatives should be performed with Equation (228) that

$$\begin{aligned}
\left(\frac{DV^1}{\partial\tau}\right)_{tr} &= \left(\frac{D}{d\tau}\frac{dr}{d\tau}\right)_{tr} = \left(\frac{dV^1}{d\tau}\right)_{tr} + \Gamma_{0i}^1 V^i V^0 + \Gamma_{1i}^1 V^i V^1 + \Gamma_{2i}^1 V^i V^2 + \Gamma_{3i}^1 V^i V^3 \\
&= \left(\frac{dV^1}{d\tau}\right)_{tr} + \Gamma_{01}^1 V^1 V^0 + \Gamma_{11}^1 V^1 V^1 + \Gamma_{21}^1 V^1 V^2 + \Gamma_{31}^1 V^1 V^3 \\
&= \left(\frac{dV^1}{d\tau}\right)_{tr} + \Gamma_{11}^1 V^1 V^1 \\
&= \frac{d^2r}{d\tau^2} + \frac{A'(r)}{2A(r)} \left(\frac{dr}{d\tau}\right)^2
\end{aligned} \tag{323}$$

$$\begin{aligned}
\left(\frac{DV^2}{\partial\tau}\right)_{tr} &= \left(\frac{D}{d\tau}\frac{rd\theta}{d\tau}\right)_{tr} = \left(\frac{dV^2}{d\tau}\right)_{tr} + \Gamma_{0i}^2 V^i V^0 + \Gamma_{1i}^2 V^i V^1 + \Gamma_{2i}^2 V^i V^2 + \Gamma_{3i}^2 V^i V^3 \\
&= \left(\frac{dV^2}{d\tau}\right)_{tr} + \Gamma_{02}^2 V^2 V^0 + \Gamma_{12}^2 V^2 V^1 + \Gamma_{22}^2 V^2 V^2 + \Gamma_{32}^2 V^2 V^3 \\
&= \left(\frac{dV^2}{d\tau}\right)_{tr} + 0
\end{aligned}$$

$$= r \frac{d^2 \theta}{d\tau^2} + \frac{d\theta}{d\tau} \frac{dr}{d\tau} \quad (324)$$

$$\begin{aligned} \left(\frac{DV^3}{\partial\tau}\right)_{tr} &= \left(\frac{D}{d\tau} \frac{r \sin\theta d\varphi}{d\tau}\right)_{tr} = \left(\frac{dV^3}{d\tau}\right)_{tr} + \Gamma_{0i}^3 V^i V^0 + \Gamma_{1i}^3 V^i V^1 + \Gamma_{2i}^3 V^i V^2 + \Gamma_{3i}^3 V^i V^3 \\ &= \left(\frac{dV^3}{d\tau}\right)_{tr} + \Gamma_{03}^3 V^3 V^0 + \Gamma_{13}^3 V^3 V^1 + \Gamma_{23}^3 V^3 V^2 + \Gamma_{33}^3 V^3 V^3 \\ &= \left(\frac{dV^3}{d\tau}\right)_{tr} + 0 \end{aligned}$$

$$= r \sin\theta \frac{d^2 \varphi}{d\tau^2} + \sin\theta \frac{d\varphi}{d\tau} \frac{dr}{d\tau} + r \cos\theta \frac{d\varphi}{d\tau} \frac{d\theta}{d\tau} \quad (325)$$

$$\begin{aligned} \left(\frac{DV^0}{\partial\tau}\right)_{tr} &= \left(\frac{D}{d\tau} \frac{dt}{d\tau}\right)_{tr} = \left(\frac{dV^0}{d\tau}\right)_{tr} + \Gamma_{0i}^0 V^i V^0 + \Gamma_{1i}^0 V^i V^1 + \Gamma_{2i}^0 V^i V^2 + \Gamma_{3i}^0 V^i V^3 \\ &= \left(\frac{dV^0}{d\tau}\right)_{tr} + \Gamma_{10}^0 V^0 V^1 \\ &= \frac{d^2 t}{d\tau^2} + \frac{B'(r)}{2B(r)} \frac{dt}{d\tau} \frac{dr}{d\tau} \end{aligned} \quad (326)$$

In comparisons of the previous two calculations, we could find subtle nuance in that they are settled by different x^μ and V^μ . But they have really given the equivalent results, in that both of them could be transformed to uniform covariant derivatives $\frac{DV}{\partial\tau}$. The latter calculation is very easy to be done. The most important is that the comparison of calculations have verified the conclusions on inequality of mixed subscript Christoffel symbols. One could easily make a checking computation for the latter calculation so that to use the simplified expression of the latter to verified the solution of the previous. That will finally indicate the errors in classical theory, as well as that in Weinberg's calculations on geodesic equations.

Many efforts [11,12,17] have been made to attempt to prove the conservative principles after the equation of geodesic equations. They attempt to show that

$$r^2 \frac{d\varphi}{d\tau} = \text{const.} \quad (327)$$

$$\left(1 - \frac{r^*}{r}\right) \frac{dt}{d\tau} = \text{const.} \quad (328)$$

are of the results of the Equation (314), and the Equation (315).

It is easy to find that these works involve with errors. In comparison on the results of $\left(\frac{DV^2}{\partial\tau}\right)_{tr}$ and $\left(\frac{DV^3}{\partial\tau}\right)_{tr}$ in previous two kinds of strategies, we will find that the two derivatives do nothing with gravity influence, and they are just come from transformation of spherical coordinates so that any doctrines after that to form angular momentum conservative principle would be lack of supports. We will make further verifications in next sections that these two equations are all of false ones. In fact, the Equation (327) is not a correct form of angular momentum, and the Equation (328) does nothing with energy. We will see that, motion trajectories could be calculated based on neither zero covariant derivatives, nor the so-called the geodesic lines.

8.3. Classical Equations of Light Ray Deflection

It is indicated in some books that the Lagrangian relates to Euler-Lagrangian equation and geodesic equation [3,4]. It is trivial to continue the discussions on whatever of the origins. I will say that the Lagrangian for light rays is absolutely correct, because we will see that it is just the expression of composition of light speed components in covariant space. It is the reason that the Lagrangian is employed for the equation of matter's trajectories in most publications. In fact, velocity composition equation could be easily employed to solve the trajectory of Newtonian problems. But it should be

highlighted that the Lagrangian equation is not proper for massive matters, which will be presented in following discussions.

Incomprehensibly, traditional solving process for Lagrangian equations seem like to do nothing with geodesic line equation and covariant differentials. On the other side, we could find that those solved trajectories all involved with contra variant angular momentum conservation, which also indicates that the solved trajectories may be not real geodesic lines.

To take the problem of the light rays passing across the Sun for granted, as shown in Figure 18.




Figure 18. Light rays passing across the Sun.

8.3.1. Classical Equations after Traditional Settings

It is expected in classical theories that for the motion of light rays, the Lagrangian is zero

$$\mathcal{L} = -(1 - \frac{r^*}{r})c^2\dot{t}^2 + (1 - \frac{r^*}{r})^{-1}\dot{r}^2 + r^2\dot{\phi}^2 = 0 \quad (329)$$

where, Schwarzschild metrics has been concerned. Thereby, derivatives by proper time have been special expressed for conveniences as $\dot{t} = \frac{dt}{d\tau}$, $\dot{r} = \frac{dr}{d\tau}$ and $\dot{\phi} = \frac{d\phi}{d\tau}$, which will also be employed in following expressions.

Two items were always set to be constant in most publications [3,4] as

$$\frac{1}{2} \frac{\partial \mathcal{L}}{\partial \dot{\phi}} = r^2 \dot{\phi} = L \quad (330)$$

$$\frac{1}{2} \frac{\partial \mathcal{L}}{\partial \dot{t}} = (1 - \frac{r^*}{r})c^2 \dot{t} = c^2 E \quad (331)$$

Because of the multiplier r^2 used, L is really an equivalent contra variant angular momentum rather than a covariant one, so that this setting seems like to insert the contra variant conservation in. It is said that the setting of L does not actually coincide with general covariance. Even so, we will find out that it is exactly wrong setting together with the setting of constant E to have given not bad results, which perform two negatives making a positive.

Then the Lagrangian would be reformed to be

$$c^2 E^2 - \dot{r}^2 - (1 - \frac{r^*}{r}) \frac{L^2}{r^2} = 0 \quad (332)$$

Setting $r' = \frac{dr}{d\phi} = \frac{\dot{r}}{\dot{\phi}}$, $u = \frac{1}{r}$, and then $r' = \frac{dr}{d\phi} = -\frac{u'}{u^2}$, there is

$$\frac{c^2 E^2}{L^2} - u'^2 - (1 - r^* u) u^2 = 0 \quad (333)$$

Derivated by ϕ , it is

$$u'' + u = \frac{3}{2} r^* u^2 \quad (334)$$

At a farthest position the right hand item goes to zero, then the equation will have a solution $u = \sin\phi/b$, where b is a distance from solar center to the asymptotic line, that could be called asymptotic distance. Replacing the very small right hand item with the solution, there is

$$u'' + u = \frac{3r^*}{2b^2} \sin^2 \varphi \quad (335)$$

There is a particular solution

$$u_1 = \frac{3r^*}{4b^2} \left(1 + \frac{1}{3} \cos 2\varphi\right) \quad (336)$$

Then the approximate solution of Equation (334) is obtained

$$u = \frac{\sin \varphi}{b} + \frac{3r^*}{4b^2} \left(1 + \frac{1}{3} \cos 2\varphi\right) \quad (337)$$

For an infinite large r , the u is infinitesimal one, there is

$$\varphi_\infty = -\frac{r^*}{b} \quad (338)$$

Observational deflection of light rays passing by the edge of the Sun is

$$\Delta = 2|\varphi_\infty| \approx 1.75'' \quad (339)$$

where, $R_\odot$ is the solar radius.

At the position $\varphi = \frac{\pi}{2}$ and $r = R$, there is

$$\frac{1}{R} = \frac{1}{b} + \frac{3r^*}{4b^2} \left(1 - \frac{1}{3}\right) \quad (340)$$

or

$$b \approx R + \frac{r^*}{2} \quad (341)$$

That will bring about contradictions for the setting of Equation (330) that at peak point there is

$$L_R = r^2 \dot{\varphi} = Rc \quad (342)$$

while at a very far point, there is

$$L_\infty = r_\infty^2 \dot{\varphi} = bc = \left(R + \frac{r^*}{2}\right)c \neq L_R \quad (343)$$

It is said that the equations have been solved incorrectly. Might as well, we would find out an inevitable solution after the inappropriate setting of Equation (330) in next section.

8.3.2. Errors Hidden in the Solving Process

It is obvious that a wrong setting has been made in Equation (332), because in gravitational field, $(1 - \frac{r^*}{r})c^2 \dot{t}$ really varies with r . In fact, the Equation (329) could be solved directly only based on the setting of Equation (330) as following.

Considering $\dot{t} = \frac{dt}{d\tau} = (1 - \frac{r^*}{r})^{-1/2}$, the Lagrangian really is

$$-c^2 + (1 - \frac{r^*}{r})^{-1} \dot{r}^2 + r^2 \dot{\varphi}^2 = 0 \quad (344)$$

or

$$-c^2 + \dot{r}^2 + r^2 \dot{\varphi}^2 = 0 \quad (345)$$

It is really the composition of light speed in covariant space. Any equation that leads to $(1 - \frac{r^*}{r})^{-1} \dot{r}^2 + r^2 \dot{\varphi}^2 \neq c^2$ is unacceptable, as that has been done in Equation (332).

Transform the equation by multiplying a $(1 - \frac{r^*}{r})$

$$(1 - \frac{r^*}{r})c^2 - \dot{r}^2 - (1 - \frac{r^*}{r})r^2 \dot{\varphi}^2 = 0 \quad (346)$$

If setting peak point radius as R and setting the angular momentum at that point $L = Rc = r^2 \dot{\varphi} = \text{const}$, with $u = 1/r$, it becomes

$$(1 - r^*u)c^2 - u'^2 R^2 c^2 - (1 - r^*u)u^2 R^2 c^2 = 0 \quad (347)$$

To be derivated by φ , it is

$$u'' + u = \frac{3}{2}r^*u^2 - \frac{r^*}{2R^2} \quad (348)$$

With $\frac{3r^*}{2b^2} \sin^2 \varphi$ instead of $\frac{3}{2}r^*u^2$, it becomes

$$u'' + u = \frac{3r^*}{2b^2} \sin^2 \varphi - \frac{r^*}{2R^2} \quad (349)$$

Because $R \approx b$, it could be solved as

$$u = \frac{\sin \varphi}{b} + \frac{r^*}{4b^2} (1 + \cos 2\varphi) \quad (350)$$

One will obtain the solution of the equation as

$$\varphi_\infty = -\frac{r^*}{2b} \quad (351)$$

At the peak point as $\varphi = \frac{\pi}{2}$, we will get the constant b that

$$\frac{1}{R} = \frac{1}{b} + \frac{r^*}{4b^2} (1 - 1) \quad (352)$$

so that

$$b = R \quad (353)$$

And the setting of Equation (330) has been well kept that

$$L_R = r^2 \dot{\varphi} = Rc \quad (354)$$

and

$$L_b = r^2 \dot{\varphi} = bc = Rc = L_R \quad (355)$$

This is really the inevitable solution for classical equations but it is not a true result for realities. We have seen that the classical equations to have been solved to an answer Equation (337) accurately up to the observation results is just caused by the wrong settings of the declared energy momentum conservations of Equation (330) and Equation (331). We will see in next sections that the angular momentum Equation (347) is not correct as well.

8.4. Momentum, Energy, and Angular Momentum Conservativeness

We have drawn the conclusion that light momentum keeps conservation in contra variant space rather than covariant space, and then of course, so does the light mass energy. In fact, apparent light speed or so-called contra variant light speed may varies in contra variant space, but light momentum and energy will not be affected by apparent speeds. In fact, neither geodesic equation nor the derivations of Lagrangian could help proving the Equation (330) and Equation (331), in that Equation (330) and Equation (331) are substantially not correct.

Considering a light ray goes a vertical distance on the Earth, one could gain the mass variation as

$$m_r = \frac{h\nu_{0(\infty)}}{c^2} \left(1 + \frac{r^*}{2r}\right) = m_\infty \left(1 + \frac{r^*}{2r}\right) \quad (356)$$

It could be called simplified equation of mass in gravitational field.

Light momentum could be expressed as

$$P = \frac{h\nu_0}{c} = m_r c \quad (357)$$

or the momentum square

$$P^2 = m_r^2 c^2 \quad (358)$$

Case in contra variant space, apparent light speed varies with position so that that speed cannot be used in expressions of light momentum directly. The invariant light speed $c = \text{const.}$ could be employed to present invariance of momentum. Equation (357) performs full variation with gravity

by m_r , which is the performance of momentum conservation. In fact, light momentum depends on frequency, just as m_r does. If we ask more for a deep reason, that should be mass energy equation.

Case in covariant space, if light momentum will also be expressed by frequency, that will vary with bases additionally.

We know the Lagrangian

$$c^2 = (1 - \frac{r^*}{r})^{-1} \dot{r}^2 + r^2 \dot{\varphi}^2 \quad (359)$$

With Lagrangian substituted in conservative momentum square, it turns to be

$$P^2 = m_r^2 [(1 - \frac{r^*}{r})^{-1} \dot{r}^2 + r^2 \dot{\varphi}^2] \quad (360)$$

or

$$P^2 = m_r^2 [(1 - \frac{r^*}{r})^{-2} (\frac{dr}{dt})^2 + (1 - \frac{r^*}{r})^{-1} r^2 (\frac{d\varphi}{dt})^2] \quad (361)$$

As we have discussed, light speed cannot be directly composed in contra variant space but can be done in covariant space. This is a reason the Lagrangian is employed in conservative momentum square.

In one source fields, the momentum vector could be discomposed to be components of centripetal and tangent

$$\mathbf{P} = \mathbf{P}_c + \mathbf{P}_t \quad (362)$$

or

$$P^2 = P_c^2 + P_t^2 \quad (363)$$

Obviously, the tangent momentum relates to tangent velocity and centripetal momentum relates to centripetal velocity. It could also be inferred that the tangent component varies with the corresponding velocity, and so does the centripetal one.

So that there must be

$$P_c^2 = m_r^2 (1 - \frac{r^*}{r})^{-1} \dot{r}^2 = m_r^2 (1 - \frac{r^*}{r})^{-2} (\frac{dr}{dt})^2 \quad (364)$$

and

$$P_t^2 = m_r^2 r^2 \dot{\varphi}^2 = m_r^2 (1 - \frac{r^*}{r})^{-1} r^2 (\frac{d\varphi}{dt})^2 \quad (365)$$

In one source field, the angular momentum conservation could be expressed as

$$L^2 = r^2 P_t^2 = m_r^2 r^4 \dot{\varphi}^2 = m_r^2 (1 - \frac{r^*}{r})^{-1} r^4 (\frac{d\varphi}{dt})^2 = \text{const.} \quad (366)$$

It should be highlighted that we are talking about the moment conservation in contra variant space. It is amazing that the angular momentum should be expressed in the form of $r^2 m_r \dot{\varphi}$ or $r^2 m_r \frac{d\varphi}{dt}$ rather than the form of $r^2 m_r \frac{d\varphi}{d\tau}$, or we have seen that contra variant light momentum could be only directly composed in covariant form. The real reason is that the invariant light speed c is just employed for the expressions by invariance.

The issues of light momentum have always been one of the controversies in physics for more than a hundred years [18]. The problem is the difficulty of assessing the light momentum in transparent materials between Minkowski's equation [19] and Abraham's equation [20]. To one's surprise, we would have made the conclusion different from both of them, after the discussions in previous sections, because of Lorentz covariance.

Nevertheless, angular momentum brings about new surprises on it. We will find that the surprises not only have been picked up from the expressions, but also hide behind the kinematics of light propagations in gravitational fields. These efforts might bring about tiny contributions for the attempt to answer the question of Einstein about 'What are light quanta?' [21] I appreciate what Leonhardt has said that light continues to surprise [22].

8.5. Revisit Equations for Light Ray Trajectory

8.5.1. Renovation and Resolution

We have recognized that it is conservative principle that really controls the solutions. In fact, light rays in gravitational field may experience mass energy variation.

As has mentioned previous, the light mass at the peak point is

$$m_R = \frac{h\nu_{0(\infty)}}{c^2} \left(1 + \frac{r^*}{2R}\right) \quad (367)$$

And it varies at position r

$$m_r = \frac{h\nu_{0(\infty)}}{c^2} \left(1 + \frac{r^*}{2r}\right) \quad (368)$$

These two equations involve with the energy conservation in contra variant space rather than that in covariant space.

For a light ray passing by a one source field, there is the angular momentum conservation as

$$L = r^2 m_r \dot{\phi} = R m_R c = \text{const.} \quad (369)$$

It should be highlighted again that the contra variant light momentum has been expressed by c and $\dot{\phi}$ which are of covariant space quantities rather than c_0^μ and $\frac{d\phi}{dt}$ of contra variant ones. In fact, it could be proved that c_0^μ and $\frac{d\phi}{dt}$ cannot be taken to form momentum conservation, if one takes efforts to have a try. That is because momentum variation depends on m_r that perform the effect of gravity, or in another words, the gravity input energy into the m_r . The invariant light speed c employed reveals that light momentum variation really depends on frequency rather than real velocity, just as that light propagates in transparent materials. That perhaps is really surprise.

Considering the Equation (367) and Equation (368), Equation (369) will lead to

$$\dot{\phi} = \frac{R m_R c}{m_r r^2} = \frac{R c}{r^2} \frac{1 + \frac{r^*}{2R}}{1 + \frac{r^*}{2r}} \quad (370)$$

Set $B = (R + \frac{r^*}{2})$. In weak field, the item $1/(1 + \frac{r^*}{2r}) \approx (1 - \frac{r^*}{2r})$, thus

$$\dot{\phi} = \frac{Bc}{r^2} \left(1 - \frac{r^*}{2r}\right) \quad (371)$$

I prefer to present the Lagrangian again

$$-c^2 + \left(1 - \frac{r^*}{r}\right)^{-1} \dot{r}^2 + r^2 \dot{\phi}^2 = 0 \quad (372)$$

Define $r' = \frac{dr}{d\phi}$, so that $\dot{r} = \frac{dr}{d\phi} \frac{d\phi}{d\tau} = r' \dot{\phi}$. There is

$$-c^2 + \left(1 - \frac{r^*}{r}\right)^{-1} r'^2 \dot{\phi}^2 + r^2 \dot{\phi}^2 = 0 \quad (373)$$

Insert the Equation (371) into Equation (373), the latter becomes

$$-c^2 + \left(1 - \frac{r^*}{r}\right)^{-1} r'^2 \left[\frac{1}{r^2} \left(1 - \frac{r^*}{2r}\right)\right]^2 B^2 c^2 + r^2 \left[\frac{1}{r^2} \left(1 - \frac{r^*}{2r}\right)\right]^2 B^2 c^2 = 0 \quad (374)$$

With $\left(1 - \frac{r^*}{2r}\right)^2 \approx 1 - \frac{r^*}{r}$, it is

$$-1 + r'^2 \frac{B^2}{r^4} + \left(1 - \frac{r^*}{r}\right) \frac{B^2}{r^2} = 0 \quad (375)$$

With $u = 1/r$, and $r' = \frac{dr}{d\phi} = -\frac{u'}{u^2}$ it turns to be

$$-\frac{1}{B^2} + u'^2 + (1 - ur^*)u^2 = 0 \quad (376)$$

We have seen the similar form of Equation (333), but this equation comes from the settings of the real conservative principles.

Derivate this equation to be

$$u'' + u = \frac{3}{2}r^*u^2 \tag{377}$$

Considering r is very big value so that the last item is very small so that the solution will be close to horizontal line at near R positions, as following equation will present

$$u'' + u = 0 \tag{378}$$

It could be solved to be

$$u = \frac{\sin\varphi}{b} \tag{379}$$

It is a horizontal line with a perpendicular distance b to the center of the Sun.

The right item of Equation (377) could be replaced with the simple solution, because the deviation is also very small. So that the equation could be reformed to be

$$u'' + u = \frac{3r^*}{2b^2}\sin^2\varphi \tag{380}$$

Case at positions far from R , the right item could be an approximation of the right item of Equation (377) to close to zero.

Once again, we could obtain the solution for the differential equation that

$$u = \frac{\sin\varphi}{b} + \frac{3r^*}{4b^2}(1 + \frac{1}{3}\cos2\varphi) \tag{381}$$

And then the deflection angle

$$\varphi_\infty = -\frac{r^*}{b} \tag{382}$$

Case $\varphi = \frac{\pi}{2}$, there is

$$\frac{1}{R} = \frac{1}{b} + \frac{r^*}{2b^2} \tag{383}$$

Because $b \gg r^*$ it becomes

$$R \approx b - \frac{r^*}{2} \tag{384}$$

or

$$b = R + \frac{r^*}{2} \tag{385}$$

To verify the momentum conservativeness that

$$L_R = m_R Rc \tag{386}$$

At a position $r \gg R$

$$L_b = m_b bc = \frac{1+\frac{r^*}{2r_\infty}}{1+\frac{r^*}{2R}} m_R bc = \frac{1}{R+\frac{r^*}{2}} R m_R bc = m_R Rc = L_R \tag{387}$$

It seems that we have got the same results as that of classical equations. But the truth is that the solution is the results after the conclusions of momentum and mass conservations which completely others to that of classical theory and at the same time the assumptions of Equation (330) and Equation (331) are thoroughly given up. It is said that the angular momentum keep conservation rather than that in previous.

The most important is that the real kinematics of light propagation has been worked out.

8.5.2. Detailed Discussions on the Coordinates of the Light Ray Trajectory

The trajectories of light ray in gravitational fields have more details behind the previous solution. More discussions may help to discover more realities.

Further discussions are going to be sponsored to make more detailed analysis for understanding of two items, positions and angles.

Item 1 is to recognize various kinds of lines. The most important line is the real light ray that is emitted from a farthest star to the observers on the Earth. This line should be curved as it goes closely to the Sun. Prolonging the straight parts of the light ray, one will gain the crossed straight lines, the asymptotic lines of light trajectory. They will be parallel to the two radial coordinate vectors r_∞ left and r_∞ right. There is another important line is the straight line from farthest star to the observer, that will present the real star direction from observer to the star.

Item 2 is to recognize those angles. φ_∞ is the second coordinate of farthest point on the light ray. Because the Earth is far enough to the Sun, the elevation angle of observer view line could be seen as φ_∞ . And because the star is very very far from the Sun, the angle between straight line to the star and horizontal line could be also seen as φ_∞ . Thus, the total deflect angle of Δ_{observe} is approximate $2\varphi_\infty$. They could be shown in Figure 19.




Figure 19. Detailed relationships for angles of light rays and view lines and coordinates.

For the coordinate of the Earth, $u = 1/A_u$, where A_u is astronomical unit, the simple solution could be used

$$\frac{1}{A_u} \approx \frac{\varphi_\oplus}{b} \quad (388)$$

where, $\varphi_\oplus$ is second coordinate of the Earth.

Thus, there is

$$\varphi_\oplus \approx \frac{b}{A_u} \quad (389)$$

Because we know that $b \approx R_\odot$, where $R_\odot$ is solar radius, so that

$$\varphi_\oplus \approx 0.004654478 \text{ rad} \quad (390)$$

The height of the horizon line to light ray

$$CE = R - A_u \sin \varphi_\oplus \quad (391)$$

It is difficult to get a not bad accurate solution. However, we can turn to discuss the value of $h = BC$ instead. For the approximate of $\varphi_\infty \approx \frac{1}{2} \Delta_{\text{observer}}$, it could be gain

$$h \approx A_u \sin \varphi_\infty \approx 635 \text{ km} \quad (392)$$

in the upper height $h = CF$.

There is a very small difference between real ray trajectory and asymptotic line, the $\delta = EF$. For the condition that light rays run from farthest position to the position they pass by the Sun, there is the equation after angular momentum conservativeness

$$\left(1 + \frac{r^*}{2r_\infty}\right)b = \left(1 + \frac{r^*}{2R}\right)R \quad (393)$$

NB, b is perpendicular length to asymptotic line, which is the moment distance for farthest positions, and only in approximate cases, it could be seen as $R + \delta$. R is peak point radius, and it need not be determined to be $R_{\odot}$ in these discussions theoretically.

Then it is easy to obtain

$$b = R + \frac{r^*}{2} \quad (394)$$

Peak difference between R and b is

$$\delta = \frac{r^*}{2} = 1.477 \text{ km} \quad (395)$$

It is more difficult to investigate such a fine distance in practice that not only because of the observational accuracy but also due to the coordinating of the peak point of light ray. The development of very-long-baseline interferometry have the capability of measuring angular separations and changes in angles as small as 10^{-4} seconds of arc [23]. That shows probabilities for the quite good accuracy for fine angle measuring. This issue perhaps cannot be solved easily.

We can study the right branch of asymptotic line with an overlapped φ_{∞} as shown in Figure 20.




Figure 20. Asymptotic line with an overlapped φ_{∞} .

There is the asymptotic line equation as

$$r \sin(\varphi + |\varphi_{\infty}|) = b \quad (396)$$

or

$$r \sin(\varphi + \frac{r^*}{b}) = b \quad (397)$$

Case $\varphi = \frac{\pi}{2}$, we will see that the top point is just a little bit higher than horizontal line

$$r = b / \sin(\frac{\pi}{2} + \frac{r^*}{b}) \quad (398)$$

Case φ is not very close to $\frac{\pi}{2}$, considering $\frac{r^*}{b}$ is very small, an approximate form of sine function is

$$\sin(\varphi + \frac{r^*}{b}) \approx \sin \varphi + \frac{r^*}{b} \cos \varphi \quad (399)$$

Thus, the deformed equation of asymptotic line could be written as

$$r = \frac{b}{\sin \varphi + r^* \cos \varphi / b} \quad (400)$$

It should be mentioned that the equation $u = \sin \varphi / b$ or $r \sin \varphi = b$ is a horizontal line with an distance b to the Sun center.

For the case that a light passes closely by the edge of the sun, there might be some influences from solar corona. It is a good idea to left a distance from the edge, for example, the position of $1.5R_{\odot}$ or even further.

8.5.3. A Wrong Treatment for Light Propagations

If the angular momentum conservativeness would be insisted for light propagations as

$$L = Rm_Rc = r^2m_r \frac{d\varphi}{dt} \quad (401)$$

It is seemingly that $\frac{d\varphi}{dt}$ presents the contra variant angular velocity of photons which is expected to correspond to contra variant angular momentum. And moreover, it could be argued to take $Rm_Rc_0^\mu$ as expressions of contra variant angular momentum, that will exactly lead to conservation break as gravitational redshift is concerned.

Now for the equation Equation (401), there is

$$\frac{d\varphi}{dt} = \frac{Rm_Rc}{m_rr^2} = \frac{Rc}{r^2} \frac{1+\frac{r^*}{2R}}{1+\frac{r^*}{2r}} \quad (402)$$

Set $B = (R + \frac{r^*}{2})$. In weak field, the item $1/(1 + \frac{r^*}{2r}) \approx (1 - \frac{r^*}{2r})$, and $(1 - \frac{r^*}{2r})^2 \approx 1 - \frac{r^*}{r}$, thus

$$\frac{d\varphi}{dt} = \frac{(1 - \frac{r^*}{2r})Bc}{r^2} \quad (403)$$

Because the Lagrangian with coordinate time is

$$-c^2 + (1 - \frac{r^*}{r})^{-2} (\frac{dr}{dt})^2 + (1 - \frac{r^*}{r})^{-1} r^2 (\frac{d\varphi}{dt})^2 = 0 \quad (404)$$

It could be transformed to be

$$c^2 - (1 - \frac{r^*}{r})^{-1} r'^2 \frac{B^2 c^2}{r^4} - \frac{B^2 c^2}{r^2} = 0 \quad (405)$$

or

$$(1 - \frac{r^*}{r}) - r'^2 \frac{B^2}{r^4} - (1 - \frac{r^*}{r}) \frac{B^2}{r^2} = 0 \quad (406)$$

or

$$(1 - r^*u) - u'^2 B^2 - (1 - r^*u)u^2 B^2 = 0 \quad (407)$$

To be derivated to be

$$u'' + u = \frac{3}{2} r^* u^2 - \frac{r^*}{2B^2} \quad (408)$$

Considering $b \approx B$, it could be solved as

$$u = \frac{\sin\varphi}{b} + \frac{r^*}{4b^2} + \frac{r^*}{4b^2} \cos 2\varphi \quad (409)$$

One will obtain that

$$\varphi_{\infty} = -\frac{r^*}{2b} \quad (410)$$

One can verify the peak point on the case of $\varphi = \pi/2$, and $r = R$, and then will gain

$$b = R \quad (411)$$

It is obvious that this solution of b cannot keep the conservation of angular momentum. The main reason is that the angular velocity defined in the angular momentum is not the direct component of light speed in Equation (404). As a comparison, that in Equation (372) is. Naturally, the solution in Equation (411) is a wrong answer for light ray propagations after a wrong setting.

8.6. Trajectories of Massive Matters

8.6.1. Discussions on Lagrangian

It is believed in classical theory that Lagrangian for massive matters have a special form [3,10] that others to that of light rays that

$$\mathcal{L} = -(1 - \frac{r^*}{r})\dot{t}^2 c^2 + (1 - \frac{r^*}{r})^{-1}\dot{r}^2 + r^2\dot{\phi}^2 = -c^2 \quad (412)$$

As has discussed that for the first item in Equation (412) $\dot{t}^2 = (dt/t\tau)^2 = (1 - \frac{r^*}{r})^{-1}$, it will naturally lead to

$$-c^2 + (1 - \frac{r^*}{r})^{-1}\dot{r}^2 + r^2\dot{\phi}^2 = -c^2 \quad (413)$$

So that

$$(1 - \frac{r^*}{r})^{-1}\dot{r}^2 + r^2\dot{\phi}^2 \equiv 0 \quad (414)$$

It is totally unacceptable that the covariant velocity results in absolute zero. But we know that the two settings of Equation (330) and Equation (331) seem to have solved the problem. Consequently, the Equation (412) will be then reformed to be

$$(1 - \frac{r^*}{r})^{-1}c^2 E^2 - (1 - \frac{r^*}{r})^{-1}\dot{r}^2 - \frac{L^2}{r^2} = c^2 \quad (415)$$

It is the traditional form of dynamic equation for massive matters. Of course, it is of fantastic expressions.

On another side, even the original Lagrangian cannot work for massive matters

$$\mathcal{L} = -c^2 + (1 - \frac{r^*}{r})^{-1}\dot{r}^2 + r^2\dot{\phi}^2 = 0 \quad (416)$$

in that for massive matters we know that the last two items are definitely unequal to c^2 in most cases.

In fact, you know there is no more appropriate dynamic equation for massive matters better than that

$$(V_0^s)^2 - (\frac{dr}{dt})^2 - r^2(\frac{d\phi}{dt})^2 = 0 \quad (417)$$

where, V_0^s is modulus of space velocity of massive matters.

The Equation (417) really performs velocity composition rather than the Equation (415) and Equation (416). The covariant velocity composition could be shown as

$$(Vs/0)^2 = \dot{\rho}^2 + r^2\dot{\phi}^2 \quad (418)$$

But it is difficult to be employed for dynamic equations, in that the velocity $Vs/0$ is not conservative quantity so that it cannot be employed in expressions of kinetic energy.

As has discussed above, it is easily to carried out the momentum equation for massive matters that

$$P^2 = m_r^2(\frac{dr}{dt})^2 + m_r^2(\frac{rd\phi}{dt})^2 \quad (419)$$

and the angular momentum equations for massive matters that

$$L = Rm_R V_0^s = r^2 m_r \frac{d\phi}{dt} \quad (420)$$

where, $V_0^s = V_0^\phi$ at the peak point.

8.6.2. Classical Solution for Planet Orbits

As the two settings adopt for the Lagrangian Equation (412), there will be

$$c^2 E^2 - \dot{r}^2 - (1 - \frac{r^*}{r})\frac{L^2}{r^2} = (1 - \frac{r^*}{r})c^2 \quad (421)$$

One could find that the main item of c^2 might sounds literally unreasonable, although the problem will not bring about great obstacles in process of calculation, because constant items in Equation (421) will be eliminated by a further derivation. Never mind, we will promptly pick up the problem in the issue of time delay in next section.

Set $u = 1/r$ and with $\dot{r} = \frac{dr}{d\phi} \frac{d\phi}{d\tau}$ and $r' = \frac{dr}{d\phi} = -\frac{u'}{u^2}$ it could be reformed as

$$\frac{c^2 E^2}{L^2} - u'^2 - (1 - r^* u)u^2 = (1 - r^* u)\frac{c^2}{L^2} \quad (422)$$

To act further derivation on it, there is

$$u'' + u = \frac{3}{2}r^*u^2 + \frac{r^*c^2}{2L^2} \quad (423)$$

For a planet orbit, $L = RV\varphi/0 \ll Rc$, so that in right side of the equation the item $\frac{3}{2}r^*u^2$ is a small quantity, where, $RV\varphi/0$ might be allowable expression in traditional settings. Zero order approximation of the equation is

$$u'' + u = \frac{r^*c^2}{2L^2} \quad (424)$$

It is the famous Binet equation. That has a solution

$$u = A(1 - e\sin\varphi) \quad (425)$$

where, $A = \frac{r^*c^2}{2L^2}$ and e is orbit eccentricity.

The item $\frac{3}{2}r^*u^2$ could be replaced by the solution approximately with eccentricity not very big

$$\frac{3}{2}r^*u^2 = \frac{3}{2}r^*A^2(1 - e\sin\varphi)^2 = \frac{3}{2}r^*A^2(1 - 2e\sin\varphi + e^2\sin^2\varphi) \quad (426)$$

Then, the Equation (423) becomes

$$u'' + u = A + \frac{3}{2}r^*A^2 - 3r^*A^2e\sin\varphi + \frac{3}{2}r^*A^2e^2\sin^2\varphi \quad (427)$$

Considering the conditions of very small eccentricity, in right side, the last item as a variable in absolute value is far less than the third item, as well as the second item as constant is far less than the first item. We would have an approximate form of the equation

$$u'' + u \approx A - 3r^*A^2e\sin\varphi \quad (428)$$

It should be noted, that there will be more conditions in which the Equation (428) might be invalid in that they may have greater eccentricity, such as some of asteroids and comets in solar system would be. In the case of great eccentricity, one must totally solve the Equation (427).

Any way, we could try to solve the split equations of Equation (428) of small eccentricity that

$$u'' + u = A \quad (429)$$

with

$$u_1 = A(1 - e\sin\varphi) \quad (430)$$

and the other equation

$$u'' + u = -3r^*A^2e\sin\varphi \quad (431)$$

with

$$u_2 = \frac{3}{2}r^*A^2e\varphi\cos\varphi \quad (432)$$

There is the general solution

$$u = u_1 + u_2 = A[1 - e\sin\varphi + \frac{3}{2}r^*Ae\varphi\cos\varphi] \quad (433)$$

where $\Delta\varphi = -\frac{3}{2}r^*Ae\varphi$ as absolute value is far less than φ . By Taylor equation or sine equation of two angles, we have

$$\sin(\varphi + \Delta\varphi) \approx \sin\varphi + \Delta\varphi\cos\varphi \quad (434)$$

So that

$$u \approx A - A\sin[(1 - \frac{3}{2}r^*A)\varphi] \quad (435)$$

This equation presents the perihelion precession that

$$\Delta\varphi = \frac{3}{2}r^*A\varphi \quad (436)$$

Thus, the precession in a revolution

$$\Delta\varphi_{2\pi} = 3r^*A \quad (437)$$

as shown in Fig. 21




Fig. 21 The trajectory of planet

8.6.3. Traditional Treatment for Close-to-Light-Speed Particles

Studies on the motion of close-to-light-speed particles in gravitational fields rarely have been seen in any publications probably due to sophisticated situations. I decide to study the issue firstly on traditional methodology is because that more faults would be found out in the process that may forcefully support the conclusions in section 7.

It is believed in traditional methodologies that the Lagrangian Equation (412) also could be applied in the dynamics of the motion of close-to-light-speed particles. So as well, the differential equation will be gained as same as Equation (423) that

$$u'' + u = \frac{3}{2}r^*u^2 + \frac{r^*c^2}{2L^2} \quad (438)$$

But it should be highlighted that the Equation (438) will be far different from Equation (423) in that the angular momentum in Equation (438) is $L = Rc$ which is far greater than that in Equation (423). Thus, one will see that the two items in Equation (438) is in same order of magnitudes while the first item in Equation (423) is a very small quantity with respect to the latter item.

Using light speed c to perform close-to-light-speed is an approximate treatment in numerical analysis.

We know that the trajectory is very close to the horizontal line $u = \sin\varphi/b$ at positions that r is not very farther than R . Otherwise at positions that r is very farther than R , the item $\frac{3}{2}r^*u^2$ will go to zero. Thus, the equation could be reformed to be

$$u'' + u = \frac{3r^*}{2b^2}\sin^2\varphi + \frac{r^*c^2}{2L^2} \quad (439)$$

The reform will not greatly influence the result of the original equation in that in very farther area $\frac{3r^*}{2b^2}\sin^2\varphi$ together with $\frac{3}{2}r^*u^2$ both are very small with respect to $\frac{r^*c^2}{2L^2}$.

There is the solution

$$u = \frac{\sin\varphi}{b} + \frac{5r^*}{4b^2}\left(1 + \frac{1}{5}\cos 2\varphi\right) \quad (440)$$

And then the deflection angle

$$\varphi_\infty = -\frac{3r^*}{2b} \quad (441)$$

Case $\varphi = \frac{\pi}{2}$, there is

$$\frac{1}{R} = \frac{1}{b} + \frac{r^*}{b^2} \quad (442)$$

Because $b \approx R \gg r^*$ it becomes

$$R \approx b - r^* \quad (443)$$

or

$$b = R + r^* \quad (444)$$

The asymptotic line will be

$$r = \frac{b}{\sin\varphi + \frac{3r^*}{2b}\cos\varphi} \quad (445)$$

To verify the issue of momentum conservativeness that

$$L_R = Rc \quad (446)$$

At a position $r \gg R$

$$L_b = bc = (R + r^*)c \neq L_R \quad (447)$$

The deflection angle seems something strange. The peak difference $\delta = r^*$ shows non-conservation for angular momentum. This result is not far beyond expectation in that this traditional

solution involves with errors. We will see more detailed problems on the trajectory in the study of corresponding time delay in next sub-sections.

8.6.4. Renovated Equations for Close-to-Light-Speed Massive Particles

We have got the conclusion that massive particles will not run with general covariance. They run with Newtonian laws. The velocity composition for close-to-light-speed massive particles really is

$$-c^2 + \left(\frac{dr}{dt}\right)^2 + r^2 \left(\frac{d\varphi}{dt}\right)^2 = 0 \quad (448)$$

The conservativeness of angular momentum is as discussed previous

$$L = R m_R c = r^2 m_r \frac{d\varphi}{dt} \quad (449)$$

One will find that the momentum defined is rare different from the Lagrangian for light rays. The constant c is not the speed of light while it really is the approximate speed of massive matter. So that the c in Equation (448) is just a velocity composition of close-to-light-speed motion. The Equation (448) and Equation (449) perform the differences of conservations between light fly and the motion of massive matter. We will see these differences in the equations of low velocity motions of massive matters in next sub-sections.

The angular velocity

$$\frac{d\varphi}{dt} = \frac{R m_R c}{m_r r^2} = \frac{R c}{r^2} \frac{1 + \frac{r^*}{2R}}{1 + \frac{r^*}{2r}} \quad (450)$$

Set $B = (R + r^*/2)$ and in weak field, the item $1/(1 + r^*/2r) \approx (1 - r^*/2r)$, and $(1 - r^*/2r)^2 \approx (1 - r^*/r)$. Thus

$$\frac{d\varphi}{dt} = \frac{1}{r^2} \left(1 - \frac{r^*}{2r}\right) B c \quad (451)$$

Together with $\frac{dr}{dt} = \frac{dr}{d\varphi} \frac{d\varphi}{dt} = r' \frac{d\varphi}{dt}$, the Equation (448) turns to be

$$c^2 - \left(1 - \frac{r^*}{r}\right) r'^2 \frac{B^2 c^2}{r^4} - \left(1 - \frac{r^*}{r}\right) \frac{B^2 c^2}{r^2} = 0 \quad (452)$$

or

$$\left(1 + \frac{r^*}{r}\right) - r'^2 \frac{B^2}{r^4} - \frac{B^2}{r^2} = 0 \quad (453)$$

With $u = \frac{1}{r'}$ and $r' = \frac{dr}{d\varphi} = -\frac{u'}{u^2}$, there is

$$(1 + r^* u) - u'^2 B^2 - u^2 B^2 = 0 \quad (454)$$

To be differentiated as

$$u'' + u = \frac{r^*}{2B^2} \quad (455)$$

Considering $b \approx B$, it could be solved as

$$u = \frac{\sin\varphi}{b} + \frac{r^*}{2b^2} \quad (456)$$

Obviously, there is

$$\varphi_\infty = -\frac{r^*}{2b} \quad (457)$$

It is easy to calculate in Equation (456) that at the peak point, $\varphi = \pi/2$, and $r = R$, so that

$$b = R + \frac{r^*}{2} \quad (458)$$

Or with angular momentum conservation there is

$$(1 + \frac{r^*}{2r_\infty})b = (1 + \frac{r^*}{2R})R \tag{459}$$

Again, we obtain

$$b = R + \frac{r^*}{2} \tag{460}$$

The asymptotic line

$$r = \frac{b}{\sin\varphi + r^* \cos\varphi / b/2} \tag{461}$$

It is said that the trajectories of light rays and close-to-light-speed particles present different φ_∞ but they present the same peak difference δ . The same peak difference is because of the result of momentum conservation.

In fact, these studies have help to create the dynamics of close-to-light-speed particles in gravitational field. By the way, one can compare the solution Equation (456) with the Equation (440) in section 8.5.3 that they have given far different result of φ_∞ and different value of b , because of errors involved in the Equation (438).

8.6.5. Renovated Equations for Un-Close-to-Light-Speed Massive Matters

Let us investigate an un-close-to-light-speed motion for massive matters from specific position R to another position r . Considering a variable dynamic mass m_r , the gravity

$$F = -\frac{GMm_r}{r^2} = -m_r c^2 \frac{r^*}{2r^2} \tag{462}$$

Dynamic energy converted from gravitational potential by a free motion from R to r is

$$E_{conv} = -\int_R^r m_\rho c^2 \frac{r^*}{2\rho^2} d\rho \tag{463}$$

and the total energy

$$E = m_r c^2 = m_R c^2 + E_{conv} \tag{464}$$

where, dynamic mass m_r and m_R correspond to possible velocities.

The function of total energy could be written as

$$m_r c^2 = m_R c^2 - \int_R^r m_\rho c^2 \frac{r^*}{2\rho^2} d\rho \tag{465}$$

Setting energy at position R as a known quantity, this equation could be derivated to be

$$m_r' = -m_r \frac{r^*}{2r^2} \tag{466}$$

It could be integrated to be

$$\ln m_r - \ln m_R = \frac{r^*}{2r} - \frac{r^*}{2R} \tag{467}$$

Then we gain the expression of variable mass that

$$m_r = m_R e^{(\frac{r^*}{2r} - \frac{r^*}{2R})} = m_\infty e^{\frac{r^*}{2r}} \tag{468}$$

where, m_R and m_r are dynamic mass corresponding to specific velocities, and m_∞ is defined as an imaginary concept corresponding to a reduction mass at a farthest position. This equation is general expression of mass variation with respect to the simplified expression in Equation (356). Particularly for massive matters in some cases the mass m_∞ could have imaginary velocity. This equation could be called general mass equation in gravitational field.

And we would like to carry out the expanded expression of gravity equation that

$$F = -m_r c^2 \frac{r^*}{2r^2} = -m_R c^2 \frac{r^*}{2r^2} e^{(\frac{r^*}{2r} - \frac{r^*}{2R})} = -m_\infty c^2 \frac{r^*}{2r^2} e^{\frac{r^*}{2r}} \tag{469}$$

One could find that the relativistic mass may not keep mass conservation any more that may surprise us, but it really has been revealed in realities. This is another kind of comprehensive physics which I will not make further discussions in this section. Mass of matter does matter [26]. The truths stay in realities.

Dynamic energy will be converted during a moving of a matter that

$$E_{conv} = m_r c^2 - m_R c^2 = m_R c^2 [e^{(\frac{r^*}{2r} - \frac{r^*}{2R})} - 1] \quad (470)$$

It should be pointed out that the expression of exponential equations of mass variation of course could be used for light ray propagations and close-to-light-speed massive particles for higher accuracy analysis case in strong fields.

For massive matters, with special relativity, the equation of dynamic energy at a position r is

$$E_k = \xi m_r V_r^2 \quad (471)$$

where, $\xi = (1 - \sqrt{1 - \frac{V_r^2}{c^2}}) \frac{c^2}{V_r^2}$, see Equation (300).

In gravitational field the dynamic energy varies with position that

$$E_k = \xi_r m_r V_r^2 = \xi_R m_R V_R^2 + E_{conv} = \xi_R m_R V_R^2 + m_R c^2 [e^{(\frac{r^*}{2r} - \frac{r^*}{2R})} - 1] \quad (472)$$

Conversion energy E_{conv} could be positive case potential release or negative case potential withdrawn. However, dynamic energy E_k will be always greater than zero, so that the variable r will be limited in some specific cases that depends on initial conditions.

NB, for the convenience of expressions, the tensors of velocities, frequencies or the components maybe not written in tensor format anymore case they may not bring about confusions for understandings, for examples, V_r and V_R refer to the velocity at position r and R .

It should be further discussed here that we have seen dynamic mass energy may come from the release of gravitational potential. That will then perform as inertial mass and gravitational mass. If in two source system, they move closer or farther will cause mass increase or lose in that we incline to realize that potential does not act as mass. We are not sure that in this condition mass conservativeness is available or not. I am inclining to say no. Maybe this discussion involves with new physics. In this section, this controversy does not really matter. This discussion just presents the issue for more focus.

We now prefer to study the conditions of irrelativistic velocities so that $\xi_r \approx \xi_R \approx 0.5$, and after Equation (468) and Equation (472), there is the velocity square

$$\begin{aligned} V_r^2 &= e^{(\frac{r^*}{2R} - \frac{r^*}{2r})} V_R^2 + 2e^{(\frac{r^*}{2R} - \frac{r^*}{2r})} [e^{(\frac{r^*}{2r} - \frac{r^*}{2R})} - 1] c^2 \\ &= e^{(\frac{r^*}{2R} - \frac{r^*}{2r})} V_R^2 + 2[1 - e^{(\frac{r^*}{2R} - \frac{r^*}{2r})}] c^2 \end{aligned} \quad (473)$$

As we have discussed in section 8.6.1, Lagrangian could no longer be employed for analysis on the motion of massive matters. After conclusions and inferences previous and further discussions on massive matter's motion in one source gravitational field, the velocity composition could be expressed as

$$-V_r^2 + \left(\frac{dr}{dt}\right)^2 + r^2 \left(\frac{d\varphi}{dt}\right)^2 = 0 \quad (474)$$

These considerations could also be performed for that of close-to-light-speed massive particles as $V_r \rightarrow c$, as Equation (448).

With angular momentum conservativeness, there is

$$r^2 m_r \frac{d\varphi}{dt} = R m_R V_R = \text{const.} \quad (475)$$

where, R is radius of the perihelion and V_R is the corresponding velocity and that is vertical to the radius.

Of course, the angular momentum conservation for lower velocity motion will be the same as that of close-to-light-speed massive matters. But they are really different from that of light rays.

So that

$$\frac{d\varphi}{dt} = \frac{R}{r^2} \frac{m_R}{m_r} V_R = \frac{R}{r^2} V_R e^{\left(\frac{r^*}{2R} - \frac{r^*}{2r}\right)} \quad (476)$$

Together with the Equation (473) and $\frac{dr}{dt} = \frac{dr}{d\varphi} \frac{d\varphi}{dt} = r' \frac{d\varphi}{dt}$, the Equation (474) becomes

$$-e^{\left(\frac{r^*}{2R} - \frac{r^*}{2r}\right)} V_R^2 - 2[1 - e^{\left(\frac{r^*}{2R} - \frac{r^*}{2r}\right)}]c^2 + (r')^2 \frac{1}{r^4} e^{\left(\frac{r^*}{R} - \frac{r^*}{r}\right)} R^2 V_R^2 + \frac{1}{r^2} e^{\left(\frac{r^*}{R} - \frac{r^*}{r}\right)} R^2 V_R^2 = 0 \quad (477)$$

or

$$-e^{\left(\frac{r^*}{2r} - \frac{r^*}{2R}\right)} V_R^2 - 2[e^{\left(\frac{r^*}{r} - \frac{r^*}{R}\right)} - e^{\left(\frac{r^*}{2r} - \frac{r^*}{2R}\right)}]c^2 + (r')^2 \frac{1}{r^4} R^2 V_R^2 + \frac{1}{r^2} R^2 V_R^2 = 0 \quad (478)$$

Setting $u = 1/r$, it is

$$-e^{\left(\frac{r^*}{2}u - \frac{r^*}{2R}\right)} V_R^2 - 2[e^{(ur^* - \frac{r^*}{R})} - e^{\left(\frac{r^*}{2}u - \frac{r^*}{2R}\right)}]c^2 + (u')^2 R^2 V_R^2 + u^2 R^2 V_R^2 = 0 \quad (479)$$

To be derivated by φ , it becomes

$$-\frac{r^*}{4} e^{\left(\frac{r^*}{2}u - \frac{r^*}{2R}\right)} V_R^2 - [r^* e^{(ur^* - \frac{r^*}{R})} - \frac{r^*}{2} e^{\left(\frac{r^*}{2}u - \frac{r^*}{2R}\right)}]c^2 + u'' R^2 V_R^2 + u R^2 V_R^2 = 0 \quad (480)$$

All indexes in exponent functions are small quantities, it is better to take the one-order approximation so that the equation becomes

$$-\frac{r^*}{4} \left(1 + \frac{r^*}{2}u - \frac{r^*}{2R}\right) V_R^2 - [r^* (1 + ur^* - \frac{r^*}{R}) - \frac{r^*}{2} (1 + \frac{r^*}{2}u - \frac{r^*}{2R})]c^2 + u'' R^2 V_R^2 + u R^2 V_R^2 = 0 \quad (481)$$

The exponent forms used here is to present primary expressions of mass energy in gravitational fields. In the conditions of strong fields, the exponent forms may take forceful effects. One could imagine that the kinematics of light or close-to-light particles could also be expressed in primary ways in strong fields.

In weak fields, those very small items in Equation (481) will be neglected. That becomes

$$-\frac{3}{4} r^{*2} c^2 u - \frac{1}{2} r^* c^2 + u'' R^2 V_R^2 + u R^2 V_R^2 = 0 \quad (482)$$

or

$$u'' + \left(1 - \frac{3}{4} \frac{r^{*2} c^2}{R^2 V_R^2}\right) u = \frac{1}{2} \frac{r^* c^2}{R^2 V_R^2} \quad (483)$$

Setting

$$\omega^2 = 1 - \frac{3}{4} \frac{r^{*2} c^2}{R^2 V_R^2} \quad (484)$$

For planets in solar system, the last item in Equation (484) is far less than 1.0, so that

$$\omega \approx 1 - \frac{3}{8} \frac{r^{*2} c^2}{R^2 V_R^2} \quad (485)$$

For planet orbits, $0 < \omega \leq 1$,

The equation could be solved as [27]

$$u = A[1 - e \sin(\omega\varphi)] \quad (486)$$

where, e is orbit eccentricity.

With this solution substituted back into Equation (483)

$$A = \frac{1}{2} \frac{r^* c^2}{R^2 V_R^2} \omega^{-2} \quad (487)$$

One will see that A in this equation is a little greater than that in Equation (435).

One-order approximation of Equation (479) is

$$-(1 + \frac{r^*}{2}u - \frac{r^*}{2R})V_R^2 - 2[(1 + ur^* - \frac{r^*}{R}) - (1 + \frac{r^*}{2}u - \frac{r^*}{2R})]c^2 + (u')^2 R^2 V_R^2 + u^2 R^2 V_R^2 = 0 \quad (488)$$

Neglecting very small items, it is

$$-V_R^2 + \frac{r^*}{R}c^2 - r^*uc^2 + (u')^2 R^2 V_R^2 + u^2 R^2 V_R^2 = 0 \quad (489)$$

With Equation (486), it becomes

$$-V_R^2 + \frac{r^*}{R}c^2 - r^*Ac^2 + r^*A\sin(\omega\varphi)c^2 + A^2\omega^2e^2[\cos(\omega\varphi)]^2 R^2 V_R^2 + (A - A\sin(\omega\varphi))^2 R^2 V_R^2 = 0 \quad (490)$$

where, e is orbit eccentricity.

Case $\varphi = 0$ there is

$$-V_R^2 + \frac{r^*}{R}c^2 - r^*Ac^2 + A^2\omega^2e^2 R^2 V_R^2 + A^2 R^2 V_R^2 = 0 \quad (491)$$

where, e is orbit eccentricity.

This equation is the velocity composition equation at the position of $\varphi = 0$. The first three items correspond to the component of minus total systematic dynamic energy, in which, the previous one is of the initial condition and the others come from potential conversion. The last item is of the component from angular dynamic energy. The fourth item is of the component from radial dynamic energy.

So that the eccentricity is

$$e \approx [\frac{(1-A^2R^2)V_R^2 - (1-AR)\frac{r^*}{R}c^2}{A^2\omega^2R^2V_R^2}]^{1/2} \quad (492)$$

Some publications have given the following expression for eccentricity

$$L^2 = a(1 - e^2)GM \quad (493)$$

where, L has been set the angular momentum, a is the long semi-major axis, G is gravitational constant and M is the mass of central source. But this is not a real solution for eccentricity, because that the a solved in Equation (486) is $a = A^{-1}(1 - e^2)^{-1}$ which is a reformation of Equation (493). The eccentricity e cannot be solved with a by the Equation (493).

We will then prefer to present the perihelion precession of

$$\Delta\varphi \approx \frac{3}{8} \frac{r^{*2}c^2}{R^2V_R^2} \varphi \approx \frac{3}{4} r^* A \varphi \quad (494)$$

or the precession per revolution

$$\Delta\varphi_{2\pi} \approx \frac{3\pi}{4} \frac{r^{*2}c^2}{R^2V_R^2} = \frac{3\pi}{2} r^* A \varphi \quad (495)$$

That could be shown as Fig. 22.




Fig. 22 The renovated trajectory of planet

The most surprising is that this solution of perihelion precession is non-general-relativistic. It is a half of the value of the classical solution. It is practicable to carry out more experiments of motions closely around the Sun to verified the conclusions of Equation (494).

One can focus on more sophisticated conditions for massive matters traveling in gravitational fields, based on the Equation (473), the Equation (480) or the Equation (483), especially for those motions in strong fields and with relativistic velocities.

Perhaps this result is the only one in which we have focused on that is in contradictions with the observations that have declared perihelion precessions of planets in solar system [17]. But some observations on PSR J0737-3039A/B [28,29] have shown quite big deviations from classical predictions, which had been expected to have perihelion precession together with geodetic precession. Nevertheless, the experiment of Gravity Probe B [30] shows that geodetic item has got accurate results while frame-dragging item has not, in which it could be recognized that the geodetic item is really the effect of special relativity. Observations on PSR 1913+16 [31] have provided more evidences on energy conversions rather than perihelion precessions. That have not been well done for the perihelion precessions of S2 and S62 [32,33] in our galaxy. It could be expected that more observations on them and so on would be sponsored to furtherly verify the result of perihelion precession.

8.6.6. Dynamics within Event Horizon

It could be predicted that a massive matter with initial velocity high enough will have opportunities to escape from a black hole.

We have known that massive matters have velocity limit of light speed. That indicates a minimum radius for circular motions in one source fields. It has been discussed in previous that vertical component of gravity would lead to Newtonian geometrical acceleration at that direction. This will then support the equation of centrifugal motion case the matter catches up to the velocity close to light speed. The equilibrium equation could be written as

$$m \frac{c^2}{r} = \frac{GMm}{r^2} \quad (496)$$

The minimum radius

$$r_{min} = \frac{GM}{c^2} = \frac{r^*}{2} \quad (497)$$

It indicates that there will be three conditions for the close-to-light-speed motion on close to $r^*/2$ circle. Condition 1 is that the matter move tangent to the circle under which the matter keep circular motion. Condition 2 is that the matter move with inner inclined direction to the tangent line at the start time, in which the matter cannot keep periodic motion and will fall to source center finally. Condition 3 is that at the initial state, the motion direction is outer inclined to tangent line, which could lead to complex results dependently.

It could be imagined that the dynamics in all the regions within event horizon and close outer space will be greatly sophisticated. I would like to make more discussions on radial escape. With the Equation (487), the total mass for a matter escape to event horizon from inner area is

$$m_r = m_0 e^{\left(\frac{r^*}{2r} - \frac{r^*}{2r^*}\right)} = m_0 e^{\left(\frac{r^*}{2r} - \frac{1}{2}\right)} \quad (498)$$

and the mass to escape to free

$$m_r = m_0 e^{\left(\frac{r^*}{2r} - \frac{r^*}{2r_\infty}\right)} = m_0 e^{\frac{r^*}{2r}} \quad (499)$$

where, m_0 is defined due to velocities after escape that have been set zero.

One can calculate the minimum escape energy for matters at different positions. A brief calculation is shown in table 4.

Table 4. Energy for matters to escape from within event horizon

from Initial positions of times of r^*	for escape to event horizon		or escape to free	
	dynamic mass	velocity/c	dynamic mass	velocity/c
$0.1r^*$	$90.017m_0$	0.9999383	$148.413m_0$	0.9999773
$0.2r^*$	$7.389m_0$	0.99080	$12.182m_0$	0.99663
$0.5r^*$	$1.649m_0$	0.795	$2.718m_0$	0.930
$0.9r^*$	$1.057m_0$	0.324	$1.743m_0$	0.819

8.7. Time Spending Problems

For a light ray passing by the Sun, the apparent light speed varies with the positions and directions on the trajectory. There should be a difference between the real time interval and that calculated via invariant light speed instead of apparent light speed. Shapiro proposed new tests of time delay of radar signals which transmitted from the Earth to pass by the edge of the Sun to another planet or satellite and then they would be reflected back to the Earth [24,25]. The observations on time delay of radar echoes would forcefully support the theory of general relativity as well as that of light ray deflect.

However, the solutions of time delay must have involved in the problems with light trajectory, that the solution process has inherited the errors in classical equations of light trajectory, so that it is necessary to make a detailed discussion to rectify.

8.7.1. Classical Solution for Radar Echoes

In classical procedure, with the assumptions of $\dot{t}(1 - \frac{r^*}{r}) = E$ and $r^2\dot{\phi} = L$, the Lagrangian could be transformed to be

$$\dot{r}^2 = c^2E^2 - (1 - \frac{r^*}{r})\frac{L^2}{r^2} \tag{500}$$

as that in Equation (332).

And also with

$$\dot{r} = \frac{dr}{dt}\dot{t} = \frac{dr}{dt}E(1 - \frac{r^*}{r})^{-1} \tag{501}$$

The Lagrangian becomes

$$(1 - \frac{r^*}{r})^{-3}(\frac{dr}{dt})^2 = c^2(1 - \frac{r^*}{r})^{-1} - \frac{1}{r^2}\frac{L^2}{E^2} \tag{502}$$

At the peak point, $\frac{dr}{dt} = 0$ so that there is

$$\frac{L^2}{E^2} = c^2R^2(1 - \frac{r^*}{R})^{-1} \tag{503}$$

Where, R is the coordinate of the peak point.

Taking it back into the Equation (502), there is

$$(1 - \frac{r^*}{r})^{-3}(\frac{dr}{dt})^2 - c^2(1 - \frac{r^*}{r})^{-1} + c^2\frac{R^2}{r^2}(1 - \frac{r^*}{R})^{-1} = 0 \tag{504}$$

or

$$(\frac{dr}{dt})^2 = c^2(1 - \frac{r^*}{r})^2 - c^2\frac{R^2}{r^2}(1 - \frac{r^*}{r})^3(1 - \frac{r^*}{R})^{-1} \tag{505}$$

Then the time spending from coordinate R to coordinate r at a very far distance could be integrated as

$$t|_R^r = \frac{1}{c}\int_R^r(1 - \frac{r^*}{r})^{-1}[1 - (1 - \frac{r^*}{r})(1 - \frac{r^*}{R})^{-1}\frac{R^2}{r^2}]^{-1/2}dr \tag{506}$$

Because $\frac{r^*}{r}$ and $\frac{r^*}{R}$ are very small, it could be written as

$$t|_R^r = \frac{1}{c}\int_R^r(1 - \frac{r^*}{r})^{-1}[1 - (1 - \frac{r^*}{r} + \frac{r^*}{R})\frac{R^2}{r^2}]^{-1/2}dr \tag{507}$$

It has an approximate solution [4] as

$$\begin{aligned} t|_R^r &\approx \frac{1}{c} \int_R^r (1 - \frac{R^2}{r^2})^{-1/2} [1 + \frac{r^*}{r} + \frac{r^*R}{2r(r+R)}] dr \\ &= \frac{1}{c} [\sqrt{r^2 - R^2} + r^* \ln(\frac{r + \sqrt{r^2 - R^2}}{R}) + \frac{r^*}{2} \sqrt{\frac{r-R}{r+R}}] \end{aligned} \quad (508)$$

8.7.2. Errors in

We have known that the assumption in classical equations is incorrect. In fact, the equation could be solved without the assumption. Such as the Lagrangian

$$-c^2 + (1 - \frac{r^*}{r})^{-1} \dot{r}^2 + r^2 \dot{\phi}^2 = 0 \quad (509)$$

With $L = Rc = r^2 \dot{\phi} = \text{const.}$ it is

$$-c^2 + (1 - \frac{r^*}{r})^{-1} \dot{r}^2 + \frac{c^2 R^2}{r^2} = 0 \quad (510)$$

And we know that

$$\dot{r} = \frac{dr}{dt} \dot{t} = \frac{dr}{dt} \frac{dt}{dr} = \frac{dr}{dt} (1 - \frac{r^*}{r})^{-1/2} \quad (511)$$

so that

$$(\frac{dr}{dt})^2 = c^2 (1 - \frac{r^*}{r})^2 - c^2 (1 - \frac{r^*}{r})^2 \frac{R^2}{r^2} \quad (512)$$

The differential relationship could be integrated that

$$t|_R^r = \frac{1}{c} \int_R^r (1 - \frac{r^*}{r})^{-1} (1 - \frac{R^2}{r^2})^{-1/2} dr \quad (513)$$

Approximate solution is

$$t|_R^r = \frac{1}{c} [\sqrt{r^2 - R^2} + r^* \ln(\frac{r + \sqrt{r^2 - R^2}}{R})] \quad (514)$$

One can find that this treatment has just brought about a little deviation from the previous. That is because the trajectory is only a little different from that one at deflect angle.

8.7.3. Renovated Equations for Radar Echoes

The revisit equation of trajectory may help to get new performances of the issue.

For the Lagrangian

$$-c^2 + (1 - \frac{r^*}{r})^{-1} \dot{r}^2 + r^2 \dot{\phi}^2 = 0 \quad (515)$$

In previous section, we have got the angular velocity expression based on energy momentum conservativeness as discussed in previous sections that $L = Rm_R c = r^2 m_r \dot{\phi}$, so that

$$\dot{\phi} = \frac{Rm_R c}{m_r r^2} = \frac{Rc}{r^2} \frac{1 + \frac{r^*}{2r}}{1 + \frac{r^*}{2r}} \quad (516)$$

Set $B = (R + \frac{r^*}{2})$ and in weak field the item $1/(1 + \frac{r^*}{2r}) \approx (1 - \frac{r^*}{2r})$, thus

$$\dot{\phi} = \frac{1}{r^2} (1 - \frac{r^*}{2r}) Bc \quad (517)$$

Then the Lagrangian becomes

$$-c^2 + (1 - \frac{r^*}{r})^{-1} \dot{r}^2 + \frac{1}{r^2} (1 - \frac{r^*}{r}) B^2 c^2 = 0 \quad (518)$$

A transformation could be made as

$$-c^2 + (1 - \frac{r^*}{r})^{-2} (\frac{dr}{dt})^2 + (1 - \frac{r^*}{r}) \frac{B^2 c^2}{r^2} = 0 \quad (519)$$

With $B = (R + \frac{r^*}{2})$ it becomes

$$-c^2 + (1 - \frac{r^*}{r})^{-2} (\frac{dr}{dt})^2 + (1 - \frac{r^*}{r})(1 + \frac{r^*}{R}) \frac{R^2 c^2}{r^2} = 0 \quad (520)$$

so that there is

$$(\frac{dr}{dt})^2 = c^2 [(1 - \frac{r^*}{r})^2 - (1 - \frac{r^*}{r})^3 (1 + \frac{r^*}{R}) \frac{R^2}{r^2}] \quad (521)$$

or

$$(dt)^2 = \frac{1}{c^2} [(1 - \frac{r^*}{r})^2 - (1 - \frac{r^*}{r})^3 (1 + \frac{r^*}{R}) \frac{R^2}{r^2}]^{-1} (dr)^2 \quad (522)$$

or

$$dt = \frac{1}{c} [(1 - \frac{r^*}{r})^2 - (1 - \frac{r^*}{r})^3 (1 + \frac{r^*}{R}) \frac{R^2}{r^2}]^{-1/2} dr \quad (523)$$

The time spending from coordinate R to coordinate r at a very far distance could be written as an integration as

$$t|_R^r = \frac{1}{c} \int_R^r (1 - \frac{r^*}{r})^{-1} (1 - \frac{R^2}{r^2} + \frac{r^* R^2}{r r^2} - \frac{r^* R^2}{R r^2})^{-1/2} dr \quad (524)$$

or

$$t|_R^r = \frac{1}{c} \int_R^r (1 + \frac{r^*}{r}) [1 - \frac{R^2}{r^2} + (\frac{r^*}{r} - \frac{r^*}{R}) \frac{R^2}{r^2}]^{-1/2} dr \quad (525)$$

By building a function

$$f(x) = x^{-1/2} \quad (526)$$

where, $x = 1 - \frac{R^2}{r^2}$. The derivative

$$f'(x) = -\frac{1}{2} x^{-3/2} \quad (527)$$

Considering $\frac{r^*}{r}$ is very small, the rear part in the integral Equation (525) could be simplified as

$$[1 - \frac{R^2}{r^2} + (\frac{r^*}{r} - \frac{r^*}{R}) \frac{R^2}{r^2}]^{-1/2} \approx (1 - \frac{R^2}{r^2})^{-\frac{1}{2}} - \frac{1}{2} (1 - \frac{R^2}{r^2})^{-3/2} (\frac{r^* R^2}{r^3} - \frac{r^* R}{r^2}) \quad (528)$$

Then the time integration will be reformed as

$$t|_R^r \approx \frac{1}{c} \int_R^r [(1 - \frac{R^2}{r^2})^{-1/2} (1 + \frac{r^*}{r}) - \frac{1}{2} (1 - \frac{R^2}{r^2})^{-3/2} (\frac{r^* R^2}{r^3} - \frac{r^* R}{r^2})] dr \quad (529)$$

where in the last item, a $(1 + \frac{r^*}{r})$ has been simplified to be 1.

The first part of the integration could be calculated to be

$$\text{part1} = \frac{1}{c} \int_R^r (1 - \frac{R^2}{r^2})^{-1/2} (1 + \frac{r^*}{r}) dr = \frac{1}{c} [\sqrt{r^2 - R^2} + r^* \ln(\frac{r + \sqrt{r^2 - R^2}}{R})] \quad (530)$$

and the last part is done as

$$\text{part2} = -\frac{1}{c} \int_R^r \frac{1}{2} (1 - \frac{R^2}{r^2})^{-\frac{3}{2}} (\frac{r^* R^2}{r^3} - \frac{r^* R}{r^2}) dr = \frac{1}{c} (\frac{r r^*}{2\sqrt{r^2 - R^2}} - \frac{r^* R}{2\sqrt{r^2 - R^2}}) \quad (531)$$

so that the total integration is

$$t|_R^r \approx \frac{1}{c} [\sqrt{r^2 - R^2} + r^* \ln(\frac{r + \sqrt{r^2 - R^2}}{R}) + \frac{r^*}{2} \sqrt{\frac{r - R}{r + R}}] \quad (532)$$

8.7.4. A Falsification on Traditional Methodology on Close-to-Light-Speed Particle Time Spending

Time delay of close-to-light-speed particles is the same issue as the trajectory. Nevertheless, we could see more problems revealed in the issue.

As Lagrangian is employed in traditional methodologies as Equation (415) with the two settings.

$$\dot{r}^2 = -(1 - \frac{r^*}{r})c^2 + c^2 E^2 - (1 - \frac{r^*}{r})\frac{L^2}{r^2} \quad (533)$$

It could be rewritten as

$$\dot{r}^2 = \frac{r^*}{r}c^2 + c^2(E^2 - 1) - (1 - \frac{r^*}{r})\frac{L^2}{r^2} \quad (534)$$

And also with the setting in traditional method as that has been applied in the Equation (501) for issue of radar echoes

$$\dot{r} = \frac{dr}{dt} \dot{t} = \frac{dr}{dt} E(1 - \frac{r^*}{r})^{-1} \quad (535)$$

it is

$$(1 - \frac{r^*}{r})^{-2} (\frac{dr}{dt})^2 E^2 = \frac{r^*}{r}c^2 + (E^2 - 1)c^2 - (1 - \frac{r^*}{r})\frac{L^2}{r^2} \quad (536)$$

At the peak point, $\frac{dr}{dt} = 0$, and $L = Rc$ so that there is

$$E^2 = 2(1 - \frac{r^*}{R}) \quad (537)$$

where, R is the coordinate of the peak point.

It could be imagined that in the traditional solution of light ray, constant E could be calculate to be an approximate of 1.0 as that in Equation (503), but in this condition it should goes to about 2.0.

Therefore, the equation is

$$2(1 - \frac{r^*}{r})^{-2} (\frac{dr}{dt})^2 (1 - \frac{r^*}{R}) = \frac{r^*}{r}c^2 + (1 - \frac{2r^*}{R})c^2 - (1 - \frac{r^*}{r})\frac{R^2 c^2}{r^2} \quad (538)$$

It could be reformed as

$$(\frac{dr}{dt})^2 = \frac{1}{2} [(1 - \frac{r^*}{r})^2 \frac{r^*}{r} (1 + \frac{r^*}{R})c^2 + (1 - \frac{r^*}{r})^2 (1 - \frac{r^*}{R})c^2 - (1 - \frac{r^*}{r})^3 (1 + \frac{r^*}{R})\frac{R^2 c^2}{r^2}] \quad (539)$$

The time spending could be drawn from the integration

$$t|_R^r = \frac{\sqrt{2}}{c} \int_R^r (1 - \frac{r^*}{r})^{-1} [\frac{r^*}{r} (1 + \frac{r^*}{R}) + (1 - \frac{r^*}{R}) - (1 - \frac{r^*}{r})(1 + \frac{r^*}{R})\frac{R^2}{r^2}]^{-1/2} dr \quad (540)$$

Some small items could be neglected that

$$t|_R^r = \frac{\sqrt{2}}{c} \int_R^r (1 - \frac{r^*}{r})^{-1} [1 + \frac{r^*}{r} - \frac{r^*}{R} - (1 - \frac{r^*}{r})(1 + \frac{r^*}{R})\frac{R^2}{r^2}]^{-1/2} dr \quad (541)$$

or

$$t|_R^r = \frac{\sqrt{2}}{c} \int_R^r (1 - \frac{r^*}{r})^{-1} [1 + \frac{r^*}{r} - \frac{r^*}{R} - (1 - \frac{r^*}{r} + \frac{r^*}{R})\frac{R^2}{r^2}]^{-1/2} dr \quad (542)$$

or

$$t|_R^r = \frac{\sqrt{2}}{c} \int_R^r (1 - \frac{r^*}{r})^{-1} [1 - \frac{R^2}{r^2} + (\frac{r^*}{r} - \frac{r^*}{R})(1 + \frac{R^2}{r^2})]^{-1/2} dr \quad (543)$$

It is a good idea to reform the items in root function that for the function

$$f(x) = x^{-1/2} \quad (544)$$

where, $x = 1 - \frac{R^2}{r^2}$ which could present main quantity of the function with respect to the rear part which just performs a higher order small item. The derivative

$$f'(x) = -\frac{1}{2}x^{-3/2} \quad (545)$$

Considering $\frac{r^*}{r}$ and $\frac{r^*}{R}$ are very small, the rear part in the integral Equation (543) could be simplified as

$$[1 - \frac{R^2}{r^2} + (\frac{r^*}{r} - \frac{r^*}{R})(1 + \frac{R^2}{r^2})]^{-1/2} \approx (1 - \frac{R^2}{r^2})^{-\frac{1}{2}} - \frac{1}{2}(1 - \frac{R^2}{r^2})^{-3/2}(\frac{r^*}{r} - \frac{r^*}{R})(1 + \frac{R^2}{r^2}) \quad (546)$$

Then the integration could be gain as

$$t|_R^r \approx \frac{\sqrt{2}}{c} \int_R^r [(1 - \frac{R^2}{r^2})^{-\frac{1}{2}}(1 + \frac{r^*}{r}) - \frac{1}{2}(1 - \frac{R^2}{r^2})^{-\frac{3}{2}}(\frac{r^*}{r} - \frac{r^*}{R}) + \frac{1}{2}(1 - \frac{R^2}{r^2})^{-\frac{3}{2}}(\frac{r^* R^2}{r^3} - \frac{r^* R}{r^2})] dr \quad (547)$$

where in the last item, a coefficient $(1 + \frac{r^*}{r})$ has been simplified to be 1.

The first part of the integration could be calculated to be

$$\begin{aligned} \text{part1} &= \frac{\sqrt{2}}{c} \int_R^r (1 - \frac{R^2}{r^2})^{-1/2} (1 + \frac{r^*}{r}) dr = \frac{\sqrt{2}}{c} \int_R^r [r(r^2 - R^2)^{-1/2} + r^*(r^2 - R^2)^{-1/2}] dr \\ &= \frac{\sqrt{2}}{c} [\sqrt{r^2 - R^2} + r^* \ln(\frac{r + \sqrt{r^2 - R^2}}{R})] \end{aligned} \quad (548)$$

while the third part is

$$\text{part3} = -\frac{\sqrt{2}}{2c} \int_R^r (1 - \frac{R^2}{r^2})^{-3/2} (\frac{r^* R^2}{r^3} - \frac{r^* R}{r^2}) dr = \frac{\sqrt{2}}{4c} \frac{r r^*}{\sqrt{r^2 - R^2}} - \frac{\sqrt{2}}{4c} \frac{r^* R}{\sqrt{r^2 - R^2}} = \frac{\sqrt{2}}{4c} \sqrt{\frac{r-R}{r+R}} \quad (549)$$

Then, the second part is

$$\text{part2} = -\frac{\sqrt{2}}{2c} \int_R^r (1 - \frac{R^2}{r^2})^{-3/2} (\frac{r^*}{r} - \frac{r^*}{R}) dr \quad (550)$$

It is better to be split to two parts for integrations

$$\begin{aligned} \text{part21} &= -\frac{\sqrt{2}}{2c} \int_R^r (1 - \frac{R^2}{r^2})^{-3/2} \frac{r^*}{r} dr = -\frac{\sqrt{2}}{2c} \int_R^r (r^2 - R^2)^{-3/2} r^2 r^* dr \\ &= -\frac{\sqrt{2}}{2c} \int_R^r (r^2 - R^2)^{-3/2} (r^2 - R^2 + R^2) r^* dr \\ &= -\frac{\sqrt{2}}{2c} \int_R^r [r^*(r^2 - R^2)^{-1/2} + (r^2 - R^2)^{-3/2} R^2 r^*] dr \\ &= -\frac{\sqrt{2}}{2c} [r^* \ln(\frac{r + \sqrt{r^2 - R^2}}{R}) - \frac{r^* R}{\sqrt{r^2 - R^2}}] \end{aligned} \quad (551)$$

The last item would be involved with an infinite problem at lower bound, but it could be solved by an offset with another infinite problem in next equation.

$$\begin{aligned} \text{part22} &= \frac{\sqrt{2}}{2c} \int_R^r (1 - \frac{R^2}{r^2})^{-3/2} \frac{r^*}{R} dr = \frac{\sqrt{2}}{2c} \int_R^r (r^2 - R^2)^{-3/2} r^3 \frac{r^*}{R} dr \\ &= \frac{\sqrt{2}}{4c} \int_R^r (r^2 - R^2)^{-3/2} (r^2 - R^2 + R^2) \frac{r^*}{R} dr \\ &= \frac{\sqrt{2}}{4c} \int_R^r [\frac{r^*}{R} (r^2 - R^2)^{-1/2} + (r^2 - R^2)^{-3/2} R r^*] d(r^2 - R^2) \\ &= \frac{\sqrt{2}}{2c} (\frac{r^*}{R} \sqrt{r^2 - R^2} - \frac{r^* R}{\sqrt{r^2 - R^2}}) \end{aligned} \quad (552)$$

The final solution could be drawn as

$$t|_R^r \approx \frac{\sqrt{2}}{c} [(1 + \frac{1}{2} \frac{r^*}{R}) \sqrt{r^2 - R^2} + \frac{1}{2c} r^* \ln(\frac{r + \sqrt{r^2 - R^2}}{R}) + \frac{3}{4c} \sqrt{\frac{r-R}{r+R}}] \quad (553)$$

It is an unacceptable solution with coefficient $\sqrt{2}$ in those items, especially in the first item the approximate $\frac{\sqrt{2}}{c} \sqrt{r^2 - R^2}$ will cause a giant time delay with respect to the expected time interval $\frac{1}{c} \sqrt{r^2 - R^2}$. That is because of the treatment in Equation (535) that the left item has got a double value by mistake. This result has revealed a very strong clue that the settings treatment of Equation (535) so as well the setting in Equation (331) involves with errors certainly.

If the Equation (331) will not be persisted in we will see that $t = (1 - \frac{r^*}{r})^{-1/2}$ by definition rather than $t = E(1 - \frac{r^*}{r})^{-1} = \sqrt{2}(1 - \frac{r^*}{r})^{-1/2}$ by Equation (537). Thus, we can repeat the calculation for the Equation (534) that

$$(1 - \frac{r^*}{r})^{-1} (\frac{dr}{dt})^2 = \frac{r^*}{r} c^2 + c^2(E^2 - 1) - (1 - \frac{r^*}{r}) \frac{L^2}{r^2} \quad (554)$$

Case $\frac{dr}{dt} = 0$, we get the same $E^2 = 2(1 - \frac{r^*}{R})$

It could be reformed as

$$\left(\frac{dr}{dt}\right)^2 = \left(1 - \frac{r^*}{r}\right) \frac{r^*}{r} c^2 + \left(1 - \frac{r^*}{r}\right) \left(1 - \frac{2r^*}{R}\right) c^2 - \left(1 - \frac{r^*}{r}\right)^2 \frac{R^2 c^2}{r^2} \quad (555)$$

Time spending could be solved by integration

$$t|_R^r = \frac{1}{c} \int_R^r \left(1 - \frac{r^*}{r}\right)^{-1} \left[\frac{r^*}{r} \left(1 + \frac{r^*}{r}\right) + \left(1 + \frac{r^*}{r}\right) \left(1 - \frac{2r^*}{R}\right) - \frac{R^2}{r^2}\right]^{-1/2} dr \quad (556)$$

It could be reformed as

$$t|_R^r = \frac{1}{c} \int_R^r \left(1 - \frac{r^*}{r}\right)^{-1} \left[1 - \frac{R^2}{r^2} + \left(\frac{2r^*}{r} - \frac{2r^*}{R}\right)\right]^{-1/2} dr \quad (557)$$

Because

$$\left[1 - \frac{R^2}{r^2} + \left(\frac{2r^*}{r} - \frac{2r^*}{R}\right)\right]^{-1/2} \approx \left(1 - \frac{R^2}{r^2}\right)^{-\frac{1}{2}} - \left(1 - \frac{R^2}{r^2}\right)^{-3/2} \left(\frac{r^*}{r} - \frac{r^*}{R}\right) \quad (558)$$

The time integration could be rewritten as

$$t|_R^r \approx \frac{1}{c} \int_R^r \left[\left(1 - \frac{R^2}{r^2}\right)^{-1/2} \left(1 + \frac{r^*}{r}\right) - \left(1 - \frac{R^2}{r^2}\right)^{-\frac{3}{2}} \left(\frac{r^*}{r} - \frac{r^*}{R}\right)\right] dr \quad (559)$$

in which,

$$\begin{aligned} \text{part1} &= \frac{1}{c} \int_R^r \left(1 - \frac{R^2}{r^2}\right)^{-1/2} \left(1 + \frac{r^*}{r}\right) dr = \frac{1}{c} \int_R^r \left[r(r^2 - R^2)^{-1/2} + r^*(r^2 - R^2)^{-1/2}\right] dr \\ &= \frac{1}{c} \left[\sqrt{r^2 - R^2} + r^* \ln\left(\frac{r + \sqrt{r^2 - R^2}}{R}\right)\right] \end{aligned} \quad (560)$$

$$\text{part2} = -\frac{1}{c} \int_R^r \left(1 - \frac{R^2}{r^2}\right)^{-3/2} \left(\frac{r^*}{r} - \frac{r^*}{R}\right) dr \quad (561)$$

To split part 2 to two parts

$$\begin{aligned} \text{part21} &= -\frac{1}{c} \int_R^r \left(1 - \frac{R^2}{r^2}\right)^{-3/2} \frac{r^*}{r} dr = -\frac{1}{c} \int_R^r (r^2 - R^2)^{-3/2} r^2 r^* dr \\ &= -\frac{1}{c} \int_R^r (r^2 - R^2)^{-3/2} (r^2 - R^2 + R^2) r^* dr \\ &= -\frac{1}{c} \int_R^r \left[r^*(r^2 - R^2)^{-1/2} + (r^2 - R^2)^{-3/2} R^2 r^*\right] dr \\ &= -\frac{1}{c} \left[r^* \ln\left(\frac{r + \sqrt{r^2 - R^2}}{R}\right) - \frac{r^* r}{\sqrt{r^2 - R^2}}\right]_R^r \end{aligned} \quad (562)$$

and

$$\begin{aligned} \text{part22} &= \frac{1}{c} \int_R^r \left(1 - \frac{R^2}{r^2}\right)^{-3/2} \frac{r^*}{R} dr = \frac{1}{c} \int_R^r (r^2 - R^2)^{-3/2} r^3 \frac{r^*}{R} dr \\ &= \frac{1}{2c} \int_R^r (r^2 - R^2)^{-3/2} (r^2 - R^2 + R^2) \frac{r^*}{R} dr^2 \\ &= \frac{1}{2c} \int_R^r \left[\frac{r^*}{R} (r^2 - R^2)^{-1/2} + (r^2 - R^2)^{-3/2} R r^*\right] d(r^2 - R^2) \\ &= \frac{1}{c} \left(\frac{r^*}{R} \sqrt{r^2 - R^2} - \frac{r^* R}{\sqrt{r^2 - R^2}}\right)_R^r \end{aligned} \quad (563)$$

The final solution could be drawn as

$$t|_R^r \approx \frac{1}{c} \left[\left(1 + \frac{r^*}{R}\right) \sqrt{r^2 - R^2} + \sqrt{\frac{r-R}{r+R}}\right] \quad (564)$$

This solution sounds more acceptable than the previous but it is still involved with errors.

8.7.5. Renovated Solution for Close-to-Light-Speed Particles

Now let us discuss another kind of no-delay time spending, the time spending of close-to-light-speed particles. The velocity composition of close-to-light-speed particles could be expressed as

$$-c^2 + \left(\frac{dr}{dt}\right)^2 + r^2 \left(\frac{d\varphi}{dt}\right)^2 = 0 \quad (565)$$

With angular momentum conservativeness as that in Equation (402), there is

$$-c^2 + \left(\frac{dr}{dt}\right)^2 + \left(1 - \frac{r^*}{r}\right) \frac{B^2 c^2}{r^2} = 0 \quad (566)$$

or

$$-c^2 + \left(\frac{dr}{dt}\right)^2 + \left(1 - \frac{r^*}{r}\right) \left(1 + \frac{r^*}{R}\right) \frac{R^2 c^2}{r^2} = 0 \quad (567)$$

so that the time spending integration could be built as

$$\begin{aligned} t|_R^r &= \frac{1}{c} \int_R^r \left[1 - \left(1 - \frac{r^*}{r}\right) \left(1 + \frac{r^*}{R}\right) \frac{R^2}{r^2}\right]^{-1/2} dr \\ &= \frac{1}{c} \int_R^r \left[1 - \frac{R^2}{r^2} - \left(\frac{r^*}{r} - \frac{r^*}{R}\right) \frac{R^2}{r^2}\right]^{-1/2} dr \\ &\approx \frac{1}{c} \int_R^r \left[\left(1 - \frac{R^2}{r^2}\right)^{-1/2} - \frac{1}{2} \left(1 - \frac{R^2}{r^2}\right)^{-3/2} \left(\frac{r^* R^2}{r^3} - \frac{r^* R}{r^2}\right)\right] dr \\ &= \frac{1}{c} \left[\sqrt{r^2 - R^2} + \frac{r^*}{2} \sqrt{\frac{r-R}{r+R}}\right] \end{aligned} \quad (568)$$

8.7.6. Equations of Time Delay

We have found that the revisit solution of time spend is the same with classical. The reality is that the classical treatment has got the same trajectory by additional settings, so that it is undoubtful to gain a same time spend with. In fact, the problem of time delay of radar echoes is just a kind of performance of light deflection. The results in these discussions are also just extensions of that in the problem of light deflection. The classical equation for time delay is to defined the difference between time interval of light rays and that of an imaginary motion of absolute light speed, and the half length of the trajectory has been coarsely set to be

$$l = \sqrt{r^2 - R^2} \quad (569)$$

So that the classical equation of half branch time delay could be calculated with Equation (532) as

$$\Delta t = \frac{1}{c} \left[r^* \ln\left(\frac{r + \sqrt{r^2 - R^2}}{R}\right) + \frac{r^*}{2} \sqrt{\frac{r-R}{r+R}} \right] \quad (570)$$

But the half branch time delay of light rays, with respect to the time interval of close-to-light-speed particles Equation (568), will be a little different

$$\Delta t \approx \frac{r^*}{c} \ln\left(\frac{r + \sqrt{r^2 - R^2}}{R}\right) \quad (571)$$

One may argue that the trajectory of close-to-light-speed particles will be different to light rays. In fact, the real length of the light trajectory also does not equal to $\sqrt{r^2 - R^2}$. One can easily take measures to work out the real length of that trajectory. I prefer to give a more accurate value than Equation (568) that could be estimated by geometrical relationship as shown in Fig. 20 while $r \gg R \approx b$ that

$$l_{\text{accurate}} = \sqrt{r^2 - R^2} + R \sin(-\varphi_\infty) \approx \sqrt{r^2 - R^2} + r^* \quad (572)$$

where, $\varphi_\infty = -r^*/b$ is deflect angle of light ray.

So that a real half branch time delay of light rays with respect to an absolute motion on the very trajectory is

$$\Delta t = \frac{1}{c} \left[r^* \ln \left(\frac{r + \sqrt{r^2 - R^2}}{R} \right) + \frac{r^*}{2} \sqrt{\frac{r-R}{r+R}} - r^* \right] \approx \frac{1}{c} \left[r^* \ln \left(\frac{r + \sqrt{r^2 - R^2}}{R} \right) - \frac{r^*}{2} \right] \quad (573)$$

We know that the time delay has been verified to be very high accurate value with respect to the classical equation, that is because the trajectory length has always been set to be $\sqrt{r^2 - R^2}$, which is indeed not accurate length of trajectory line.

If consider the close-to-light-speed particles, the real length of its trajectory line could be estimated to be

$$l_p = \sqrt{r^2 - R^2} + R \sin(-\varphi_\infty) = \sqrt{r^2 - R^2} + \frac{r^*}{2} \quad (574)$$

where, $\varphi_\infty = -r^*/(2b)$ is deflect angle of particles.

With Equation (568), the half branch time delay of close-to-light-speed particles is

$$\Delta t_p = 0 \quad (575)$$

The zero time delay is because the particles keep approximate velocity during flying on the trajectory. This result is a certainty of velocity conservation for massive particles.

Something different from trajectory investigation, the test of time delay of light and close-to-light-speed particle propagation might allow a quite big separation to the Sun edge thereby to provide not bad accuracy. Furthermore, it could be expected to sponsor experiments to emit light rays and massive particles on a straight line from a point not very close to the Sun at the same time for time delay verification.

8.8. Comparative Research between Numerical Solutions and Algebraic Solutions

8.8.1. The Invalidity of Newtonian Second Law in Light Propagation

It is the most interesting that perhaps there is the probability to carry out new numerical method to calculate the light trajectories or close-to-light-speed motions in gravitational field, which could be called ballistic trajectory method. Considering that the gravity component parallel to the motion trajectory will not bring about changes to further motion, we could only consider the calculation on motion variation due to vertical component of gravity so that to determine a differential coordinate on the trajectory. Thus, in the way incremental, the trajectory could be solved at last.

Firstly, it is proficient to take a motion of close-to-light-speed massive particle in one source field into discussion, as shown in Figure 23.




Figure 23. A diagram of ballistic method for a close-to-light-speed motion.

It is believed that for a massive particle in weak field, the vertical deviation could be solved by Newtonian second law

$$\Delta s = \frac{1}{2} g_v \Delta t^2 \quad (576)$$

where, g_v is the component of gravity vertical to the velocity of the particle.

The coordinates at position A are known to be (r_A, φ_A) or (x_A, y_A) . As well, the trajectory will have a direction with angle α_A . Then, the distance in a time interval Δt the particle travelling is

$$AB = c \Delta t \quad (577)$$

The deviation angle of AB to AC is

$$\delta \approx \Delta s / AB \quad (578)$$

and

$$\gamma = \alpha_A + \delta \quad (579)$$

Thus, the coordinates at position B could be calculated as

$$x_B = x_A + AB \cos \gamma \quad (580)$$

$$y_B = y_A + AB \sin \gamma \quad (581)$$

or

$$r_B = \sqrt{x_B^2 + y_B^2} \quad (582)$$

$$\varphi_B = \arccos \frac{x_B}{r_B} \quad (583)$$

It should be highlighted that the angle γ is not the direction angle α_B . The α_B could be calculated by angular momentum conservation that

$$(1 + \frac{r^*}{2R})R = (1 + \frac{r^*}{2r_B})r_B \sin \beta_B \quad (584)$$

where, $\beta_B = \alpha_B + \varphi_B$.

Now back to the fly of light ray from position A to position B. It might be imagined to calculate the deviation Δs also by Newtonian second law. That sounds naturally to see the photons as light speed particles with dynamic mass so that to deviate in the same way as massive particles do. Unfortunately, it is impossible in that that kind of calculation will lead to the result very close to that of massive particles. It seems like that the Newtonian second law sounds invalid in light propagation even at the vertical direction. Perhaps one can propose a double gravity for a simulation on light propagation with respect to that of massive particles and it could be imagined that the results would be of not bad precisions. But the truth is that they are really not of a same issue in that they would have different asymptotic distance b . An interesting clue is that light rays are un-accelerable even at the direction vertical to the motion. Maybe that could be employed to interpret the invalidity of Newtonian second law.

Anyway, there is still an opportunity to explore the ballistic trajectory method for them. That is to perform by trial methodologies. We can imagine that after point A there will be a wave front after Huygens' postulation, then, some points on it may be selected for further considerations. With angular momentum conservativeness and Lagrangian, one can calculate the velocity and angle α_B , as well as coordinates as shown in Figure 24. Thus, the deviations of the points could be estimated to help for further trials. I have made more efforts to try but not done, although I still believe the probabilities.




Figure 24. Probability of ballistic trajectory method for light deviation.

8.8.2. Comparisons of Numerical Solutions and Algebraic Solutions

To make comparisons with numerical and algebraic solutions is not only a kind of further verification but also a further support to the conclusion of energy moment conservativeness, and at last, a surprise of light propagation. As has discussed previous, ballistic trajectory method could be developed to calculate the trajectories of close-to-light massive particles. It could be applied by

dividing trajectory to finite segments and calculating coordinates with Equation (576) to Equation (584) step by step. The invalidity of Newtonian second law for light propagation is actual another kind of support to the inferences of kinematics.

Notwithstanding, difference method for all differential equations could also be employed for more comparisons. Difference method has shown great advantages in scientific calculations and many excellent schemes have been developed to serve for more complex requirements.

For the equation of light rays

$$u'' + u = \frac{3}{2}r^*u^2 \tag{585}$$

The simple central difference scheme could be suggested that

$$u'' = \frac{u(\varphi+\Delta\varphi)-2u(\varphi)+u(\varphi-\Delta\varphi)}{(\Delta\varphi)^2} \tag{586}$$

Thus, the difference equation could be built as

$$\frac{u(\varphi+\Delta\varphi)-2u(\varphi)+u(\varphi-\Delta\varphi)}{(\Delta\varphi)^2} + u(\varphi) = \frac{3}{2}r^*u^2(\varphi) \tag{587}$$

Pre-exercises show that if step intervals defined by $\Delta\varphi < 0.1^\circ$, the calculations would get quite good accuracy.

As for the equation of close-to-light massive particles, the differential equation

$$u'' + u = \frac{r^*}{2b^2} \tag{588}$$

could also be built simply as

$$\frac{u(\varphi+\Delta\varphi)-2u(\varphi)+u(\varphi-\Delta\varphi)}{(\Delta\varphi)^2} + u(\varphi) = \frac{r^*}{2b^2} \tag{589}$$

Thus, we would sponsor a comparison with the analytical solutions, difference method solutions and the ballistic trajectory method solutions mentioned previously, together with the asymptotic lines and horizontal line for references.

Numerical methods of difference and ballistic were carried in office computer and others were completed by the calculator of my mobile phone, in that the phone calculator could provide 8 floating point precisions more than that of the Fortran software in the computer.

It is shown in Table 5 that difference method and ballistic method still perform not bad precisions in the results of upper parts of the trajectories. Of course, both of the two methods are of step by step arithmetic so as to accumulate quite amount of deviations at the rear parts of the trajectories. Anyway, the two numerical methods are reliable, that greatly supports the inferences of kinematics.

It is easy to develop more optimal schemes for difference method and the way of optimal secant seeking instead of the tangent seeking for ballistic method or any other effective technologies to improve the precisions of numerical analyses. That will confirm the conclusions and inferences that have been drawn previous.

Table 5. Comparison of the analytical solutions, difference method solutions and the ballistic trajectory method solutions.

coordinate (degree)	coordinate radius (m) of solution of light ray by method of			coordinate radius (m) of solution of close-to-light-speed massive particle by method of				coordinate radius (m) of horizontal line
	analytical	difference	asymptotic line	analytical	difference	ballistic	asymptotic line	
90	695,500,000	695,500,000	695,501,477	695,500,000	695,500,000	695,500,000	695,501,477	695,501,477
80	706,229,140	706,229,166	706,230,180	706,229,186	706,229,216	706,229,226	706,230,444	706,230,708
70	740,135,345	740,135,367	740,136,069	740,135,540	740,135,570	740,135,590	740,136,641	740,137,212
60	803,093,469	803,093,487	803,093,961	803,093,961	803,093,991	803,094,020	803,094,945	803,095,929
50	907,909,142	907,909,153	907,909,463	907,910,181	907,910,213	907,910,251	907,911,080	907,912,699
40	1,082,002,548	1,082,002,549	1,082,002,744	1,082,004,645	1,082,004,679	1,082,004,738	1,082,005,482	1,082,008,219
30	1,390,992,617	1,390,992,592	1,390,992,723	1,390,997,047	1,390,997,085	1,390,997,170	1,390,997,839	1,391,002,954
20	2,033,486,508	2,033,486,403	2,033,486,554	2,033,497,655	2,033,497,694	2,033,497,838	2,033,498,416	2,033,510,279
10	4,005,136,911	4,005,136,314	4,005,136,922	4,005,184,405	4,005,184,395	4,005,184,750	4,005,185,149	4,005,233,377
0.26	153,123,772,603	153,122,660,623	153,123,772,604	153,195,382,906	153,195,143,112	153,195,397,701	153,195,383,644	153,267,061,697
0.01	3,890,278,441,952	3,889,557,122,200	3,890,278,441,977	3,937,035,389,754	3,936,873,242,381	3,937,033,549,851	3,937,035,390,479	3,984,929,947,953
0	163,784,893,532,900	162,515,766,173,612	163,784,893,516,163	327,569,787,065,801	326,450,054,739,264	327,616,498,942,582	327,569,787,031,587	∞

8.9. Additional Discussions

8.9.1. On Geodesic Line, Inertial Motion, and General Covariance

We have discussed those derivatives theoretically in section 4, section 5 and section6 that in covariant space the acceleration of a massive matter could not be really calculate to be zero and even revisit light frequency shift will perform double of that of contra variant space. There is only the light speed that always serve the general covariance as the result the acceleration of a light ray may perform nonvanishing in covariant space in any conditions. These conclusions would have then challenged the theory of geodesic lines, because geodesic lines ask for thoroughly nonvanishing solutions for covariant derivatives. We also have proved that geodesic equation cannot be really employed for the solutions of massive matter’s motions as well as the motions of light rays in section 8.

Now it is the time to make additional discussions on the tips between matter’s trajectories, inertial motions, and geodesic lines.

General covariance asks for matters in covariant space to perform inertial motions just as they perform in no gravity fields. That requires trajectories of matters stay on the lines of geodesic lines in gravitational fields.

Geodesic lines are of the so-called straight lines of a covariant space that present the geometrical properties of the space. But inertial motions must ask for more. For example, an inertial trajectory must serve for any matters to run on, either the massive matters or light rays. The more important is that inertial trajectory must be run on by any matters with any velocities with same direction. It could be said that inertial motions share trajectories.

Back to the massive matter’s trajectories, it is no doubt that any two matters with different velocities in gravitational field with same initial positions and directions, will fly apart, so that to fly on two trajectories, just as matter’s motions in solar system. These trajectories of course correspond to two lines in covariant space. It is said that the matters with different velocities may not share trajectory in covariant space so that the motions are not inertial motions in covariant space and the trajectories are not geodesic lines.

As for light rays, due to the consequence of invariant light speed, they will come up to zero accelerations in covariant space and we might have seen that light rays could share trajectories in different frequencies, that really performs general covariance. But the trajectories of light rays are not that of real inertial motions or geodesic lines yet, in that light rays do not share trajectories with massive matters.

The motions of close-to-light-speed particles are something special, in that they would have zero acceleration approximately just as light rays, but they do not share trajectories with light rays and un-close-to-light-speed particles as have discussed.

8.9.2. On 4-dimensional and 3-dimensional Velocities

We all know that vector composition could be expressed in vector summation as well as in way of component square, by the Pythagorean theorem.

For example 4-dimensional velocity in contra variant space as

$$\mathbf{V}_0^\uparrow = (V_0^0, V_0^1, V_0^2, V_0^3) \quad (590)$$

could have a composition of

$$\mathbf{V}_0^\uparrow = \mathbf{V}_0^0 + \mathbf{V}_0^1 + \mathbf{V}_0^2 + \mathbf{V}_0^3 \quad (591)$$

where $\mathbf{V}_0^\mu$ is defined the component vector at the direction μ which has a value V_0^μ .

or the form of

$$-(V_0^\uparrow)^2 = -(V_0^0)^2 + (V_0^1)^2 + (V_0^2)^2 + (V_0^3)^2 \quad (592)$$

where $V_0^0 = \frac{cdt}{dt} = c$, and $V_0^\uparrow$ is module of $\mathbf{V}_0^\uparrow$. Only in the condition that matter's velocity catches up close to light speed c , the modules may be close to zero.

As in covariant space, there is

$$\mathbf{V}^\uparrow/0 = (V0/0, V1/0, V2/0, V3/0) \quad (593)$$

That would have a composition of

$$\mathbf{V}^\uparrow/0 = \mathbf{V}0/0 + \mathbf{V}1/0 + \mathbf{V}2/0 + \mathbf{V}3/0 \quad (594)$$

or the form of

$$-(V^\uparrow/0)^2 = -(V0/0)^2 + (V1/0)^2 + (V2/0)^2 + (V3/0)^2 \quad (595)$$

where $V0/0 = \frac{cd\tau}{d\tau} = c$, and $(V^\uparrow/0)^2$ could have a value uncertain to compare with c^2 so that the setting of $\mathcal{L} = -c^2$ is really a fault treatment.

For light rays, the contra variant velocity will be composed to be nonvanishing

$$\mathbf{c}_0^\uparrow = \mathbf{c}_0^0 + \mathbf{c}_0^1 + \mathbf{c}_0^2 + \mathbf{c}_0^3 \quad (596)$$

or

$$-(c_0^\uparrow)^2 = -(c_0^0)^2 + (c_0^1)^2 + (c_0^2)^2 + (c_0^3)^2 \neq 0 \quad (597)$$

where $c_0^0 = \frac{cdt}{dt} = c$.

But their velocity composition in covariant space there is

$$\mathbf{c}^\uparrow/0 = \mathbf{c}0/0 + \mathbf{c}1/0 + \mathbf{c}2/0 + \mathbf{c}3/0 \quad (598)$$

or

$$-(c^\uparrow/0)^2 = -(c0/0)^2 + (c1/0)^2 + (c2/0)^2 + (c3/0)^2 = 0 \quad (599)$$

where $c0/0 = \frac{cd\tau}{d\tau} = c$, and because the summation of the last 3 items is c^2 we will then gain zero value of $c^\uparrow/0$ just as the Lagrangian has been defined.

We have seen that 4-dimensional velocity does not always make sense. It is not a bad idea to return to the expressions of 3-dimensional velocity in some conditions that we call them true velocities.

$$\mathbf{V}_0^s = \mathbf{V}_0^1 + \mathbf{V}_0^2 + \mathbf{V}_0^3 \quad (600)$$

where, $\mathbf{V}_0^s$ is total quantity of the true velocity.

or the form of

$$(V_0^s)^2 = (V_0^1)^2 + (V_0^2)^2 + (V_0^3)^2 < c^2 \quad (601)$$

In some cases $(V_0^s)^2$ could be greater than $(c_0^s)^2$.

As in covariant space, there is

$$\mathbf{V}s/0 = \mathbf{V}1/0 + \mathbf{V}2/0 + \mathbf{V}3/0 \quad (602)$$

or

$$(Vs/0)^2 = (V1/0)^2 + (V2/0)^2 + (V3/0)^2 \quad (603)$$

where, $(Vs/0)^2$ could have a value less than or equal to or great than c^2 .

For light rays, the contra variant velocity will be composed to be

$$\mathbf{c}_0^s = \mathbf{c}_0^1 + \mathbf{c}_0^2 + \mathbf{c}_0^3 \quad (604)$$

or

$$(c_0^s)^2 = (c_0^1)^2 + (c_0^2)^2 + (c_0^3)^2 \leq c^2 \quad (605)$$

And in covariant space there is

$$cs/0 = c1/0 + c2/0 + c3/0 \quad (606)$$

or

$$(cs/0)^2 = (c1/0)^2 + (c2/0)^2 + (c3/0)^2 = c^2 \quad (607)$$

It could be found that in some cases, matter's contra variant velocity may be greater than light contra variant velocity that

$$(V_0^s)^2 \geq (c_0^s)^2 \quad (608)$$

or

$$(Vs/0)^2 \geq (cs/0)^2 \quad (609)$$

8.9.3. On Light Momentum and Massive Matter's Momentum

As have discussed, momentum and energy refer to that issues which will keep conservation during whole movement, so much as we cannot confirm that they are only valid in contra variant space. We have seen that for light rays, frequencies could be employed to perform momentum and energy, but for massive matters, there will be something different. It is based on physical realities that the conservative quantities be defined. Therefore, I will not discuss any transformed forms of momentum and energy, so far as to deal them as tensors. Owing to mass energy equation, matter's total mass could be seen to be equivalent to total energy. These treatments will not do any harms to our understandings on realities. One of the reasons is that conservative quantities are of something invariant. For example, as light speed c is used as invariant value in some cases, it could be treated as a scalar. Momentum and energy keep conservation except inputs or outputs.

As we talk about light momentum of square forms, the modulus of 4-dimensional velocities $(c_0^1)^2$ in Equation (597) and $(c \uparrow/0)^2$ in Equation (599) cannot be selected for momentum equation, in that $(c \uparrow/0)^2$ is zero and $(c_0^1)^2$ is close to zero in weak fields. As for the modulus of 3-dimensional velocities $(c_0^T)^2$ in Equation (605), it varies in gravitational fields. So that $(cs/0)^2 \equiv c^2$ in Equation (607) is the only choice.

For square momentum of massive matters, the modulus of 4-dimensional velocities $(V_0^1)^2$ in Equation (592) and $(V \uparrow/0)^2$ in Equation (595) also cannot be employed because of similar reasons. The $(Vs/0)^2$ in Equation (603) is not a conservative quantity, as has been proved previous, while $(V_0^s)^2$ in Equation (601) is so that the latter could be employed for momentum expression.

As has mentioned light momentum will be performed in special forms that

$$P^2 = m^2[(c1/0)^2 + (c2/0)^2 + (c3/0)^2] \quad (610)$$

or

$$\mathbf{P} = \mathbf{P}_1 + \mathbf{P}_2 + \mathbf{P}_3 = mc1/0 + mc2/0 + mc3/0 \quad (611)$$

The most surprising is that the vector of momentum would have direction others to the tangent line of trajectory. For a light ray propagating in one source fields, the momentum direction will coincide with trajectory at peak point or be asymptotically close to the trajectory in very fast area as shown in Figure 25.




Figure 25. Light momentum vectors on the trajectory

The surprising forms of light momentum encourages me to make an assumption that a light ray refracted from a media to another would have equivalent light momentums.

Momentum of massive matters could be expressed as

$$P^2 = m^2[(V_0^1)^2 + (V_0^2)^2 + (V_0^3)^2] \quad (612)$$

or

$$\mathbf{P} = \mathbf{P}_1 + \mathbf{P}_2 + \mathbf{P}_3 = m\mathbf{V}_0^1 + m\mathbf{V}_0^2 + m\mathbf{V}_0^3 \quad (613)$$

The vector of momentum of massive matters will always coincide with trajectory in gravitational fields.

9. Relativistic Release and Relativistic Frequency Shift

It is predicted that in one source field, the inflow of matters may cause relativistic release. Sole matters moving into the source center for a separation is of course exactly a kind of inflow. But I want to point out that the inflows of accretions of quasars and active galactic nuclei are of normal performances in celestial evolutions. As the matter of facts, they have not been well investigated, so that it is necessary to cast a few concentrations on.

9.1. Dynamics of Accretions of Quasars and Active Galactic Nuclei

9.1.1. Galactic Accretions and Planet Rings

The discovery of quasars and active galactic nuclei is of the greatest advancing of the astronomy in the 1960s [34,35]. Quasars and active galactic nuclei are of the most spectacular extragalactic objects not only because of their extraordinary giant radiations [36] but also due to the complicated spectra and very high redshifts. The emission spectra of quasars and active galactic nuclei are really non-thermal continuum [37]. Those issued models for quasars and active galactic nuclei in the past decades, such as the so-called standard model [38] or unified model [39], have not really interpreted the mechanism of accretion inflow and energy release. This is the reason that I want to sponsor a brief discussion on the dynamics of accretions of quasars and active galactic nuclei.

Galactic accretion disks exactly experience resemblant kinematics with planet ring systems in that both of them are of surrounding dusts revolving the center source, loosely inside the Roche limit [40]. I am going to sponsor a study firstly with respect to the planet ring systems so that to renovate those recognitions upon the evolution and fueling mechanism of galactic accretions.

Planet rings are not static structures [41,42]. Their evolutions involve with numerous internal and external processes in which Keplerian shear acts the key role. It causes rings to spread [43] in all process. As we have seen, Saturn rings usually have that striking refined structures.

In the condition that ring particles are small enough, the electromagnet interactions will then overcome the inertial forces so that to control the motion within the rings. Such particles or together with gases getting to form colloid, will be as a whole to perform fluid behaviors so as to bear pressure. Under the driving effects from outer boundaries, a fluid ring may go bolder cubically because pressure increasing helps particles to move deviate vertically to the ring plane, such as the Halo ring of Jupiter [44]. Spectrum observations predict that the particles in Halo ring are less than submicron. Optical observations also show that Halo ring seems blue and gray with respect to that the main rings show red color [44]. It is because that the fine particles scatter more blue rays, just like our sky does, while coarse particles scatter more red.

Firstly, I want to make a discussion based on a fluid ring in the condition that the center velocity equals to Keplerian velocity.

9.1.2. The Dynamic Models of Fluid Rings

For Keplerian motions of matters around center object, there is

$$mr\omega_k^2 = \frac{GMm}{r^2} \quad (614)$$

where, m is mass of a particle, G is gravitational constant, M is the mass of center source, r is distance to the center and $\omega_k = \frac{v_k}{r}$ is angular velocity while v_k is tangent velocity.

The tangent velocity and angular velocity will increase as radius getting smaller that

$$v_k = \frac{(GM)^{0.5}}{r^{0.5}} \text{ and } \omega_k = \frac{(GM)^{0.5}}{r^{1.5}} \quad (615)$$

Keplerian kinetic energy could be expressed as

$$E_k = \frac{1}{2}mv_k^2 = \frac{1}{2}m \frac{GM}{r} \quad (616)$$

Keplerian motion determines constant angular momentum for a particle at specific position.

$$L_k = rmv_k = mr^{0.5}(GM)^{0.5} \quad (617)$$

Namely, there is Keplerian shear at the specific position, and the shearing rate can be derived to be

$$\frac{dv_k}{dr} = -\frac{(GM)^{0.5}}{2r^{1.5}} \text{ and } \frac{d\omega_k}{dr} = -\frac{3(GM)^{0.5}}{2r^{2.5}} \quad (618)$$

I do not think that the dynamics of planet rings is very easy to be built in that the evolutions of the rings must be rarely sophisticated. I prefer to suggest three simplified models that may reveal a little bit of probabilities.

Model 1: Central Keplerian motion with driving front and resistant front

It is imagined that a ring keeps Keplerian velocity at its center position while outer part has velocity lower than Keplerian velocity so that to form driving force, and inner part has velocity higher than Keplerian velocity so that to form resistant force. In this way, the driving force and resistant force could help to form fluid pressure to bolder the ring at vertical dimension as shown in Figure 26. That really others to the flat rings as that of Saturn.




Figure 26. Fluid ring in one source field.

For example, in the outer part of a fluid ring, if the velocity is lower to form driving force, the pressure could be calculated as

$$p = \int_{r_o}^r (\rho \frac{GM}{x^2} - \rho x \omega^2) dx \quad (619)$$

where, ρ is density of fluid, r_o is radius from source center to the outer edge of fluid ring, and the angular velocity ω is less than Keplerian angular velocity ω_k of the very point so that to give driving force. If a linear function is assumed for an element of outer part

$$\omega^2 = (\omega_k^2 - \frac{r-r_c}{r_o-r_c} \Delta\omega_o^2) \quad (620)$$

where, r_c refers to radius of pressure center and $\Delta\omega_o^2$ represents the angular velocity square difference with respect to ω_k^2 , which is the Keplerian angular velocity square at the very position r .

This equation presents an active falling of the outer part particles because of losing of energy and velocity.

Concern that at every position there is the relationship

$$\rho \frac{GM}{r^2} = \rho r \omega_k^2 \quad (621)$$

The Equation (619) becomes

$$p = \int_{r_o}^r \rho x \frac{x-r_c}{r_o-r_c} \Delta\omega_o^2 dx \quad (622)$$

Giving the pressure of inner edge and outer edge equal to zero, and constant density, and $r_o - r_c \ll r_c$, there will be an approximate solution of center pressure

$$p_c \approx \frac{1}{2} \rho r_c (r_o - r_c) \Delta\omega_o^2 = \frac{1}{2} \rho r_c b \Delta\omega_o^2 \quad (623)$$

where, $b = r_o - r_c$ represents the outer radius of the fluid ring.

On another side, the resistant force determined by the higher velocity of inner parts, with velocity as

$$\omega^2 = (\omega_k^2 + \frac{r_c-r}{r_c-r_i} \Delta\omega_i^2) \quad (624)$$

In which ω^2 is greater than ω_k^2 so that to form a passive resistant force.

There will be the center pressure

$$p_c \approx \frac{1}{2} \rho r_c a \Delta\omega_i^2 \quad (625)$$

where, $a = r_c - r_i$ represents inner radius of fluid ring.

On the direction vertical to the plane, it is easily to calculate [36] that

$$p_c \approx \int_0^h \rho r \omega_{ck}^2 \frac{y}{r} dy = \frac{\rho h^2}{2} \omega_{ck}^2 \quad (626)$$

where, h represents vertical height from top or the bottom to the center of the ring, y is the distance to pressure center and ω_{ck} is Keplerian angular velocity at pressure center.

Thus, the relationships between semi axes are

$$a \approx \frac{b \Delta\omega_i^2}{\Delta\omega_o^2} \quad (627)$$

$$a \approx \frac{h^2 \omega_{ck}^2}{r_c \Delta\omega_i^2} \quad (628)$$

$$b \approx \frac{h^2 \omega_{ck}^2}{r_c \Delta\omega_o^2} \quad (629)$$

The outer edge of the ring may be called driving front, depending on a random falling of outer particles, that forms an ambiguous boundary. The inner edge of the ring comes from passive driving, may be called driven front, so that to form a sharp edge as seen in Halo ring [44], just like curved water surface in an accelerated container. Of course, fluid rings also can be flattened and split to be refined structures so that to perform more complicated than Halo ring.

Model 2: Driving condition with no shear motion in inner part

It is assumed that the driving force will form a pressure gradient in inner part that just fit the gravity difference coming from Keplerian velocity difference so that to form no shear motion as shown in Figure 27.




Figure 27. Driving condition with no shear motion in inner part.

(Note, the curved line of ring shape is an imaginary result probable, concerning variable density).

There will be

$$\omega_{ik} = \omega_i = \omega_c \quad (630)$$

It is said that any differential slice in the inner part may be accelerated by outer squeezing to catch up the velocity ω_{ik} .

That indicates an increase of the angular velocity square to meet the force equilibrium

$$\Delta\omega^2 = \omega_{ik}^2 - \omega_k^2 = \frac{GM}{r_i^3} - \frac{GM}{r^3} \quad (631)$$

Thus the ring pressure could be written as

$$p = \int_{r_i}^r \rho x \Delta\omega^2 dx = \int_{r_i}^r \rho x \left(\frac{GM}{r_i^3} - \frac{GM}{x^3} \right) dx \quad (632)$$

Assume a constant density, the central pressure is about

$$\begin{aligned} p_c &\approx \rho \frac{GM}{r_i^2} (r_c - r_i) - \rho \frac{GM}{r_i r_c} (r_c - r_i) \\ &\approx \rho \frac{GM}{r_i^2} \left(1 - \frac{r_i}{r_c}\right)^2 \end{aligned} \quad (633)$$

That will lead to a special state that inner part of the ring has no Keplerian shear. But this state may need strict conditions and may be destroyed by further evolution in a long term.

Model 3: Momentum conversion and ring split

After then, I want to discuss the split of a ring under the condition of developed Keplerian shear. If a fluid ring is well Keplerian sheared, it could be predicted that the ring will go thinner. We know that there may be Keplerian shear at any position in the ring. But we can make an analysis on the angular momentum conversion in the middle point which could then at last split the ring to two parts. For convenience, it could be defined as two rings originally, the ring i and the ring $i+1$ as shown in Figure 28. They will have their own Keplerian velocity points at r_i and r_{i+1} . In a time interval, the two rings will complete quite amount of angular momentum conversion because of the Keplerian shear, which is just like viscous friction acts in the interface that makes the conversion.




Figure 28. Angular momentum conversion between neighboring rings.

Setting the mass of rings $m_i = m_{i+1} = m$, the relative angular momentum difference between two neighboring rings could be estimated as

$$rm\Delta v = -m \frac{(GM)^{0.5}}{2r^{0.5}} \Delta r \quad (634)$$

where $\Delta r = r_{i+1} - r_i$ and $r \approx 0.5(r_{i+1} + r_i)$

The exchange of angular momentum may be expressed as

$$\Delta L = -\beta m \frac{(GM)^{0.5}}{2r^{0.5}} \Delta r \quad (635)$$

where, momentum exchange ratio $0 \leq \beta \leq 1$.

It is said that after a Keplerian shear, the outer one should get increase of angular momentum and the inner one should get that decline. For the reasons of kinetic mechanism, both the inner ring and outer ring will be driven apart from original position. If inner ring goes a difference δr , the difference of Keplerian angular momentum could be calculated as

$$\delta L_k = m \frac{(GM)^{0.5}}{2r^{0.5}} \delta r \quad (636)$$

Let δL_k equals to ΔL , it is obtained that

$$\delta r = \beta \Delta r \quad (637)$$

For the inner ring, it goes to shrink by falling a (minus) δr separation so that to form a new state. And for the outer ring, it expands by going up to a (positive) δr separation.

We have seen that this is not a whole dynamic analysis because that the pressure in the ring has also taken the effect of angular momentum conversion directly, but that does no harm for the conclusion that Keplerian shear, i.e., the viscous shear in the ring, leads to angular momentum conversion and rings split. One can make detailed study with Navier Stokes equations or the renovated ones.

A process of ring spreading around a source center is an irreversible process with entropy production.

9.1.3. Shearing Dissipation in Fluid Rings

This issue is carried out to estimate the shearing dissipation so that to discuss the evolution by Keplerian shear. Of course, the evolution would involve together with momentum conversion and ring split. But in the way of theoretical methodology, it is better to be investigated in two steps, the angular momentum conversion by Keplerian shear, and the energy momentum conservation during rings split.

Angular momentum conversion must be done across the interface between the two rings, so that the first step analysis is reasonable. I prefer to focus on the states before and after momentum conversion of two neighboring rings as has been shown in Figure (25). Before conversion, the neighboring rings have been separated by $\Delta r = r_{i+1} - r_i$, ($r_{i+1} > r_i$), with velocities

$$v_i = \frac{(GM)^{0.5}}{r_i^{0.5}} \text{ and } v_{i+1} = \frac{(GM)^{0.5}}{r_{i+1}^{0.5}} \quad (638)$$

The difference between v_i and v_{i+1} could be written after Equation (618)

$$\Delta v = -\frac{(GM)^{0.5}}{2r^{1.5}} \Delta r \quad (639)$$

where, $r \approx 0.5(r_{i+1} + r_i)$, Δv is defined as minus quantity while Δr as plus. Namely,

$$v_{i+1} = v_i + \Delta v \quad (640)$$

Considering ring mass equals to each other, two conditions would be taken into study.

As angular momentum conversion happens, there are

$$v_{i,after} = v_i + \beta \Delta v \text{ and } v_{i+1,after} = v_i + \Delta v - \beta \Delta v \quad (641)$$

As Δv has been defined minus, the inner ring is really slowed down and the outer ring is accelerated. Because of $0 \leq \beta \leq 1.0$, kinetic energy may not keep conservation. With original kinetic energy of

$$E_{before} = \frac{1}{2} m v_i^2 + \frac{1}{2} m (v_i + \Delta v)^2 \quad (642)$$

and the kinetic energy after conversion of

$$E_{after} = \frac{1}{2} m (v_i + \beta \Delta v)^2 + \frac{1}{2} m (v_i + \Delta v - \beta \Delta v)^2 \quad (643)$$

Thus, there is the dissipation energy without considering potential variations

$$Dissipation_1 = E_{before} - E_{after} = \beta m \Delta v^2 - \beta^2 m \Delta v^2 \quad (644)$$

Considering the Equation (639), it turns to be

$$Dissipation_1 = \beta(1 - \beta) m \frac{GM}{4r^3} \Delta r^2 \quad (645)$$

Then go to the second step that after the first step of angular momentum conversion, the outer part of inner ring loses momentum so that provides driving force to drive the inner ring to inflow and the inner part of outer ring will be accelerated to higher momentum so that to squeeze the outer

ring to outflow. This evolution will perform a little sophisticated in that the inflow ring should be accelerated by gravity and the outflow ring should be slow down by gravity. We know that the inflow of a ring will cause half potential release surplus with respect to the Keplerian energy requirement and the outflow will experience potential withdraw. And also, we know that the gravity is perpendicular to statistic velocity of the ring. These issues could be solved in fluid flow anyway, even though that must involve with the evolution of vortexes.

Now we are going to check the energy conservation of final state. After the inner ring inflows a separation δr and the outer ring outflows a separation δr , they come to a new state of Keplerian balance. Taking the inner ring for granted, before Keplerian shear, its total energy involves with Keplerian kinetic energy and potential

$$E_{i,before} = \frac{1}{2} m \frac{GM}{r_i} + 0 \quad (646)$$

After Keplerian shear and inflowing a separation δr , total energy becomes

$$E_{i,after} = \frac{1}{2} m \frac{GM}{r_i - \delta r} - m \frac{GM}{r_i^2} \delta r \quad (647)$$

and for outer ring, the energy before

$$E_{i+1,before} = \frac{1}{2} m \frac{GM}{r_{i+1}} + 0 \quad (648)$$

The energy of final state

$$E_{i+1,after} = \frac{1}{2} m \frac{GM}{r_{i+1} + \delta r} + m \frac{GM}{r_{i+1}^2} \delta r \quad (649)$$

Total energy difference before and after the two steps could be estimated by

$$\begin{aligned} & E_{i,before} - E_{i,after} + E_{i+1,before} - E_{i+1,after} \\ &= \left(\frac{1}{2} m \frac{GM}{r_i} - \frac{1}{2} m \frac{GM}{r_i - \delta r} \right) + \left(\frac{1}{2} m \frac{GM}{r_{i+1}} - \frac{1}{2} m \frac{GM}{r_{i+1} + \delta r} \right) + \left(\frac{GM}{r_i^2} \delta r - \frac{GM}{r_{i+1}^2} \delta r \right) \end{aligned} \quad (650)$$

The first two items could be transformed as

$$\begin{aligned} & \left(\frac{1}{2} m \frac{GM}{r_i} - \frac{1}{2} m \frac{GM}{r_i - \delta r} \right) \\ &= \frac{1}{2} m \frac{GM}{r_i} - \frac{1}{2} m GM \left(\frac{1}{r_i} + \frac{1}{r_i^2} \delta r + \frac{1}{r_i^3} \delta r^2 \right) \\ &= GMm \left(-\frac{1}{2r_i^2} \delta r - \frac{1}{2r_i^3} \delta r^2 \right) \end{aligned} \quad (651)$$

The middle two items could be transformed as

$$\begin{aligned} & \left(\frac{1}{2} m \frac{GM}{r_{i+1}} - \frac{1}{2} m \frac{GM}{r_{i+1} + \delta r} \right) \\ &= \frac{1}{2} m \frac{GM}{r_{i+1}} - \frac{1}{2} m GM \left(\frac{1}{r_{i+1}} - \frac{1}{r_{i+1}^2} \delta r + \frac{1}{r_{i+1}^3} \delta r^2 \right) \\ &= GMm \left(\frac{1}{2r_{i+1}^2} \delta r + \frac{1}{2r_{i+1}^3} \delta r^2 \right) \\ &= GMm \left[\frac{1}{2(r_i + \Delta r)^2} \delta r + \frac{1}{2r_{i+1}^3} \delta r^2 \right] \\ &= GMm \left[\frac{1}{2r_i^2} \left(1 - \frac{2\Delta r}{r_i} \right) \delta r + \frac{1}{2r_{i+1}^3} \delta r^2 \right] \end{aligned} \quad (652)$$

so that the previous four items are

$$-GMm \frac{\Delta r \delta r}{r_i^3} = -GMm \beta \frac{\Delta r^2}{r_i^3} \quad (653)$$

The last two items are

$$\begin{aligned}
& \left(\frac{GMm}{r_i^2} \delta r - \frac{GMm}{r_{i+1}^2} \delta r \right) \\
&= \frac{GMm}{r_i^2} \delta r - \frac{GMm}{(r_i + \Delta r)^2} \delta r \\
&= \frac{GMm}{r_i^2} \delta r - \frac{GMm}{r_i^2} \delta r \left(1 - \frac{2\Delta r}{r_i} \right) \\
&= 2 \frac{GMm}{r_i^3} \Delta r \delta r \\
&= 2GMm\beta \frac{\Delta r^2}{r_i^3}
\end{aligned} \tag{654}$$

The energy difference is total energy dissipation

$$\begin{aligned}
Dissipation &= E_{i,before} - E_{i,after} + E_{i+1,before} - E_{i+1,after} \\
&= \beta GMm \frac{\Delta r^2}{r_i^3}
\end{aligned} \tag{655}$$

One can calculate the gravitational potential release of a single ring inflowing, although it does not sound reasonable for a comparison with dissipations because there is another ring outflowing. Anyway, that is

$$dW = m \frac{GM}{r^2} \delta r = \beta m \frac{GM}{r^2} \Delta r \tag{656}$$

Thus, we get the ratio of dissipation and potential release

$$\frac{Dissipation}{dW} = \frac{\Delta r}{r} \tag{657}$$

Or the ratio of dissipation-1 and potential release

$$\frac{Dissipation_1}{dW} = \frac{(1-\beta)\Delta r}{4r} \tag{658}$$

where $\beta \neq 0$.

Ring evolution dynamics indicates that the inner half of an accretion inclines to inflow and outer half inclines to outflow by and large. Additionally, the process of vortex evolution would dissipate, even though the two dissipations are not independent.

$$Dissipation_2 = Dissipation - Dissipation_1 = \frac{1}{4}\beta(3 + \beta)m \frac{GM}{r^3} \Delta r^2 \tag{659}$$

It shows that the dissipation in vortex evolution might be greater than that in momentum exchange between rings.

We know that inter-ring separation Δr would be far less than the radius r of relative rings. It is said that in the condition of Keplerian shear, the dissipative energy is a higher rank infinitesimal quantity of that of potential release of inflow, no matter that inner ring potential release will support the outer ring potential withdrawing. Classical quasars could have luminosities of 10^{45} erg s⁻¹ to 10^{46} erg s⁻¹ [45]. It is unimaginable that those releases completely come from gravitational potential energy.

9.1.4. On Particle Rings

For solid particle rings, it is difficult to make a simple analysis on angular momentum conversions and on ring's evolutions. In fact, any head-to-head collisions, and partial collisions of edge-by-edge that might form additional rotations, between two particles, might cause great changes to their trajectories. It could be imagined that for numerous particles in neighboring rings, assuming very small velocity difference between any two Keplerian collision particles, we can predict quasi fluid behavior for Keplerian shear acting on neighboring rings, that in a whole, Keplerian shear help to make angular momentum conversion and ring split. One can also write down those equations of

energy momentum just as that of fluid rings. But all in all, particle rings cannot form pressure so that they will only spread as a very thin disk with perfect and refined structures in the equator plane, unlike the fluid rings, which could form a bold 3-dimensional torus structures. I believe that numerical method could be expected for quasi fluid behavior simulation for Keplerian shear of particle rings, as the effects of the rings of Saturn have shown us.

9.2. Relativistic Release

Any efforts to conceiving the mechanism of relativistic release would encounter great difficulties, because of two reasons that relativistic release has not been well recognized and we still know less about the intrinsic structures of fundamental particles that might be revealed relativistic emissions. But on the other hand, tremendous of observations have shown incredible probabilities for intrinsic emissions and correlated pseudo redshifts, as well as intrinsic redshift of absorptions, which are so intensely corresponding to the inference of relativistic release that I cannot help to make a try.

A direct realization on this issue is that the conservation of electromagnetic momentum and invariance of electromagnetic wave speed will lead to the variation of proficiency of light moment transportation, then the interactions or so-called the forces between particles, especially that in intrinsic structures. The most complicated is that these variations would be anisotropic so that the dynamics must be even profound and comprehensive.

Despite the complexities of relativistic release for a massive matter inflow into the center source, I suggest a mathematic analysis based on an assumption of continuum release. It is still not a true expression of realities, but that may help for more understandings on that topic.

The apparent light speed c_0^1 could be employed to describe the concept of equivalent state in gravitational fields. It may be a state after relativistic release, although this state should be not really stable because there should be energy deficiency in the other two dimensions.

One can discuss the momentum exchange efficiency under the conditions of different apparent light speeds so that to talk about the variation of equivalent state. Another important clue that indicates equivalent state is the fine structure constant. We know that fine structure constant determines the dimensions of condensed matters, of course that may determine the dimensions of intrinsic structures. We can imagine that the variation of apparent light speed could bring about the variation of fine structure constant so that to bring about the variation of the energy of equivalent state.

I don't think all of energy of a matter would take part in the relativistic release, for example, the most fundamental particles may be expected to be made of quasi photons, in common recognitions, they might not be split anymore. Thus, we would rather define the concept of releasable mass $\tilde{m}$. I cannot give an estimation whether the maximum releasable mass should be up to $0.9m_0$ or not, but it could be believed to be quite amount, where m_0 is total rest mass of original equivalent state, which corresponding to the state that the matters do not experience relativistic release.

Now for a center source field, equivalent state of matters is proposed to be

$$\tilde{E} = \tilde{m}(c_0^1)^2 \quad (660)$$

Matters located at different positions in the gravitational field perform different equivalent states. As a matter goes a separation dr along the radial direction, the expression of exceeding energy is

$$d\tilde{E} = 2\tilde{m}c_0^1(c_0^1)'dr \quad (661)$$

As a matter goes from the farthest point to the position of radius r , the relativistic released energy could be integrated to be

$$\tilde{E}_{released} = \int_r^\infty d\tilde{E} = \tilde{m}c^2 - \tilde{m}(c_0^r)^2 = \tilde{m}c^2[1 - (1 - \frac{r^*}{r})^2] \quad (662)$$

Thereafter, we could also define a corresponding concept of the relativistic residual energy that, at a position, matters keep an amount of energy for subsequent releases,

$$\tilde{E}_{residual} = \tilde{m}c^2(1 - \frac{r^*}{r})^2 \quad (663)$$

It seems that the residual energy may go zero as matters reaches r^* . It is just a solution of theoretical model. In practice, relativistic release will be intermittent, and matters perhaps to experience flattenizations as they close to r^* so that to bring about instabilities. As results, residual energy may not go zero at the end in practice. This effort is just to manage to perform kind of probabilities of relativistic release.

It can be calculated that most part of energy is kept before $10r^*$

$$\tilde{E}_{residual}(10r^*) = 0.81\tilde{m}c^2 \quad (664)$$

Now it is naturally to define the release rate by derivative of release energy along the radial direction to the center source. Released energy could be the integration of release distribution function from a position farthest to the position r or the minus inversely

$$\tilde{E}_{released} = \int_{\infty}^r e dr = \int_r^{\infty} -e dr \quad (665)$$

On the other hand, the residual energy could also be defined as an integration of release distribution from position r^* to position r or the minus form

$$\tilde{E}_{residual} = \int_{r^*}^r e dr = - \int_r^{r^*} e dr \quad (666)$$

Thus, the release distribution could be derivated to be

$$e = 2\tilde{m}c^2(1 - \frac{r^*}{r})\frac{r^*}{r^2} \quad (667)$$

It is easy to calculate that the peak value locates at the position

$$r_{max} = \frac{3}{2}r^* \quad (668)$$

with the maximum release distribution

$$e_{max} = \frac{8}{27}\frac{\tilde{m}c^2}{r^*} \quad (669)$$

One can calculate the release intensity of accretion inflow. Setting equivalent mass for every single ring in an accretion, the luminosity could be derived as

$$l = 2\tilde{\rho}^*vc^2(1 - \frac{r^*}{r})\frac{r^{*2}}{r^3} \quad (670)$$

where, $\tilde{\rho}^*$ is matter's assuming density at the position r^* and v is inflow velocity.

Suppose the accretion spreading sufficiently, it can be calculated that the most luminous area is at the position

$$r_{lmax} = \frac{4}{3}r^* \quad (671)$$

As well as maximum luminosity

$$l_{max} = \frac{27}{128}\frac{m\tilde{\rho}^*vc^2}{r^*} \quad (672)$$

Release distribution and accretion luminosity could be shown in Figure 29.




Figure 29. Release distribution and accretion luminosity.

The Equation (671) represents intensity of release at a specific position in an accretion disk. It is revealed that the peak luminosity should not happen at the edge of horizon event, but at a little outer position, just as that was shown in the event-horizon-scale images of M87 taken by the Event Horizon Telescope Collaboration with wavelength of 1.3mm [46].

9.3. Relativistic Emission Lines and Relativistic Redshift

Eyeing on the stimulated release of electrons in atoms, we could carry out a concept of exceeding ratio for relativistic release of an energy structure.

$$\sigma = \frac{\tilde{E}_{excited} - \tilde{E}_{ground}}{\tilde{E}_{ground}} \quad (673)$$

where $\tilde{E}_{excited}$ and $\tilde{E}_{ground}$ are energy of excited state and ground state of specific structure. Exceeding ratios for relativistic release may be more different from that of stimulated release of electrons, probably, they might be quite small or big. We know less about that.

In one source field, exceeding ratio would be variable with position as the so-called equivalent state has shown.

$$\sigma = \sigma_{\infty} \left(1 - \frac{r^*}{r}\right)^2 \quad (674)$$

Thus, frequencies of specific emission rays could be written as

$$\nu_r = \nu_{\infty} \left(1 - \frac{r^*}{r}\right)^2 \quad (675)$$

Case $r \gg r^*$ it could be simplified to be

$$\nu_r = \nu_{\infty} \left(1 - \frac{2r^*}{r}\right) \quad (676)$$

These emissions of course include but not limit to electron transition emissions that we are more familiar to that could be easily certificated comparing to stimulated release on the Earth. This kind of frequency will be quite different with which we have known well in weak field, in that it looks like redshifted after photons reach the Earth. But it is just a pseudo redshift because the emissions have not experienced real redshift at the emission time. The fact is that it is only verified to be redshifted by comparing to the spectrum characteristics of a matter in weak fields. In this case, it is just redshift seemingly. It could be called relativistic redshift.

In practice, emission lines will experience gravitational redshift to arrive a farthest position. The Equation (468) could also be employed to interpret light mass energy variation in gravitational field, so that the detectable frequency could be expressed as

$$\nu_{r \rightarrow d} = \nu_{\infty} \left(1 - \frac{r^*}{r}\right)^2 e^{-\frac{r^*}{2r}} \quad (677)$$

If a detector is at a position close to the Earth, one should furtherly calculate the gravitational frequency shift in gravitational field of the Earth. That will not be presented here.

Case $r \gg r^*$, the Equation (677) could be simplified to be

$$\nu_{r \rightarrow d} \approx \nu_{\infty} \left(1 - \frac{r^*}{r}\right)^2 \left(1 + \frac{r^*}{2r}\right)^{-1}$$

$$= \nu_{\infty}(1 - \frac{r^*}{r})^{2.5} \approx \nu_{\infty}(1 - 2.5 \frac{r^*}{r}) \tag{678}$$

Propose a definition of relativistic frequency shift based on frequency

$$z_{rel} = \frac{\nu_{\infty}-\nu_r}{\nu_r} = (1 - \frac{r^*}{r})^{-2} - 1 \tag{679}$$

Case $r \gg r^*$ it could be simplified to be

$$z_{rel} = \frac{\nu_{\infty}-\nu_r}{\nu_r} = \frac{2r^*}{r} \tag{680}$$

It is an intrinsic frequency shift rather than cosmological redshift. Note that ν_r keep invariant during propagation process if neglect the effect of gravitational redshift. ν_{∞} is the comparative frequency of equivalent radiation in no gravity condition.

Detectable redshift should be calculated as

$$z_{r \rightarrow d} = \frac{\nu_{\infty}-\nu_{r \rightarrow d}}{\nu_{r \rightarrow d}} = (1 - \frac{r^*}{r})^{-2} e^{\frac{r^*}{2r}} - 1 \tag{681}$$

Case $r \gg r^*$ it could be simplified to be

$$z_{r \rightarrow d} = (1 - \frac{r^*}{r})^{-2.5} - 1 \approx \frac{2.5r^*}{r} \tag{682}$$

Relativistic frequency shift and detectable frequency shift could be calculated for comparison in Table 6.

Table 6. Relativistic and detectable frequency shifts at different positions.

Emission positions	Relativistic frequency shifts	Detectable frequency shifts
1.3r*	17.778	26.585
1.4r*	11.250	16.508
1.5r*	8.000	11.561
2.0r*	3.000	4.136
3.0r*	1.250	1.658
10.0r*	0.235	0.298
20.0r*	0.108	0.136

The highest redshift we have observed on quasars is up to 16.4 [47,48]. It could be expected that higher redshift would be seen to be up to more than 20.0 in recent future. Perhaps higher redshifts have already been observed but not certificated.

Perhaps, up to now, we have found the probable way by which the accretions of quasars and active galactic nuclei release energy. It has been pointed out that the giant radiations of quasars and active galactic nuclei are not of thermal continuum. It has ever been believed that the radiations come from the energy of gravitational potential conversion [36], but we know that the energy of gravitational potential conversion must release radiation by the way of thermal black-body radiation. This will then lead to a contradiction. And in previous section, the dissipation of potential conversion of an accretion has been proved to be an infinitesimal of total conversion, that furtherly deduces the probabilities of potential origin of accretion radiation.

It is carried out that the detectable redshifts are just the property of relativistic emissions, which obviously rise against the theory of cosmological origin. The distribution statistics of observational high redshifts [49,50] might have indicated the redshifts are of property of the emissions rather than that of cosmic reasons.

Theoretically, it will be rarely probable that emission lines of multiple redshifts might be verified in some special observations in that the mechanism implies the probabilities although some of them may be ambiguous perhaps one of them dominates others as has been discussed. In fact, observations on frequency shift and line width and some other aspects of emissions might have present the performances of multiple redshifts. Celestial observations have always brought us surprises, multiple redshift emission might be the new one.

9.4. Broad Lines and Narrow Lines

In the emissions of quite amount of continuous inflow matters, the exciting width that the inflow involved sustains at radial direction may cause specific continuous emission distribution. As a result, that could be certificated to be a broadened line in spectrum diagram. For a massive accretion around a galactic nucleus, the Kepler shear may be employed to interpret the mechanism of inflow of a ring. If there is an inflow of a ring at position r with exciting width of Δr , the emission frequency might be distributed from ν_r to $\nu_{r+\Delta r}$. Considering the effect of gravitational redshift, one can get a line width expressed by frequency after Equation (677) that

$$w_\nu = \nu_{r+\Delta r \rightarrow d} - \nu_{r \rightarrow d} = \left[\left(1 - \frac{r^*}{r+\Delta r}\right)^2 e^{-\frac{r^*}{2(r+\Delta r)}} - \left(1 - \frac{r^*}{r}\right)^2 e^{-\frac{r^*}{2r}} \right] \nu_\infty \quad (683)$$

Case $r \gg r^*$ and $r \gg \Delta r$ it could be simplified to be

$$w_\nu \approx \left(1 - \frac{2.5r^*}{r+\Delta r} - 1 + \frac{2.5r^*}{r}\right) \nu_\infty = \left(\frac{1}{r} - \frac{1}{r+\Delta r}\right) 2.5r^* \nu_\infty \approx \frac{2.5\Delta r r^*}{r^2} \nu_\infty \quad (684)$$

or by wave length

$$\begin{aligned} w_\lambda &= \lambda_r - \lambda_{r+\Delta r} = c/\nu_{r \rightarrow d} - c/\nu_{r+\Delta r \rightarrow d} \\ &= \left[\left(1 - \frac{r^*}{r}\right)^{-2} e^{\frac{r^*}{2r}} - \left(1 - \frac{r^*}{r+\Delta r}\right)^{-2} e^{\frac{r^*}{2(r+\Delta r)}} \right] \lambda_\infty \end{aligned} \quad (685)$$

Case $r \gg r^*$ and $r \gg \Delta r$ it could be simplified to be

$$w_\lambda \approx \left(1 + \frac{2.5r^*}{r} - 1 - \frac{2.5r^*}{r+\Delta r}\right) \lambda_\infty \approx \frac{2.5\Delta r r^*}{r^2} \lambda_\infty \quad (686)$$

which could also be amount to a so-called Doppler velocity by frequency as

$$v_{D\nu} = \frac{w_\nu}{\nu_r} c \quad (687)$$

or Doppler velocity by wave length as

$$v_{D\lambda} = \frac{w_\lambda}{\lambda_r} c \quad (688)$$

where, c is light speed.

Case $r \gg r^*$ it could be simplified to be

$$v_{D\nu} \approx v_{D\lambda} \quad (689)$$

These equations show that in the regions near to r^* , relativistic emissions should have broader line widths, and in the regions far off the r^* , relativistic emissions should have more narrow line widths. Calculations on line widths of various conditions will be presented in Table 7.

Table 7. Emission line widths at different positions with specific exciting widths.

Emission positions	Exciting widths	w_v/v_∞	v_{Dv} km/s	w_λ/λ_∞	$v_{D\lambda}$ km/s
1.5 r^*	0.1 r^*	0.02327	87681	2.8408	67850
1.5 r^*	0.05 r^*	0.01158	43634	1.5949	38093
3.0 r^*	0.1 r^*	0.01433	11425	0.09752	11006
3.0 r^*	0.05 r^*	0.007239	5772	0.05018	5663
10.0 r^*	0.1 r^*	0.002079	809	0.003492	807
10.0 r^*	0.05 r^*	0.001044	406	0.001756	406
20.0 r^*	0.1 r^*	0.0005706	194	0.0007360	194
20.0 r^*	0.05 r^*	0.0002860	97	0.0003690	97
100.0 r^*	0.1 r^*	0.00002455	7.6	0.00002582	7.6
100.0 r^*	0.05 r^*	0.00001228	3.8	0.00001291	3.8

This result forcefully matches the observational dimensions of broad line regions and narrow line regions. It is said that line widths depend on inflow exciting width, but in totally analysis, they highly relate to the inflow positions. The regions close to r^* may emit lines broader than that of farther regions. Observations show that the sizes of broad line regions are estimated within 0.01 or 0.1pc and line widths are about 1000 kms⁻¹ to 10000 kms⁻¹, while the sizes of narrow line regions are about 0.1pc to 1kpc [49,51,52]. In fact, those certificated narrow lines incline to have been certificated as lower redshift lines. For radio emissions, some of them were observed to have compact cores and extended components [51]. The compact cores could be interpreted that equivalent state in inner areas of accretions would have lower equivalent energy to release lower frequency emissions, while some of the giant extended components of emission pictures might be the jet outflow who emit radio lines.

Relativistic release must perform in sophisticated conditions. The most probable condition is that the release in one of the rings could overwhelming all the others, especially that of the most inner ring. Thus, in most cases, releases of an active galactic nucleus might be certificated by the emissions of the sole ring. One can find that most of narrow lines have been certificated lower redshift, such as NGC 4151 with $z=0.0033$ [53] and MCG-5-23-16 with $z=0.00849$ [54] have narrow Fe K α lines. But on the other hand, their spectra both involve with broad lines [55], that indicate controversies in emission line certifications.

The variability, asymmetry, wave length shift and so much as line broadening of emissions of quasars and active nuclei [50] indicate more sophisticated dynamic conditions than imagine.

9.5. Relativistic Absorption

Most of quasars and active galactic nuclei have giant accretion bodies spreading around their equatorial planes. Case a pulse of light ray goes through some parts of an accretion body, that light ray may experience different absorptions due to their passing positions. If a pulse of light crosses several separate rings or blocks of an accretion, it might experience multiple absorptions. Absorption frequencies depend on equivalent state of the matters that have experienced relativistic release, so that the absorption frequencies could be expressed as

$$\nu_r = \nu_\infty(1 - \frac{r^*}{r})^2 \tag{690}$$

and the detectable frequency is

$$\nu_{r \rightarrow d} = \nu_\infty(1 - \frac{r^*}{r})^2 e^{-\frac{r^*}{2r}} \tag{691}$$

where, ν_r is the frequency of absorption line in that the light rays cross throughout a ring located at a position of r , and ν_∞ is the absorption frequency case that structure would absorb in no-gravity fields. One can of course calculate the widths of absorption lines just like that of emission lines.

$$w_v = [(1 - \frac{r^*}{r+\Delta r})^2 e^{-\frac{r^*}{2(r+\Delta r)}} - (1 - \frac{r^*}{r})^2 e^{-\frac{r^*}{2r}}] \nu_\infty \tag{692}$$

or

$$w_\lambda = [(1 - \frac{r^*}{r})^{-2} e^{\frac{r^*}{2r}} - (1 - \frac{r^*}{r+\Delta r})^{-2} e^{\frac{r^*}{2(r+\Delta r)}}] \lambda_\infty \quad (693)$$

where, Δr is absorption width of a ring that the light ray passes across.
or expressed by Doppler velocity in frequency as

$$v_{D\nu} = \frac{w_\nu}{\nu_r} c \quad (694)$$

or Doppler velocity in wave length as

$$v_{D\lambda} = \frac{w_\lambda}{\lambda_r} c \quad (695)$$

But more different from emissions, absorption spectra must have more narrow lines and lower redshift than emissions, because the absorption rings are mostly at more outer regions with respect to the inner shining emission source, otherwise they would be difficult to be detected.

As has been discussed above, the inflow of inner rings of accretion may act as key role of entire emissions. So, we can image that brighter inner rings emit light rays, and on our view line, they go through some surrounding rings to be absorbed. It is said that, absorbing dusts might be the matters located at surrounding outer positions rather than those so-called insert bodies far away from the emission areas. This image may lead to the so-called relative blueshift, with respect to that of emission lines. Of course, there might be still seldom absorptions happening in inner areas so that we will also observe relative redshift absorption lines occasionally. Given a separation Δr between emission position and absorption position more exterior, the emission frequency could be calculated as

$$\nu_{em \rightarrow d} = \nu_\infty (1 - \frac{r^*}{r_{em}})^2 e^{-\frac{r^*}{2r_{em}}} \quad (696)$$

with redshift of the line

$$z_{em \rightarrow d} = \frac{\nu_\infty - \nu_{em \rightarrow d}}{\nu_{em \rightarrow d}} \quad (697)$$

and absorption frequency varies because a separation Δr between absorption and emission positions

$$\nu_{ab \rightarrow d} = \nu_\infty (1 - \frac{r^*}{r_{ab}})^2 e^{-\frac{r^*}{2(r_{ab} + \Delta r)}} \quad (698)$$

so that the redshift of absorption line is

$$z_{ab \rightarrow d} = \frac{\nu_\infty - \nu_{ab \rightarrow d}}{\nu_{ab \rightarrow d}} \quad (699)$$

And then, there is a blueshift of an absorption line

$$\Delta z = z_{ab \rightarrow d} - z_{em \rightarrow d} = (1 - \frac{r^*}{r_{ab}})^{-2} e^{\frac{r^*}{2(r_{ab} + \Delta r)}} - (1 - \frac{r^*}{r_{em}})^{-2} e^{\frac{r^*}{2r_{em}}} \quad (700)$$

Given $r_{em} = 2.5r^*$ and $\Delta r = 0.5r^*$, it can be calculated that $z_{em} = 2.39$ and $z_{ab} = 1.66$. Here, the absorption lines may show a blueshift of 0.73 with respect to the emission lines.

It was found that there are both absorption broad line region and absorption narrow line region [56] in single system in which those absorption lines are all corresponding to that of emission lines.

Case a continuum spectrum of light rays passes across multiple rings and arrive at an observatory on the Earth, one can then get a set of multiple frequency shift of absorptions. It is because absorptions only depend on the positions in accretions. This does interpret the multiple redshift absorption lines in tremendous of observations [49–51] in those quasars and active galactic nuclei. Especially for that of so-called Lyman-alpha forest [57], a series of regular Lyman-alpha absorption lines, with descending order of redshifts, queue up amazingly at the left side of the great main emission line.

There is a special condition for a matter out flowing from center source, for example the center jet outflow. When a matter at ground state moves a separation toward outer direction, its structural energy should experience energy deficiency in that the energy of equivalent state gets increasing. We

then see that the matter is in the state lack of energy, and generally, the state should be kept until surrounding environment happens to emit amount of specific light rays across, so as the result, the matter gets absorptions. It is said that, this kind of absorptions would be far different from normal relativistic absorptions as that have been discussed above, so that they could be called the super relativistic absorption. As for the mentioned absorption energy, it depends on the position it getting absorbing as well as the position it has ever been in ground state. Note that this kind of absorptions really distinguish from the general relativistic absorptions in that the super relativistic absorption energy depends on the difference between the final emission position and specific absorption position, while the general relativistic absorption is due to property of structure state. It could be calculated as

$$\Delta v = v_{ab} - v_{gr} \quad (701)$$

where, $v_{ab} = (1 - \frac{r^*}{r_{ab}})^2 e^{\frac{-r^*}{2r_{ab}}} v_{\infty}$, $v_{gr} = (1 - \frac{r^*}{r_{gr}})^2 e^{\frac{-r^*}{2r_{gr}}} v_{\infty}$, r_{ab} is the radius at absorption position, and r_{gr} is the radius of the position of equivalent ground state before outflow. This kind of absorption might need greater absorption energy than that of general relativistic absorption. Therefore, it should be called the super relativistic absorption.

To the observer at a farthest position, the absorption line will also perform special frequency shift that

$$z_{super} = \frac{v_{\infty} - \Delta v}{\Delta v} = \frac{1}{(1 - \frac{r^*}{r_{ab}})^2 e^{\frac{-r^*}{2r_{ab}}} - (1 - \frac{r^*}{r_{gr}})^2 e^{\frac{-r^*}{2r_{gr}}}} - 1 \quad (702)$$

Case the jet flow particles of an active galactic nucleus are assigned in a broad area, it could cause a broader absorption pit for a crossing light ray. One can calculate the average redshift and the width of absorption pit.

Additionally speaking on the topic of jet flow, massive particles may be easily accelerated up and keep approaching absolute light speed so as to escape from inner event horizon. On the other hand, apparent light speed may be at lower level in the nearby r^* area so that to form pseudo super luminal motions. A high pseudo super luminal motion may cause Cherenkov radiation, which could generally show polarized propagation. That might also be the reasons for some high energy light rays. One can imagine that the events of γ -ray burst detection delay in observations on SN1987A [58] and GW170817 [59] might be interpreted by pseudo super luminal propagations of neutrinos and gravitational waves.

Acknowledgement: Great many thanks to whom that I sent them emails for communications years ago early after part of the topics were developed. Thereafter, these researches have been completed in a very long time. I appreciate anyone who would like to download and read this article in that it will be a hard work of weeks or longer to make complicated calculations and comprehensive understandings. And I will highly appreciate them who are going to feedback comments and peer reviews that may help for more communications and cooperation.

References

1. A. Einstein, Die Grundlage der allgemeinen Relativitätstheorie, Annalen der Phys., 49,769(1916)
2. J. Jost, Riemannian Geometry and Geometric Analysis, Springer –Verlag Berlin Heidelberg, Sixth Edition 2011, (1965)
3. T. P. Cheng, Relativity, Gravitation, and Cosmology, Oxford University Press (2005)
4. N. Straumann, General Relativity, Springer Science + Business Media Dordrecht (2004, 2013)
5. L. P. Eisenhart, Riemannian geometry, Princeton University Press, Eighth Printing 1997, (1925)
6. П.К.Рашевский, Riemannian Geometry and Tensor Analysis, Technique and Theory Press, (1953)
7. C. Huang, General Relativity Lecture Sheet, China Science Press (2023)
8. A. Einstein, Zur Elektrodynamik bewegter Körper, Ann. Physik, 17,891(1905)
9. H. Minkowski, Nachr. Ges. Wiss. Göttingen Math.-Phys. Kl. 53 (1908)

10. Y. Yu, General Relativity, Peking University Press (1997)
11. H. Jin, Introduction to General Theory of Relativity, Zhejiang University Press (2022)
12. B. Chen, General Relativity, Peking University Press (2018)
13. C. E. Weatherburn, An Introduction to Riemannian Geometry and the Tensor Calculus, Cambridge University Press, (1942)
14. R. Pound, G. Rebka, Gravitational Red-shift in Nuclear Resonance, Physical Review Letters. 3 (9): 439–441 (1959)
15. R. Vessot, M. Levine; E. Mattison et al., Test of Relativistic Gravitation with a Space-Borne Hydrogen Maser, Physical Review Letters. 45 (26): 2081–2084(1980)
16. T. Do, et al., Relativistic Redshift of the Star S0-2 Orbiting the Galactic Center Supermassive Black Hole, Science. 365 (6454) (2019)
17. S. Weinberg, Gravitation and Cosmology, Principles and Applications of the General Theory of Relativity, John Wiley (1972)
18. R. Peierls, More Surprises in Theoretical Physics, Princeton Univ. Press (1991)
19. H. Minkowski, Nachr. Ges. Wiss. Gött. Math.-Phys. Kl. 53 (1908)
20. M. Abraham, Rend. Circ. Matem. Palermo 28, 1 (1909)
21. A. Einstein, A centenary volume letter to M. Besso, 1951(p.138)
22. U. Leonhardt, Optics: Momentum in an Uncertain Light, Nature, 444:823–824 (2006)
23. M. Froeschle, F. Miguard, F. Arenon, in Proceedings of the Hubblecos Venice 1997 Symposium, ESA Publications Division, Noordwijk (1997)
24. I. I. Shapiro, Fourth Test of General Relativity. Physical Review Letters. 13 (26): 789–791 (1964)
25. I. I. Shapiro, M. E. Ash, R. P. Ingalls etc., Fourth Test of General Relativity: New Radar Result. Physical Review Letters. 26 (18): 1132–1135 (1971)
26. J. Roche, What is mass? European Journal of Physics, 26:1–18(2005).
27. Q. Xu, Z. Liu, Ordinary Differential Equations Analytical Methods and Numerical Methods, Publishing House of Electronics Industry, China (2022)
28. B. B. Perera, M. A. McLaughlin, M. Kramer etc., The Evolution of PSR J0737-3039B and a Model for Relativistic Spin Precession, the Astrophysical Journal, 721: 1193-1205(2010)
29. P. Landry, B. Kumar, Constraints on the Moment of Inertia of PSR J0737-3039A from GW170817, arXiv: 1807. 04727v2 [gr-qc] 10 Oct 2018
30. C. W. F. Everitt etc., Gravity Probe B: Final Result of a Space Experiment to Test General Relativity, Phys. Rev. Lett., 106(22): 221101(2011)
31. J.H. Taylor, L.A. Fowler, and P.M. McCulloch, Measurements of General Relativistic Effects in the Binary Pulsar PSR1913+16, Nature, 277:437 (1978)
32. R. Abuter, A. Amorim, N. Anugu etc., Detection of the Gravitational Redshift in the Orbit of the Star S2 Near the Galactic Centre Massive Black Hole, Astronomy & Astrophysics, 615, L15:1-10(2018)
33. Victor M.L., Riccardo D.M., Ivan M. etc., Future prospects for measuring 1PPN parameters using observations of S2 and S62 at the Galactic Center, arXiv: 2410. 22864v2 [astro-ph.GA] 28 Jan 2025
34. E. E. Salpeter, Accretion of interstellar matter by massive objects. Astrophys. J., 140, 796–800, doi:10.1086/147973 (1964)
35. Y. B. Dovich, and I. D. Novikov, The radiation of gravity waves by bodies moving in the field of a collapsing star. Sov. Phys. Dokl., 9, 246 (1964)
36. S. Kato, J. Fukue, S. Mineshige, Black-hole Accretion Disks, Kyoto University Press (2008)
37. R. D. Blandford, H. Netzer, L. Woltjer, Active galactic nuclei, Saas-Fee Advanced Course 20, Lecture Notes 1990, Swiss Society for Astrophysics and Astronomy, Springer-Verlag Berlin Heidelberg (1990)
38. N. I. Shakura, R. A. Sunyaev, Black holes in binary systems. Observational appearance. Astron. Astrophys., 24, 337–355 (1973)
39. S. L. Shapiro, A. P. Lightman, and D. M. Eardley: Astrophys. J. 204 187 (1976)
40. B. Sicardy, Dynamics of planetary rings, Lect. Notes Phys. 682, 183–200 (2006)
41. J. C. Maxwell, On the Stability of the Motions of Saturn's Rings, Cambridge and London: MacMillan and Company. Reprinted in Scientific Papers of James Clerk Maxwell, Vol. 1, Cambridge University Press (1890)

42. S. Charnoz, A. Croda, R. Hyodo, Rings in the Solar System: a Short Review, arxiv, 201805.08963 (2018)
43. L. W. Esposito, Composition, Structure, Dynamics, and Evolution of Saturn's Rings, *Annu. Rev. Earth Planet. Sci.* 38,383-410 (2010)
44. E. D. Miner, R. R. Wessen, J. N. Cuzzi, Planetary Ring Systems, Praxis Publishing Ltd, Chichester, UK (2007)
45. B. Lee, R. Chary, Insights into Physical Conditions and Magnetic Fields from High Redshift Quasars, arxiv: 2207. 07290v1 [astro-ph.GA] 15 Jul 2022
46. Event Horizon Telescope Collaboration, First M87 event horizon telescope results. I. The shadow of the supermassive black hole, *ApJ*, 875, L1, 17 (2019)
47. C.T. Donnan, D.J. Mcleod, J.S. Dulop ect., The Evolution of the Galaxy UV Luminosity Function at Reshift $z \approx 8-15$ from Deep JWST and Ground-based Near-infrared Imaging, arxiv: 2207. 12356v3 [astro-ph.GA] 24 Nov 2022
48. L.Y. Yung, R.S. Somerville, S.L. Finkelstein ect., Are the Ultra-high-redshift Galaxies at $z > 10$ Surprising in the Context of Standard Galaxy Formation Models? arxiv: 2304. 04348v2 [astro-ph.GA] 11 Nov 2023
49. K. Huang, Quasars and Active Galactic Nuclei, China Science and Technology Press, (2005)
50. B. M. Peterson, An introduction to active galactic nuclei, Cambridge University Press (1997)
51. V. Beckmann, Chris shrader, Active galactic nuclei, WILEY-VCH Verlag GmbH & Co. KGaA, Boschstr. 12, 69469, Weinheim, Germany (2012)
52. D. E. Osterbrock, Emission-line spectra and the nature of active galactic nuclei, proceedings of a conference held at the Georgia State University, Springer-Verlag, 1-18 (1987)
53. M.C. Bentz, P.R. Williams, T. Treu, The Broad Line Region and Black Hole Mass of NGC 4151, arxiv: 2206. 03513v1 [astro-ph.GA] 7 Jun 2022
54. R. Serafinelli, A. Marinucci, A.D. Rosa etc., A Remarkably Stable Accretion Disk in the Sefert Galaxy, arxiv: 2309. 06092v1 [astro-ph.HE] 12 Sep 2023
55. V. Liu, A. Zoghbi, J.M. Miller, Dectection of Asymmetry in the Narrow Fe K α Line in MCG-5-23-16 with Chandra, arxiv: 2312. 16354v2 [astro-ph.HE] 4 Jan 2024
56. H. Netzer, The Physics and Evolution of Active Galactic Nuclei, Cambridge University Press (2013)
57. R. Lynds, The Absorption-line spectrum of 4c 05.34, *Astrophysical Journal*, vol. 164, p.L73 (1971)
58. K. Hirata et al., Observation of a Neutrino Burst from the Supernova SN1987a, *Phys. Rev. Lett.*, 58:1490-1493(1987)
59. B. P. Abbott et al., GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral, *Phys. Rev. Lett.*, 119:161101(2017)

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.