

Hypothesis

Not peer-reviewed version

The Energy-Deficit Hypothesis of Autism: Linking Parental TNF- α -Mediated Diseases to Offspring Autism Risk via Mitochondrial Dysfunction

[Byul Kang](#) *

Posted Date: 31 December 2025

doi: 10.20944/preprints202512.2808.v1

Keywords: autism spectrum disorder; TNF- α ; mitochondrial dysfunction; synaptic pruning; energy metabolism; normal-tension glaucoma; psoriasis; type 1 diabetes; neuroinflammation; eye contact avoidance



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Hypothesis

The Energy-Deficit Hypothesis of Autism: Linking Parental Normal-Tension Glaucoma, Type 1 Diabetes, and Psoriasis to Offspring Autism Risk via TNF- α -Mediated Mitochondrial Dysfunction

Byul Kang

Independent Researcher, Republic of Korea; Republic of Korea; halococoa@gmail.com

Abstract

Background: Autism spectrum disorder (ASD) affects approximately 1-2% of children worldwide, yet its etiology remains incompletely understood. Emerging evidence suggests that parental autoimmune diseases significantly increase offspring autism risk, with psoriasis (OR 1.59), type 1 diabetes (OR 1.49-2.36), and rheumatoid arthritis (OR 1.51) showing particularly strong associations. **Hypothesis:** I propose that autism is fundamentally an immune-metabolic disorder characterized by TNF- α -mediated mitochondrial dysfunction leading to cerebral energy deficiency. This energy deficit impairs two critical processes: (1) synaptic pruning during neurodevelopment, and (2) real-time social cognition including gaze processing and emotion recognition. The primary mechanism involves TNF- α pathway dysregulation—through genetic inheritance from parents with autoimmune diseases such as psoriasis, type 1 diabetes, and rheumatoid arthritis, and/or through direct fetal exposure to elevated maternal TNF- α during pregnancy. Additionally, for cases without parental autoimmune history, I propose a speculative secondary mechanism: mitonuclear immune conflict, where paternal immune genes may partially recognize maternal mitochondria as non-self, generating endogenous TNF- α . **Novel Prediction:** Based on this framework, I predict that parents with normal-tension glaucoma (NTG)—another TNF- α -mediated condition characterized by neurodegeneration independent of intraocular pressure—will show elevated prevalence of autistic offspring. This prediction has not been tested in any published study. **Implications:** This hypothesis unifies disparate observations about autism pathophysiology and suggests that anti-inflammatory interventions targeting the TNF- α pathway may have therapeutic potential, particularly when administered early in neurodevelopment.

Keywords: autism spectrum disorder; TNF- α ; mitochondrial dysfunction; synaptic pruning; energy metabolism; normal-tension glaucoma; psoriasis; type 1 diabetes; neuroinflammation; eye contact avoidance

1. Introduction

Autism spectrum disorder (ASD) is a neurodevelopmental condition characterized by deficits in social communication and interaction, restricted interests, and repetitive behaviors. Despite decades of research, the fundamental biological mechanisms underlying autism remain elusive. While genetic factors contribute substantially to autism risk, environmental and immunological factors increasingly appear to play critical roles.

A growing body of evidence indicates that parental autoimmune diseases significantly increase the risk of autism in offspring. Meta-analyses have demonstrated that family history of autoimmune disease is associated with a 28-50% increased risk of autism. Importantly, specific autoimmune conditions mediated by tumor necrosis factor-alpha (TNF- α) show particularly robust associations with offspring autism risk.

In this hypothesis paper, I propose that autism is fundamentally an **immune-metabolic disorder** characterized by TNF- α -mediated mitochondrial dysfunction. I argue that the resulting cerebral energy deficiency impairs two critical processes: synaptic pruning during neurodevelopment and real-time social cognitive processing. This framework explains core autism symptoms from a unified energetic perspective and generates testable predictions.

2. Evidence for Parental Autoimmune Disease-Autism Association

2.1. Epidemiological Evidence

Multiple large-scale epidemiological studies have established associations between parental autoimmune diseases and offspring autism risk. Table 1 summarizes key findings from major studies.

Table 1. Parental Autoimmune Disease and Offspring Autism Risk.

Parental Disease	Odds Ratio	95% CI	Key Reference
Psoriasis	1.59	1.21-2.10	Wu et al. 2015
Type 1 Diabetes (T1D)	1.49-2.36	1.21-4.12	JAMA 2018, IJE 2023
Rheumatoid Arthritis	1.51	1.14-2.00	Keil et al. 2010
Hypothyroidism	1.64	1.16-2.32	Atladóttir et al. 2009
Any Autoimmune Disease	1.28-1.50	1.11-1.75	Wu et al. 2015 Meta

Note: All listed conditions are TNF- α -mediated autoimmune/inflammatory disorders. CI = Confidence Interval.

2.2. The TNF- α Common Denominator

A critical observation is that all parental diseases strongly associated with offspring autism risk share a common pathogenic mechanism: dysregulation of TNF- α signaling. TNF- α is a master pro-inflammatory cytokine that plays central roles in:

- **Psoriasis:** TNF- α drives keratinocyte proliferation and inflammatory cascade; anti-TNF biologics are first-line therapy
- **Type 1 Diabetes:** TNF- α directly induces β -cell apoptosis and promotes autoimmune destruction of pancreatic islets
- **Rheumatoid Arthritis:** TNF- α orchestrates synovial inflammation and joint destruction; anti-TNF therapy revolutionized treatment
- **Normal-Tension Glaucoma:** TNF- α mediates retinal ganglion cell death independent of intraocular pressure elevation

3. TNF- α and Mitochondrial Dysfunction: The Mechanistic Link

3.1. Direct Effects of TNF- α on Mitochondrial Function

TNF- α exerts profound inhibitory effects on mitochondrial function through multiple mechanisms. Table 2 summarizes the key pathways by which TNF- α impairs cellular energy production.

Table 2. TNF- α Effects on Mitochondrial Function.

Mechanism	Effect on Energy Metabolism
ETC Complex I Inhibition	Blocks electron transfer at the first step of oxidative phosphorylation
ETC Complex III Inhibition	Disrupts cytochrome bc1 complex function
Cytochrome c Oxidase (COX)	Reduces terminal electron transfer and oxygen consumption
Membrane Depolarization	Collapses mitochondrial membrane potential ($\Delta\Psi_m$), halting ATP synthesis
PDH Suppression	Inhibits pyruvate dehydrogenase, blocking glucose entry into TCA cycle
ROS Overproduction	Increases reactive oxygen species, causing oxidative damage to mitochondrial components

Mechanism	Effect on Energy Metabolism
Warburg Effect Induction	Shifts metabolism to inefficient aerobic glycolysis (2 vs 36 ATP per glucose)

Note: ETC = Electron Transport Chain; PDH = Pyruvate Dehydrogenase; TCA = Tricarboxylic Acid Cycle; ROS = Reactive Oxygen Species.

3.2. Rapid Neurotoxicity of TNF-α

Critically, TNF-α-induced mitochondrial dysfunction occurs rapidly in neurons. Studies using pathophysiologically relevant concentrations demonstrate:

- Reduction in mitochondrial basal respiration within **1.5 hours** of TNF-α exposure
- Decreased ATP production preceding neuronal cell death
- Effects mediated specifically through TNF-R1 receptor signaling
- Cascade involving caspase-8 activation, membrane potential collapse, and cytochrome c release

4. The Energy-Deficit Hypothesis of Autism

I propose that chronic TNF-α elevation, whether inherited genetically or transmitted during pregnancy, leads to persistent mitochondrial dysfunction and cerebral energy deficiency. This energy deficit manifests in two critical domains that explain core autism symptoms.

4.1. Impaired Synaptic Pruning

The Energy Cost of Synaptic Pruning: The developing brain undergoes massive synaptic pruning, eliminating approximately 50% of synapses from infancy to adolescence. This process is extraordinarily energy-intensive because:

- Microglia actively phagocytose synapses, requiring substantial ATP
- The infant brain consumes **40% of total body energy**—far exceeding adult proportions
- Complement cascade activation and autophagy pathways require ATP

Evidence of Pruning Deficits in Autism: Postmortem studies reveal striking differences in synaptic density between autistic and neurotypical brains:

Table 3. Synaptic Pruning in Neurotypical vs Autistic Brains.

Parameter	Neurotypical	Autism
Synaptic density reduction (childhood→adolescence)	~50%	~16%
Dendritic spine density	Normal	Elevated
mTOR pathway activity	Normal	Hyperactive
Autophagy function	Normal	Impaired

Source: Tang et al. 2014, Neuron (Columbia University study).

Consequences of Excess Synapses: The failure to prune synapses results in:

- **Local over-connectivity:** Excess short-range connections creating "neural noise"
- **Long-distance under-connectivity:** Insufficient resources for developing major "highway" connections between brain regions
- **Reduced signal-to-noise ratio:** Difficulty filtering relevant from irrelevant information
- **Sensory overload:** Heightened sensitivity due to failure to attenuate sensory inputs

4.2. Impaired Social Cognition and Gaze Avoidance

The Energy Demands of Social Processing: Social cognition—including face recognition, gaze processing, and emotion interpretation—is among the most computationally and energetically demanding brain functions. It requires simultaneous activation of:

- Fusiform Face Area (FFA): Face identity processing

- Superior Temporal Sulcus (STS): Gaze direction and biological motion
- Amygdala: Emotional salience and threat detection
- Prefrontal Cortex: Social context integration and decision-making

Eye Contact as Energy Conservation: I propose that gaze avoidance in autism represents an *adaptive energy conservation strategy*. Direct evidence supports this interpretation:

Table 4. Self-Reported Experiences of Eye Contact in Autism.


Experience Category	Representative Quote
Energy Exertion	"Eye contact feels like I'm using up a lot of energy. Maximum 2-6 seconds."
Audiovisual Integration Failure	"I cannot listen to someone while making eye contact at the same time."
Cognitive Trade-off	"When I focus on eye contact, I can't process what's being said."
Recovery Requirement	"The longer I maintain eye contact, the more recovery time I need afterward."

Source: Trevisan et al. 2017, PLOS ONE - Qualitative analysis of first-hand accounts.

Neural Evidence: Functional neuroimaging studies demonstrate that in autism, eye contact triggers **amygdala hyperactivation**, suggesting heightened metabolic demand. Gaze avoidance thus serves to reduce this hyperarousal and conserve limited neural energy for other cognitive tasks.

5. Integrated Pathophysiological Model

Figure 1 presents the unified model linking parental TNF-α-mediated diseases to offspring autism through mitochondrial dysfunction and energy deficiency.






Figure 1. Integrated Pathophysiological Model of Autism as an Immune-Metabolic Disorder. Figure 1 Legend: The model illustrates how parental TNF- α pathway dysregulation leads to offspring autism through two converging pathways: developmental (impaired synaptic pruning) and real-time (impaired social cognition). Both pathways result from ATP deficiency secondary to mitochondrial dysfunction. ETC = Electron Transport Chain; ROS = Reactive Oxygen Species.

6. The Mitonuclear Immune Conflict Hypothesis: An Endogenous Source of TNF- α

While the preceding sections describe how TNF- α -mediated mitochondrial dysfunction leads to autism, an important question remains: what about cases where parents have no autoimmune disease? I propose that mitonuclear immune conflict may represent an endogenous source of TNF- α that activates the same pathogenic pathway.

6.1. The Gap in the TNF- α Hypothesis

The TNF- α energy deficit hypothesis explains autism risk in offspring of parents with autoimmune diseases such as psoriasis, type 1 diabetes, and rheumatoid arthritis. However, autism also occurs in families with no history of autoimmune disease. Additionally, studies show that a substantial proportion of autistic individuals exhibit mitochondrial dysfunction biomarkers without carrying classical mitochondrial disease mutations.

This raises a critical question: if TNF- α -mediated mitochondrial dysfunction is central to autism pathophysiology, what is the source of TNF- α in cases without parental autoimmune disease?

6.2. The Unique Inheritance Pattern of Mitochondria

Mitochondria possess a unique inheritance pattern. Mitochondrial DNA (mtDNA) is inherited **exclusively from the mother**—paternal mitochondria are actively eliminated from the fertilized egg. Every mitochondrion in an individual's body carries only maternal genetic information.

In contrast, the nuclear genome—including genes governing immune function and self/non-self recognition—is inherited from **both parents**. This creates an asymmetry: the immune system is shaped by both parental genomes, but the mitochondria it must tolerate are exclusively maternal.

6.3. The Conflict Hypothesis: Paternal Immune Genes vs. Maternal Mitochondria

I hypothesize that in some individuals, paternally inherited immune genes may fail to fully recognize maternal mitochondria as "self." This could result in:

- **Immune misrecognition:** The paternal contribution to immune recognition machinery (HLA genes, innate immune pathways) may be calibrated to recognize mitochondrial signatures that differ from those inherited from the mother.

- **Chronic immune attack:** The immune system may mount persistent inflammatory responses against the individual's own mitochondria, treating them as partially foreign.
- **Endogenous TNF- α production:** This chronic immune activation would result in sustained TNF- α release—activating the same pathogenic cascade described in previous sections, even without external TNF- α exposure from parental autoimmune disease.

6.4. Two Pathways to the Same Outcome

The mitonuclear immune conflict hypothesis does not replace the parental autoimmune disease hypothesis—it complements it by providing a second pathway to TNF- α -mediated mitochondrial dysfunction:

Table 5. Two Pathways to TNF- α -Mediated Mitochondrial Dysfunction.

	Pathway 1: External	Pathway 2: Internal
Source of TNF- α	Parental autoimmune disease	Mitonuclear immune conflict
Mechanism	Genetic inheritance + fetal exposure during pregnancy	Paternal immune attack on maternal mitochondria
Parental disease required?	Yes	No
Final common pathway	TNF- α elevation → Mitochondrial dysfunction → Energy deficit → Autism	

Note: Both pathways converge on the same final mechanism of TNF- α -mediated mitochondrial dysfunction.

This framework explains why:

- Parental autoimmune disease increases autism risk (Pathway 1)
- Autism also occurs without parental autoimmune disease (Pathway 2)
- Only a subset of children with autoimmune parents develop autism (variable mitonuclear compatibility may be protective or additive)

6.5. Testable Predictions

The mitonuclear immune conflict hypothesis generates testable predictions:

- Anti-mitochondrial antibodies or mitochondria-targeted immune markers may be elevated in autistic individuals without parental autoimmune history
- Inflammatory cytokines including TNF- α may be elevated even in autism cases without parental autoimmune disease
- Specific HLA haplotype combinations from parents may show associations with autism risk

7. Novel Prediction: Normal-Tension Glaucoma and Autism

A critical test of this hypothesis involves normal-tension glaucoma (NTG). NTG is a neurodegenerative condition affecting retinal ganglion cells (RGCs) that shares key pathophysiological features with both the TNF- α -mediated autoimmune diseases associated with autism and with autism itself.

7.1. NTG as a TNF- α -Mediated Condition

Evidence supporting TNF- α involvement in NTG:

- Elevated TNF- α levels in aqueous humor and serum of NTG patients
- TNF- α directly induces RGC apoptosis via TNF-R1 signaling
- Anti-TNF therapy shows protective effects in animal models
- NTG frequently co-occurs with systemic inflammatory conditions (e.g., psoriasis)
- Disease progression occurs despite normal intraocular pressure, implicating IOP-independent mechanisms

7.2. The Untested Hypothesis

I predict that parents with NTG will have elevated prevalence of autistic offspring compared to the general population.

To my knowledge, *no published study has examined the association between parental NTG and offspring autism risk*. This represents a critical gap in the literature and a testable prediction of this hypothesis.

Table 6. TNF-α-Mediated Conditions and Autism Association Studies.

Parental Condition	TNF-α Role	Autism Association Studied?
Psoriasis	Central	Yes (OR 1.59)
Type 1 Diabetes	Central	Yes (OR 1.49-2.36)
Rheumatoid Arthritis	Central	Yes (OR 1.51)
Normal-Tension Glaucoma	Central	NO STUDIES EXIST

Note: The absence of studies on parental NTG and offspring autism represents a critical research gap and a testable prediction of the present hypothesis.

8. Therapeutic Implications

The energy-deficit hypothesis suggests several therapeutic approaches:

8.1. Anti-TNF-α Interventions

Existing anti-TNF biologics (etanercept, infliximab, golimumab, adalimumab) have proven efficacy in TNF-α-mediated diseases. In type 1 diabetes, golimumab preserved β-cell function in a phase 2 trial (NEJM 2020). Similar approaches might be considered for autism prevention in high-risk pregnancies, though significant safety and ethical considerations would need to be addressed.

8.2. Mitochondrial Support

Interventions supporting mitochondrial function may provide benefit:

- **Coenzyme Q10:** Essential electron carrier in ETC
- **L-Carnitine:** Facilitates fatty acid transport into mitochondria
- **NAD+ Precursors (NR, NMN):** Support ETC function and cellular energy production
- **B Vitamins:** Cofactors for mitochondrial enzymes

8.3. Early Identification

Screening for parental TNF-α-mediated diseases (psoriasis, T1D, RA, NTG) could identify pregnancies at elevated autism risk, enabling earlier monitoring and potentially earlier intervention.

9. Limitations and Future Directions

Limitations: This hypothesis paper synthesizes existing evidence but does not present new experimental data. The proposed mechanisms, while supported by converging lines of evidence, require direct experimental validation.

Future Directions: Key studies needed include: (1) Epidemiological investigation of parental NTG and offspring autism risk; (2) Longitudinal studies of mitochondrial function in infants at high autism risk; (3) Clinical trials of mitochondrial support interventions; (4) Mechanistic studies of TNF-α effects on synaptic pruning in animal models.

10. Conclusion

I propose that autism spectrum disorder can be understood as an immune-metabolic disorder characterized by TNF- α -mediated mitochondrial dysfunction leading to cerebral energy deficiency. This energy deficit impairs synaptic pruning during development and compromises real-time social cognitive processing, explaining core autism symptoms from a unified mechanistic perspective.

The hypothesis generates a novel, testable prediction: that parents with normal-tension glaucoma—a TNF- α -mediated neurodegenerative condition not previously linked to offspring autism—will show elevated prevalence of autistic children. Confirmation of this prediction would provide strong support for the broader hypothesis.

If validated, this framework has important implications for autism prevention and treatment, suggesting that anti-inflammatory and mitochondrial support interventions may have therapeutic potential, particularly when administered early in neurodevelopment.

References

1. Wu S, Ding Y, Wu F, et al. Family history of autoimmune diseases is associated with an increased risk of autism in children: A systematic review and meta-analysis. *Neurosci Biobehav Rev.* 2015;55:322-332.
2. Xiang AH, Wang X, Martinez MP, et al. Maternal Type 1 Diabetes and Risk of Autism in Offspring. *JAMA.* 2018;320(1):89-91.
3. Persson M, et al. Maternal type 1 diabetes, pre-term birth and risk of autism spectrum disorder. *Int J Epidemiol.* 2023;52(2):377-386.
4. Keil A, Daniels JL, Forssen U, et al. Parental autoimmune diseases associated with autism spectrum disorders in offspring. *Epidemiology.* 2010;21(6):805-808.
5. Atladóttir HO, Pedersen MG, Thorsen P, et al. Association of family history of autoimmune diseases and autism spectrum disorders. *Pediatrics.* 2009;124(2):687-694.
6. Tang G, Gudsnuik K, Kuo SH, et al. Loss of mTOR-dependent macroautophagy causes autistic-like synaptic pruning deficits. *Neuron.* 2014;83(5):1131-1143.
7. Trevisan DA, Roberts N, Lin C, Birmingham E. How do adults and teens with self-declared Autism Spectrum Disorder experience eye contact? *PLOS ONE.* 2017;12(11):e0188446.
8. Dalton KM, Nacewicz BM, Johnstone T, et al. Gaze fixation and the neural circuitry of face processing in autism. *Nat Neurosci.* 2005;8(4):519-526.
9. Morris G, Berk M. The many roads to mitochondrial dysfunction in neuroimmune and neuropsychiatric disorders. *BMC Med.* 2015;13:68.
10. Quattrin T, Haller MJ, Steck AK, et al. Golimumab and Beta-Cell Function in Youth with New-Onset Type 1 Diabetes. *N Engl J Med.* 2020;383(21):2007-2017.
11. Chen Y, et al. The role of the immune system in autism spectrum disorder. *Neuropsychopharmacology.* 2017;42(1):284-298.
12. Rossignol DA, Frye RE. Mitochondrial dysfunction in autism spectrum disorders: a systematic review and meta-analysis. *Mol Psychiatry.* 2012;17(3):290-314.
13. Nakazawa T, et al. Tumor necrosis factor- α mediates oligodendrocyte death and delayed retinal ganglion cell loss in a mouse model of glaucoma. *J Neurosci.* 2006;26(49):12633-12641.
14. Deng W, et al. Rapid mitochondrial dysfunction mediates TNF- α -induced neurotoxicity. *J Neurochem.* 2015;132(4):443-451.
15. Ashwood P, et al. Immunological cytokine profiling identifies TNF- α as a key molecule dysregulated in autistic children. *Mol Psychiatry.* 2017;22(6):809-816.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.