

Article

Not peer-reviewed version

A Pilot Study of Family Integrated Care (FICare) in Critically Ill Preterm and Term Infants in the NICU: FICare Plus

[Najmus Sehr Ansari](#) ^{*} , [Linda S. Franck](#) , Christopher Tomlinson , Anna Colucci , [Karel O'Brien](#)

Posted Date: 26 June 2023

doi: 10.20944/preprints202306.1768.v1

Keywords: Family integrated care; critically ill neonates; safety; feasibility



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

A Pilot Study of Family Integrated Care (FICare) in Critically Ill Preterm and Term Infants in the NICU: FICare Plus

Najmus Sehr Ansari ^{1,2*}, Linda S. Franck ³, Christopher Tomlinson ^{1,2}, Anna Colucci ⁴ and Karel O'Brien ^{1,4}

¹ Department of Pediatrics, University of Toronto, Toronto, ON M5S 1A1, Canada

² Division of Neonatology, The Hospital for Sick Children, Toronto, ON, M5G 1X8, Canada

³ School of Nursing, University of California, San Francisco CA 94143, USA

⁴ Department of Pediatrics, Mount Sinai Hospital, Toronto, ON M5G 1X5, Canada

* Correspondence: sehr.ansari@uhn.ca Tel: +1 647 821 5381

Abstract: Family Integrated Care (FICare) is associated with improved developmental outcomes and decreased parental mental health risks in stable preterm infants. However, less is known about its application in critically ill infants who are at greater risk for adverse outcomes. The objective of this study was to assess the safety and feasibility of implementation of FICare in critically ill infants in the first few weeks of life. In this prospective cohort study, 41 critically ill infants were enrolled: 17 in observational group receiving standard care (SC) and 24 in the FICare Plus group. Resources were developed for staff and parents to support earlier engagement in infant care. Outcomes were assessed using standardized questionnaires. The t-test or Wilcoxon Rank-Sum test were used to compare continuous variables, while Chi-square or Fisher exact test were used for categorical variables, respectively. Baseline maternal and infant characteristics were similar in both groups. The tools and procedures developed for FICare Plus successfully supported greater engagement in the care of their infant with no increase in adverse events and no increase in parent stress. Preliminary measures of infant efficacy were similar in both groups. Total anxiety scores were high among all parents at enrolment {83.1 (67-94) vs 77.3 (66-86); p-value 0.4}. However, the scores prior to discharge were lower in FICare Plus group {78 (71-90) vs 63 (52-74.5); p-value 0.02}. This pilot study showed that it is feasible and safe to implement family integrated care in critically ill infants.

Keywords: family integrated care; critically ill neonates; safety; feasibility

1. Introduction

Family Integrated Care (FICare) involves engaging parents in their infants' care and supporting them to be active participants in all aspects of infant care planning and caregiving, in partnership with the health care professionals [1]. Studies conducted in different countries and health systems have shown improved neonatal outcomes and enhanced psychological and emotional well-being in parents. [2-4]. Clinical trials have shown improved weight gain, shorter hospitalization, lower rates of infection and higher rate of breastfeeding at discharge in infants who received family integrated care [5,6]. Furthermore, there is growing body of evidence which suggests that FICare improves long term neurodevelopmental outcomes in infants [7,8].

The original studies of FICare enrolled preterm infants born less than or equal to 35 weeks gestation, who were stable on non-invasive ventilation at the time of enrolment although many were born very preterm [1]. Investigators have also reported the outcomes of implementing FICare in level 2 neonatal intensive care units (NICU) [9]. However, limited evidence is available about the feasibility of implementing FICare with families of infants who are critically ill at the time of enrolment. Only one study has investigated FICare in critically ill preterm neonates in the United States [10]. Although there were no group differences in outcomes commonly measured in prior studies of stable preterm infants, there were no adverse effects and several FICare components (rounds, peer parent mentors

and group parent classes) were associated with improved outcomes compared with standard family-centered care [10]. Another study implemented a modified FICare program NICU-wide in a community level hospital, including some term and near-term infants. In this study, nurses reported positive views about the program and parents had reduced stress and improved readiness for discharge after implementation of the program [2].

The technologically advanced environment of neonatal intensive care units can often be very daunting and stressful for parents [11,12]. This stress is further heightened by their infants' fragility and their limited ability to participate as a caregiver, which may adversely impact their confidence to acquire normal parenting skills [13–15]. Although critically ill infants may have greater gains from FICare, there was a concern especially from our family advisors, that implementing FICare in this population might cause unintended distress to parents. Given the potential benefits for critically ill infants and their families, we set out to adapt FICare to accommodate the needs of this vulnerable population and to pilot this adapted model, termed FICare Plus, to assess the safety and feasibility of its implementation.

2. Materials and Methods

Study Design

This was a prospective cohort study, pre-post implementation, of an augmented FICare program for critically ill infants of all gestational ages, termed as FICare Plus. It was conducted in two tertiary care NICUs, Mount Sinai Hospital (Site A) and SickKids Hospital (Site B) in Toronto, Canada. Site A is a tertiary care perinatal center with more than 1100 infant admissions per year to the NICU. Of these infants, 13% are extremely low birth weight (ELBW) (<1000 grams) infants. Site B is a quaternary care referral center with an average of 800 admissions/year to the NICU. Approximately one third of these infants require surgery during the neonatal period. In addition to serving different infant populations, these two NICUs have different levels of experience with FICare. Site A has over 10 years of experience, whereas Site B has not implemented FICare as their standard of care.

To assess the safety and feasibility of implementation of FICare Plus in critically ill infants, data was collected in a sequential manner, pre and post implementation. This design was considered most feasible for this study as families could not be randomized to different models of care within the same unit. The study protocol, questionnaires and the educational toolkit developed for this study were approved by the research ethics board at both hospitals.

Participants

Infants eligible for this study included: ELBW infants on invasive positive pressure ventilation for >48 hours after birth, infants with surgical necrotizing enterocolitis or bowel perforation, infants with tracheoesophageal fistula or esophageal atresia. Families were approached while the infant was <2 weeks of age. For both cohorts, families were asked to commit to being present at the hospital for 4 hours per day, for a minimum of 5 days/week. For parents of multiples, families were only asked to commit to the same time commitment even though their time might be divided between their infants. Infants were excluded if they were receiving palliative care, were deemed unlikely to survive either due to critical illness or other life-threatening congenital anomaly or were born to parents with inability to participate either due to health, social or language issues. Parents in the FICare Plus group were encouraged to attend and participate in daily morning rounds. They were encouraged to engage in their infant's care under nursing supervision. Written informed consent was obtained from all enrolled participants. Parents in both groups were asked to complete questionnaires at specific time points including their demographic information.

Procedures

Pre-implementation: In the development phase of this study, teams were formed at both sites which included parents, bedside nurse/s, respiratory therapist, social worker, physicians, and an educator to plan and execute this project at their respective sites. The nursing staff, veteran parents

(through Canadian Premature Babies Foundation) and current parents were surveyed at both hospitals to determine how parents could safely and meaningfully be involved in the care of their critically ill babies. The information gathered from these surveys was reviewed in a joint meeting of both site teams. The teams identified critical opportunities for creating a connection with families at both sites and went on to develop educational resources for parents and staff (will be uploaded to the FICare website (<https://familyintegratedcare.com>)). From the parent survey, it was identified that parents might have difficulty attending group classes in the immediate days following admission of their infant to the NICU. In response, a toolkit was developed for both sites by consensus, which included a parent booklet and online educational resources including videos for parents to access at their own time. It was also recognized that staff might need additional education so that they could better support parents particularly when their infant was just admitted to the hospital. In response, an on-line staff educational booster toolkit was developed, comprising of short videos providing information about the importance of parent engagement in infant care, parent reflections on their experience in the NICU, and coaching staff about how best to support parents (see <https://familyinteratedcare.com>). A detailed inventory of potential parent and nurse responsibilities for FICare Plus group was created and reviewed at both sites, recognizing that there could be some site-specific accommodations depending on the needs of the infant. While the resources for implementation were being developed, participants were enrolled in the standard care (SC) cohort at both sites.

Implementation: Once recruitment to the SC group was completed, FICare Plus booster training was provided to >90% of nursing staff at both sites. The NICU policies and procedures at both sites were reviewed and revised to enable greater parent participation from admission. Between Jan 1, 2021, to July 1, 2021, parents were enrolled in the FICare Plus group. Parents were oriented to the tools provided for their self-education by a specially trained FICare Plus nurse educator. Parents were supported by their bedside nurse to participate in their infants' care and were taught ways to interact with their infants which may optimize their development. They were also provided with information about coping strategies and stress -reducing activities, in addition to the routine support provided by the social worker. Enrolment was maintained to ensure equal representation of infants in the observational and the implementation cohorts.

The research coordinators at both sites ensured that all enrolled families were able to access the educational materials provided, were being offered to participate in their infant's care and were supported to participate in rounds. Since families in both cohorts were recruited during the COVID-19 pandemic, research coordinators had limited access to both NICU's and therefore had limited ability to approach families for enrolment. A single coordinator approached all families for all the concurrent studies in the unit and then referred those who were interested in hearing more about a specific study to the respective coordinator. Our FICare Plus study coordinators were also unable to have face-to-face contact with families. They provided support and encouragement to families to complete the data collection through phone calls. In addition, peer support, which has been a cornerstone of FICare implementation in our other studies, was not provided over this time period. The parent ongoing education sessions, which were usually standard at site A, were provided inconsistently and virtually rather than in person.

Data Collection

The primary aim of this study was to assess if it was feasible to support greater parent engagement in taking care of critically ill infants given the tools and procedures developed for FICare. The measures of this outcome included 1) the percentage of approached parents who enrolled in the pilot study 2) parent access to and evaluation of on-line education modules and educational materials 3) the parents' perceptions of their engagement in their infants care as indicated on their parent surveys. Secondary outcomes included the safety of the intervention as measured by 1) the number of reported adverse events related to increased parental involvement in their infants care 2) the acceptability of FICare Plus as reported by the multidisciplinary team on their surveys 3) whether there was any difference between the staff responses at both sites.

Preliminary measures of efficacy were collected pre and post implementation and included both infant and parent measures. The infant measures were the type of feeding at discharge (tube feeding, exclusive breast milk feeding, exclusive formula feeding or a combination of formula/breast milk), weight at discharge and discharge destination. Parent measures included measures of parent stress and anxiety, parent perceived competence in parenting and family centered care. Parent anxiety was measured by the State -Trait Anxiety Inventory (STAI) [16] at enrolment, day 21, prior to discharge and one week post discharge. The STAI is a 40-item questionnaire with scores ranging from 1 (lowest anxiety) to 4 (highest anxiety) for each item with cumulative scores ranging from 40-160. Parental stress was assessed using the Parental Stressor Scale: NICU (PSS: NICU) [17] which is a validated scale to assess parental stress during NICU hospitalization of their infant; the average overall scores range from 1-5 in order of severity. Parental perceived parenting self-efficacy was measured using Perceived Parenting Self- Efficacy Tool [18]. The score range in this validated tool is 20-80, with higher scores indicating higher perceived self-efficacy. Parents' NICU experience was assessed using the Family Centered Care Survey (FCC) [19]. Their response was rated on a 7-point scale with higher values indicating a more positive experience. Linked parent and infant demographic data was extracted from the Canadian Neonatal Network database.

Statistical analysis:

The data analysis was based on an intention-to-treat basis. The measures of both the primary and secondary outcomes are presented with calculated descriptive statistics. Comparisons by intervention group were performed using the t-test or Wilcoxon Rank-Sum test for continuous variables, and Chi-square or Fisher exact test for categorical variables. For the group comparisons, p-value <0.05 was considered statistically significant. All analyses were performed using SAS version 9.4 (Cary, NC).

3. Results

Between January 1, 2020, and December 31, 2020, a total of 115 infants were assessed at both hospitals for eligibility in the SC cohort (Figure 1). Of the 36 families who were approached, 17 parent -infant dyads (47.2%) were enrolled in the SC cohort, between the two sites. Between January 1, 2021, and July 31, 2021, a total of 245 infants were assessed for eligibility in the FICare Plus cohort (Figure 2). Of the 64 families approached, 24 parent infant dyads (37.5%) were enrolled in the FICare Plus group, between the two sites.




Figure 1. Flow diagram of families included in Standard Care cohort.




Figure 2. Flow diagram of families included in FICare Plus cohort.

Baseline parent characteristics were similar in both groups (Table 1) except caesarean delivery, which was more common in FICare Plus group (75% vs 35.2%; p- value 0.01). For infants, the median

age at enrolment was 8 days which was similar in both groups {8 (5,19) vs 8 (5.5,13); p-value 0.8}. No significant difference was observed in gestational age, birth weight, major morbidity, and clinical course in infants between the two groups (Table 2).

Table 1. Baseline parental characteristics.

	Standard Care (17)	FICare Plus (24)	p-value
Relationship, % (n/N)			0.99
Mother	82.35 (14/17)	83.3 (20/24)	
Father	17.65 (3/17)	16.6 (4/24)	
English as first language, % (n/N)	52.94 (9/17)	70(14/20)	0.29
Education level (Bachelor or above), % (n/N)	35.29(6/17)	63.16(12/19)	0.09
Employment (or self-employment), % (n/N)	82.35(14/17)	83.3(20/24)	0.09
Prenatal Care (yes), % (n/N)	52.94(9/17)	62.5(15/24)	0.16
Parental age (years), median (IQR)	30 (26, 33)	31.5 (29.5, 36.5)	0.12
Parental self-rated health status (0-100), median (IQR)	97 (77.5, 100)	97 (85.5, 100)	0.77
Singleton conception % (n/N)	70.59(12/17)	83.33(20/24)	0.45
Primigravida mothers, % (n/N)	29.41(5/17)	20.83(5/24)	0.71
Cesarean delivery, % (n/N)	35.29(6/17)	75(18/24)	0.01
Abortions, % (n/N)	41.18(7/17)	58.33(14/24)	0.27
Out born delivery, % (n/N)	64.71(11/17)	41.67(10/24)	0.14

Table 2. Baseline infant characteristics and their clinical parameters.

	Standard Care	FICare Plus	p-value
Number of infants	17	24	
Gestational age (weeks), median (IQR)	26.05 (24.2, 29)	28.9 (25.5, 32.1)	0.15
Age at enrolment (days), median (IQR)	8 (5,19)	8 (5.5-13)	0.8
Birth weight (kg), median (IQR)	0.71 (0.62, 0.82)	0.93 (0.74, 1.63)	0.07
Sex, % (n/N)	52.94(9/17)	50(12/24)	0.85
Apgar score<7 at 1 min, % (n/N)	71.43(10/14)	50(12/24)	0.19
SNAP II score>20, % (n/N)	35.29(6/17)	20.83(5/24)	0.47
Intestinal Perforation, % (n/N)	11.76(2/17)	8.33(2/24)	0.99
Surgical intervention, % (n/N)	35.29(6/17)	37.5(9/24)	0.89
Sepsis, % (n/N)	35.29(6/17)	16.67(4/24)	0.27
Esophageal atresia	17.65(3/17)	29.17(7/24)	0.47
Invasive ventilation days, median (IQR)	6 (1, 44)	3 (0, 6)	0.09
NIPPV ventilation days, median (IQR)	4 (0, 14)	3 (0, 18.5)	0.96
CPAP days, median (IQR)	15 (6, 37)	21 (12, 30)	0.75

All FICare Plus families received a link to the on-line educational resources and the FICare Plus parent education booklet. The online educational module was accessed sixty-five times by 24 families who were enrolled in FICare Plus. Sixteen of the 24 families (66%) completed the parent feedback survey. All sixteen parents indicated that they had accessed the on-line educational materials and they agreed or strongly agreed with the statement that the educational materials provided the right kind and right amount of information. Twelve of the sixteen parents responded that they found the information booklet useful.

Of the 25 families who consented to participate in the FICare cohort, 24 families continued their participation throughout their hospital stay. Sixteen FICare Plus families completed the discharge surveys. All FICare Plus families indicated that they enjoyed taking part in the study. The results of the parent feedback survey are presented in Table 3. Despite the COVID restrictions, parents in both cohorts indicated that they were able to spend more than 4 hours/day with their baby. Of note, all parents in the FICare Plus cohort indicated that they felt ready to interact with their baby when invited to do so, felt confident being involved in their baby's care and 87.5% (14/16) of parents indicated that they felt part of their baby's care team and were able to participate in development of a care plan.

Table 3. Parent feedback survey.

	Standard Care	FICare Plus
Able to spend > 4 hours/day with my baby % n/N	90.91 (10/11)	93.75 (15/16)
Feel confident being involved in my baby's care % n/N	90.91 (10/11)	100 (16/16)
Feel like I know my baby % n/N	100 (11/11)	100 (16/16)
Felt ready to interact with my baby when invited to do so % n/N	63.64 (7/11)	100 (16/16)
Found participating on rounds helpful % n/N	90.91 (10/11)	100 (16/16)
Felt part of my baby's care team % n/N	81.82 (9/11)	87.5 (14/16)
Was able to participate in development of care plan % n/N	63.64 (7/11)	87.5 (14/16)
Had enough information on baby's plan of care at discharge % n/N	80 (8/10)	85.71 (12/14)

The staff survey was completed by 177 members of the multidisciplinary team involved at both sites {Site A n=101/200 (50%), Site B n=76/200 (38%)}. 70% of the respondents indicated that they found the booster education kit helpful. The survey responses indicated that the teams at both sites found FICare Plus acceptable, and it was generally well adopted as presented in Table 4. Of note, there was a significant difference noted between the responses from staff at both sites regarding how satisfying they found the experience of taking care of families enrolled in FICare Plus. In addition, significant differences were noted in the staff survey responses regarding the barriers experienced in implementing FICare between the two sites. More respondents from Site B indicated that inadequate staffing (70 vs 47%) and limited space to accommodate families (86 vs 37%) were barriers to FICare Plus implementation. In addition, staff at both sites reported a different role in their pattern of communication with families; more staff in site A indicated that the bedside nurse would act as a liaison between the family and the clinical team (60%) than at Site B (45%).

Table 4. Staff feedback survey.

	Site A	Site B	p-value
I became more of a mentor to the parents rather than a direct care provider			0.4
Very/Somewhat true % n/N	56.44(57/101)	56.58(43/76)	
Neutral % n/N	32.67(33/101)	26.32(20/76)	
Somewhat untrue/Not at all true % n/N	10.89(11/101)	17.11(13/76)	
I felt I was taking care of the entire family not just the baby			0.88
Very/Somewhat true % n/N	83.17(84/101)	85.9(67/78)	
Neutral % n/N	10.89(11/101)	8.97(7/78)	
Somewhat untrue/Not at all true % n/N	5.94(6/101)	5.13(4/78)	
I found taking care of families in the FICare Plus model more challenging than taking care of SC families			0.86
Very/Somewhat true % n/N	32.67(33/101)	28.95(22/76)	

Neutral % n/N	31.68(32/101)	34.21(26/76)
Somewhat untrue/Not at all true % n/N	35.64(36/101)	36.84(28/76)
I found taking care of families in the FICare Plus model more satisfying		0.028
Very/Somewhat true % n/N	69.31(70/101)	53.25(41/77)
Neutral % n/N	21.78(22/101)	40.26(31/77)
Somewhat untrue/Not at all true % n/N	8.91(9/101)	6.49(5/77)
I have seen a positive effect on the parent-infant relationship		0.95
Very/Somewhat true % n/N	81.19(82/101)	81.25(65/80)
Neutral % n/N	15.84(16/101)	15(12/80)
Somewhat untrue/Not at all true % n/N	2.97(3/101)	3.75(3/80)
I have seen a positive effect of this model on the parent healthcare team relationship		0.55
Very/Somewhat true % n/N	78(78/100)	70.89(56/79)
Neutral % n/N	18(18/100)	24.05(19/79)
Somewhat untrue/Not at all true % n/N	4(4/100)	5.06(4/79)
I feel overall that this model of care allows for greater scope on a nurse's role in the NICU		0.08
Very/Somewhat true % n/N	66.34(67/101)	61.54(48/78)
Neutral % n/N	22.77(23/101)	34.62(27/78)
Somewhat untrue/Not at all true % n/N	10.89(11/101)	3.85(3/78)

Safety:

No untoward events in infants, such as accidental extubation, removal of intravenous lines, displacements of nasogastric/orogastric tubes, dislodgement of umbilical catheters, mishandling of equipment or parental emotional breakdown was observed during the study in FICare Plus group.

Preliminary measures of efficacy

The results of the parent measures of efficacy are presented in Tables 3. The measures of efficacy for infants as indicated by clinical outcomes at discharge (Table 5), were similar in both groups. Specifically, there was no difference in the feeding outcome.

Table 5. Clinical outcomes at discharge.

	Standard Care	FiCare Plus	p-value
Number of infants	17	24	
Gavage, %(n/N)	60(9/15)	70.83(17/23)	0.49
Breast milk only, %(n/N)	53.33 (8/15)	60.87 (14/23)	0.64
Formula milk only, %(n/N)	6.67 (1/15)	8.70 (2/23)	0.99
Both Breast and Formula and milk, %(n/N)	33.33 (5/15)	30.43 (7/23)	0.99
Weight at discharge(kg), median (IQR)	3.24 (2.03, 3.65)	2.64 (1.71, 2.92)	0.22
Discharge destination, %(n/N)			0.26
Home	23.53(4/17)	12.5(3/24)	
Inpatient area	29.41(5/17)	16.67(4/24)	
Other hospitals (NICU or Community)	35.29(6/17)	66.67(16/24)	
Died before discharge	11.76(2/17)	4.17(1/24)	

The total anxiety scores among parents were similar at enrolment between the two groups {87(67-94) vs.70.5 (66-86); p-value 0.22} (Table 6). However, the scores prior to discharge were lower (less anxiety) in FiCare Plus group as compared to the SC group {78 (71-90) vs.63 (52-74.5); p-value 0.02}. This difference did not persist post-discharge. No difference was observed in parental perception of family centered care (Table 7) and their Perceived Parenting Self-efficacy scores between FiCare Plus and SC groups (Table 8). Similarly, the mean parental stress scores (Table 9) were similar in both groups at enrolment (3.1 vs 2.8; p-value 0.31) and on day 21 (3.3 vs 3; p-value 0.3).

Table 6. Total anxiety scores.

	Standard Care	FiCare Plus	p-value
At enrollment, median (IQR)	87 (67, 94)	70.5 (66, 86)	0.22
At Day 21, median (IQR)	88.5 (71, 103)	78.5 (63, 91)	0.3
Prior to Discharge, median (IQR)	78 (71, 90)	63 (52, 74.5)	0.029
Post Discharge, median (IQR)	73 (65, 88)	70.5 (49, 77)	0.35

Table 7. Perception of family centered care.

	Standard Care	FiCare Plus	p-value
At enrollment, median (IQR)	5.56 (4.89, 6.56)	6.22(5.78, 7)	0.12
Prior to Discharge, median (IQR)	5.11 (3.89, 5.89)	5.33 (4.89, 5.78)	0.68

Table 8. Perceived parenting self-efficacy scores.

	Standard Care	FiCare Plus	p-value
At day 21, mean (SD)	58.13 (10.86)	62.44 (6.33)	0.18
Prior to Discharge, mean (SD)	65.7 (8.04)	65.69 (7.07)	0.99
Post discharge, mean (SD)	66.44 (7.84), 9	66.67 (6.41), 8	0.94

Table 9. Parental stress scores.

	Standard Care	FICare Plus	p-value
At enrollment, mean (SD)	3.10 (0.96)	2.81 (0.74)	0.31
At day 21, mean (SD)	3.32 (0.93)	3.0 (0.62)	0.30

4. Discussion

The results of this prospective cohort study suggest that it is feasible and safe to implement FICare in critically ill infants as defined in this study. The tools and procedures provided to staff and families for implementation of FICare Plus resulted in an increased parental perception of their ability to be a part of their infants' care team, specifically around feeling ready to interact with their baby and being a participant in developing care plans. Our data also suggests that such engagement may have had the positive impact of decreasing parental anxiety prior to discharge. The specific concerns around the risk of increasing parent stress or increasing the risk of safety concerns for the infant were not identified in this pilot study, recognizing that the sample size was very small.

The reflection of health care practitioners on their experience of FICare Plus was generally positive, with a recognition of the importance of the parent role in their infants' care. Interestingly, in the hospital with more complex surgical patients, health care practitioners provided a neutral response to the question about whether they found this model of care more satisfying than SC. This merits further exploration. There may be specific characteristics of the staff who chose to work in a surgical NICU that orients their care differently or it may just reflect less familiarity with this model of care and comfort with the skills it requires.

Implementation of FICare in stable term and preterm infants has shown remarkable success across the world. In a multi-center, randomized controlled trial conducted in 26 tertiary care NICUs, implementation of FICare resulted in improved weight gain in infants, higher rates of exclusive breast feeding at discharge and decreased parental stress and anxiety [5]. In another RCT in 11 tertiary care NICUs in China, Hei et al. reported better neonatal outcomes such as shorter hospitalization, shorter duration of oxygen supplementation, reduction in nosocomial infections and less antibiotic use in FICare group infants [6]. In this context, we extended this model of care to both surgical and non-surgical high-risk neonates.

FICare Plus is an adaptation of FICare which is tailored to address the needs of parents of critically ill infants early in their hospital course. Through the engagement of parents and multidisciplinary staff, we developed a specific FICare Plus toolkit including educational resources for staff and parents. We aimed to promote close physical contact between parents and their ill infants, even when they required specialized care due to the severity of their illness. An important task in this regard was to train and support parents to get familiar with the special needs of their infants, such as invasive and non-invasive respiratory support, ostomy bags, catheters, feeding tubes, etc. Although there was a possibility that the expectation placed on parents to perform these tasks might distress them, it was interesting to note that parents felt comfortable in learning and performing these skills when invited to do so as described in their feedback survey. No difference was noted in parental stress on day 21 (3.3 vs 3; p-value 0.3), though this might be due to the small sample size of our study. A recent case-control study by Kubicka et al supported these findings. They reported that parental stress scores were significantly lower in parents in the FICare group as compared to those in the control group and that the more parents were involved in their infants' care, the lower were their stress scores [2].

In our study, no adverse events such as accidental extubation, removal of intravenous catheters, displacement of nasogastric tubes or parental emotional breakdown were observed in the FICare Plus group. This is consistent with the findings of a recent clinical trial conducted in three NICUs in California in which preterm infants of gestational age < 33 weeks with complex medical needs were enrolled. No parent related adverse events were observed in the FICare group in this study [10].

Our findings are also in keeping with those of other recent studies in this population of infants. Moreno-Sanz and her colleagues carried out a study in an advanced NICU which is a referral center for critically ill infants requiring surgeries, extracorporeal membrane oxygenation (ECMO) and other

complex procedures. Of the 91 infants enrolled, 74.7% were preterm, 18.7% had complex surgeries and 6.6% had other morbidities. All families enrolled in this FICare study completed the educational module and training and participated regularly in their infants' care, except one who withdrew voluntarily. Assessment of parental perception about implementation of FICare by both qualitative and quantitative measures (FICare satisfaction questionnaire) showed that parents felt confident and safe to participate in their infants' care [20]. Similarly, the report from the study conducted in three large neonatal intensive care units (NICU) including two quaternary care NICU's in California, demonstrated that FICare could be implemented in infants with complex care needs, including those who required surgery. It also showed that implementation of FICare in critically ill preterm babies resulted in improved weight gain and reduction in nosocomial infection [10].

Another interesting finding of our study is that parental anxiety scores prior to discharge were significantly lower in FICare Plus group as compared to the control group {63 (52-74.5) vs 78 (71-90); p-value 0.02}. This is similar to the finding in a multicenter, multinational, cluster- randomized control trial in which parents in the FICare group had lower mean anxiety scores as compared to those in the control cohort {70.8 vs 74.2 ; p-value 0.004}[5]. This could be accredited to the fact that they had the opportunity to directly observe and participate in their infants' care along with the medical team. This may have given them a feeling of self-efficacy and confidence over their parenting abilities resulting in less anxiety. Reduced parental anxiety is a finding of added importance in our study, especially in the context of critically ill infants. There is convincing evidence available in literature which suggests that parental anxiety influences parenting behaviors with a cumulative long-term negative impact on their child's development [21-23]. This suggests that FICare Plus has the potential to improve not only the short -term outcomes, but also provide long-term benefits to infants and their families.

We acknowledge the limitations of our study. As a pilot study with a small sample size, the statistical power was limited. Another limitation of our study is that we only included families who could receive education in English, even if not their primary language. These findings may therefore not be generalized to our entire NICU population. As FICare is being adapted in many different cultures and situations, further information may become available.

The implications of this study are manifold with the potential to impact health outcomes, clinical practice, health policy and cost of care. The parents' comfort with knowing and interacting with their infants even when critically ill may give them confidence and expertise to take better care of their infants. This could result in improved short- and long-term health outcomes for infants including reduced length of hospital stay. In addition, they may be able to care better after discharge home, with the potential for decreased post-discharge morbidity, reduced health care utilization and re-hospitalization. All these may reduce the cost of NICU care, and thus have important implications for both policy development and resource allocation decisions.

5. Conclusions

This pilot study shows that it is feasible and safe to implement family integrated care in critically ill infants. Future quality improvement initiatives may provide further information about the processes required for its sustainability and refinement.

Author Contributions: KOB, AC, NSA, LF, and CT conceptualized and designed the study. KOB, NSA and AC coordinated and interpreted the initial analysis. NSA drafted the initial manuscript, which was reviewed and revised by KOB, LF, AC and CT for important intellectual content. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by MSH-UHN AMO Innovation fund.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Review Board of Mount Sinai Hospital (REB # 18-0076E, dated June12, 2018) and Sick Kids Hospital ((REB # 1000059541, dated February 14, 2019).

Informed Consent Statement: Informed consent was obtained from parents of all infants involved in the study.

Data Availability Statement: The data presented in this study is not publicly available due to privacy restrictions.

Acknowledgments: We would like to acknowledge Xiang Ye for his contribution in performing the statistical analysis as well as Eugene Woojin Yoon for reviewing the data analysis.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. O'Brien K, Bracht M, Macdonell K, McBride T, Robson K, O'Leary L, Christie K, Galarza M, Dicky T, Levin A, Lee SK. A pilot cohort analytic study of Family Integrated Care in a Canadian neonatal intensive care unit. *BMC Pregnancy Childbirth.* 2013;13 Suppl 1(Suppl 1):S12. doi: 10.1186/1471-2393-13-S1-S12. Epub 2013 Jan 31. PMID: 23445639; PMCID: PMC3561192.
2. Kubicka Z, Fiascone J, Williams D, Zahr E, Ditzel A, Perry D, Rousseau T, Lacy M, Arzuaga B. Implementing modified family integrated care in a U.S. neonatal intensive care unit: nursing perspectives and effects on parents. *J Perinatol.* 2023 Apr;43(4):503-509. doi: 10.1038/s41372-023-01601-y. Epub 2023 Jan 10. PMID: 36627393; PMCID: PMC9838294.
3. Hei M, Gao X, Gao X, Nong S, Zhang A, Zhang Q, Lee SK. Is family integrated care in neonatal intensive care units feasible and good for preterm infants in China: study protocol for a cluster randomized controlled trial. *Trials.* 2016 Dec;17(1):1-8.
4. He SW, Xiong YE, Zhu LH, Lv B, Gao XR, Xiong H, Wang H, Shi HR, Latour JM. Impact of family integrated care on infants' clinical outcomes in two children's hospitals in China: a pre-post intervention study. *Ital J Pediatr.* 2018 Jun 5;44(1):65. doi: 10.1186/s13052-018-0506-9. PMID: 29871689; PMCID: PMC5989344.
5. O'Brien K, Robson K, Bracht M, Cruz M, Lui K, Alvaro R, da Silva O, Monterrosa L, Narvey M, Ng E, Soraisham A, Ye XY, Mirea L, Tarnow-Mordi W, Lee SK; FICare Study Group and FICare Parent Advisory Board. Effectiveness of Family Integrated Care in neonatal intensive care units on infant and parent outcomes: a multicentre, multinational, cluster-randomised controlled trial. *Lancet Child Adolesc Health.* 2018 Apr;2(4):245-254. doi: 10.1016/S2352-4642(18)30039-7. Epub 2018 Feb 8. Erratum in: *Lancet Child Adolesc Health.* 2018 Aug;2(8):e20. PMID: 30169298.
6. Hei M, Gao X, Li Y, Gao X, Li Z, Xia S, Zhang Q, Han S, Gao H, Nong S, Zhang A, Li J, Wang Y, Ye XY, Lee SK. Family Integrated Care for Preterm Infants in China: A Cluster Randomized Controlled Trial. *J Pediatr.* 2021 Jan;228:36-43.e2. doi: 10.1016/j.jpeds.2020.09.006. Epub 2020 Sep 6. PMID: 32898578.
7. Feldman R, Rosenthal Z, Eidelman AI. Maternal-preterm skin-to-skin contact enhances child physiologic organization and cognitive control across the first 10 years of life. *Biol Psychiatry.* 2014;75(1):56-64. doi:10.1016/j.biopsych.2013.08.012.
8. Church PT, Grunau RE, Mirea L, Petrie J, Soraisham AS, Synnes A, Ye XY, O'Brien K. Family Integrated Care (FICare): Positive impact on behavioural outcomes at 18 months. *Early Hum Dev.* 2020 Dec;151:105196. doi: 10.1016/j.earlhumdev.2020.105196. Epub 2020 Sep 19. PMID: 32987227.
9. Benzies KM, Aziz K, Shah V, Faris P, Isaranuwatchai W, Scotland J, Larocque J, Mrklas KJ, Naugler C, Stelfox HT, Chari R, Soraisham AS, Akierman AR, Phillipos E, Amin H, Hoch JS, Zanoni P, Kurilova J, Lodha A; Alberta FICare Level II NICU Study Team. Effectiveness of Alberta Family Integrated Care on infant length of stay in level II neonatal intensive care units: a cluster randomized controlled trial. *BMC Pediatr.* 2020 Nov 28;20(1):535. doi: 10.1186/s12887-020-02438-6. PMID: 33246430; PMCID: PMC7697372.
10. Franck LS, Gay CL, Hoffmann TJ, Kriz RM, Bisgaard R, Cormier DM, Joe P, Lothe B, Sun Y. Neonatal outcomes from a quasi-experimental clinical trial of Family Integrated Care versus Family-Centered Care for preterm infants in U.S. NICUs. *BMC Pediatr.* 2022 Nov 22;22(1):674. doi: 10.1186/s12887-022-03732-1.
11. Murphy M, Shah V, Benzies K. Effectiveness of Alberta Family-Integrated Care on Neonatal Outcomes: A Cluster Randomized Controlled Trial. *J Clin Med.* 2021 Dec 14;10(24):5871. doi: 10.3390/jcm10245871. PMID: 34945163; PMCID: PMC8708302.
12. Als H, McAnulty GB. The Newborn Individualized Developmental Care and Assessment Program (NIDCAP) with Kangaroo Mother Care (KMC): Comprehensive Care for Preterm Infants. *Curr Womens Health Rev.* 2011 Aug;7(3):288-301. doi: 10.2174/157340411796355216. PMID: 25473384; PMCID: PMC4248304.

13. Waddington C, van Veenendaal NR, O'Brien K, Patel N; International Steering Committee for Family Integrated Care. Family integrated care: Supporting parents as primary caregivers in the neonatal intensive care unit. *Pediatr Investig.* 2021 Jun 18;5(2):148-154. doi: 10.1002/ped4.12277. PMID: 34179713; PMCID: PMC8212757
14. Bouet KM, Claudio N, Ramirez V, García-Fragoso L. Loss of parental role as a cause of stress in the neonatal intensive care unit. *Bol Asoc Med P R.* 2012 Jan-Mar;104(1):8-11. PMID: 22788072
15. Warre R, O'Brien K, Lee SK. Parents as the primary caregivers for their infant in the NICU: benefits and challenges. *NeoReviews.* 2014 Nov;15(11): e472-7.
16. Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA. Manual for the state-trait anxiety inventory. Palo Alto: Consulting Psychologists Press, 1983.
17. Miles MS, Funk SG, Carlson J. Parental Stressor Scale: neonatal intensive care unit. *Nurs Res* 1993; 42: 148-52.
18. Barnes CR, Adamson-Macedo EN. Perceived maternal parenting self-efficacy (PMP S-E) tool: development and validation with mothers of hospitalized preterm neonates. *J Adv Nurs.* 2007;60(5):550-560
19. Raiskila S, Lehtonen L, Tandberg BS, et al. Parent and nurse perceptions on the quality of family-centered care in 11 European NICUs. *Austr Crit Care.* 2016;29(4):201-209.
20. Moreno-Sanz B, Montes MT, Antón M, Serrada MT, Cabrera M, Pellicer A. Scaling Up the Family Integrated Care Model in a Level IIIC Neonatal Intensive Care Unit: A Systematic Approach to the Methods and Effort Taken for Implementation. *Front Pediatr.* 2021 Jun 9;9:682097. doi: 10.3389/fped.2021.682097. PMID: 34178899; PMCID: PMC8219911.
21. Delobel-Ayoub M, Arnaud C, White-Koning M, Casper C, Pierrat V, Garel M, Burguet A, Roze JC, Matis J, Picaud JC, Kaminski M, Larroque B; EPIPAG Study Group. Behavioral problems and cognitive performance at 5 years of age after very preterm birth: the EPIPAG Study. *Pediatrics.* 2009 Jun;123(6):1485-92. doi: 10.1542/peds.2008-1216. PMID: 19482758.
22. Singer LT, Fulton S, Davillier M, Koshy D, Salvator A, Baley JE. Effects of infant risk status and maternal psychological distress on maternal-infant interactions during the first year of life. *J Dev Behav Pediatr.* 2003 Aug;24(4):233-41. doi: 10.1097/00004703-200308000-00003. PMID: 12915795.
23. Treyvaud K, Anderson VA, Lee KJ, Woodward LJ, Newnham C, Inder TE, Doyle LW, Anderson PJ. Parental mental health and early social-emotional development of children born very preterm. *J Pediatr Psychol.* 2010 Aug;35(7):768-77. doi: 10.1093/jpepsy/jsp109. Epub 2009 Dec 2. PMID: 19955253.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.