

Article

Not peer-reviewed version

Runaway Climate Across the Wider Caribbean and Eastern Tropical Pacific in the Anthropocene: Threats to Coral Reef Conservation, Restoration, and Social-Ecological Resilience

[Edwin A. Hernández-Delgado](#)^{*} and Yanina M. Rodriguez-Gonzalez

Posted Date: 25 April 2025

doi: 10.20944/preprints202504.2137.v1

Keywords: Anthropocene; Caribbean; climate change; coral reefs; eastern tropical Pacific; surface(2m) temperature; sea surface temperature; runaway climate



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Article

Runaway Climate Across the Wider Caribbean and Eastern Tropical Pacific in the Anthropocene: Threats to Coral Reef Conservation, Restoration, and Social-Ecological Resilience

Edwin A. Hernández-Delgado * and Yanina M. Rodríguez-González

Sociedad Ambiente Marino, San Juan, Puerto Rico; edwinhernandez@sampr.org, yaninarodriguez@sampr.org

* Correspondence: edwinhernandez@sampr.org

Abstract: Marine heatwaves (MHWs) are increasingly affecting tropical seas, causing mass coral bleaching and mortality in the wider Caribbean (WC) and eastern tropical Pacific (ETP). This leads to significant coral loss, reduced biodiversity, and impaired ecological functions. Climate models forecast a troubling future for Latin American coral reefs, but downscaled projections for the WC and ETP remain limited. Understanding regional temperature thresholds that threaten coral reef futures and restoration efforts is critical. Our goals included analyzing historical trends in July-August-September-October (JASO) temperature anomalies and exploring future projections at subregional and country levels. From 1940 to 2023, JASO air and ocean temperature anomalies showed significant increases. Projections indicate that even under optimistic scenario 4.5, temperatures may exceed the +1.5°C air threshold beyond pre-industrial levels by the 2040s and the +1.0°C ocean threshold beyond historical annual maximums by the 2030s, resulting in severe coral bleaching and mortality. Business-as-usual scenario 8.5 suggests conditions will become intolerable for coral conservation and restoration by the 2030s, with decadal warming trends largely surpassing historical rates, under unbearable conditions for corals. Immediate development of regional and local adaptive coral reef conservation and restoration plans, along with climate change adaptation and mitigation strategies, is essential to provide time for optimistic scenarios to materialize.

Keywords: Anthropocene; Caribbean; climate change; coral reefs; eastern tropical Pacific; surface (2m) temperature; sea surface temperature; runaway climate

1. Introduction

The Intergovernmental Panel on Climate Change (IPCC) Sixth Assessment Report characterizes marine heatwaves (MHWs) as periods during which water temperatures are abnormally warm for the time of year, relative to historical temperatures, with such extreme warmth persisting for days to months [1]. These events can occur anywhere in the ocean and span areas up to thousands of kilometers. A MHW is defined as a discrete, prolonged event of anomalously warm sea surface temperatures (SST) that persists for five or more consecutive days, during which the SST exceeds the 90th percentile relative to a 30-year historical baseline for that region and time of year [2]. MHWs have surged in tropical seas over recent decades [3], with 2023 and 2024 setting numerous temperature records [4,5]. Sea surface warming trends have doubled per decade [6], projecting widespread coral population declines as temperatures exceed +2°C [7]. Recent prolonged MHWs have caused significant coral reef habitat modifications, leading to long-term consequences for ecological functions and resilience [8,9]. Local stressors synergize with rising temperatures to kill corals [10] by increasing their vulnerability to bleaching and diseases [11–15], while also heightening genetic susceptibility to warming [16]. These rapidly changing conditions signal the potential onset of a runaway climate scenario, with threatening unprecedented impacts across the Wider Caribbean (WC) and Eastern Tropical Pacific (ETP).

Severe regional extreme events may trigger tipping points in both ecosystems and human systems due to climate drivers [17]. Interactions among tipping elements beyond +2°C could result in unprecedented cascading impacts [18–20], potentially leading to runaway climate conditions characterized by permanent MHWs. MHWs significantly affect marine ecosystems, particularly in tropical regions, impacting even highly migratory species and top predators [21]. Varying in duration and intensity, MHWs are particularly concerning in tropical and subtropical waters. Their frequency has increased more than 20-fold due to anthropogenic climate change [22].

Elevated SST during prolonged MHWs cause corals to expel their endosymbiotic algae, leading to mass bleaching [23–26] (**Figure 1**). This not only jeopardizes coral health but also diminishes their nutrient acquisition [27]. Extended bleaching can result in high coral mortality rates, severely impacting reef structure and biodiversity [28,29]. Many marine species may migrate to cooler waters, disrupt local ecosystems and alter species composition [30], which can result in loss of biodiversity and changes in predator-prey dynamics. Vulnerable species face heightened extinction risks due to their inability to adapt to rapid temperature changes and chronic MHW conditions [31–37], ultimately reshaping reef functionality [38]. Projections indicate that by 2080, coral bleaching will likely begin on most reefs in spring, with year-round bleaching risks remaining high for some low-latitude reefs, regardless of global greenhouse gas (GHG) mitigation efforts [39]. Consequently, understanding regional SST changes across the WC and ETP has become increasingly crucial.




Figure 1. a) Example of severe bleaching and recent mass mortality on a remnant patch of endangered Elkhorn coral (*Acropora palmata*) in Puerto Rico, 2024; b) NOAA Coral Reef Watch 5km Degree Heating Week (DHW) year-to-date maximum (as of Nov. 24, 2024), showing an unprecedented accumulation of over 20 DHWs across the eastern Caribbean and the spatial extent of the 2024 marine heatwave.

Warmer waters significantly impact primary productivity, particularly phytoplankton growth—the foundation of the marine food web [40–42]. Changes in phytoplankton dynamics can cascade through trophic levels, altering tropical coastal productivity and fisheries [43–45]. MHWs disrupt nutrient cycling, affecting the availability of essential nutrients and dissolved oxygen for marine life [46–48]. They can also impact seagrass community structure and ecosystem processes [49]. Changes in fish behavior, migration, and breeding patterns due to rising temperatures can lead to declines in fish stocks, affecting local fisheries and dependent communities. Long-term consequences, such as reef flattening from sea surface warming and stronger hurricanes, may result in further habitat homogenization and loss of fish diversity [50–52]. This decline in fisheries can severely impact the economies of coastal communities reliant on fishing [53]. Additionally, warmer SST can facilitate pathogen growth, increasing disease outbreaks in marine species, especially corals and fish [54–56]. The degradation of marine ecosystems due to increasing MHWs compromises essential services, such as coastal protection, tourism, and recreational opportunities. Communities dependent on marine resources may face food insecurity as fish populations decline and ecosystems are disrupted. Overall, MHWs pose a significant threat to tropical marine ecosystems, with far-reaching implications for

biodiversity, fisheries, and coastal livelihoods. Addressing runaway climate change and mitigating its impacts are critical for enhancing ecosystem resilience and protecting these communities.

Runaway climate refers to a scenario in which climate change accelerates uncontrollably, resulting in severe and potentially irreversible alterations to the Earth's climate system. This phenomenon is marked by feedback loops that exacerbate warming, such as GHG release from melting permafrost and the loss of reflective ice cover. It typically occurs when climate tipping points are exceeded, causing changes in large parts of the climate system—known as tipping elements—to become self-perpetuating beyond a certain warming threshold [57]. This process creates a self-reinforcing cycle where initial warming triggers further warming, pushing the climate system into a state of extreme change [58,59], with potentially long-term dire global consequences [60]. Runaway climate can manifest when specific thresholds or tipping points are crossed, leading to abrupt climate shifts that are challenging to reverse.

Rising SST and record accumulated degree heating weeks (DHWs) during 2023 and 2024 resulted in widespread catastrophic coral bleaching, diminishing coral health, biodiversity, and in the long-term project to affect the structural integrity of reef ecosystems. Coral reefs support a vast array of marine life, and their collapse can lead to significant habitat loss for numerous species. Increased SST and habitat changes can facilitate the spread of invasive species, further stressing native marine life. Runaway climate impacts phytoplankton productivity, which is essential for marine food webs. Variations in nutrient availability and SST can disrupt phytoplankton growth and distribution, affecting higher trophic levels. Changes in fish populations due to temperature shifts and habitat loss can lead to declines in fisheries, jeopardizing food security for coastal communities. Additionally, elevated CO₂ atmospheric concentration can lead to higher absorption by oceans, resulting in ocean acidification. This negatively impacts calcareous organisms, such as corals and shellfish, compromising their ability to build and maintain structures. Acidification can disrupt the functions and services of marine ecosystems, including nutrient cycling and habitat provision. Runaway climate also contributes to rising sea levels, threatening coastal habitats like mangroves, seagrasses, and salt marshes. These ecosystems are vital for coastal protection and biodiversity, making their conservation and restoration increasingly complex and expensive, particularly in small island developing states (SIDS) and least developed countries [61].

Runaway climate poses severe risks to tropical marine ecosystems, leading to degradation, loss of biodiversity, and altered ecological dynamics. The impacts extend beyond marine life; human communities that rely on healthy oceans for their livelihoods and well-being are also affected. More intense storms can increase coastal erosion and habitat destruction, further threatening marine and coastal social-ecological systems. Changes in temperature gradients disrupt ocean currents, affecting the distribution of nutrients and dissolved oxygen, as well as marine life migration patterns. The increased frequency of extreme weather events—such as marine heatwaves, coral bleaching, mass coral mortality, hypoxic/anoxic events, and harmful algal blooms—can cause significant disruptions to marine ecosystems, including habitat destruction and increased sedimentation, leading to ecological collapses. The bleaching events of 2023 and 2024 resulted in catastrophic coral mortality, affecting nursery-raised and out-planted colonies across various locations across the WC and ETP, decimating coral populations and reducing biodiversity and habitat complexity. The rising frequency and greater spatial and temporal extent of extreme events can limit the success of coral restoration projects, as newly restored corals struggle to survive repeated stress events, hindering their growth and survival. In the long term, exceeding critical tipping points that lead to runaway climate can incur profound socioeconomic costs [62], increasing confidence in physical and socioeconomic impacts of compound extreme events [63]. Estimates indicate over \$3.1 billion in indirect losses of ecosystem services over multiple years [64]. Additionally, studies suggest a regional or global increase in the number and intensity of hurricanes due to climate change [65,66], making large economic shocks more frequent and exacerbating runaway climate threats. Addressing climate change and its drivers is essential to prevent a runaway climate scenario and to safeguard marine ecosystems for future generations.

Coral reefs provide essential services, including coastal protection, fisheries, and tourism. Their degradation results in increased coastal erosion, loss of fish habitats, and a decline in tourism revenues. This decline significantly impacts fish populations, leading to reduced catches and affecting local livelihoods, which can result in food insecurity and economic losses for communities [67]. As major tourist attractions, coral reefs' degradation leads to fewer tourist arrivals, causing further economic challenges for communities reliant on tourism. Weakening coral reefs make coastal areas more susceptible to storm surges and erosion, exacerbating the impacts of hurricanes and sea-level rise (SLR).

Recent climate model projections from the IPCC indicate a troubling future of global-scale warming for Latin American coral reefs. However, downscaled climate projections for the WC and ETP are limited. There is an urgent need to understand the projected spatio-temporal scales of future temperature thresholds that could jeopardize coral reef futures and the success of restoration efforts in the Anthropocene.

Our primary goals include: 1. Understanding historical trends: We aim to analyze historical changes in July-August-September-October (JASO) air temperature and sea surface temperature (SST) across the region; 2. Exploring fluctuations: We explore subregional and country-level spatio-temporal fluctuations in projected JASO temperature thresholds that could threaten coral reefs in the future; 3. Local-level recommendations: We discuss important recommendations at the local level to adapt to projected climate changes. To achieve these goals, we used the European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5 climate database to reconstruct historical climate trends across 30 different countries and subregions of the WC and ETP, focusing on the spatio-temporal variation from 1940 to 2023 for surface (2m) air temperature and SST. We also employed climate models from the CMIP6 SSP245 and SSP585 ensemble averages to project future regional warming trends from 1851 to 2100, based on two contrasting scenarios defined by the IPCC [1]. This information will be critical for developing adaptive regional and local-scale coral reef restoration plans, as well as for formulating environmental planning and sustainable development strategies to adapt to projected changes.

Addressing the increased challenges posed by projected extreme weather events and runaway climate changes across the WC and ETP throughout the Anthropocene necessitates a deeper, downscaled approach. This involves focusing on regional, subregional, and national scales to better understand the projected impacts on coral reef futures. It also requires a profound understanding of the spatio-temporal patterns of projected increases in surface (2m) temperature anomalies beyond +1.5°C compared to pre-industrial times, as well as projected increases over +1.0°C above mean annual maximums in SST. Key aspects include: 1. Critical Thresholds: Gaining insights into these critical temperature thresholds is essential for understanding the limits to which coral reefs can adapt and survive; 2. Adaptive Management: Enhanced comprehension of temperature impacts will improve our ability to plan and implement adaptive management approaches tailored to local conditions; 3. Coral Restoration Planning: Understanding these patterns will inform effective coral restoration strategies, identifying specific needs and interventions required for recovery; 4. Increased Funding Strategies: Awareness of the urgency surrounding these temperature thresholds can help secure increased local and regional funding for resilience-building initiatives; and 5. International Cooperation: Promoting collaboration among WC and ETP countries will be vital for enhancing the resilience of Latin American coral reefs and supporting the communities that depend on them. By deepening our understanding of these dynamics, we can better prepare for future challenges and work towards sustainable solutions for coral reef conservation and restoration and community livelihoods.

2. Materials and Methods

This study aimed to analyze historical changes in JASO air temperature and SST anomalies across the WC and ETP region. Regional, subregional and country-level spatio-temporal fluctuations in projected JASO temperature thresholds that could threaten coral reefs in the future were also

explored under two contrasting climate scenarios. The Climate Reanalyzer tool (<https://climatoreanalyzer.org>) served as the primary source for historical JASO surface (2m) temperature and SST anomaly (SST-a) data across 30 countries and subregions in the WC and ETP. Polygons were specifically drawn for each country or subregion to extract historical data sets (**Figure 2, Table S1**). Data were derived from the ECMWF ERA5 climate database [68], which is the fifth generation of ECMWF atmospheric reanalysis, covering the global climate since January 1940. JASO temperature anomaly data were obtained from 1940 to 2023 from each selected area. To project future climate scenarios, we utilized the International Coupled Model Intercomparison Project 6 (CMIP6) developed by the World Climate Research Programme (WCRP), which provides a new set of climate scenarios [69]. These scenarios encompass various socio-economic developments and pathways of atmospheric GHG concentrations. For this project, two scenarios were selected: 1. SSP245: This is the optimistic model, representing a medium pathway of future GHG emissions. It assumes that climate protection measures are being implemented, with an additional radiative forcing of 4.5 W/m² by the year 2100; 2. SSP585: This business-as-usual model represents the upper boundary of climate scenarios, predicting an additional radiative forcing of 8.5 W/m² by the year 2100. It reflects a scenario where no significant climate protection measures are taken. These models and datasets are critical for understanding historical trends and projecting future impacts on coral reef ecosystems in the region.

Climate models CMIP6 SSP245 and CMIP6 SSP585 ensemble averages were utilized to project future JASO warming trends at regional, subregional, and country levels, using historical data and modeling from 1851 to 2100 under both optimistic (4.5 W/m²) and business-as-usual (8.5 W/m²) scenarios based on IPCC criteria [1]. Decadal variation analysis was conducted using a one-way permutational analysis of variance (PERMANOVA) on normalized annual temperature anomalies (based on the 1951-1980 climatology) to assess temporal variations in JASO surface (2m) temperature and SST-a [70]. Linear regression was employed to quantify the relationship between JASO surface (2m) temperature and SST-a using historical datasets from 1940 to 2023 to parameterize the model, organized by country and subregion. Individual regressed formulas were used to calculate future SST-a trends based on projected surface (2m) temperature anomalies under both scenarios.

Table S1 summarizes the study locations by subregion, including coordinates of sampling polygons and the linear regression results between normalized JASO surface (2m) temperature and SST-a for the period of 1940 to 2023, using the 1951-1980 climatology as a baseline. A one-way analysis of similitude (ANOSIM) was implemented to test significant variations between observed and modeled temperatures from 2015 to 2024, contrasting observed versus modeled JASO temperature trends [71]. All multivariate tests used 9,999 permutations and normalized temperature anomalies. Principal components analysis (PCA) was used to project decadal variation in temperature anomalies [71]. Projected temperature anomaly variation between 1851 and 2100 under scenarios 4.5 and 8.5 at the country level were summarized in the Supplementary section (**Figures S1–S8**). This comprehensive approach allowed for a nuanced understanding of temperature trends and their potential impacts on coral reef futures in the face of climate change.




Figure 2. Spatial distribution of studied subregions: 1. Western Caribbean (WCa); 2. Southwestern Caribbean (SWC); 3. Southern Caribbean (SCa); 4. Eastern Caribbean (ECa); 5. Northern Caribbean (NCa); 6. Northwestern Caribbean (NWC); 7. Northeastern Tropical Pacific (NETP); 8. Southeastern Tropical Pacific (SETP). Image source: Google Earth.

3. Results

3.1. Regional Temperature Anomaly Patterns 1940-2023

The analysis of regional spatio-temporal trends in JASO temperature anomalies between 1940 and 2023 revealed significant warming patterns. There was a significant increase in JASO surface (2m) temperature anomalies ($p<0.0001$) (**Figure 3, Table 1**). There was a regional average of -0.1767°C during the 1940s and of $+0.9680^{\circ}\text{C}$ during the 2020s, or an average warming trend of $\sim 0.1431^{\circ}\text{C}$ per decade. Pairwise analyses showed that regional surface (2m) temperature anomalies showed a significant increase during the 1960s, a marginal increase during the 1970s and a irreversible increase since the 1980s (**Table 1**).




Figure 3. Spatio-temporal variation in decadal JASO regional mean surface (2m) temperature anomaly across the wider Caribbean and eastern Pacific regions between 1940 and 2023.

Table 1. Summary of a one-way PERMANOVA test of the spatio-temporal variation in JASO regional normalized surface (2m) temperature anomaly.

Source ¹	df	SS	MS	Pseudo-F	p
Region	8	452.3	56.5	20.03	<0.0001
Residual	75	211.7	2.8		
Pairwise groups	t	p			
1940 vs 1950	1.44	0.1507			
1940 vs 1960	3.43	0.0003			
1940 vs 1970	1.77	0.0586			
1940 vs 1980	2.82	0.0045			
1940 vs 1990	4.84	<0.0001			
1940 vs 2000	8.14	<0.0001			
1940 vs 2010	10.58	<0.0001			
1940 vs 2020	7.36	0.0012			

¹ Based on 9,999 permutations; df= Degrees of freedom; SS= Sum of squares; MS= Mean squares; Data= Pseudo-F statistic, p value.

A significant increase in SST anomalies (SST-a) was also observed ($p<0.0001$) (**Figure 4, Table 2**). A a regional mean SST-a of $+0.0259^{\circ}\text{C}$ during the 1940s and of $+0.7071^{\circ}\text{C}$ during the 2020s was documented, or an average warming trend of $\sim 0.0916^{\circ}\text{C}$ per decade. Pairwise analysis showed a significant increase in regional SST-a during the 1970s, and then a irreversible increase after the 2000s.




Figure 4. Spatio-temporal variation in decadal JASO regional mean sea surface temperature anomaly (SST-a) across the wider Caribbean and eastern Pacific regions between 1940 and 2023.

Table 2. Summary of a one-way PERMANOVA test of the spatio-temporal variation in JASO regional normalized sea surface temperature anomaly.

Source ¹	df	SS	MS	Pseudo-F	p
Region	8	324.6	40.6	8.97	<0.0001
Residual	75	339.4	4.5		
Pairwise groups	t	p			
1940 vs 1950	0.73	0.5969			
1940 vs 1960	1.01	0.3599			

1940 vs 1970	2.58	0.0018
1940 vs 1980	1.19	0.2391
1940 vs 1990	0.86	0.4941
1940 vs 2000	3.72	0.0002
1940 vs 2010	4.23	0.0002
1940 vs 2020	4.59	0.0011

¹ Based on 9,999 permutations; df= Degrees of freedom; SS= Sum of squares; MS= Mean squares; Data= Pseudo-F statistic, p value.

3.2. Country-Level Temperature Anomaly Patterns 1940-2023

All 30 countries and subregions analyzed exhibited a significant increase ($p<0.0001$) in JASO surface (2m) temperature anomalies (**Table 3, Figure 5**). The temperature anomaly across the western Caribbean countries showed a mean rate of increase of 0.1723°C/decade, 0.1311°C/decade across the southwestern Caribbean, 0.1482°C/decade across the southern Caribbean, and 0.1140°C/decade across the eastern Caribbean. The temperature anomaly across the northern Caribbean showed a mean rate of increase of 0.1243°C/decade, a mean rate of 0.1119°C/decade across the northwestern Caribbean, 0.1935°C/decade across the northeastern tropical Pacific, and 0.1494°C/decade across the southeastern tropical Pacific. The data indicate a consistent warming trend across all subregions, with profound implications for the conservation and restoration of coral reefs.

Table 3. Summary of a one-way PERMANOVA test of the spatio-temporal variation in JASO country-level normalized surface (2m) temperature anomaly.

Country ¹	Pseudo-F	p	Country	Pseudo-F	P	Country	Pseudo-F	P
			Leeward Is.			Bahamas		
			Windward			Florida		
México	25.15	<0.0001	Is.	18.24	<0.0001	N. Méx. Pacific	15.97	<0.0001
Belize/Guatemala	40.68	<0.0001	US/British	16.28	<0.0001	S. Méx. Pacific	14.78	<0.0001
Honduras	32.07	<0.0001	Vis	15.67	<0.0001	Guatemala	10.61	<0.0001
Nicaragua	20.41	<0.0001	Puerto Rico	18.82	<0.0001	Pacific	26.59	<0.0001
Costa Rica	18.85	<0.0001	Dom. Rep.	19.51	<0.0001	El Salv./Hond.	45.28	<0.0001
Panamá	17.70	<0.0001	Haiti	19.54	<0.0001	Pac	17.21	<0.0001
San Andrés	16.40	<0.0001	Jamaica	20.01	<0.0001	Nicaragua	5.60	<0.0001
Colombia	19.06	<0.0001	Gr.	19.96	<0.0001	Pacific	10.05	<0.0001
Venezuela	25.40	<0.0001	Cayman	19.61	<0.0001	Costa Rica	12.64	<0.0001
ABC Islands	17.60	<0.0001	Cuba	13.81	<0.0001	Pacific	19.18	<0.0001
			Turks-Caicos			Panamá Pacific		
						Colombia		
						Pacific		

¹ Based on 9,999 permutations; Data= Pseudo-F statistic, p value; d.f.=8,75.

Significant observed changes in JASO surface (2m) temperature anomalies across Belize/Guatemala, Colombia, Venezuela, the southern Mexican Pacific, Guatemala Pacific, El Salvador/Honduras Pacific, and the Colombian Pacific became permanent after the decade of 1960. In the case of Panamá, Aruba, Bonaire and Curaçao (ABC Islands), Puerto Rico, Haiti, the northern Mexican Pacific, Nicaragua Pacific, Costa Rica Pacific, and the Panamá Pacific, significant increases in JASO surface (2m) temperature anomalies became permanent during the 1980s. Elsewhere, increases became significant during the 1990s, except for South Florida in the 2000s.





Figure 5. Spatio-temporal variation in country-level decadal means of JASO surface (2m) temperature anomalies: (a) Western Caribbean, (b) Southwestern Caribbean, (c) Southern Caribbean, (d) Eastern Caribbean, (e) Northern Caribbean, (f) Northwestern Caribbean, (g) Northeastern tropical Pacific, (h) Southeastern tropical Pacific. Country acronyms described in Table S1.

All 30 countries and subregions analyzed also exhibited a significant increase ($p=0.0123$ to $p<0.0001$) in JASO SST-a (Table 4, Figure 6). SST-a across the western Caribbean exhibited a mean rate of increase of $0.0878^{\circ}\text{C}/\text{decade}$. The southwestern Caribbean averaged $0.0775^{\circ}\text{C}/\text{decade}$, $0.0894^{\circ}\text{C}/\text{decade}$ across the southern Caribbean, and $0.0831^{\circ}\text{C}/\text{decade}$ the eastern Caribbean. SST-a across the northern Caribbean increased at a rate of $0.0863^{\circ}\text{C}/\text{decade}$, $0.0954^{\circ}\text{C}/\text{decade}$ across the northwestern Caribbean, $0.0928^{\circ}\text{C}/\text{decade}$ across northeastern tropical Pacific, and $0.0690^{\circ}\text{C}/\text{decade}$ across the southeastern tropical Pacific.

Table 4. Summary of a one-way PERMANOVA test of the spatio-temporal variation in JASO country-level normalized sea surface temperature anomaly.

Country ¹	Pseudo-F	p	Country	Pseudo-F	P	Country	Pseudo-F	P
			Leeward Is.			Bahamas		
			Windward			Florida		
México	14.85	<0.0001	Is.	11.71	<0.0001	N. Méx. Pacific	8.39	<0.0001
Belize/Guatemala	6.39	<0.0001	US/British	12.09	<0.0001	S. Méx. Pacific	12.64	<0.0001
Honduras	9.94	<0.0001	Vis	11.58	<0.0001	Guatemala	4.03	0.0005
Nicaragua	10.31	<0.0001	Puerto Rico	11.43	<0.0001	Pacific	9.22	<0.0001
Costa Rica	11.51	<0.0001	Dom. Rep.	8.84	<0.0001	El Salv./Hond.	9.55	<0.0001
Panamá	11.80	<0.0001	Haiti	8.33	<0.0001	Pac	6.12	<0.0001
San Andrés	8.88	<0.0001	Jamaica	10.26	<0.0001	Nicaragua	3.46	0.0017
Colombia	6.86	<0.0001	Gr.	11.29	<0.0001	Pacific	2.72	0.0123
Venezuela	9.07	<0.0001	Cayman	10.07	<0.0001	Costa Rica	3.46	0.0021
ABC Islands	7.36	<0.0001	Cuba	5.64	0.0002	Pacific	4.21	0.0003
			Turks-Caicos			Panamá Pacific		
						Colombia Pacific		

¹ Based on 9,999 permutations; Data= Pseudo-F statistic, *p* value; d.f.=8,75.






Figure 6. Spatio-temporal variation in country-level decadal means of JASO sea surface temperature anomalies: (a) Western Caribbean, (b) Southwestern Caribbean, (c) Southern Caribbean, (d) Eastern Caribbean, (e) Northern Caribbean, (f) Northwestern Caribbean, (g) Northeastern tropical Pacific, (h) Southeastern tropical Pacific. Country acronyms described in Table S1.

The data show a consistent upward trend in SST-a across all regions, indicating significant warming through 1940-2023. The northwestern Caribbean experienced the highest mean rate of increase at 0.0954°C/decade, while the southeastern tropical Pacific had the lowest at 0.0690°C/decade. These warming trends have critical implications for marine ecosystems, particularly coral reefs, and highlight the need for effective climate adaptation and management strategies.

Significant observed changes in JASO SST-a across the Bahamas and southern Florida became permanent after the decade of 1990. Elsewhere, increases in JASO SST-a became significant during the 2000s, except for the northeastern Mexican Pacific and Costa Rica Pacific which became significant during the 2010s, and the Panamá Pacific and Colombian Pacific, which became significant during the 2020s.

3.3. Modeled Historical Variation in Temperature Anomalies 1851-2100 – Scenario 4.5

There was a highly significant decadal-scale increase ($p < 0.0001$) in JASO surface (2m) temperature anomaly projections across each of the eight different subregions in the WC and ETP for the period from 1851 to 2100 under the optimistic scenario 4.5, compared to pre-industrial temperatures of the 1850s (**Figure 7, Table 5**). Pairwise PERMANOVA analyses revealed significant permanent increases in temperature anomalies across the northeastern tropical Pacific after the 1930s, southern, northern and northwestern Caribbean after the 1940s, western and eastern Caribbean after the 1950s, southeastern tropical Pacific across the 1960s, and across the southwestern Caribbean after the 1970s.




Figure 7. Projected spatio-temporal variation in decadal JASO regional mean 2m temperature anomaly across the wider Caribbean and eastern tropical Pacific regions between 1851 and 2100 based on Scenario 4.5.

The analysis indicates a clear trend of increasing temperature anomalies across all subregions over the decades, with each region experiencing significant changes at different times. These findings highlight the ongoing impacts of climate change in these regions and underscore the urgency for adaptive management and conservation strategies to mitigate these effects on marine ecosystems, even under the optimistic scenario 4.5.

Table 5. Summary of a one-way PERMANOVA test of the spatio-temporal variation in modeled JASO normalized surface (2m) temperature and sea surface temperature anomalies 1851-2100 under scenario 4.5, and projected onsetting of +1.5°C beyond pre-industrial JASO 2m temperature anomalies and of +1.0°C beyond maximum JASO SST anomalies.

Region ¹	Pseudo-F	p	Pseudo-F	p	2m temp ≥1.5°C	SST ≥1.0°C
	2m temp		SST			
W Caribbean	152.9	<0.0001	158.5	<0.0001	2080	2060
SW Caribbean	110.5	<0.0001	109.8	<0.0001	2060	2050
S Caribbean	145.0	<0.0001	138.5	<0.0001	2060	2060
E Caribbean	106.2	<0.0001	106.2	<0.0001	2030	2020
N Caribbean	148.4	<0.0001	147.2	<0.0001	2050	2040
NW Caribbean	210.5	<0.0001	209.6	<0.0001	2040	2030
NE Tropical Pacific	74.4	<0.0001	80.2	<0.0001	2040	2040
SE Tropical Pacific	60.6	<0.0001	61.2	<0.0001	2060	2050

¹ Based on 9,999 permutations; Data= Pseudo-F statistic, *p* value; d.f.=8,75.

The projected trends in JASO surface (2m) temperature and SST anomalies under scenarios 4.5 and 8.5 indicate significant warming across the WC and ETP (**Figures S1–S8**). Modeled JASO surface (2m) temperature anomalies across the western Caribbean averaged -0.2981°C during the 1850s, +1.1063°C during the 2020s, and were projected to rise to +2.2623°C during the 2090s under scenario 4.5, an average increase of 0.1170°C/decade until the 2020s and a projected increase of 0.1651°C/decade between the 2020s and 2090s (**Figure 7, Table 6**). The southwestern Caribbean average anomaly was -0.2918°C in the 1850s, +1.1131°C in the 2020s, and was projected to increase to +2.1057°C in the 2090s, an average of 0.1171°C/decade until the 2020s, and a projection of 0.1418°C/decade between the 2020s and 2090s. There was an average of -0.4090°C in the 1850s and

+1.0517°C in the 2020s across the southern Caribbean, with a projected rise of +2.2318°C by the 2090s, which represents an average rise of 0.1217°C/decade until the 2020s and a projected shift of 0.1686°C/decade between the 2020s and 2090s. Modeled JASO surface (2m) temperature anomalies across the eastern Caribbean averaged -0.3334°C during the 1850s, +1.1159°C during the 2020s, and were projected to rise to +2.0885°C during the 2090s under scenario 4.5, an average increase of 0.1208°C/decade until the 2020s and a projected increase of 0.1389°C/decade between the 2020s and 2090s

Table 6. Summary of observed and modeled decadal changes in JASO surface (2m) temperature anomaly under scenario 4.5. Units: °C/decade.

Region ¹	1850-2020	2020-2100	2000-2020	2020-2050	2050-2100
W Caribbean	0.1170	0.1651	0.2704	0.2064	0.1342
SW Caribbean	0.1171	0.1418	0.3038	0.1784	0.1144
S Caribbean	0.1217	0.1686	0.2309	0.2337	0.1198
E Caribbean	0.1208	0.1389	0.3033	0.1817	0.1069
N Caribbean	0.1204	0.1535	0.2832	0.2018	0.1172
NW Caribbean	0.1197	0.1654	0.2708	0.2196	0.1248
NE Tropical Pacific	0.1245	0.2014	0.1447	0.2938	0.1322
SE Tropical Pacific	0.1158	0.1577	0.1960	0.2246	0.1076
Regional average	0.1196	0.1616	0.2504	0.2175	0.1196

The projected trends in JASO surface (2m) temperature anomalies under scenario 4.5 indicate significant warming also across the northern Caribbean, with -0.3329°C during the 1850s, +1.1115°C during the 2020s, and a projected rise to +2.1857°C during the 2090s, or an average increase of 0.1204°C/decade until the 2020s, and a projected shift of 0.1535°C/decade between the 2020s and 2090s (**Figure 7, Table 6**). The northwestern Caribbean averaged -0.2733°C in the 1850s, +1.1629°C in the 2020s, and had a projected rise of +2.3208°C by the 2090s, an average of 0.0798°C/decade until the 2020s, and a projected rise of 0.1654°C/decade between the 2020s and 2090s. An average of -0.5379°C in the 1850s and of +0.9555°C in the 2020s was observed in the northeastern tropical Pacific, with a projected rise of +2.3655°C by the 2090s, an average of 0.0830°C/decade until the 2020s and a projected rise to 0.2014°C/decade between the 2020s and 2090s. The southeastern tropical Pacific averaged -0.4475°C in the 1850s, +0.9415°C in the 2020s, and had a projected rise of +2.0457°C by the 2090s, an average of 0.0772°C/decade until the 2020s and a projected increase of 0.1577°C/decade between the 2020s and 2090s.

All regions exhibit significant warming trends, with the northeastern tropical Pacific showing the highest projected increase of 0.2014°C/decade between the 2020s and 2090s. These trends are critical for understanding the potential impacts on marine ecosystems, emphasizing the need for proactive climate adaptation measures in these vulnerable regions.

The modeled trends in JASO surface (2m) temperature anomalies reveal significant regional increases across the WC and ETP, particularly in recent decades, as well as concerning warming projections, even under optimistic scenario 4.5. Regional average increase between 1850 and 2020 was 0.1196°C/decade, however the sub-period of 2000-2020 averaged 0.2504°C per decade (**Table 6**). This rate is 2.5 times the global average. The projected average temperature anomaly rise for 2020-2100 under scenario 4.5 is 0.1616°C/decade (1.6 times the global average), but with a projected average of 0.2175°C/decade for the sub-period of 2020-2050, and 0.1196°C/decade for the sub-period of 2050-2100. The data indicate a marked projected increase in regional temperature anomalies, with the most significant rise occurring in the 2000-2020 period. The projections under scenario 4.5 suggest continued warming, particularly in the first half of the 21st century, emphasizing the need for urgent climate action and adaptation strategies in the region.

The results from the pairwise PERMANOVA analysis and principal component analysis (PCA) provide important insights into projected changes in JASO surface (2m) temperature anomalies

across the WC and ETP under scenario 4.5. Significant increases in temperature anomalies were projected across the southern Caribbean by the 2030s ($t=2.32$, $p=0.0326$). Significant increases in temperature anomaly by the 2040s are projected to occur across the western ($t=3.35$, $p=0.0008$), southwestern ($t=2.78$, $p=0.0059$), eastern ($t=2.28$, $p=0.0233$), northern ($t=3.27$, $p=0.0005$), and northwestern Caribbean ($t=4.04$, $p<0.0001$), and across the northeastern tropical Pacific ($t=4.17$, $p=0.0006$) and the southeastern tropical Pacific ($t=2.80$, $p=0.0056$). PCA showed that after the 1990s, JASO surface (2m) temperature anomalies are projected to exceed a $+1.5^{\circ}\text{C}$ anomaly compared to pre-industrial temperatures, even under the optimistic scenario 4.5 (**Figure 8**). The analyses indicate a concerning trend of increasing temperature anomalies across all assessed subregions even under an optimistic scenario, with significant increments anticipated as early as the 2030s and 2040s.

The analysis of JASO SST-a projections also reveals significant rising trends ($p<0.0001$) across all eight subregions of WC and ETP for the period of 1851 to 2100, compared to pre-industrial temperatures of the 1850s (**Figure 9, Table 5**). Permanent increases in SST-a were documented by the 1930s across the northeastern tropical Pacific, by the 1940s across the southern, northern and northwestern Caribbean, by the 1950s across the western and eastern Caribbean, by the 1960s across the southeastern tropical Pacific, and by the 1970s across the southwestern Caribbean. The findings illustrate a clear trend of increasing SST anomalies across all subregions, indicating significant warming over the decades. This warming has important implications for marine ecosystems, influencing species distribution, coral health, and overall biodiversity.




Figure 8. Principal component analysis (PCA) of the projected mean decadal variation in JASO 2m temperature anomaly across the wider Caribbean and eastern Pacific regions between 1851 and 2100 based on Scenario 4.5. This solution explains 99.9% of the observed decadal variation.

The projected trends in JASO SST-a under scenario 4.5 reflect significant warming across various subregions. Observed JASO SST-a across the western Caribbean averaged -0.1796°C in the 1850s and $+0.6141^{\circ}\text{C}$ in the 2020s, with a projected rise of $+1.2549^{\circ}\text{C}$ by the 2090s, and average increase of $0.0661^{\circ}\text{C}/\text{decade}$ by the 2020s and a projected shift of $0.0915^{\circ}\text{C}/\text{decade}$ between the 2020s and 2090s (**Figure 9, Table 7**). The southwestern Caribbean averaged -0.1678°C during the 1850s and $+0.7661^{\circ}\text{C}$ in the 2020s, with a projection of $+1.4264^{\circ}\text{C}$ during the 2090s, an average rise of $0.0778^{\circ}\text{C}/\text{decade}$ until the 2020s and a projection of $0.0943^{\circ}\text{C}/\text{decade}$ between the 2020s and 2090s. The SST-a across the southern Caribbean averaged -0.1709°C in the 1850s and $+0.6266^{\circ}\text{C}$ in the 2020s, with a projection of $+1.2540^{\circ}\text{C}$ by the 2090s, an average rise of $0.0665^{\circ}\text{C}/\text{decade}$ until the 2020s and a projection of $0.0896^{\circ}\text{C}/\text{decade}$ between the 2020s and 2090s. Average SST-a was -0.3209°C in the 1850s and $+1.0931^{\circ}\text{C}$ in the 2020s across the eastern Caribbean, with a projection of $+2.0421^{\circ}\text{C}$ by the 2090s, with an average increase of $0.1178^{\circ}\text{C}/\text{decade}$ until the 2020s and a projection of $0.1356^{\circ}\text{C}/\text{decade}$ between the 2020s and 2090s.




Figure 9. Projected spatio-temporal variation in decadal JASO regional mean sea surface temperature anomaly across the wider Caribbean and eastern tropical Pacific regions between 1851 and 2100 based on Scenario 4.5.

The modeled trends in JASO SST-a under scenario 4.5 also indicate notable warming across other subregions (**Figure 9, Table 7**). The northern Caribbean averaged -0.2736°C during the 1850s and $+0.8151^{\circ}\text{C}$ during the 2020s, with a projected rise of $+1.6236^{\circ}\text{C}$ by the 2090s, an average increase of $0.0907^{\circ}\text{C/decade}$ until the 2020s and a projected increase of $0.1155^{\circ}\text{C/decade}$ between the 2020s and 2090s. Average JASO SST-a was -0.2758°C in the 1850s and $+0.9123^{\circ}\text{C}$ in the 2020s across the northwestern Caribbean, with a projection of $+1.8674^{\circ}\text{C}$ by the 2090s, an average rise of $0.0990^{\circ}\text{C/decade}$ until the 2020s and a projection of $0.1364^{\circ}\text{C/decade}$ by the 2090s. JASO SST-a across the northeastern tropical Pacific averaged -0.3592°C in the 1850s and $+0.5461^{\circ}\text{C}$ in the 2020s, with projections of $+1.4121^{\circ}\text{C}$ by the 2090s, an average increment of $0.0754^{\circ}\text{C/decade}$ until the 2020s and a projection of $0.1237^{\circ}\text{C/decade}$ by the 2090s. The southeastern tropical Pacific averaged -0.2862°C in the 1850s and $+0.6035^{\circ}\text{C}$ in the 2020s, with a projection of $+1.3082^{\circ}\text{C}$ by the 2090s, an average rise of $0.0741^{\circ}\text{C/decade}$ until the 2020s and a projection of $0.1007^{\circ}\text{C/decade}$ by the 2090s.

Table 7. Summary of observed and modeled decadal changes in JASO sea surface temperature anomaly under scenario 4.5. Units: $^{\circ}\text{C/decade}$.

Region ¹	1850-2020	2020-2100	2000-2020	2020-2050	2050-2100
W Caribbean	0.0661	0.0915	0.1550	0.1149	0.0740
SW Caribbean	0.0778	0.0943	0.2039	0.1183	0.0764
S Caribbean	0.0665	0.0896	0.1378	0.1266	0.0619
E Caribbean	0.1178	0.1356	0.2958	0.1774	0.1042
N Caribbean	0.1155	0.1155	0.2142	0.1517	0.0883
NW Caribbean	0.0990	0.1364	0.2256	0.1805	0.1034
NE Tropical Pacific	0.0754	0.1237	0.0925	0.1813	0.0806
SE Tropical Pacific	0.0741	0.1007	0.1259	0.1427	0.0692
Regional average	0.0865	0.1109	0.1813	0.1492	0.0823

Even under scenario 4.5, all subregions show significant warming trends, with projections indicating increases in SST-a becoming more pronounced as we approach the end of the century. The eastern, northwestern and northern Caribbean exhibit the highest projected increases in mean SST-a between the 2020s and 2090s, highlighting it as a critical area for monitoring. These temperature increases are likely to have substantial effects on marine ecosystems, necessitating proactive climate strategies to mitigate potential impacts on biodiversity, fisheries and livelihoods.

Mean regional average increase in JASO SST-a was $0.0865^{\circ}\text{C}/\text{decade}$ for the period of 1850-2020, but $0.1813^{\circ}\text{C}/\text{decade}$ for the sub-period of 2000-2020, which was 1.8 times the global ocean average (**Table 7**). The projected regional average for the period of 2020-2100 under scenario 4.5 is $0.1109^{\circ}\text{C}/\text{decade}$. However, for the sub-period of 2020-2050 the projection is $0.1492^{\circ}\text{C}/\text{decade}$, while for the sub-period of 2050-2100 is $0.0823^{\circ}\text{C}/\text{decade}$. This suggests a significant overshoot beyond average ocean warming trends and then a gradual moderate leveling, but not a reversal in warming trends.

When JASO SST-a during the 2020s was compared to projected changes under scenario 4.5, a significant increase was projected by the 2040s across the western ($t=3.39$, $p=0.0010$), southwestern ($t=2.74$, $p=0.0071$), southern ($t=3.45$, $p=0.0016$), eastern ($t=2.28$, $p=0.0244$), northern ($t=3.25$, $p=0.0007$) and northwestern Caribbean ($t=4.01$, $p=0.0002$), and across the northeastern ($t=4.29$, $p=0.0002$) and the southeastern tropical Pacific ($t=2.80$, $p=0.0070$). PCA showed that following the 1990s, SST anomaly is also projected to curve well beyond the $+1.0^{\circ}\text{C}$ anomaly when compared to historical maximum JASO temperatures, even under optimistic scenario 4.5 (**Figure 10**). This variation is projected to respond to generalized increases in SST-a across all subregions.




Figure 10. Principal component analysis (PCA) of the projected mean decadal variation in JASO sea surface temperature anomaly across the wider Caribbean and eastern Pacific regions between 1851 and 2100 based on Scenario 4.5. This solution explains 99.9% of the observed variation.

3.4. Modeled Historical Variation in Temperature Anomalies 1851-2100 – Scenario 8.5

There was a highly significant decadal scale increase ($p < 0.0001$) in JASO surface (2m) temperature anomaly projections across each of the eight different WC and ETP subregions for the period of 1851 to 2100 under business-as-usual scenario 8.5 when compared to pre-industrial temperatures of the 1850s (**Figure 11, Table 8**).

Table 8. Summary of a one-way PERMANOVA test of the spatio-temporal variation in modeled JASO normalized surface (2m) temperature and sea surface temperature anomalies 1851-2100 under scenario 8.5, and projected onsetting of +1.5°C beyond pre-industrial JASO 2m temperature anomalies and of +1.0°C beyond maximum JASO SST anomalies.

Region ¹	Pseudo-F	p	Pseudo-F	p	2m temp ≥1.5°C	SST ≥1.0°C
	<u>2m temp</u>		<u>SST</u>			
W Caribbean	350.0	<0.0001	356.9	<0.0001	2030	2040
SW Caribbean	231.6	<0.0001	227.7	<0.0001	2030	2040
S Caribbean	299.0	<0.0001	263.6	<0.0001	2030	2040
E Caribbean	208.5	<0.0001	208.6	<0.0001	2030	2020
N Caribbean	302.9	<0.0001	300.3	<0.0001	2030	2030
NW Caribbean	385.0	<0.0001	385.8	<0.0001	2030	2030

NE Tropical Pacific	161.5	<0.0001	171.6	<0.0001	2020	2030
SE Tropical Pacific	129.7	<0.0001	131.5	<0.0001	2030	2040

¹ Based on 9,999 permutations; Data= Pseudo-F statistic, *p* value; d.f.=8,75.

Pairwise PERMANOVA analyses showed a significant permanent increase in temperature anomaly after the 1930s across the southern Caribbean and the northeastern tropical Pacific, after the 1940s across the northern and northwestern Caribbean, after the 1950s across the western and eastern Caribbean, and across the southeastern tropical Pacific, and after the 1970s across the southwestern Caribbean.

The modeled JASO surface (2m) temperature anomaly under scenario 8.5 averaged -0.2981°C across the western Caribbean during 1850s, +1.1561°C during the 2020s, and +4.1161°C during the 2090s (**Figure 11, Table 9**), an average increase of 0.1212°C/decade until the 2020s, and a projected increase of 0.4229°C/decade between the 2020s and 2090s. Temperature anomaly averaged -0.2918°C in the 1850s across the southwestern Caribbean, +1.0504°C in the 2020s, and +3.6753°C in the 2090s, an average increase of 0.1119°C/decade until the 2020s, and a projected increase of 0.3750°C/decade between the 2020s and 2090s. The southern Caribbean averaged -0.4090°C in the 1850s and +1.0808°C in the 2020s, with a projection of +4.1277°C by the 2090s, an average of 0.1242°C/decade until the 2020s, and a projection of 0.4353C/decade between the 2020s and 2090s. The eastern Caribbean averaged -0.3334°C during the 1850s and +1.0634°C during the 2020s, with a projected rise of +3.5575°C by the 2090s, an average of 0.1164°C/decade until the 2020s, and a projection of 0.3563°C/decade between the 2020s and 2090s.



Figure 11. Projected spatio-temporal variation in decadal JASO regional mean 2m temperature anomaly across the wider Caribbean and eastern tropical Pacific regions between 1851 and 2100 based on scenario 8.5.

Mean JASO surface (2m) temperature anomaly across the northern Caribbean averaged -0.3329°C during the 1850s and +1.1302°C during the 2020s, with projections of +3.8079°C during the 2090s, averaging 0.1219°C/decade until the 2020s and a projection of 0.3825°C/decade between the 2020s and 2090s under scenario 8.5. Temperature anomaly across the northwestern Caribbean averaged -0.2733°C in the 1850s and +1.1664°C in the 2020s, with projected +4.0969°C during the 2090s, averaging 0.1200°C/decade until the 2020s, and a projection of 0.4178°C/decade between the 2020s and 2090s. The northeastern tropical Pacific averaged -0.5000°C in the 1850s and +1.1309°C in the 2020s, with projected +4.8434°C during the 2090s, averaging 0.1359°C/decade until the 2020s, and a projection of 0.5304°C/decade between the 2020s and 2090s. JASO surface (2m) temperature anomaly averaged -0.4475°C in the 1850s across the southeastern tropical Pacific and +1.0079°C in the

2020s, with projected +4.1014°C by the 2090s, averaging 0.1213°C/decade until the 2020s and a projection of 0.4419°C/decade between the 2020s and 2090s.

Mean regional average increase in JASO surface (2m) temperature anomaly was 0.1216°C/decade for the period of 1850-2020, but 0.2597°C/decade for the sub-period of 2000-2020, which was 2.6 times the global average (Table 9). Projected average for the period of 2020-2100 under scenario 8.5 is 0.4203°C/decade, or 4.2 times the global average. However, for the sub-period of 2020-2050 the projection is 0.3440°C/decade, while for the sub-period of 2050-2100 is 0.4776°C/decade.

Table 9. Summary of observed and modeled decadal changes in JASO surface (2m) temperature anomaly under scenario 8.5. Units: °C/decade.

Region ¹	1850-2020	2020-2100	2000-2020	2020-2050	2050-2100
W Caribbean	0.1212	0.4229	0.2748	0.3521	0.4756
SW Caribbean	0.1119	0.3750	0.2724	0.3225	0.4144
S Caribbean	0.1242	0.4353	0.2455	0.3513	0.4983
E Caribbean	0.1164	0.3563	0.2770	0.3148	0.3874
N Caribbean	0.1219	0.3825	0.2925	0.3387	0.4154
NW Caribbean	0.1200	0.4178	0.2726	0.3575	0.4645
NE Tropical Pacific	0.1359	0.5304	0.2134	0.3833	0.6406
SE Tropical Pacific	0.1213	0.4419	0.2293	0.3318	0.5245
Regional average	0.1216	0.4203	0.2597	0.3440	0.4776

Pairwise PERMANOVA analysis showed that when JASO surface (2m) temperature anomaly during the 2020s was compared to projected changes under a business-as-usual scenario 8.5, a significant increase was projected by the 2030s across the western (t=2.51, p=0.0250), southwestern (t=2.31, p=0.0342), southern (t=2.81, p=0.0125), eastern (t=2.27, p=0.0381), northern (t=2.39, p=0.0284) and northwestern Caribbean (t=2.93, p=0.0077), and across the northeastern tropical Pacific (t=2.41, p=0.0272), and by the 2040s across the southeastern tropical Pacific (t=2.83, p=0.0139). PCA showed that following the 2000s, JASO surface (2m) temperature anomaly is projected to increase well beyond the +1.5°C anomaly when compared to pre-industrial temperatures under business-as-usual scenario 8.5 (Figure 12). This variation is predicted to respond to generalized increases in JASO 2m temperature anomaly across all subregions.

The analyses reveal a worrying trend of increasing temperature anomalies under business-as-usual scenario 8.5, with significant warming expected as early as the 2030s. The projections highlight the urgency of addressing climate change impacts, as sustained increases in temperature anomalies could lead to severe ecological and socio-economic consequences in the affected regions. This information underscores the need for robust climate policies and adaptive strategies to mitigate the effects of rising temperatures on marine ecosystems and communities.

There was also a highly significant decadal scale increase (p<0.0001) in JASO SST-a projections across each of the eight different WC and ETP subregions for the period of 1851 to 2100 under business-as-usual scenario 8.5 when compared to pre-industrial temperatures of the 1850s (Figure 13). Pairwise PERMANOVA analyses showed a significant permanent increase in JASO SST-a after the 1930s across the northeastern tropical Pacific subregion, and after the 1940s across the southern, northern and northwestern Caribbean subregions. A permanent increase in SST anomaly was also observed after the 1950s across the western, southwestern, and eastern Caribbean, and across the southeastern tropical Pacific.




Figure 12. Principal component analysis (PCA) of the projected mean decadal variation in JASO 2m temperature anomaly across the wider Caribbean and eastern Pacific regions between 1851 and 2100 based on scenario 8.5. This solution explains 100% of the observed variation.

The modeled JASO SST-a under scenario 8.5 averaged -0.1796°C across the western Caribbean during 1850s and $+0.6325^{\circ}\text{C}$ during the 2020s, with projected $+2.2839^{\circ}\text{C}$ by the 2090s (**Figure 13**, **Table 10**), an average rise of $0.0677^{\circ}\text{C}/\text{decade}$ until the 2020s, and a projection of $0.2359^{\circ}\text{C}/\text{decade}$ between the 2020s and 2090s. SST-a averaged -0.1678°C in the 1850s across the southwestern Caribbean and $+0.7229^{\circ}\text{C}$ in the 2020s, and a projection of $+2.4593^{\circ}\text{C}$ during the 2090s, averaging $0.0742^{\circ}\text{C}/\text{decade}$ until the 2020s, and $0.2481^{\circ}\text{C}/\text{decade}$ between the 2020s and 2090s. An average anomaly of -0.1709°C was observed during the 1850s in the southern Caribbean and $+0.6181^{\circ}\text{C}$ during the 2020s, with a projection of $+2.2473^{\circ}\text{C}$ by the 2090s, averaging $0.0658^{\circ}\text{C}/\text{decade}$ until the 2020s, and $0.2327^{\circ}\text{C}/\text{decade}$ between the 2020s and 2090s. Mean JASO SST-a was -0.3209°C in the 1850s across the eastern Caribbean and $+1.0418^{\circ}\text{C}$ in the 2020s, with a projection of $+3.4757^{\circ}\text{C}$ during the 2090s, averaging $0.1136^{\circ}\text{C}/\text{decade}$ until the 2020s, and a projection of $0.3477^{\circ}\text{C}/\text{decade}$ between the 2020s and 2090s.




Figure 13. Projected spatio-temporal variation in decadal JASO regional mean sea surface temperature anomaly across the wider Caribbean and eastern tropical Pacific regions between 1851 and 2100 based on scenario 8.5.

Table 10. Summary of observed and modeled decadal changes in JASO sea surface temperature anomaly under scenario 8.5. Units: $^{\circ}\text{C}/\text{decade}$.

Region ¹	1850-2020	2020-2100	2000-2020	2020-2050	2050-2100
W Caribbean	0.0677	0.2359	0.1643	0.1979	0.2645
SW Caribbean	0.0742	0.2481	0.1823	0.2139	0.2737

S Caribbean	0.0658	0.2327	0.1335	0.1902	0.2647
E Caribbean	0.1136	0.3477	0.2701	0.2982	0.3781
N Caribbean	0.0919	0.2879	0.2209	0.2549	0.3127
NW Caribbean	0.0992	0.3459	0.2267	0.2947	0.3843
NE Tropical Pacific	0.0821	0.3233	0.1325	0.2366	0.3884
SE Tropical Pacific	0.0779	0.2843	0.1487	0.2105	0.3396
Regional average	0.0841	0.3021	0.1849	0.2371	0.3258

The northern Caribbean averaged a mean JASO SST-a of -0.2736°C during the 1850s and +0.8287°C during the 2020s, with projected +2.8440°C during the 2090s, an average rise of 0.0919°C/decade until the 2020s and a projection of 0.2879°C/decade between the 2020s and 2090s under scenario 8.5. The northwestern Caribbean averaged -0.2758°C in the 1850s and +0.9144°C in the 2020s, with projections of +3.3358°C during the 2090s, an average increase of 0.0992°C/decade until the 2020s and a projection of 0.3459°C/decade between the 2020s and 2090s. There was an average JASO SST-a of -0.3380°C in the 1850s across the northeastern tropical Pacific and +0.6473°C in the 2020s, with projections of +2.9107°C by the 2090s, averaging 0.0821°C/decade until the 2020s and a projection of 0.3233°C/decade between the 2020s and 2090s. The southeastern tropical Pacific averaged -0.2862°C in the 1850s and +0.6491°C in the 2020s, with projected +2.6390°C during the 2090s, averaging 0.0779°C/decade until the 2020s, and a projection of 0.2843°C/decade between the 2020s and 2090s.

Mean regional average increase in JASO SST-a was 0.0841°C/decade for the period of 1850-2020, but 0.1849°C/decade for the sub-period of 2000-2020, which was 1.8 times the global ocean average (Table 10). Projected average for the period of 2020-2100 under scenario 8.5 is 0.3021°C/decade. However, for the sub-period of 2020-2050 the projection is 0.2371°C/decade, while for the sub-period of 2050-2100 is 0.3258°C/decade.

Pairwise PERMANOVA analysis showed that when JASO SST-a during the 2020s was compared to projected change under scenario 8.5, a significant increase was expected by the 2030s across the western Caribbean (t=2.59, p=0.0203), southwestern Caribbean (t=2.22, p=0.0371), southern Caribbean (t=2.60, p=0.0169), eastern Caribbean (t=2.27, p=0.0381), northern Caribbean (t=2.38, p=0.0305), northwestern Caribbean (t=2.91, p=0.0082), and the northeastern tropical Pacific (t=2.60, p=0.0200), and by the 2040s across the southeastern tropical Pacific (t=2.81, p=0.0132). PCA also showed that following the 2000s, SST-a is also likely ascend well beyond the +1.0°C anomaly when compared to historical maximum JASO SST-a under business-as-usual scenario 8.5 (Figure 14).




Figure 14. Principal component analysis (PCA) of the projected mean decadal variation in JASO sea surface temperature anomaly across the wider Caribbean and eastern Pacific regions between 1851 and 2100 based on scenario 8.5. This solution explains 100% of the observed variation.

There is a significant upward trend in SST-a scenario 8.5, with notable changes expected as early as the 2030s. The projected warming underscores the urgent need for climate action and coral reef conservation and restoration adaptative strategies, as rising temperatures could severely impact

marine ecosystems, biodiversity, and coastal communities. This information highlights the importance of monitoring and addressing climate change effects to mitigate potential adverse outcomes across the WC and ETP.

3.5. Observed vs. Modeled Variation in JASO Temperature Anomalies in 2015-2024

The analysis of observed versus modeled deviations in JASO surface (2m) and SST anomalies for the decade spanning 2015 to 2024 across the WC and ETP regions revealed significant findings. An ANOSIM test showed that the observed surface (2m) temperature anomaly was significantly higher than modeled anomalies across the WC and ETP regions ($R=0.128$, $p=0.0015$), with a particular deviation from scenario 4.5 ($R=0.274$, $p=0.0050$) (**Table 11**). No significant deviation from scenario 8.5, or between scenarios 4.5 and 8.5 were documented. Observed SST-a were significantly higher than modeled anomalies ($R=0.152$, $p=0.0030$), with significantly higher observations than modeled scenarios 4.5 ($R=0.292$, $p=0.0010$) and 8.5 ($R=0.119$, $p=0.0260$). No significant difference between scenarios 4.5 and 8.5 was observed.

Table 11. Summary of a one-way ANOSIM test of the observed vs. modeled regional variation in JASO normalized surface (2m) temperature and sea surface temperature anomalies in 2015-2024 under scenarios 4.5 and 8.5.

Region ¹	Global R	p	Obs vs 4.5 R	p	Obs vs 8.5 R	p	4.5 vs 8.5 R	p
Surface (2m) temperature	0.128	0.0015	0.274	0.0050	0.076	0.1050	0.024	0.2710
Sea surface temperature	0.152	0.0030	0.292	0.0010	0.119	0.0260	0.021	0.2930

¹ Based on 9,999 permutations; Data= R statistic, *p* value.

The results indicate that observed temperature anomalies in the WC and ETP regions are significantly higher than those projected by the models, particularly under scenario 4.5. The lack of significant differences between scenarios 4.5 and 8.5 suggests that both scenarios may not fully capture the observed extremes. These findings highlight potential shortcomings in climate models and the need for further refinement to better predict future temperature anomalies, which are critical for understanding climate change impacts on marine ecosystems and regional climates.

Subregional patterns of observed vs. modeled deviations in JASO surface (2m) temperature anomaly during 2015-2024 were determined. Observed anomalies were significantly higher than modeled outcomes across the western Caribbean ($R=0.083$, $p=0.0480$), with significant deviations from scenario 4.5 ($R=0.198$, $p=0.0030$) (**Table 12**). Observed JASO surface (2m) temperature anomaly was also significantly different from scenario 4.5 projections across the southwestern Caribbean ($R=0.182$, $p=0.0190$), across the eastern Caribbean ($R=0.179$, $p=0.0080$), with significant deviations from scenarios 4.5 ($R=0.324$, $p=0.0050$) and 8.5 ($R=0.072$, $p=0.0011$), and across the northern Caribbean ($R=0.083$, $p=0.0590$), but significantly higher than projected scenario 4.5 ($R=0.198$, $p=0.0150$). Observed JASO surface (2m) temperature anomaly was significantly different from scenario 4.5 projections across the NW Caribbean ($R=0.198$, $p=0.0260$), and across the northeastern tropical Pacific ($R=0.098$, $p=0.0340$), with significant deviations from scenarios 4.5 ($R=0.167$, $p=0.0210$) and 8.5 ($R=0.185$, $p=0.0150$). No significant difference between observed and modeled temperature anomaly was observed either across the southern Caribbean or across the southeastern tropical Pacific. No significant differences were observed between scenarios 4.5 and 8.5 for 2015-2024 across any of the subregions.

Table 12. Summary of a one-way ANOSIM test of the observed vs. modeled subregional variation in JASO normalized surface (2m) temperature anomaly 2015-2024 under scenarios 4.5 and 8.5.

Region ¹	Global R	p	Obs vs 4.5 R	p	Obs vs 8.5 R	p	4.5 vs 8.5 R	p
W Caribbean	0.083	0.0480	0.198	0.0030	0.055	0.1660	-0.007	0.4480
SW Caribbean	0.064	0.0960	0.182	0.0190	-0.010	0.4630	0.019	0.2940
S Caribbean	-0.008	0.5150	0.023	0.2830	-0.038	0.7010	-0.011	0.4690
E Caribbean	0.179	0.0080	0.324	0.0050	0.072	0.0011	0.107	0.0890
N Caribbean	0.083	0.0590	0.198	0.0150	0.032	0.2450	-0.005	0.4150
NW Caribbean	0.076	0.0700	0.198	0.0260	0.050	0.1790	-0.024	0.5760
NE Tropical Pacific	0.098	0.0340	0.167	0.0210	0.185	0.0150	-0.060	0.8930
SE Tropical Pacific	0.010	0.3330	0.032	0.2440	0.031	0.2430	-0.023	0.5170

¹ Based on 9,999 permutations; Data= R statistic, *p* value.

Subregional patterns of observed vs. modeled deviations in JASO SST-a during 2015-2024 were also quantified. Observed anomalies were significantly higher than modeled outcomes across the western Caribbean (R=0.130, *p*=0.0050), with significant deviations from scenarios 4.5 (R=0.254, *p*=0.0010) and 8.5 (R=0.131, *p*=0.0210) (**Table 13**). It was also significantly higher than modeled projections across the southwestern Caribbean (R=0.099, *p*=0.0290), with significant deviations from scenario 4.5 (R=0.193, *p*=0.0080), across the eastern Caribbean (R=0.199, *p*=0.0020), with significant deviations from scenarios 4.5 (R=0.331, *p*=0.0050) and 8.5 (R=0.132, *p*=0.0310), across the northern Caribbean (R=0.134, *p*=0.0080), with significant deviations from scenarios 4.5 (R=0.239, *p*=0.0090) and 8.5 (R=0.128, *p*=0.0470), and across the northwestern Caribbean (R=0.085, *p*=0.0470), with a significant deviation from scenario 4.5 (R=0.211, *p*=0.0150). No significant deviation between observed and both modeled scenarios was observed across the southern Caribbean, the northeastern and the southern tropical Pacific.

Table 13. Summary of a one-way ANOSIM test of the observed vs. modeled subregional variation in JASO normalized sea surface temperature anomaly 2015-2024 under scenarios 4.5 and 8.5.

Region ¹	Global R	p	Obs vs 4.5 R	p	Obs vs 8.5 R	p	4.5 vs 8.5 R	p
W Caribbean	0.130	0.0050	0.254	0.0010	0.131	0.0210	0.015	0.4900
SW Caribbean	0.099	0.0290	0.193	0.0080	0.063	0.1420	0.022	0.2840
S Caribbean	0.027	0.2390	0.076	0.0670	0.011	0.3710	-0.018	0.5290
E Caribbean	0.199	0.0020	0.331	0.0050	0.132	0.0310	0.108	0.0700
N Caribbean	0.134	0.0080	0.239	0.0090	0.128	0.0470	-0.002	0.3880
NW Caribbean	0.085	0.0470	0.211	0.0150	0.066	0.1260	-0.021	0.5600
NE Tropical Pacific	0.058	0.0910	0.115	0.0520	0.106	0.0730	-0.062	0.9100
SE Tropical Pacific	-0.015	0.5350	0.032	0.2170	-0.040	0.7390	-0.028	0.5320

¹ Based on 9,999 permutations; Data= R statistic, *p* value.

4. Discussion

4.1. Spatio-Temporal Variation in JASO Temperature Anomalies 1940-2023

Significant increases in mean JASO surface (2m) temperature anomalies (0.1431°C/decade) and JASO SST-a (0.0916°C/decade) were observed from 1940 to 2023 across the WC and ETP. Decadal increases in surface temperature anomalies varied by subregion, ranging from 0.1119°C/decade in the northwestern Caribbean to 0.1935°C/decade in the northeastern tropical Pacific. These warming trends were 1.1 to 1.9 times the global average. There was notable spatio-temporal variability among countries, with significant increases recorded in the 1940s and 1960s in Belize/Guatemala, Colombia, Venezuela, and ETP subregions. Additionally, significant variations occurred between the 1940s and

1980s in Panamá, Aruba-Bonaire-Curaçao, Puerto Rico, Haiti, and Panamá Pacific. From the 1940s to the 1990s, significant temperature variations were noted in México, Costa Rica, the Leeward and Windward Islands, and the Bahamas, while changes from the 1940s to the 2000s were significant in south Florida.

Decadal increases in JASO SST-a from 1940 to 2023 varied by subregion, ranging from 0.0690°C/decade in the southeastern tropical Pacific to 0.0954°C/decade in the northwestern Caribbean, reflecting rates from just below average to average global ocean warming. Significant increases were recorded in the Bahamas and south Florida from the 1940s to 1990s, in the northeastern Mexican Pacific and Costa Rica Pacific from the 1940s to 2010s, and in the Panamá and Colombian Pacific from the 1940s to 2020s. Additionally, significant increases were observed in other regions from the 1940s to 2000s.

These results indicate a consistent warming trend across all regions from 1940 to 2023, with decadal warming rates matching or exceeding global averages. Significant coral bleaching and mortality have occurred on regional and global scales, jeopardizing coral reef conservation and restoration efforts. These changes underscore the urgent need for adaptive management and conservation strategies to protect coral reefs and their associated ecosystems.

4.2. Spatio-Temporal Variation in JASO Temperature Anomalies 1851-2100 – Scenario 4.5

A significant increase in JASO surface (2m) temperature anomalies is projected from 1851 to 2100 under optimistic scenario 4.5. Increases were noted after the 1930s in the northeastern tropical Pacific, after the 1940s in the southern, northern, and northwestern Caribbean, after the 1950s in the western and eastern Caribbean, after the 1960s in the southeastern tropical Pacific, and after the 1970s in the southwestern Caribbean. The average increase across the region was 0.1196°C/decade from the 1850s to 2020s, rising to 0.2504°C/decade in the 2000s-2020s. Projected increases for 2020s-2100s under scenario 4.5 average 0.1616°C/decade, with 0.2175°C/decade for 2020s-2050s and 0.1196°C/decade for 2050s-2100s. This indicates a likely overshoot of JASO surface temperature anomalies lasting three to four decades, with warming trends persisting regardless of scenario. Projected increases compared to the 2020s baseline suggest significant warming by the 2030s in the southern Caribbean and by the 2040s elsewhere. Even under scenario 4.5, these trends will quickly exceed the +1.5°C threshold above pre-industrial levels, emphasizing a narrow window for regional action, mitigation, and adaptation.

A significant increase in JASO SST-a is also projected from 1851 to 2100 even under optimistic scenario 4.5, aligning with surface (2m) temperature warming trends. The average increase in SST-a was 0.0865°C/decade from the 1850s to 2020s, rising to 0.1813°C/decade in the 2000s-2020s. Projected increases for 2020s-2100s under scenario 4.5 average 0.1109°C/decade, with 0.1492°C/decade for 2020s-2050s and 0.0823°C/decade for 2050s-2100s. This suggests a moderate overshoot of JASO SST-a lasting three to four decades, with trends continuing in line with current global patterns. Comparison to the 2020s baseline indicates significant increases by the 2040s across all subregions. Even under scenario 4.5, expected SST warming trends will exceed the +1.0°C threshold above maximum annual temperatures, likely leading to recurrent mass coral bleaching and mortality events by the 2040s, threatening coral reef conservation and restoration efforts in the WC and ETP. This underscores the limited window for regional action, mitigation, and adaptation.

4.3. Spatio-Temporal Variation in JASO Temperature Anomalies 1851-2100 – Scenario 8.5

JASO surface (2m) temperature anomalies are projected to show significant increases from 1851 to 2100 under a business-as-usual scenario (8.5). Noteworthy increases were observed from the 1850s to the 1930s in the southern Caribbean and northeastern tropical Pacific, after the 1940s in the northern and northwestern Caribbean, after the 1950s in the western and eastern Caribbean, and after the 1970s in the southwestern Caribbean. Under scenario 8.5, the average rate of increase in JASO surface (2m) temperature anomaly was 0.1216°C/decade from the 1850s to 2020s, escalating to 0.2597°C/decade in the 2000s-2020s. Projected increases for 2020s-2100s average 0.4203°C/decade, with 0.3440°C/decade for 2020s-2050s and 0.4776°C/decade for 2050s-2100s. This indicates a

significant overshoot of JASO surface (2m) temperature anomalies, with projected warming rates 4.2 times higher than the global average. Comparing predicted JASO surface temperature anomalies to the 2020s baseline suggests significant increases by the 2040s in the southeastern tropical Pacific and by the 2030s elsewhere. These trends under scenario 8.5 will rapidly exceed the +1.5°C threshold above pre-industrial levels, stressing a minimal temporal window for regional action, mitigation, and adaptation.

A significant increase in JASO SST-a is projected from 1851 to 2100 under a business-as-usual scenario (8.5). Prominent increases were observed from the 1850s to the 1930s in the northeastern tropical Pacific, after the 1940s across the southern, northern, and northwestern Caribbean, and after the 1950s elsewhere. The average rate of increase in SST-a across the region was 0.0841°C/decade from the 1850s to 2020s, rising to 0.1849°C/decade in the 2000s-2020s. Projected increases for 2020s-2100s under scenario 8.5 average 0.3021°C/decade, with 0.2371°C/decade for 2020s-2050s and 0.3258°C/decade for 2050s-2100s. This indicates a significant overshoot of JASO SST-a of 2-3X the global average. When compared to the 2020s baseline, significant increases are expected by the 2040s in the western, southwestern, and southern Caribbean, as well as the southeastern tropical Pacific. Remarkable warming is also projected by the 2030s in the northern and northwestern Caribbean, and the northeastern tropical Pacific, with increases anticipated before the end of the 2020s in the eastern Caribbean. These projected SST-a warming trends under scenario 8.5 will promptly exceed the +1.0°C threshold above maximum annual temperatures, leading to recurrent mass coral bleaching and mortality events as early as the 2030s. This risks coral reef conservation and restoration attempts in the WC and ETP, underlining an extremely narrow window for regional action, mitigation, and adaptation.

4.4. Factors Explaining Regional SST Warming Trends

Several natural climate teleconnections can influence sea surface warming across the WC and ETP. These teleconnections are large-scale climate patterns that may affect weather and climate variability over extensive areas. Key teleconnections affecting these regions include the El Niño–Southern Oscillation (ENSO). ENSO is one of the most influential teleconnections, characterized by periodic warming (El Niño) or cooling (La Niña) of SST in the central and eastern Pacific [72]. Warmer SSTs in the ETP often result in weakening of the trade winds and SST increase in the WC due to weakened mixing and increasing ocean heat content (OHC). This can reduce rainfall and increase drought risk in the WC. Cooler SST in the ETP during La Niña is typically associated with lower wave heights [73], and warmer-than-normal SST in the WC, potentially leading to increased rainfall and hurricane activity due to reduced windshear across the WC [74,75].

The North Atlantic Oscillation (NAO) is another significant factor and refers to fluctuations in atmospheric pressure between the Icelandic Low and the Azores High in the North Atlantic. NAO also affects the ocean through changes in OHC, gyre circulations, mixed layer depth fluctuations, salinity variability, high latitude deep water formation and sea ice cover [76], therefore strongly influencing marine ecosystem processes. The interaction between NAO and ENSO dynamics can significantly influence WC climate and SST [77]. A positive NAO phase strengthens the trade winds, which can cool the SST in the WC by enhancing evaporation and ocean mixing. A negative NAO phase weakens the trade winds, allowing SST to warm up, leading to enhanced mass coral bleaching risk, and potentially intensifying tropical cyclone development in the region. The interaction of NAO and ENSO dynamics can also influence rainfall regional rainfall patterns [78,79], potentially leading to enhanced sediment-laden and nutrient-loaded runoff pulses to coastal environments, further stressing out corals.

The Atlantic Multidecadal Oscillation (AMO) is a long-term oscillation of SST in the North Atlantic, with phases lasting several decades. AMO dynamics correlates with other indices and can have a significant influence on WC SST [80], although evidence of this correlation is ambiguous. During the warm phase of the AMO, the WC tends to experience warmer conditions, enhanced risk of coral bleaching and mortality [81], and greater hurricane activity and regional rainfall [82]. The

cool phase typically brings cooler SST and reduced hurricane activity. Significant negative relationships between star coral, *Orbicella faveolata*, skeletal density and mean SST, maximum SST, AMO, and accumulated degree heating months have been documented [83], suggesting that AMO may increasingly affect coral growth under projected warming trends.

The Madden-Julian Oscillation (MJO) is an intra-seasonal climate pattern characterized by a large-scale eastward-moving disturbance of clouds, rainfall, and winds across the tropics, with cycles of around 30–60 days. It can influence SST variability across the WC and ETP by modulating convection and wind patterns [84]. It can increase or suppress rainfall and tropical cyclone activity, affecting SST in those regions over short timescales.

The Pacific Decadal Oscillation (PDO) is a long-term fluctuation in SST in the North Pacific, with warm and cool phases that can last for decades [85]. A warm phase of the PDO can lead to increased SST in the ETP, which might enhance the influence of El Niño on SST variability in the ETP [86]. A cool phase typically has the opposite effect, cooling SST and interacting with La Niña conditions to reduce temperatures in the ETP, though potentially enhancing SST warming across the WC, augmenting coral bleaching risk.

The Intertropical Convergence Zone (ITCZ) is a belt of low pressure that circles the Earth near the equator, where trade winds converge, and it is associated with convective rainfall and cloud formation. Variations in the position of the ITCZ can directly influence SST patterns in the tropical Atlantic and ETP by altering cloud cover, precipitation, and wind patterns [87]. A more northerly ITCZ position can enhance SST warming in the WC.

The Caribbean Low-Level Jet (CLLJ) is a strong wind current that flows from east to west across the WC, typically peaking during summer. The intensity of the CLLJ affects evaporation and ocean mixing, which can either cool or warm SST, affecting regional OHC and hurricane formation [88]. A stronger jet generally cools SST in the WC, while a weaker jet allows SST to warm and accumulate OHC more easily [89,90]. Interannual variability in OHC is also related with ENSO events which modulate the CLLJ [91], and may influence SST and tropical cyclogenesis [92], potentially affecting coral bleaching risk.

These complex teleconnections do not act independently; they often interact in compounded ways, creating compound spatio-temporal patterns of SST variability across the WC and ETP. For example, ENSO and the AMO may simultaneously influence the WC's climate, with one pattern amplifying or moderating the other's effects. Understanding these natural teleconnections is key to predicting climate variability, including hurricane activity and droughts, across the region, and mass coral bleaching and mortality events. However, projected regional and global warming trends, despite the scenario, suggest that conditions like a combination of teleconnections leading to strong warming are likely to be dominant across the region through the Anthropocene. Such complex interactions may also trigger potential regional runaway climate effects with paramount consequences for WC and ETP coral reefs and coastal human communities.

4.5. Regional Consequences of Projected Warming Trends

Our findings are consistent with previous studies that have documented strong warming patterns across all subregions of the WC, particularly since the 1970s [93]. The only decades warmer than the 1970s were the 1960s for the northern Caribbean, the 1950s for the western Caribbean and the 1940s for the eastern and southern Caribbean [93]. But historical warming patterns have been geographically consistent across the region and projected warming trends are widespread and concerning under both optimistic and business-as-usual scenarios. Numerous studies have documented trends of increasing atmospheric temperature anomalies of ~0.10-0.12°C/decade across extensive geographic areas on global or regional scales throughout the 20th century to present [94–102]. However, warming rates of up to 0.16°C/decade were documented during the summer months in China between 1900 and 2006 [103]. Similar findings were observed in Israel [104]. Warming rates of up to 0.29°C/decade were projected for the period of 2011-2040 and of up to 0.14°C/decade for the period of 2061-2090 under scenario 4.5, and rates of up to 0.43°C/decade were anticipated for the

period of 2011-2040 and of up to $0.72^{\circ}\text{C}/\text{decade}$ for the period of 2061-2090 under scenario 8.5 in China [105]. Areas over the permafrost are projected to warm at even faster rates [106]. These findings highlight that warming trends during the early 21st century have accelerated when compared to historical trends.

Global oceans have also warmed at a general rate of $0.10^{\circ}\text{C}/\text{decade}$ [1]. The WC basin warmed at a rate of $0.04^{\circ}\text{C}/\text{decade}$ between 1871 and 2020, and at rates of $0.17^{\circ}\text{C}/\text{decade}$ between 1981 and 2020, and $0.20^{\circ}\text{C}/\text{decade}$ between 1994 and 2020 [107]. Other accounts have estimated WC sea surface warming trends of $0.108^{\circ}\text{C}/\text{decade}$ for the combined periods of 1906-1969 and 1972-2005, and of $0.118^{\circ}\text{C}/\text{decade}$ for the sub-period of 1972-2005 [108]. However, The Gulf of Mexico warmed at rates of up to $0.32^{\circ}\text{C}/\text{decade}$ during the period of 2000 to 2021, and the south Florida estuaries up to $0.42^{\circ}\text{C}/\text{decade}$ during the same period [109]. Areas across the southwestern Pacific warmed 0.10 - $0.20^{\circ}\text{C}/\text{decade}$ between 1982 and 2016 [110]. Coral proxy paleoclimate reconstructions have also shown significant WC SST warming trends through the 20th century, particularly after the 1950s [77,111]. SST-a was projected to increase by a factor of $+1.92$ to $+3.01^{\circ}\text{C}$ by 2071-2100 in contrast to the 1976-2005 climatology [112]. Similar projections were found for the WC and ETP [113].

Observations of increasing atmospheric and ocean temperatures across the WC since at least the 1980s have strongly correlated with the AMO [80,108]. The AMO is a north Atlantic SST signal that influences decadal scale variability in Caribbean precipitation [114] and SST [115]. The AMO signal was more pronounced on temperature during the end of the summer (i.e., JASO) than for other periods of the year [80]. However, recent evidence regarding the role of AMO as a main driver of sea surface warming trends is increasingly ambiguous, exhibiting high spatio-temporal variability and not always correlated to ocean warming [116–121], so it may not necessarily directly affect observed temperature patterns. There is evidence suggesting that AMO oscillations respond to mixed anthropogenic GHG and sulfate aerosols forcing, with long-term modeling evidence pointing out at the influence of volcanic forcing [122]. Observed sea surface warming trends have accelerated during recent decades through a complex combination of mechanisms, raising major concerns for the socio-economic and ecological resiliency of SIDS and of low-lying areas of numerous developing countries across the WC and ETP.

An increase of $0.10^{\circ}\text{C}/\text{decade}$ in global temperatures is considered a critical threshold in climate change due to its significant cumulative impact over time and the associated risks of surpassing internationally agreed-upon temperature limits. While $0.10^{\circ}\text{C}/\text{decade}$ may seem modest, over a century, this rate results in a total increase of 1°C . Such a rise contributes to the intensification of climate-related phenomena, including more frequent and prolonged MHWs, altered precipitation patterns, rapid hurricane intensification, increased SST and OHC, and accelerated SLR. Model projections point out an increasing risk of surpassing regional critical temperature thresholds within only a decade or two. Higher decadal rates of increase will rapidly reach or exceed the 1°C threshold above annual maximum SST. Reaching or exceeding a total increase of 1°C above annual maximum temperatures will result in continuously reaching hot spot temperatures which will lead to mass coral bleaching after 8 weeks and potential mass coral mortalities beyond 12 weeks. These changes can disrupt coastal ecosystems, but also agriculture, food and water security, and human societies.

There is mounting evidence that under current projections, warming trends may lead to runaway climate conditions, which would lead to temperature anomalies of $+2$ to $+5^{\circ}\text{C}$ by 2100 [123–128], with outcomes depending on projected pathways [129]. Other modeling efforts have suggested that temperature anomalies may range from $+1.6$ to $+6.9^{\circ}\text{C}$ by 2100, depending on global emissions pathways, but with likely conditions of reaching $+4^{\circ}\text{C}$ during the early 2060s under business-as-usual scenarios [130]. These changes may lead to important spatio-temporal variations in storm tracks and in storm intensity at both hemispheres [131], to significant spatio-temporal variability in river flows [132], altered fisheries productivity [133–136], declining agriculture [137–139] and to significant soil warming, which may alter temperature/moisture-sensitive soil processes and productivity [140]. In the long-term, this might jeopardize water and food security on regional to global scales, and will lead to enhanced carbon loss from soils, which would further contribute to runaway climates [141–

143]. Furthermore, climate models project substantial changes across the WC, including significant warming and drying trends [144,145], increasing threats by SLR [146], and projected degradation of coastal ecosystems, their ecological functions and productivity [67,147]. Islands may also experience increased impacts on tropical forests by stronger hurricanes [148] and potential runaway climate effects, with some models predicting warming beyond 4.5°C [137]. This might have potential adverse impacts on future regional socio-economic development [149]. México and Central America will also be affected by warming and a decrease in precipitation [150,151] and runoff [152], with adverse impacts to agriculture [153], fisheries and livestock [137], and a decline in tropical forest cover and productivity [154].

Under such projected conditions, despite the global emissions scenario, conditions for the conservation and restoration of coral reefs will be severely compromised as early as the 2030s or 2040s, which warrant the need for rapid action. The 2015 Paris Agreement aims to limit global warming to well below 2°C above pre-industrial levels, with efforts to cap the increase at 1.5°C [1]. Exceeding these thresholds significantly heightens the risk of severe climate impacts, particularly on coral reef ecosystems. Current observations indicate that global temperatures have already risen by approximately 1 to 1.5°C since the late 19th century. This study has shown that continuing at a warming rate of 0.10°C/decade or higher could lead to rapidly surpassing the 1.5°C threshold within the 2040s even under optimistic scenario 4.5, and within the 2030s under business-as-usual scenario 8.5, intensifying the urgency for mitigation and adaptation efforts. A sustained increase of 0.10°C/decade or higher is critical because it contributes to the cumulative rise in global temperatures, rapidly bringing us closer to, and potentially beyond, thresholds that are associated with severe and potentially irreversible, runaway climate conditions. Recognizing and addressing this rate of change is essential for effective climate policy and action.

4.6. SOS for Coral Reefs – The Urgency for Adaptive Coral Conservation and Restoration Strategies

Projected warming trends across the WC and ETP, regardless of the climate scenario, will very likely result in annual mass coral bleaching and mortality, adversely affecting coral conservation and restoration efforts as soon as the 2030s or earlier. This represents an important warning signal to promote the rapid development and implementation of regional and country-specific adaptive coral restoration strategies. Creating an adaptive coral restoration plan for the WC and ETP in the face of projected climate change, sea surface warming, and extreme events requires a multifaceted approach. Critical considerations include the following components.

4.6.1. Select and cultivate climate-resilient coral species. This requires prioritizing the selection and cultivation of coral species that have shown resilience to higher temperatures and acidification [155–157], and that have shown adaptive molecular and physiological survival mechanisms [158–163], enhancing genetic diversity in restoration efforts to increase the adaptive potential of coral populations across regional and sub-regional scales, and improving strategies for site selection [164,165].

4.6.2. Increase restoration efficiency, focusing on scale and cost-effectiveness of deployment. The use of cost-effective approaches is critical to enhance the scale of interventions in developing countries and SIDS, where resources are most likely limited [166–169].

4.6.3. Advanced coral propagation techniques. Micro-fragmentation [170,171], larval rearing [172], methods to enhance coral recruitment, growth and survival [169], genotyping [173], gene banking [174], assisted gene flow and cryopreservation [175,176], and rapid growth techniques are vital to accelerating the growth of slow-growing coral species and implementing larval rearing and settlement techniques to boost natural recruitment processes.

4.6.4. Ensure restoration of endangered coral species proceeds within a population-genetics management context. Develop a population genetics management plan to promote enhanced genetic diversity and disease resistance in future coral restoration efforts [170,177].

4.6.5. Dynamic site selection. Use predictive modeling to identify sites less likely to be affected by future climate impacts (i.e., deeper water refugia, sites under stronger circulation and

oxygenation, sites with remnant populations of rare, threatened or endangered corals with population $\lambda > 1$). Rotation of restoration sites, such as the Reef of Hope initiative [178] should be based on real-time monitoring of environmental conditions (i.e., stronger circulation, cooler temperatures) to maximize coral survival rates [61].

4.6.6. Enhanced monitoring and rapid response. Implement consistent ecological [179], environmental DNA (eDNA) [180], and epigenetics monitoring [181,182], as well as remote sensing technologies [183–185] to track coral health and develop early warning systems to identify stress [162,186,187]. Establish teams for rapid intervention during extreme events to protect and stabilize affected coral populations (i.e., implementing shadowing techniques, rotating coral nurseries to deeper waters or areas of strong circulation, collecting fragments or colonies of highly-depleted species and placed in upland nurseries for future propagation and restoration).

4.6.7. Thermal tolerance and pre-stress conditioning. Expose corals to controlled sub-lethal stress levels to build resilience against future temperature spikes (i.e., through restoration rotation strategies) [188,189]. Explore the potential of assisted evolution techniques to enhance thermal tolerance in corals [190–193].

4.6.8. Support a holistic approach to coral reef ecosystem restoration [194,195]. It is critical to consider the integration of restoring fish assemblages [61] and herbivory [196–198]. The integration of multifaceted cost-effective nature-based strategies to enhance coastal resilience and provide protection against wave energy, runup and coastal erosion is paramount under projected stronger storm swells and SLR [199–202].

4.6.9. Integration of local and traditional knowledge. Involve local communities and stakeholders in restoration efforts, leveraging traditional ecological knowledge and practices, and providing capacity building, training, hands on experiences and resources to local stakeholders to empower them in coral conservation and restoration activities. The integration of community-based solutions has been paramount in achieving restoration success in numerous examples [203–208].

4.6.10. Restoration of adjacent ecosystems. Restore and protect mangrove forests, seagrass beds and other green coastal infrastructure that support coral reef health and provide additional coastal protection, climate change adaptation and mitigation, and socio-economic benefits [200]. Implement holistic/integrated coastal zone management plans that address the health of all coastal ecosystems, not just coral reefs.

4.6.11. Innovation in restoration techniques. Develop artificial and hybrid reef structures that can serve as nurseries and habitat for coral and other marine life, and depending on the material, location, design and configuration, these reef structures can also provide long-term protection against shoreline erosion [200–202,209]. Employ eco-engineering approaches to design reef structures that enhance coral resilience and biodiversity [210–213].

4.6.12. Climate-ready infrastructure. Use natural, artificial and hybrid structures to promote wave energy and runup attenuation and protect reefs from storm damage [191–202,214–218]. Experiment with shading devices and localized cooling systems to protect corals during heatwaves [219,220].

4.6.13. Policy, funding, international cooperation, and governance. Advocate for dedicated funding streams from international climate funds, NGOs, and recurrent governmental sources to support long-term restoration projects. Foster regional cooperation and knowledge exchange among WC and ETP nations to enhance collective restoration efforts. Ensure coral restoration and conservation are integrated into national and regional climate adaptation and biodiversity strategies and policies [221] to foster improved governance and sustainable success [222–224].

4.6.14. Adaptive management and continuous learning. Implement adaptive management frameworks that allow for flexibility and continuous improvement of restoration strategies based on new scientific insights and monitoring data [211,225]. Continuously support research to develop innovative restoration techniques and improve understanding of coral resilience mechanisms. Promote, support and strengthen community-based restoration efforts.

4.6.15. Develop and promote the use of standardized terms and metrics for coral reef restoration. Using uniform strategies tailored to local realities and needs, similar terms and a set of standard operating procedures and metrics is vital to improve outcome comparativeness among projects [169,179].

4.6.16. Support coral reef restoration practitioners working in diverse geographic locations. Supporting projects across WC and ETP developing countries and SIDS requires international cooperation strategies, novel partnerships among regional governments, private sectors, hospitality industry, tourism businesses, creating local cooperatives and other strategic partnerships involving base communities, fishing villages, etc. It will also require recognizing and providing access to colonial states (i.e., Puerto Rico, U.S. Virgin Islands) to international funding sources.

4.6.17. Public awareness and education. Develop public awareness campaigns and outreach programs to highlight the importance of coral reefs in the context of climate change and the need for adapting restoration strategies. Integrate coral reef conservation and restoration into school curricula and community education programs to build a culture of stewardship from a young age.

A regional adaptive coral restoration plan for the WC and ETP must be comprehensive, incorporating resilient species selection, advanced propagation techniques, dynamic site management, enhanced monitoring, rapid response strategies, community-based capacity building, and providing access to recurrent funding sources. Integrating local knowledge, restoring adjacent ecosystems, and fostering regional collaboration are also crucial. Securing funding, building public awareness, and continuously adapting strategies based on new data will enhance the success and sustainability of coral restoration efforts in the face of mounting climate challenges.

4.7. Timely Adaptation and Mitigation Strategies for Developing Nations and Small Island Developing States Across the WC and ETP

Regardless of the climate scenario, projected warming trends across the WC and ETP warrant the rapid development and implementation of climate change adaptation and mitigation strategies to minimize the risk of runaway climate impacts in the future. The following list provides potential alternative solutions for developing nations and SIDS.

4.7.1. Promote national-regional lobbying, activism and advocacy actions for change. These will play crucial roles in advocating for the protection of the environment and promoting sustainable practices, including influencing policy and legislation, raising awareness, encouraging corporate responsibility, supporting renewable energy, agroecology practices, mobilizing communities (grassroots, empowering individuals), protecting and restoring biodiversity, addressing climate change, and water and food security (advocating for action, raising public consciousness), promoting environmental justice (addressing inequities, community support), research and innovation, workforce development, and building global-regional-national networks, such as a Latin American Coral Restoration Network (international collaboration, shared best practices and lessons learned).

4.7.2. Promote sustainable development. A participatory/fair ocean economy is vital to promote integration, equity and a fair inclusion of base communities and local businesses in a participatory ocean economy. A massive reforestation campaign is instrumental to manage microclimates, promote greener cities, improve water catchment, reduce runoff and lead to food and water security, while protecting coastal ecosystems from excessive sediment and nutrient loads and pollution. In addition, greener and sustainable hospitality and food and beverage industries are paramount to promote fresher operations, reduce their carbon footprint, mitigate and compensate environmental impacts, promote sustainable food and seafood consumption, promote consumption of local markets and food sources, reduce waste production, eliminate single-use plastics, and promote and support coastal ecosystem restoration, reforestation, agroecological practices, composting, recycling and other sustainable practices.

4.7.3. Water resource management. Rainwater harvesting practices are becoming increasingly important. Implement systems to capture and store rainwater, reducing dependence on overexploited freshwater sources. Promote reforestation for enhanced aquifer recharge. Explore

renewable energy-powered desalination technologies to provide potable water, especially in arid regions and SIDS.

4.7.4. Integrated waste management systems. Solid waste management is a critical concern across the WC and ETP nations and will require a multifaceted approach. Inappropriate waste disposal is a mounting source of GHG emissions and should be a top priority in the bucket list of recommended actions for climate adaptation and mitigation practices. Encourage practices that minimize waste generation at the source. This includes promoting reusable products, reducing packaging, and implementing policies that discourage single-use plastics. Implement public awareness campaigns to educate residents and businesses about the benefits of reducing waste and how to implement waste-reducing practices. Establish efficient recycling programs that target common recyclable materials such as paper, plastics, glass, and metals. Provide accessible recycling bins and regular collection services, and economic or tax incentives to businesses that achieve meaningful recycling levels. Promote organic waste composting at both household and community levels, provide incentives to the hospitality and food beverage industries that achieve successful composting practices. This can significantly reduce the volume of waste sent to landfills while producing valuable compost for gardening and landscaping. Implement an integrated/holistic waste management system that combines waste reduction, recycling, composting, and safe disposal methods. This approach ensures that all waste types are managed effectively. Engage local communities in the planning and implementation of waste management strategies, and through the creation of community-owned waste management businesses and cooperatives to ensure they meet the specific needs and capacities of local communities.

4.7.5. Ecosystem-based adaptation. Invest in restoring mangroves, wetlands, sand dunes, seagrasses and coral reefs, which provide natural barriers against storm surges and erosion while supporting biodiversity. Promote sustainable agricultural practices that integrate trees with crops and livestock, enhancing resilience to climate-related shocks, and promoting water and food security. Implement widespread reforestation programs to promote greener cities, enhance runoff control, manage microclimates, enhance carbon sequestration and further promote water and food security.

4.7.6. Restore shallow coral reef resilience. Promote coral gardening and transplantation to restore resilient coral species to degraded areas. This can include using coral nurseries to grow coral in controlled conditions before reintroducing them to the wild (i.e., promoting restoration rotation strategies). Focus on enhancing genetic diversity in coral populations to increase resilience to diseases and sea surface warming. Establish long-term monitoring programs to assess changes in coral health, biodiversity, the impact of invasive species, and changing environmental conditions. These data can inform innovative regional management strategies. Invest in research to understand the mechanisms of coral resilience and identify species and genotypes that are more adaptable to changing conditions, as well as quantify coral demographic performance under variable environmental conditions.

4.7.7. Sustainable fisheries management. Implement community-based, local management practices that involve communities in decision-making, helping to ensure sustainable fish stocks and livelihoods [61]. Establish marine protected areas (MPAs) under participatory, cooperative management models to protect critical habitats and replenish fish populations, supporting both ecological health and local economies. Integrate coral, seagrass and mangrove restoration strategies with community-based sustainable fisheries management to enhance coastal essential fish habitats.

4.7.8. Sustainable tourism. Non-sustainable massive tourism across the WC and ETP is critical for the regional economy, but can generate significant adverse impacts to natural resources [226]. Sustainable tourism practices are essential for preserving the natural and cultural heritage of the WC and ETP while promoting economic development. Promote tourism development practices that eliminate the displacement and gentrification of base communities. Support eco-certification programs for tourism operators that meet environmental and sustainability standards and equity. Encourage activities that emphasize the natural environment; while ensuring they are conducted responsibly to minimize impacts and reduce carbon footprints. Involving local communities in tourism planning and decision-making and management to ensure their needs and perspectives are

considered. Promote cultural tourism that highlights local traditions, crafts, and folklore, providing economic benefits to communities while preserving cultural identity. Encourage green building practices, such as green roofs, rain gardens, and the use of sustainable materials and energy-efficient designs in new tourism developments, including hotels and resorts. Invest in public transportation options and promote non-motorized transport (bicycles, walking) when feasible to reduce carbon footprints. Implement recycling initiatives in tourist areas to minimize waste generation and promote responsible waste disposal practices. Encourage businesses to reduce single-use plastics by providing alternatives and promoting reusable products. Support the establishment and effective management of MPAs and coastal natural reserves to conserve biodiversity and support sustainable fishing and tourism practices. Implement measures to protect local wildlife and habitats, including regulations against poaching and habitat destruction, and by supporting local breeding of rare and endangered species, including corals, and by restoring degraded habitats. Develop educational programs for tourists that promote awareness of local ecosystems, cultural heritage, and sustainable practices. Provide education, capacity-building and hands-on training for tourism operators and local businesses on sustainable practices, conservation and restoration efforts, and customer engagement. Provide incentives and encourage tourists to visit during off-peak seasons to reduce overcrowding and pressure on local resources. Market green-certified, eco-friendly accommodations, businesses and activities that prioritize local food consumption and sustainability. Develop and monitor sustainability indicators to assess the environmental, social, and economic impacts of tourism. Implement systems for feedback from tourists and local communities to inform ongoing improvements in sustainable tourism practices. Foster regional collaboration among WC and ETP nations to share best practices, resources, and strategies for sustainable tourism. Encourage partnerships between government, NGOs, and the private sector to develop and promote local, subregional and regional standardized sustainable tourism initiatives. Develop strategies to enhance the climate resilience of tourism infrastructure and ecosystems, ensuring long-term sustainability in the face of climate change. Implement disaster preparedness plans for tourism operators to ensure safety and minimize impact during extreme weather events.

4.7.9. Renewable energy development. Invest in solar and wind energy, and in hydroelectric if viable river sources are available, leveraging abundant natural resources to reduce reliance on fossil fuels and enhance energy security. Develop decentralized energy systems (i.e., microgrids) to provide reliable electricity to remote communities, reducing vulnerability to climate impacts. Avoid costly/irreversible socio-economic and environmental tradeoffs (i.e., impacts on critical agricultural lands, protected areas, watersheds, etc.), maximizing the use of existing built infrastructure.

4.7.10. Climate resilient infrastructure. Promote the use of sustainable materials and designs that enhance resilience to climate impacts, such as elevated structures in flood-prone areas, green roofs, rain gardens, passive cooling structures, etc. Upgrade and maintain transportation infrastructure to withstand extreme weather events, ensuring access to essential services.

4.7.11. Disaster risk reduction. Invest in technology and training for early warning systems to prepare communities for extreme weather events, minimizing loss of life and property. Foster community engagement in disaster planning, response strategies and preparedness programs, ensuring local knowledge is incorporated.

4.7.12. Education and capacity building of different societal sectors. Develop educational, hands-on training initiatives focused on sustainable practices, climate adaptation, and resilience-building within communities. Promote local research and development initiatives to innovate climate adaptation and mitigation strategies tailored to specific regional, subregional and local challenges.

By adopting many of these recommendations, the WC and ETP can promote and enhance socio-economic and environmental sustainability and promote critical conservation and restoration practices on their coastal ecosystems, which should enhance social-ecological resilience to projected climate changes.

5. Conclusions

MHWs are increasingly affecting tropical seas, causing mass coral bleaching and mortality across the WC and ETP. This leads to significant coral loss, reduced biodiversity, and impaired ecological functions. Climate models forecast a troubling future for Latin American coral reefs. From 1940 to 2023, JASO air and ocean temperature anomalies showed significant decadal increases. Projections indicate that even under optimistic scenario 4.5, temperatures may exceed the +1.5°C air threshold beyond pre-industrial levels by the 2040s and the +1.0°C ocean threshold beyond historical annual maximums by the 2030s, resulting in severe coral bleaching and mortality. Business-as-usual scenario 8.5 suggests SST-a will become intolerable for coral conservation and restoration by 2030, with decadal warming trends largely surpassing historical rates, leading to unbearable conditions for corals. Given these alarming trends, immediate and coordinated climate action across the WC and ETP is critical. This includes not only the development of regional, sub-regional, and local adaptive coral reef conservation and restoration plans that should incorporate novel strategies but also robust multifaceted climate change adaptation and mitigation strategies to minimize losing socio-economic and ecological resilience of human coastal communities. By prioritizing climate action now, we can create a more resilient marine environment and human coastal communities, providing the necessary time for optimistic scenarios to materialize and ensure the survival of these vital ecosystems for future generations.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org, Table S1. Summary of study locations by region, with coordinates of sampling polygons and the linear regression results between normalized JASO 2m temperature and sea surface temperature (SST) anomalies for the period of 1940 to 2023 on each location, using the 1951-1980 climatology as baseline. Figure S1. Projection of 2m temperature and SST anomalies across the Mesoamerican Barrier System (MSA), western Caribbean, into the year 2100 under scenarios 4.5 and 8.5: (a) México, (b) Belize/Guatemala; (c) Honduras, (d) Nicaragua. Figure S2. Projection of 2m temperature and SST anomalies across the Caribbean Central America into the year 2100 under scenarios 4.5 and 8.5: (a) Costa Rica, (b) Panamá; (c) San Andrés-Providencia, Colombia. Figure S3. Projection of 2m temperature and SST anomalies across the southern Caribbean into the year 2100 under scenarios 4.5 and 8.5: (a) Colombia, (b) Venezuela; (c) ABC Islands (Aruba, Bonaire, Curaçao). Figure S4. Projection of 2m temperature and SST anomalies across the eastern Caribbean into the year 2100 under scenarios 4.5 and 8.5: (a) Leeward Islands, (b) Windward Islands; (c) U.S./British Virgin Islands (UBV). Figure S5. Projection of 2m temperature and SST anomalies across the northern Caribbean into the year 2100 under scenarios 4.5 and 8.5: (a) Puerto Rico, (b) Dominican Republic, (c) Haiti, (d) Jamaica. Figure S6. Projection of 2m temperature and SST anomalies across the northwestern Caribbean into the year 2100 under scenarios 4.5 and 8.5: (a) Grand Cayman, (b) Cuba, (c) Turks & Caicos, (d) Bahamas, (e) Florida. Figure S7. Projection of 2m temperature and SST anomalies across the eastern Pacific into the year 2100 under scenarios 4.5 and 8.5: (a) Northern México, (b) Southern México, (c) Guatemala, (d) El Salvador/Honduras. Figure S8. Projection of 2m temperature and SST anomalies across the eastern Pacific into the year 2100 under scenarios 4.5 and 8.5: (a) Nicaragua, (b) Costa Rica, (c) Panamá, (d) Colombia.

Author Contributions: Conceptualization, E.A.H.D. and Y.M.R.G.; methodology, E.A.H.D.; validation, E.A.H.D.; formal analysis, E.A.H.D.; investigation, E.A.H.D. and Y.M.R.G.; resources, E.A.H.D. and Y.M.R.G.; data curation, E.A.H.D.; writing—original draft preparation, E.A.H.D. and Y.M.R.G.; writing—review and editing, E.A.H.D. and Y.M.R.G.; visualization, E.A.H.D.; supervision, E.A.H.D.; project administration, E.A.H.D. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data supporting reported results can be found at the Supplementary Materials section in "Hernandez&Rodriguez_data.xlsx" file.

Acknowledgments: This publication was the first peer-reviewed contribution of Sociedad Ambiente Marino's Climate Science Division. The study was fully conducted under in kind contributions by both authors.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. IPCC. Climate change 2021: The physical science basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. **2021**, Cambridge University Press, Cambridge. www.ipcc.ch/report/ar6/wg1/
2. Hobday, A.J.; Alexander, L.V.; Perkins, S.E.; Smale, D.A.; Straub, S.C.; Oliver, E.C.; Benthuyssen, J.A.; Burrows, M.T.; Donat, M.G.; Feng, M.; Holbrook, N.J. A hierarchical approach to defining marine heatwaves. *Progr. Oceanogr.* **2016**, *141*, 227-338.
3. Dutheil, C.; Lal, S.; Lengaigne, M.; Cravatte, S.; Menkès, C.; Receveur, A.; Börgel, F.; Gröger, M.; Houlbreque, F.; Le Gendre, R.; Mangolte, I. The massive 2016 marine heatwave in the Southwest Pacific: An "El Niño–Madden-Julian Oscillation" compound event. *Sci. Adv.* **2024**, *10*(41), eadp2948.
4. Hoegh-Guldberg, O.; Skirving, W.; Dove, S.G.; Spady, B.L.; Norrie, A.; Geiger, E.F.; Liu, G.; De La Cour, J.L.; Manzello, D.P. Coral reefs in peril in a record-breaking year. *Science*. **2023**, *382*(6676), 1238-1240.
5. Goessling, H.F.; Rackow, T.; Jung, T. Recent global temperature surge intensified by record-low planetary albedo. *Science*. **2025**, *387*(6729), 68-73.
6. Mejias-Rivera, C.L.; Courtney, T.A. Ocean warming, heat stress, and coral bleaching in Puerto Rico. *Caribb. J. Sci.* **2024**, *54*(1), 132-149.
7. Lachs, L.; Bozec, Y.M.; Bythell, J.C.; Donner, S.D.; East, H.K.; Edwards, A.J.; Golbuu, Y.; Gouezo, M.; Guest, J.R.; Humanes, A.; Riginos, C. Natural selection could determine whether *Acropora* corals persist under expected climate change. *Science*. **2024**, *386*(6727), 1289-1294, eadl6480.
8. Baum, J.K.; Claar, D.C.; Tietjen, K.L.; Magel, J.M.; Maucieri, D.G.; Cobb, K.M.; McDevitt-Irwin, J.M. Transformation of coral communities subjected to an unprecedented heatwave is modulated by local disturbance. *Science Adv.* **2023**, *9*(14), eabq5615.
9. Wang, S.; Foster, A.; Lenz, E.A.; Kessler, J.D.; Stroeve, J.C.; Anderson, L.O.; Turetsky, M.; Betts, R.; Zou, S.; Liu, W.; Boos, W.R. Mechanisms and impacts of Earth system tipping elements. *Rev. Geophys.* **2023**, *61*(1), e2021RG000757.
10. Donovan, M.K.; Burkepile, D.E.; Kratochwill, C.; Shlesinger, T.; Sully, S.; Oliver, T.A.; Hodgson, G.; Freiwald, J.; van Woesik, R. Local conditions magnify coral loss after marine heatwaves. *Science*. **2021**, *372*(6545), 977-980.
11. Wiedenmann, J.; D'Angelo, C.; Smith, E.G.; Hunt, A.N.; Legiret, F.E.; Postle, A.D.; Achterberg, E.P. Nutrient enrichment can increase the susceptibility of reef corals to bleaching. *Nature Clim. Change*. **2013**, *3*(2), 160-164.
12. Vega Thurber, R.L.; Burkepile, D.E.; Fuchs, C.; Shantz, A.A.; McMinds, R.; Zaneveld, J.R. Chronic nutrient enrichment increases prevalence and severity of coral disease and bleaching. *Global Change Biol.* **2014**, *20*(2), 544-554.
13. Burkepile, D.E.; Shantz, A.A.; Adam, T.C.; Munsterman, K.S.; Speare, K.E.; Ladd, M.C.; Rice, M.M.; Ezzat, L.; McIlroy, S.; Wong, J.C.; Baker, D.M. Nitrogen identity drives differential impacts of nutrients on coral bleaching and mortality. *Ecosystems*. **2020**, *23*, 798-811.
14. Fernandes de Barros Marangoni, L.; Ferrier-Pagès, C.; Rottier, C.; Bianchini, A.; Grover, R. Unravelling the different causes of nitrate and ammonium effects on coral bleaching. *Sci. Rept.* **2020**, *10*(1), 11975.
15. Lesser, M.P. Eutrophication on coral reefs: what is the evidence for phase shifts, nutrient limitation and coral bleaching. *BioScience*. **2021**, *71*(12), 1216-1233.
16. Starko, S.; Fifer, J.E.; Claar, D.C.; Davies, S.W.; Cunning, R.; Baker, A.C.; Baum, J.K. Marine heatwaves threaten cryptic coral diversity and erode associations among coevolving partners. *Sci. Adv.* **2023**, *9*(32), eadf0954.

17. Romanou, A.; Hegerl, G.; Seneviratne, S.; Abis, B.; Bastos, A.; Conversi, A.; Landolfi, A.; Kim, H.; Lerner, P.; Mekus, J.; Otto-Bliesner, B. Extreme events contributing to tipping elements and tipping points. *Surv. Geophys.* **2024**, 1-46. <https://doi.org/10.1007/s10712-024-09863-7>
18. Wunderling, N.; Donges, J.F.; Kurths, J.; Winkelmann, R. Interacting tipping elements increase risk of climate domino effects under global warming. *Earth Syst. Dynam. Disc.* **2021**, 12, 1-21. <https://doi.org/10.5194/esd-12-601-2021>
19. Wunderling, N.; Winkelmann, R.; Rockström, J.; Loriani, S.; Armstrong McKay, D.I.; Ritchie, P.D.; Sakschewski, B.; Donges, J.F. Global warming overshoots increase risks of climate tipping cascades in a network model. *Nature Clim. Change.* **2023**, 13(1): 75-82.
20. Wunderling, N.; von der Heydt, A.S.; Aksenov, Y.; Barker, S.; Bastiaansen, R.; Brovkin, V.; Brunetti, M.; Couplet, V.; Kleinen, T.; Lear, C.H.; Lohmann, J. Climate tipping point interactions and cascades: a review. *Earth Syst. Dynam.* **2024**, 15(1), 41-74.
21. Braun, C.D.; Lezama-Ochoa, N.; Farchadi, N.; Arostegui, M.C.; Alexander, M.; Allyn, A.; Bograd, S.J.; Brodie, S.; Crear, D.P.; Curtis, T.H.; Hazen, E.L. Widespread habitat loss and redistribution of marine top predators in a changing ocean. *Science Adv.* **2023**, 9(32), eadi2718.
22. Laufkötter, C.; Zscheischler, J.; Frölicher, T.L. High-impact marine heatwaves attributable to human-induced global warming. *Science.* **2020**, 369(6511), 1621-1625.
23. Dalton, S.J.; Carroll, A.G.; Sampayo, E.; Roff, G.; Harrison, P.L.; Entwistle, K.; Huang, Z.; Salih, A.; Diamond, S.L. Successive marine heatwaves cause disproportionate coral bleaching during a fast phase transition from El Niño to La Niña. *Sci. Total Env.* **2020**, 715, 136951.
24. Asner, G.P.; Vaughn, N.R.; Martin, R.E.; Foo, S.A.; Heckler, J.; Neilson, B.J.; Gove, J.M. Mapped coral mortality and refugia in an archipelago-scale marine heat wave. *Proc. Natl. Acad. Sci. USA*, **2022**, 119(19), e2123331119.
25. Shlesinger, T.; van Woesik, R. Oceanic differences in coral-bleaching responses to marine heatwaves. *Sci. Total Env.* **2023**, 871, 162113.
26. Goreau, T.J.; Hayes, R.L. 2023 Record marine heat waves: coral reef bleaching HotSpot maps reveal global sea surface temperature extremes, coral mortality, and ocean circulation changes. *Oxford Open Clim. Change.* **2024**, 4(1), kgae005.
27. Godinot, C.; Houlbrèque, F.; Grover, R.; Ferrier-Pagès, C. Coral uptake of inorganic phosphorus and nitrogen negatively affected by simultaneous changes in temperature and pH. *PLoS One.* **2011**, 6(9), e25024.
28. Hughes, T.P.; Rodrigues, M.J.; Bellwood, D.R.; Ceccarelli, D.; Hoegh-Guldberg, O.; McCook, L.; Moltschanivskyj, N.; Pratchett, M.S.; Steneck, R.S.; Willis, B. Phase shifts, herbivory, and the resilience of coral reefs to climate change. *Current Biol.* **2007**, 17(4), 360-365.
29. Leggat, W.P.; Camp, E.F.; Suggett, D.J.; Heron, S.F.; Fordyce, A.J.; Gardner, S.; Deakin, L.; Turner, M.; Beeching, L.J.; Kuzhiumparambil, U.; Eakin, C.M. Rapid coral decay is associated with marine heatwave mortality events on reefs. *Curr. Biol.* **2019**, 29(16), 2723-2730.
30. Anderson, J.J.; Gurarie, E.; Bracis, C.; Burke, B.J.; Laidre, K.L. Modeling climate change impacts on phenology and population dynamics of migratory marine species. *Ecol. Mod.* **2013**, 264, 83-97.
31. Alvarez-Filip, L.; Estrada-Saldívar, N.; Pérez-Cervantes, E.; Molina-Hernández, A.; González-Barrios, F.J. A rapid spread of the stony coral tissue loss disease outbreak in the Mexican Caribbean. *PeerJ.* **2019**, 7, e8069.
32. Bernal-Sotelo, K.; Acosta, A.; Cortés, J. Decadal change in the population of *Dendrogyra cylindrus* (Scleractinia: Meandrinidae) in old providence and St. Catalina Islands, Colombian Caribbean. *Front. Mar. Sci.* **2019**, 5, 513.
33. Chan, A.N.; Lewis, C.L.; Neely, K.L.; Baums, I.B. Fallen pillars: The past, present, and future population dynamics of a rare, specialist coral-algal symbiosis. *Front. Mar. Sci.* **2019**, 6, 218.
34. Estrada-Saldívar, N.; Molina-Hernández, A.; Pérez-Cervantes, E.; Medellín-Maldonado, F.; González-Barrios, F.J.; Alvarez-Filip, L. Reef-scale impacts of the stony coral tissue loss disease outbreak. *Coral Reefs.* **2020**, 39, 861-866.

35. Estrada-Saldívar, N.; Quiroga-García, B.A.; Pérez-Cervantes, E.; Rivera-Garibay, O.O.; Alvarez-Filip, L. Effects of the stony coral tissue loss disease outbreak on coral communities and the benthic composition of Cozumel reefs. *Front. Mar. Sci.* **2021**, *8*, 632777.
36. Jones, N.P.; Kabay, L.; Semon Lunz, K.; Gilliam, D.S. Temperature stress and disease drives the extirpation of the threatened pillar coral, *Dendrogyra cylindrus*, in southeast Florida. *Sci. Rept.* **2021**, *11*(1), 14113.
37. Cavada-Blanco, F.; Croquer, A.; Vermeij, M.; Goergen, L.; Rodriguez, R. *Dendrogyra cylindrus*: extinction risk assessment. *The Red List of Threatened Species.* **2022**, Dec 22.
38. Alvarez-Filip, L.; González-Barrios, F.J.; Pérez-Cervantes, E.; Molina-Hernández, A.; Estrada-Saldívar, N. Stony coral tissue loss disease decimated Caribbean coral populations and reshaped reef functionality. *Comm. Biol.* **2022**, *5*(1), 440.
39. Mellin, C.; Brown, S.; Cantin, N.; Klein-Salas, E.; Mouillot, D.; Heron, S.F.; Fordham, D.A. Cumulative risk of future bleaching for the world's coral reefs. *Sci. Adv.* **2024**, *10*(26), eadn9660.
40. Hays, G.C.; Richardson, A.J.; Robinson, C. Climate change and marine plankton. *Trends Ecol. Evol.* **2005**, *20*(6), 337-344.
41. Pesce, M.; Critto, A.; Torresan, S.; Giubilato, E.; Santini, M.; Zirino, A.; Ouyang, W.; Marcomini, A. Modelling climate change impacts on nutrients and primary production in coastal waters. *Sci. Total Env.* **2018**, *628*, 919-937.
42. Raven, J.A.; Beardall, J. Influence of global environmental change on plankton. *J. Plankton Res.* **2021**, *43*(6), 779-800.
43. Möllmann, C.; Müller-Karulis, B.; Kornilovs, G.; St. John, M.A. Effects of climate and overfishing on zooplankton dynamics and ecosystem structure: regime shifts, trophic cascade, and feedback loops in a simple ecosystem. *ICES J. Mar. Sci.* **2008**, *65*(3), 302-310.
44. Kirby, R.R.; Beaugrand, G.; Lindley, J.A. Synergistic effects of climate and fishing in a marine ecosystem. *Ecosystems.* **2009**, *12*, 48-61.
45. Pratchett, M.S.; Munday, P.L.; Graham, N.A.; Kronen, M.; Pinca, S.; Friedman, K.; Brewer, T.D.; Bell, J.D.; Wilson, S.K.; Cinner, J.E.; Kinch, J.P. Vulnerability of coastal fisheries in the tropical Pacific to climate change. In: Bell, J.D.; Johnson, J.E.; Hobday, A.J. (eds) *Vulnerability of Tropical Pacific Fisheries and Aquaculture to Climate Change. Secretariat of the Pacific Community, Noumea, New Caledonia.* **2011**, Chapter 9, 167-185.
46. Hayashida, H.; Matear, R.J.; Strutton, P.G. Background nutrient concentration determines phytoplankton bloom response to marine heatwaves. *Global Change Biol.* **2020**, *26*(9), 4800-4811.
47. Wyatt, A.M.; Resplandy, L.; Marchetti, A. Ecosystem impacts of marine heat waves in the Northeast Pacific. *Biogeosciences.* **2022**, *19*(24), 5689-5705.
48. Kauppi, L.; Göbeler, N.; Norkko, J.; Norkko, A.; Romero-Ramirez, A.; Bernard, G. Changes in macrofauna bioturbation during repeated heatwaves mediate changes in biogeochemical cycling of nutrients. *Front. Mar. Sci.* **2023**, *9*, 1070377.
49. Serrano, O.; Arias-Ortiz, A.; Duarte, C.M.; Kendrick, G.A.; Lavery, P.S. Impact of marine heatwaves on seagrass ecosystems. In: *Ecosystem Collapse and Climate Change*, **2021**, 345-364. Cham: Springer International Publishing.
50. Alvarez-Filip, L.; Dulvy, N.K.; Gill, J.A.; Côté, I.M.; Watkinson, A.R. Flattening of Caribbean coral reefs: region-wide declines in architectural complexity. *Proc. Royal Soc. B: Biol. Sci.* **2009**, *276*(1669), 3019-3025.
51. Alvarez-Filip, L.; Paddack, M.J.; Collen, B.; Robertson, D.R.; Côté, I.M. Simplification of Caribbean reef-fish assemblages over decades of coral reef degradation. *PLoS One.* **2015**, *10*(4), e0126004.
52. Hernández-Delgado, E.A.; Alejandro-Camis, P.; Cabrera-Beauchamp, G.; Fonseca-Miranda, J.S.; Gómez-Andújar, N.X.; Gómez, P.; Guzmán-Rodríguez, R.; Olivo-Maldonado, I.; Suleimán-Ramos, S.E. Stronger hurricanes and climate change in the Caribbean Sea: Threats to the sustainability of endangered coral species. *Sustainability.* **2024**, *16*(4), 1506.
53. Allison, E.H.; Perry, A.L.; Badjeck, M.C.; Adger, W.N.; Brown, K.; Conway, D.; Halls, A.S.; Pilling, G.M.; Reynolds, J.D.; Andrew, N.L.; Dulvy, N.K. Vulnerability of national economies to the impacts of climate change on fisheries. *Fish Fisher.* **2009**, *10*(2), 173-196.

54. Burge, C.A.; Eakin, C.M.; Friedman, C.S.; Froelich, B.; Hersherberger, P.K.; Hofmann, E.E.; Petes, L.E.; Prager, K.C.; Weil, E.; Willis, B.L.; Ford, S.E. Climate change influences on marine infectious diseases: implications for management and society. *Ann. Rev. Mar. Sci.* **2014**, *6*(1), 249-277.
55. Tracy, A.M.; Pielmeier, M.L.; Yoshioka, R.M.; Heron, S.F.; Harvell, C.D. Increases and decreases in marine disease reports in an era of global change. *Proc. Royal Soc. B.* **2019**, *286*(1912), 20191718.
56. Byers, J.E. Marine parasites and disease in the era of global climate change. *Ann. Rev. Mar. Sci.* **2021**, *13*(1), 397-420.
57. Armstrong McKay, D.I.; Staal, A.; Abrams, J.F.; Winkelmann, R.; Sakschewski, B.; Loriani, S.; Fetzer, I.; Cornell, S.E.; Rockström, J.; Lenton, T.M. Exceeding 1.5°C global warming could trigger multiple climate tipping points. *Science*. **2022**, *377*(6611), eabn7950.
58. Lenton, T.M.; Held, H.; Kriegler, E.; Hall, J.W.; Lucht, W.; Rahmstorf, S.; Schellnhuber, H.J. Tipping elements in the Earth's climate system. *Proc. Natl. Acad. Sci. USA* **2008**, *105*(6), 1786-93.
59. Lenton, T.M. Environmental tipping points. *Ann. Rev. Environ. Res.* **2013**, *38*(1), 1-29.
60. Goldblatt, C.; Watson, A.J. The runaway greenhouse: implications for future climate change, geoengineering and planetary atmospheres. *Phil. Trans. Royal Soc. A: Math., Phys. Eng. Sci.* **2012**, *370*(1974), 4197-4216.
61. Hernández-Delgado, E.A.; Laureano, R. Bringing back the fish: Sustainable success of community-based restoration of Elkhorn coral (*Acropora palmata*) in Vega Baja, Puerto Rico (2008-2023). *Sustainability*. **2024**, *16*, 5985.
62. Dietz, S.; Rising, J.; Stoerk, T.; Wagner, G. Economic impacts of tipping points in the climate system. *Proc. Natl. Acad. Sci. USA*. **2021**, *118*(34): e2103081118.
63. Kopp, R.E.; Hayhoe, K.; Easterling, D.R.; Hall, T.; Horton, R.; Kunkel, K.E.; LeGrande, A.N. Potential surprises—Compound extremes and tipping elements. Climate Science Special Report: Fourth National Climate Assessment, D.J. Wuebbles et al., Eds., Vol. I, US Global Change Research Program, **2017**, 411-429.
64. Smith, K.E.; Burrows, M.T.; Hobday, A.J.; Sen Gupta, A.; Moore, P.J.; Thomsen, M.; Wernberg, T.; Smale, D.A. Socioeconomic impacts of marine heatwaves: Global issues and opportunities. *Science*. **2021**, *374*(6566), eabj3593.
65. Emanuel, K.A. Downscaling CMIP5 climate models shows increased tropical cyclone activity over the 21st century. *Proc. Natl. Acad. Sci. USA*, **2013**, *110*(30), 12219-12224, doi:10.1073/pnas.1301293110.
66. Knutson, T.R.; Sirutis, J.J.; Vecchi, G.A.; Garner, S.; Zhao, M.; Kim, H.-S.; Bender, M.; Tuleya, R.E.; Held, I.M.; Villarini, G. Dynamical downscaling projections of twenty-first-century Atlantic Hurricane activity: CMIP3 and CMIP5 model-based scenarios, *J. Climate*. **2013**, *26*(17), 6591-6617, doi:10.1175/JCLI-D-12-00539.1.
67. Hernández-Delgado, E.A. The emerging threats of climate change on tropical coastal ecosystem services, public health, local economies and livelihood sustainability of small islands: Cumulative impacts and synergies. *Mar. Poll. Bull.* **2015**, *101*(1), 5-28.
68. Hersbach, H.; Bell, B.; Berrisford, P.; Biavati, G.; Horányi, A.; Muñoz Sabater, J.; Nicolas, J.; Peubey, C.; Radu, R.; Rozum, I.; Schepers, D.; Simmons, A.; Soci, C.; Dee, D.; Thépaut, J.-N. ERA5 monthly averaged data on single levels from 1940 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS), **2023**, 10.24381/cds.f17050d7 (Accessed on 01-10-2024).
69. Eyring, V.; Bony, S.; Meehl, G.A.; Senior, C.A.; Stevens, B.; Stouffer, R.J.; Taylor, K.E. Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geosci. Model Dev.* **2016**, *9*(5), 1937-1958.
70. Anderson, M.J.; Gorley, R.N.; Clarke, K.R. PERMANOVA+ for PRIMER: Guide to Software and Statistical Methods; Massey University: Auckland, New Zealand; PRIMER-e Ltd.: Plymouth, UK, **2008**.
71. Clarke, K.R.; Gorley, R.N.; Somerfield, P.J.; Warwick, R.M. Change in Marine Communities: An Approach to Statistical Analysis and Interpretation, 3rd ed.; PRIMER-E: Plymouth, UK, **2014**.
72. Philander, S.G.; Rasmusson, E.M. The southern oscillation and El Niño. *Adv. Geophys.* **1985**, *28*, 197-215.
73. Vega, M.J.; Alvarez-Silva, O.; Restrepo, J.C.; Ortiz, J.C.; Otero, L.J. Interannual variability of wave climate in the Caribbean Sea. *Ocean Dynam.* **2020**, *70*, 965-976.

74. Tartaglione, C.A.; Smith, S.R.; O'Brien, J.J. ENSO impact on hurricane landfall probabilities for the Caribbean. *J. Clim.* **2003**, *16*(17), 2925-2931.
75. Klotzbach, P.J. The influence of El Niño–Southern Oscillation and the Atlantic multidecadal oscillation on Caribbean tropical cyclone activity. *J. Clim.* **2011**, *24*(3), 721-731.
76. Hurrell, J.W.; Deser, C. North Atlantic climate variability: the role of the North Atlantic Oscillation. *J. Mar. Syst.* **2010**, *79*(3-4), 231-244.
77. Hetzinger, S.; Pfeiffer, M.; Dullo, W.C.; Garbe-Schönberg, D.; Halfar, J. Rapid 20th century warming in the Caribbean and impact of remote forcing on climate in the northern tropical Atlantic as recorded in a Guadeloupe coral. *Palaeogeog., Palaeoclimatol., Palaeoecol.* **2010**, *296*(1-2), 111-124.
78. Jury, M.; Malmgren, B.A.; Winter, A. Subregional precipitation climate of the Caribbean and relationships with ENSO and NAO. *J. Geophys. Res.: Atmospheres.* **2007**, *112*(D16), D16107.
79. Moron, V.; Gouirand, I.; Taylor, M. Weather types across the Caribbean basin and their relationship with rainfall and sea surface temperature. *Clim. Dynam.* **2016**, *47*, 601-21.
80. Stephenson, T.S.; Vincent, L.A.; Allen, T.; Van Meerbeeck, C.J.; McLean, N.; Peterson, T.C.; Taylor, M.A.; Aaron-Morrison, A.P.; Auguste, T.; Bernard, D.; Boekhoudt, J.R.I.; Blenman, R.C.; Braithwaite, G.C.; Brown, G.; Butler, M.; Cumberbatch, C.J.M.; Etienne-Leblanc, S.; Lake, D.E.; Martin, D.E.; McDonald, J.L.; Ozoria Zaruela, M.; Porter, A.O.; Santana Ramirez, M.; Tamar, G.A.; Roberts, B.A.; Sallons Mitro, S.; Shaw, A.; Spence, J.M.; Winter, A.; Trotman, A.R. Changes in extreme temperature and precipitation in the Caribbean region, 1961–2010. *Int. J. Climatol.* **2014**, *34*, 2957–2971.
81. Simonti, A.L.; Eastman, J.R. 2005 Caribbean mass coral bleaching event: A sea surface temperature empirical orthogonal teleconnection analysis. *J. Geophys. Res.: Oceans.* **2010**, *115*(C11), C11009.
82. Vimont, D.J.; Kossin, J.P. The Atlantic meridional mode and hurricane activity. *Geophys. Res. Lett.* **2007**, *34*(7), L07709.
83. Lizcano-Sandoval, L.D.; Marulanda-Gómez, Á.; López-Victoria, M.; Rodríguez-Ramírez, A. Climate change and Atlantic Multidecadal Oscillation as drivers of recent declines in coral growth rates in the Southwestern Caribbean. *Front. Mar. Sci.* **2019**, *6*, 38.
84. Jury, M.R. MJO influence in the Caribbean. *Theor. Appl. Climatol.* **2020**, *139*(3), 1559-1567.
85. Mantua, N.J.; Hare, S.R. The Pacific decadal oscillation. *J. Oceanogr.* **2002**, *58*, 35-44.
86. Nidheesh, A.G.; Lengaigne, M.; Vialard, J.; Izumo, T.; Unnikrishnan, A.S.; Cassou, C. Influence of ENSO on the Pacific decadal oscillation in CMIP models. *Clim. Dynam.* **2017**, *49*, 3309-3326.
87. Pierrehumbert, R.T. Thermostats, radiator fins, and the local runaway greenhouse. *J. Atm. Sci.* **1995**, *52*(10), 1784-1806.
88. Wang, C. Variability of the Caribbean low-level jet and its relations to climate. *Climate Dynam.* **2007**, *29*, 411-422.
89. Wang, C.; Lee, S.K. Atlantic warm pool, Caribbean low-level jet, and their potential impact on Atlantic hurricanes. *Geophys. Res. Lett.* **2007**, *34*(2), L02703.
90. Li, H.; Misra, V. Thirty-two-year ocean–atmosphere coupled downscaling of global reanalysis over the Intra-American Seas. *Climate Dynam.* **2014**, *43*, 2471-2489.
91. Montoya-Sánchez, R.A.; Devis-Morales, A.; Bernal, G.; Poveda, G. Seasonal and interannual variability of the mixed layer heat budget in the Caribbean Sea. *J. Mar. Syst.* **2018**, *187*, 111-127.
92. Rajasree, V.P.; Cao, X.; Ramsay, H.; Ocasio, K.M.; Kilroy, G.; Alvey, III, G.R.; Chang, M.; Nam, C.C.; Fudeyasu, H.; Teng, H.F.; Yu, H. Tropical cyclogenesis: Controlling factors and physical mechanisms. *Trop. Cyclone Res. Rev.* **2023**, *12*(3), 165-181.
93. Jones, P.D.; Harpham, C.; Harris, I.; Goodess, C.M.; Burton, A.; Centella-Artola, A.; Taylor, M.A.; Bezanilla-Morlot, A.; Campbell, J.D.; Stephenson, T.S.; Joslyn, O. Long-term trends in precipitation and temperature across the Caribbean. *Int. J. Climatol.* **2016**, *36*(9), 3314-3333.
94. Zhao, L.; Xu, J.; Powell, A.M.; Jiang, Z. Uncertainties of the global-to-regional temperature and precipitation simulations in CMIP5 models for past and future 100 years. *Theor. Appl. Climatol.* **2015**, *122*, 259-270.
95. Zhou, C.; Wang, K. Land surface temperature over global deserts: Means, variability, and trends. *J. Geophys. Res.: Atmospheres.* **2016**, *121*(24), 14344-14357.

96. Sun, X.; Ren, G.; Xu, W.; Li, Q.; Ren, Y. Global land-surface air temperature change based on the new CMA GLSAT data set. *Sci. Bull.* **2017**, *62*(4), 236-238.
97. Sun, C.; Jiang, Z.; Li, W.; Hou, Q.; Li, L. Changes in extreme temperature over China when global warming stabilized at 1.5°C and 2.0°C. *Sci. Rep.* **2019**, *9*, 14982.
98. Sun, X.; Ren, G.; Ren, Y.; Lin, W.; Zhang, P.; Zhang, S.; Xue, X. Asian climate warming since 1901: observation and simulation. *Clim. Res.* **2023**, *91*, 67-82.
99. Duan, J.; Li, L.; Chen, L.; Zhang, H. Time-dependent warming amplification over the Tibetan Plateau during the past few decades. *Atm. Sci. Lett.* **2020**, *21*(10), e998.
100. Nita, I.A.; Sfîcă, L.; Voiculescu, M.; Birsan, M.V.; Micheu, M.M. Changes in the global mean air temperature over land since 1980. *Atm. Res.* **2022**, *279*, 106392.
101. Ren, G.; Zhan, Y.; Ren, Y.; Wen, K.; Zhang, Y.; Sun, X.; Zhang, P.; Zheng, X.; Qin, Y.; Zhang, S.; He, J. Observed changes in temperature and precipitation over Asia, 1901-2020. *Clim. Res.* **2023**, *90*, 31-43.
102. Gautam, R.; Pathak, B.; Bhuyan, P.K.; Borgohain, A.; Kundu, S.S. Long-term trend analysis of surface temperature over North-East India and adjoining regions based on CRU and ERA5 reanalysis. *J. Earth Syst. Sci.* **2024**, *133*(3), 141.
103. Li, Q.; Dong, W.; Li, W.; Gao, X.; Jones, P.; Kennedy, J.; Parker, D. Assessment of the uncertainties in temperature change in China during the last century. *Chinese Sci. Bull.* **2010**, *55*, 1974-1982.
104. Yosef, Y.; Aguilar, E.; Alpert, P. Detecting and adjusting artificial biases of long-term temperature records in Israel. *Int. J. Climatol.* **2018**, *38*(8), 3273-3289.
105. You, Q.; Min, J.; Fraedrich, K.; Zhang, W.; Kang, S.; Zhang, L.; Meng, X. Projected trends in mean, maximum, and minimum surface temperature in China from simulations. *Global Planet Change.* **2014**, *112*, 53-63.
106. Hu, G.; Zhao, L.; Wu, T.; Wu, X.; Park, H.; Li, R.; Zhu, X.; Ni, J.; Zou, D.; Hao, J.; Li, W. Continued warming of the permafrost regions over the Northern Hemisphere under future climate change. *Earth's Future.* **2022**, *10*(9), e2022EF002835.
107. Bove, C.B.; Mudge, L.; Bruno, J.F. A century of warming on Caribbean reefs. *PloS Climate.* **2022**, *1*(3), e0000002.
108. Antuña-Marrero, J.C.; Otterå, O.H.; Robock, A.; Mesquita, M.D. Modelled and observed sea surface temperature trends for the Caribbean and Antilles. *Int. J. Climatol.* **2016**, *36*(4), 1873-1886.
109. Shi, J.; Hu, C. South Florida estuaries are warming faster than global oceans. *Env. Res. Lett.* **2022**, *18*(1), 014003.
110. Shears, N.T.; Bowen, M.M. Half a century of coastal temperature records reveal complex warming trends in western boundary currents. *Sci. Rept.* **2017**, *7*(1), 1-9.
111. Alpert, A.E.; Cohen, A.L.; Oppo, D.W.; DeCarlo, T.M.; Gaetani, G.A.; Hernández-Delgado, E.A.; Winter, A.; Gonneea, M.E. Twentieth century warming of the tropical Atlantic captured by Sr-U paleothermometry. *Paleoceanography.* **2017**, *32*(2), 146-160.
112. Bustos Usta, D.F.; Torres Parra, R.R. Ocean and atmosphere changes in the Caribbean Sea during the twenty-first century using CMIP5 models. *Ocean Dynam.* **2021**, *71*(6), 757-777.
113. Karmalkar, A.V.; Taylor, M.A.; Campbell, J.; Stephenson, T.; New, M.; Centella, A.; Benzanilla, A.; Charlery, J. A review of observed and projected changes in climate for the islands in the Caribbean. *Atmósfera.* **2013**, *26*(2), 283-309.
114. Enfield, D.B.; Mestas-Núñez, A.M.; Trimble, P.J. The Atlantic multidecadal oscillation and its relation to rainfall and river flows in the continental US. *Geophys. Res. Lett.* **2001**, *28*(10), 2077-2080.
115. Alexander, M.A.; Kilbourne, K.H.; Nye, J.A. Climate variability during warm and cold phases of the Atlantic Multidecadal Oscillation (AMO) 1871–2008. *J. Mar. Syst.* **2014**, *133*, 14-26.
116. Delworth, T.L.; Zhang, R.; Mann, M.E. Decadal to centennial variability of the Atlantic from observations and models. *Ocean Circ.: Mech. Impacts* **2007**, *173*, 131-148.
117. Kavvada, A.; Ruiz-Barradas, A.; Nigam, S. AMO's structure and climate footprint in observations and IPCC AR5 climate simulations. *Clim. Dynam.* **2013**, *41*, 1345-1364.
118. Mann, M.E.; Steinman, B.A.; Miller, S.K. On forced temperature changes, internal variability, and the AMO. *Geophys. Res. Lett.* **2014**, *41*(9), 3211-3219.

119. Steinman, B.A.; Mann, M.E.; Miller, S.K. Atlantic and Pacific multidecadal oscillations and Northern Hemisphere temperatures. *Science*. **2015**, *347*(6225), 988-991.
120. Mann, M.E.; Steinman, B.A.; Miller, S.K. Absence of internal multidecadal and interdecadal oscillations in climate model simulations. *Nat. Comm.* **2020**, *11*(1), 49.
121. Zhang, W.; Mei, X.; Geng, X.; Turner, A.G.; Jin, F.F. A nonstationary ENSO–NAO relationship due to AMO modulation. *J. Clim.* **2019**, *32*(1), 33-43.
122. Mann, M.E.; Steinman, B.A.; Brouillette, D.J.; Miller, S.K. Multidecadal climate oscillations during the past millennium driven by volcanic forcing. *Science*. **2021**, *371*(6533), 1014-1019.
123. Houghton, J. The science of global warming. *Int. Sci. Rev.* **2001**, *26*(4), 247-257.
124. Sanderson, M.G.; Hemming, D.L.; Betts, R.A. Regional temperature and precipitation changes under high-end ($\geq 4^{\circ}\text{C}$) global warming. *Phil. Trans. Royal Soc. A: Math., Phys. Eng. Sci.* **2011**, *369*(1934), 85-98.
125. Donnelly, C.; Greuell, W.; Andersson, J.; Gerten, D.; Pisacane, G.; Roudier, P.; Ludwig, F. Impacts of climate change on European hydrology at 1.5, 2 and 3 degrees mean global warming above preindustrial level. *Clim. Change*. **2017**, *143*, 13-26.
126. Raftery, A.E.; Zimmer, A.; Frierson, D.M.; Startz, R.; Liu, P. Less than 2°C warming by 2100 unlikely. *Nature Clim. Change*. **2017**, *7*(9), 637-641.
127. Wang, X.; Jiang, D.; Lang, X. Climate change of 4°C global warming above pre-industrial levels. *Adv. Atm. Sci.* **2018**, *35*, 757-770.
128. Pielke, Jr. R.; Burgess, M.G.; Ritchie, J. Plausible 2005-2050 emissions scenarios project between 2 and 3 degrees C of warming by 2100. *Env. Res. Lett.* **2022**, *17*, 024027.
129. Barring, L.; Strandberg, G. Does the projected pathway to global warming targets matter? *Env. Res. Lett.* **2018**, *13*(2), 024029.
130. Betts, R.A.; Collins, M.; Hemming, D.L.; Jones, C.D.; Lowe, J.A.; Sanderson, M.G. When could global warming reach 4°C ? *Phil. Trans. Royal Soc. A: Math., Phys. Eng. Sci.* **2011**, *369*(1934), 67-84.
131. Chang, E.K.; Guo, Y.; Xia, X. CMIP5 multimodel ensemble projection of storm track change under global warming. *J. Geophys. Res.: Atmospheres*. **2012**, *117*(D23), D23118.
132. Arnell, N.W.; Gosling, S.N. The impacts of climate change on river flow regimes at the global scale. *J. Hydrol.* **2013**, *486*, 351-364.
133. Allison, E.H.; Perry, A.L.; Badjeck, M.C.; Adger, N.; Brown, K.; Conway, D.; Halls, A.S.; Pilling, G.M.; Reynolds, J.D.; Andrew, N.L.; Dulvy, N.K. Vulnerability of national economies to the impacts of climate change on fisheries. *Fish Fisheries*. **2009**, *10*(2), 173-196.
134. Brander, K. Impacts of climate change on fisheries. *J. Mar. Syst.* **2010**, *79*(3-4), 389-402.
135. Hollowed, A.B.; Barange, M.; Beamish, R.J.; Brander, K.; Cochrane, K.; Drinkwater, K.; Foreman, M.G.; Hare, J.A.; Holt, J.; Ito, S.I.; Kim, S. Projected impacts of climate change on marine fish and fisheries. *ICES J. Mar. Sci.* **2013**, *70*(5), 1023-1037.
136. Erauskin-Extramiana, M.; Arrizabalaga, H.; Hobday, A.J.; Cabré, A.; Ibaibarriaga, L.; Arregui, I.; Murua, H.; Chust, G. Large-scale distribution of tuna species in a warming ocean. *Global Change Biol.* **2019**, *25*(6), 2043-2060.
137. Reyer, C.P.; Adams, S.; Albrecht, T.; Baarsch, F.; Boit, A.; Canales Trujillo, N.; Carlsburg, M.; Coumou, D.; Eden, A.; Fernandes, E.; Langerwisch, F. et al. Climate change impacts in Latin America and the Caribbean and their implications for development. *Reg. Environ. Change*. **2017**, *17*, 1601-1621.
138. Prager, S.D.; Ríos, A.R.; Schiek, B.; Almeida, J.S.; González, C.E. Climate change vulnerability and economic impacts in the agricultural sector in Latin America and the Caribbean. IDB Technical Note IDB-TN-01915. **2020**, Inter-American Development Bank (IDB); International Center for Tropical Agriculture (CIAT). Cali, Colombia.
139. Lachaud, M.A.; Bravo-Ureta, B.E.; Ludena, C.E. Economic effects of climate change on agricultural production and productivity in Latin America and the Caribbean (LAC). *Agricult. Econ.* **2022**, *53*(2), 321-332.
140. Soong, J.L.; Phillips, C.L.; Ledna, C.; Koven, C.D.; Torn, M.S. CMIP5 models predict rapid and deep soil warming over the 21st century. *J. Geophys. Res.: Biogeosciences*. **2020**, *125*(2), e2019JG005266.

141. Brien, R.J.; Phillips, O.L.; Feldpausch, T.R.; Gloor, E.; Baker, T.R.; Lloyd, J.; López-González, G.; Monteagudo-Mendoza, A.; Malhi, Y.; Lewis, S.L.; Vázquez Martínez, R. Long-term decline of the Amazon carbon sink. *Nature*. **2015**, *519*(7543), 344-348.
142. Crowther, T.W.; Todd-Brown, K.E.; Rowe, C.W.; Wieder, W.R.; Carey, J.C.; Machmuller, M.B.; Snoek, B.L.; Fang, S.; Zhou, G.; Allison, S.D.; Blair, J.M. et al. Quantifying global soil carbon losses in response to warming. *Nature*. **2016**, *540*(7631), 104-108.
143. Qin, Y.; Xiao, X.; Wigneron, J.P.; Ciais, P.; Brandt, M.; Fan, L.; Li, X.; Crowell, S.; Wu, X.; Doughty, R.; Zhang, Y. et al. Carbon loss from forest degradation exceeds that from deforestation in the Brazilian Amazon. *Nature Clim. Change*. **2021**, *11*(5), 442-448.
144. Neelin, J.D.; Münnich, M.; Su, H.; Meyerson, J.E.; Holloway, C.E. Tropical drying trends in global warming models and observations. *Proc. Natl. Acad. Sci. USA*. **2006**, *103*(16), 6110-6115.
145. Cashman, A.; Nurse, L.; John, C. Climate change in the Caribbean: the water management implications. *The J. Env. Dev.* **2010**, *19*(1), 42-67.
146. Taylor, M.A.; Stephenson, T.S.; Chen, A.A.; Stephenson, K.A. Climate change and the Caribbean: Review and response. *Caribb. Studies*. **2012**, *40*(2), 169-200.
147. Hernández-Delgado, E.A. Climate change impacts on Caribbean coastal ecosystems: Emergent ecological and environmental geography challenges. In: *Routledge Handbook of Latin America and the Environment*, **2023**, 36-50. Routledge.
148. Lugo, A.E. Effects and outcomes of Caribbean hurricanes in a climate change scenario. *Sci. Total Env.* **2000**, *262*(3), 243-251.
149. Bishop, M.L.; Payne, A. Climate change and the future of Caribbean development. *The J. Dev. Studies*. **2012**, *48*(10), 1536-1553.
150. Karmalkar, A.V.; Bradley, R.S.; Díaz, H.F. Climate change in Central America and Mexico: regional climate model validation and climate change projections. *Clim. Dynam.* **2011**, *37*, 605-629.
151. Imbach, P.; Chou, S.C.; Lyra, A.; Rodrigues, D.; Rodriguez, D.; Latinovic, D.; Siqueira, G.; Silva, A.; Garofolo, L.; Georgiou, S. Future climate change scenarios in Central America at high spatial resolution. *PLoS One*. **2018**, *13*(4), e0193570.
152. Hidalgo León, H.G.; Amador Astúa, J.A.; Alfaro Martínez, E.J.; Quesada, B. Hydrological climate change projections for Central America. *J. Hydrol.* **2013**, *495*, 94-112.
153. Harvey, C.A.; Saborio-Rodríguez, M.; Martínez-Rodríguez, M.R.; Viguera, B.; Chain-Guadarrama, A.; Vignola, R.; Alpizar, F. Climate change impacts and adaptation among smallholder farmers in Central America. *Agric. Food Sec.* **2018**, *7*(1), 1-20.
154. Lyra, A.; Imbach, P.; Rodríguez, D.; Chou, S.C.; Georgiou, S.; Garofolo, L. Projections of climate change impacts on Central America tropical rainforest. *Clim. Change*. **2017**, *141*, 93-105.
155. Caruso, C.; Hughes, K.; Drury, C. Selecting heat-tolerant corals for proactive reef restoration. *Front. Mar. Sci.* **2021**, *8*, 632027.
156. Quigley, K.M. Breeding and selecting corals resilient to global warming. *Ann. Rev. Anim. Biosci.* **2024**, *12*(1), 209-332.
157. Humanes, A.; Lachs, L.; Beauchamp, E.; Bukorou, L.; Buzzoni, D.; Bythell, J.; Craggs, J.R.; de la Torre Cerro, R.; Edwards, A.J.; Golbuu, Y.; Martínez, H.M. Selective breeding enhances coral heat tolerance to marine heatwaves. *Nature Comm.* **2024**, *15*(1), 8703.
158. Palumbi, S.R.; Barshis, D.J.; Traylor-Knowles, N.; Bay, R.A. Mechanisms of reef coral resistance to future climate change. *Science*. **2014**, *344*(6186), 895-898.
159. Smith, E.G.; Hazzouri, K.M.; Choi, J.Y.; Delaney, P.; Al-Kharafi, M.; Howells, E.J.; Aranda, M.; Burt, J.A. Signatures of selection underpinning rapid coral adaptation to the world's warmest reefs. *Sci. Adv.* **2022**, *8*(2), eabl7287.
160. Lachs, L.; Donner, S.D.; Mumby, P.J.; Bythell, J.C.; Humanes, A.; East, H.K.; Guest, J.R. Emergent increase in coral thermal tolerance reduces mass bleaching under climate change. *Nature Comm.* **2023**, *14*(1), 4939.
161. Selmoni, O.; Bay, L.K.; Exposito-Alonso, M.; Cleves, P.A. Finding genes and pathways that underlie coral adaptation. *Trends Gen.* **2024**, *40*(3), 213-227.

162. Nunn, B.L.; Brown, T.; Timmins-Schiffman, E.; Mudge, M.C.; Riffle, M.; Axworthy, J.B.; Dilworth, J.; Kenkel, C.D.; Zaneveld, J.; Rodrigues, L.J.; Padilla-Gamiño, J.L. Protein signatures predict coral resilience and survival to thermal bleaching events. *Comm. Earth Environ.* **2025**, *6*(1), 191.
163. Huang, W.; Chen, J.; Yang, E.; Meng, L.; Feng, Y.; Chen, Y.; Huang, Z.; Tan, R.; Xiao, Z.; Zhou, Y.; Xu, M. Heat-tolerant subtropical *Porites lutea* may be better adapted to future climate change than tropical one in the South China Sea. *Sci. Total Environ.* **2025**, *962*, 178381.
164. Baums, I.B.; Baker, A.C.; Davies, S.W.; Grottoli, A.G.; Kenkel, C.D.; Kitchen, S.A.; Kuffner, I.B.; LaJeunesse, T.C.; Matz, M.V.; Miller, M.W.; Parkinson, J.E. Considerations for maximizing the adaptive potential of restored coral populations in the western Atlantic. *Ecol. Appl.* **2019**, *29*(8), e01978.
165. Shaver, E.C.; McLeod, E.; Hein, M.Y.; Palumbi, S.R.; Quigley, K.; Vardi, T.; Mumby, P.J.; Smith, D.; Montoya-Maya, P.; Muller, E.M.; Banaszak, A.T. A roadmap to integrating resilience into the practice of coral reef restoration. *Global Change Biol.* **2022**, *28*(16), 4751-4764.
166. Montoya Maya, P.H.; Smit, K.P.; Burt, A.J.; Frias-Torres, S. Large-scale coral reef restoration could assist natural recovery in Seychelles, Indian Ocean. *Nature Conserv.* **2016**, *16*, 1-7.
167. Doropoulos, C.; Vons, F.; Elzinga, J.; Ter Hofstede, R.; Salee, K.; Van Koningsveld, M.; Babcock, R.C. Testing industrial-scale coral restoration techniques: harvesting and culturing wild coral-spawn slicks. *Front. Mar. Sci.* **2019**, *6*, 658.
168. McLeod, I.M.; Hein, M.Y.; Babcock, R.; Bay, L.; Bourne, D.G.; Cook, N.; Doropoulos, C.; Gibbs, M.; Harrison, P.; Lockie, S.; van Oppen, M.J. Coral restoration and adaptation in Australia: The first five years. *Plos One.* **2022**, *17*(11), e0273325.
169. Vardi, T.; Hoot, W.C.; Levy, J.; Shaver, E.; Winters, R.S.; Banaszak, A.T.; Baums, I.B.; Chamberland, V.F.; Cook, N.; Gulko, D.; Hein, M.Y. et al. Six priorities to advance the science and practice of coral reef restoration worldwide. *Rest. Ecology.* **2021**, *29*(8), e13498.
170. Page, C.A.; Muller, E.M.; Vaughan, D.E. Microfragmenting for the successful restoration of slow growing massive corals. *Ecol. Eng.* **2018**, *123*, 86-94.
171. Tortolero-Langarica, J.A.; Rodríguez-Troncoso, A.P.; Cupul-Magaña, A.L.; Rinkevich, B. Micro-fragmentation as an effective and applied tool to restore remote reefs in the Eastern Tropical Pacific. *Int. J. Env. Res. Public Health.* **2020**, *17*(18), 6574.
172. Banaszak, A.T.; Marhaver, K.L.; Miller, M.W.; Hartmann, A.C.; Albright, R.; Hagedorn, M.; Harrison, P.L.; Latijnhouwers, K.R.; Mendoza Quiroz, S.; Pizarro, V.; Chamberland, V.F. Applying coral breeding to reef restoration: best practices, knowledge gaps, and priority actions in a rapidly-evolving field. *Rest. Ecol.* **2023**, *31*(7), e13913.
173. Blanco-Pimentel, M.; Kenkel, C.D.; Kitchen, S.A.; Calle-Triviño, J.; Baums, I.B.; Cortés-Useche, C.; Morikawa, M.K. Overcoming barriers to reef restoration: field-based method for approximate genotyping of *Acropora cervicornis*. *Rest. Ecol.* **2024**, *32*(3), e14073.
174. Hobbs, R.J.; O'Brien, J.K.; Bay, L.K.; Severati, A.; Spindler, R.; Henley, E.M.; Quigley, K.M.; Randall, C.J.; van Oppen, M.J.; Carter, V.; Zuchowicz, N. A decade of coral biobanking science in Australia-transitioning into applied reef restoration. *Front. Mar. Sci.* **2022**, *9*, 960470.
175. Daly, J.; Zuchowicz, N.; Nuñez Lendo, C.I.; Khosla, K.; Lager, C.; Henley, E.M.; Bischof, J.; Kleinhans, F.W.; Lin, C.; Peters, E.C.; Hagedorn, M. Successful cryopreservation of coral larvae using vitrification and laser warming. *Sci. Rept.* **2018**, *8*(1), 15714.
176. Hagedorn, M.; Page, C.A.; O'Neil, K.L.; Flores, D.M.; Tichy, L.; Conn, T.; Chamberland, V.F.; Lager, C.; Zuchowicz, N.; Lohr, K.; Blackburn, H. Assisted gene flow using cryopreserved sperm in critically endangered coral. *Proc. Natl. Acad. Sci. USA.* **2021**, *118*(38), e2110559118.
177. Brown, A.L.; Anastasiou, D.E.; Schul, M.; MacVittie, S.; Spiers, L.J.; Meyer, J.L.; Manfrino, C.; Frazer, T.K. Mixtures of genotypes increase disease resistance in a coral nursery. *Sci. Rept.* **2022**, *12*(1), 19286.
178. Bowden-Kerby, A. Coral-focused climate change adaptation and restoration based on accelerating natural processes: Launching the "Reefs of Hope" paradigm. *Oceans* **2022**, *4*(1), 13-26.
179. Goergen, E.A.; Schopmeyer, S.; Moulding, A.L.; Moura, A.; Kramer, P.; Viehman, T.S. Coral reef restoration monitoring guide: Methods to evaluate restoration success from local to ecosystem scales. NOAA Technical Memorandum NOS NCCOS 279, **2020**, Silver Spring, MD. 145 pp. doi: 10.25923/xndz-h538

180. Alexander, J.B.; Bunce, M.; White, N.; Wilkinson, S.P.; Adam, A.A.; Berry, T.; Stat, M.; Thomas, L.; Newman, S.J.; Dugal, L.; Richards, Z.T. Development of a multi-assay approach for monitoring coral diversity using eDNA metabarcoding. *Coral Reefs*. **2020**, *39*, 159-171.
181. Rodríguez-Casariago, J.A.; Mercado-Molina, A.E.; García-Souto, D.; Ortiz-Rivera, I.M.; Lopes, C.; Baums, I.B.; Sabat, A.M.; Eirin-López, J.M. Genome-wide DNA methylation analysis reveals a conserved epigenetic response to seasonal environmental variation in the staghorn coral *Acropora cervicornis*. *Front. Mar. Sci.* **2020**, *7*, 560424.
182. Eirin-López, J.M.; Putnam, H. Marine environmental epigenetics. *Front. Mar. Sci.* **2021**, *8*, 685075.
183. Xu, J.; Zhao, D. Review of coral reef ecosystem remote sensing. *Acta Ecologica Sinica*. **2014**, *34*(1): 19-25.
184. Hedley, J.D.; Roelfsema, C.M.; Chollett, I.; Harborne, A.R.; Heron, S.F.; Weeks, S.; Skirving, W.J.; Strong, A.E.; Eakin, C.M.; Christensen, T.R.; Ticzon, V. Remote sensing of coral reefs for monitoring and management: a review. *Remote Sensing*. **2016**, *8*(2), 118.
185. Foo, S.A.; Asner, G.P. Scaling up coral reef restoration using remote sensing technology. *Front. Mar. Sci.* **2019**, *6*, 79.
186. Eakin, C.M.; Lough, J.M.; Heron, S.F. Climate variability and change: monitoring data and evidence for increased coral bleaching stress. In: *Coral Bleaching: Patterns, Processes, Causes and Consequences*, **2009** (pp. 41-67). Berlin, Heidelberg: Springer Berlin Heidelberg.
187. Flower, J.; Ortiz, J.C.; Chollett, I.; Abdullah, S.; Castro-Sanguino, C.; Hock, K.; Lam, V.; Mumby, P.J. Interpreting coral reef monitoring data: A guide for improved management decisions. *Ecol. Indicators*. **2017**, *72*, 848-869.
188. Bellantuono, A.J.; Granados-Cifuentes, C.; Miller, D.J.; Hoegh-Guldberg, O.; Rodríguez-Lanetty, M. Coral thermal tolerance: tuning gene expression to resist thermal stress. *PloS One*. **2012**, *7*(11), e50685.
189. Page, C.E.; Leggat, W.; Heron, S.F.; Fordyce, A.J.; Ainsworth, T.D. High flow conditions mediate damaging impacts of sub-lethal thermal stress on corals' endosymbiotic algae. *Conserv. Physiol.* **2021**, *9*(1), coab046.
190. van Oppen, M.J.; Oliver, J.K.; Putnam, H.M.; Gates, R.D. Building coral reef resilience through assisted evolution. *Proc. Natl. Acad. Sci. USA*. **2015**, *112*(8), 2307-2313.
191. van Oppen, M.J.; Oakeshott, J.G. A breakthrough in understanding the molecular basis of coral heat tolerance. *Proc. Natl. Acad. Sci. USA*. **2020**, *117*(46), 28546-28548.
192. Humanes, A.; Beauchamp, E.A.; Bythell, J.C.; Carl, M.K.; Craggs, J.R.; Edwards, A.J.; Golbuu, Y.; Lachs, L.; Martínez, H.M.; Palmowski, P.; Paysinger, F. An experimental framework for selectively breeding corals for assisted evolution. *Front. Mar. Sci.* **2021**, *8*, 669995.
193. DeFilippo, L.B.; McManus, L.C.; Schindler, D.E.; Pinsky, M.L.; Colton, M.A.; Fox, H.E.; Tekwa, E.W.; Palumbi, S.R.; Essington, T.E.; Webster, M.M. Assessing the potential for demographic restoration and assisted evolution to build climate resilience in coral reefs. *Ecol. Appl.* **2022**, *32*(7), e2650.
194. Boström-Einarsson, L.; Babcock, R.C.; Bayraktarov, E.; Ceccarelli, D.; Cook, N.; Ferse, S.C.; Hancock, B.; Harrison, P.; Hein, M.; Shaver, E.; Smith, A. Coral restoration—A systematic review of current methods, successes, failures and future directions. *PloS One*. **2020**, *15*(1), e0226631.
195. Watt-Pringle, R.; Razak, T.B.; Jompa, J.; Ambo-Rappe, R.; Kostaman, A.N.; Smith, D.J. Coral reef restoration in Indonesia: lessons learnt from the world's largest coral restoration nation. *Biodiv. Conserv.* **2024**, *33*(10), 2675-707.
196. Cano, I.; Sellares-Blasco, R.I.; Lefcheck, J.S.; Villalpando, M.F.; Croquer, A. Effects of herbivory by the urchin *Diadema antillarum* on early restoration success of the coral *Acropora cervicornis* in the central Caribbean. *J. Exp. Mar. Biol. Ecol.* **2021**, *539*, 151541.
197. Williams, S.M. The reduction of harmful algae on Caribbean coral reefs through the reintroduction of a keystone herbivore, the long-spined sea urchin *Diadema antillarum*. *Rest. Ecol.* **2022**, *30*(1), e13475.
198. Butler, M.J. IV; Durán, A.; Feehan, C.J.; Harborne, A.R.; Hykema, A.; Patterson, J.T.; Sharp, W.C.; Spadaro, A.J.; Wijers, T.; Williams, S.M. Restoration of herbivory on Caribbean coral reefs: are fishes, urchins, or crabs the solution? *Front. Mar. Sci.* **2024**, *11*, 1329028.

199. Viehman, T.S.; Reguero, B.G.; Lenihan, H.S.; Rosman, J.H.; Storlazzi, C.D.; Goergen, E.A.; Canals Silander, M.F.; Groves, S.H.; Holstein, D.M.; Bruckner, A.W.; Carrick, J.V. et al. Coral restoration for coastal resilience: Integrating ecology, hydrodynamics, and engineering at multiple scales. *Ecosphere*. **2023**, *14*(5), e4517.
200. Hernández-Delgado, E.A. Coastal restoration challenges and strategies for small island developing states in the face of sea level rise and climate change. *Coasts*. **2024**, *4*(2), 235-286.
201. Storlazzi, C.D.; Reguero, B.G.; Alkins, K.C.; Shope, J.B.; Gaido-Lassarre, C.; Viehman, T.S.; Beck, M.W. Hybrid coral reef restoration can be a cost-effective nature-based solution to provide protection to vulnerable coastal populations. *Sci. Adv.* **2025**, *11*(3), eadn4004.
202. Familkhalili, R.; Storlazzi, C.D.; Nemeth, M.; Viehman, S. Assessing the effect of coral reef restoration location on coastal flood hazard along the San Juan Coastline, Puerto Rico. *Front. Mar. Sci.* **2025**, *12*, 1528460.
203. dela Cruz, D.W.; Rinkevich, B.; Gomez, E.D.; Yap, H.T. Assessing an abridged nursery phase for slow growing corals used in coral restoration. *Ecol. Eng.* **2015**, *84*, 408-415.
204. Frey, J.B.; Berkes, F. Can partnerships and community-based conservation reverse the decline of coral reef social-ecological systems? *Int. J. Commons*. **2014**, *8*(1), 26-46.
205. Hernández-Delgado, E.A.; Mercado-Molina, A.E.; & Suleimán-Ramos, S.E. Multi-disciplinary lessons learned from low-tech coral farming and reef rehabilitation practices. I. Best management practices. 213-243. In: Duque-Beltrán, C.; & E. Tello-Camacho (Eds.), *Corals in a Changing World*, **2018**, InTech Publ.
206. Hernández-Delgado, E.A.; Mercado-Molina, A.E.; Suleimán-Ramos, S.E.; Lucking, M.A. Multi-disciplinary lessons learned from low-tech coral farming and reef rehabilitation practices. II. Coral demography and social-ecological benefits. 245-268. In: Duque-Beltrán, C.; & E. Tello-Camacho (Eds.), *Corals in a Changing World*, **2018**, InTech Publ.
207. Hein, M.Y.; Birtles, A.; Willis, B.L.; Gardiner, N.; Beeden, R.; Marshall, N.A. Coral restoration: Socio-ecological perspectives of benefits and limitations. *Biol. Conserv.* **2019**, *229*, 14-25.
208. Bayraktarov, E.; Banaszak, A.T.; Montoya Maya, P.; Kleypas, J.; Arias-González, J.E.; Blanco, M.; Calle-Triviño, J.; Charuvi, N.; Cortés-Useche, C.; Galván, V.; García Salgado, M.A. et al. Coral reef restoration efforts in Latin American countries and territories. *PLoS One*. **2020**, *15*(8), e0228477.
209. Norris, B.K.; Storlazzi, C.D.; Pomeroy, A.W.; Reguero, B.G. Optimizing infragravity wave attenuation to improve coral reef restoration design for coastal defense. *J. Mar. Sci. Eng.* **2024**, *12*(5), 768.
210. Rinkevich, B. Climate change and active reef restoration—ways of constructing the “reefs of tomorrow”. *J. Mar. Sci. Eng.* **2015**, *3*(1), 111-127.
211. Rinkevich, B. The active reef restoration toolbox is a vehicle for coral resilience and adaptation in a changing world. *J. Mar. Sci. Eng.* **2019**, *7*(7), 201.
212. Rinkevich, B. Ecological engineering approaches in coral reef restoration. *ICES J. Mar. Sci.* **2021**, *78*(1), 410-420.
213. Kim, T.; Baek, S.; Kwon, Y.; Lee, J.; Cha, S.M.; Kwon, S. Improved coastal erosion prevention using a hybrid method with an artificial coral reef: large-scale 3D hydraulic experiment. *Water*. **2020**, *12*(10), 2801.
214. Roelvink, F.E.; Storlazzi, C.D.; Van Dongeren, A.R.; Pearson, S.G. Coral reef restorations can be optimized to reduce coastal flooding hazards. *Front. Mar. Sci.* **2021**, *8*, 653945.
215. Brathwaite, A.; Clua, E.; Roach, R.; Pascal, N. Coral reef restoration for coastal protection: Crafting technical and financial solutions. *J. Env. Mgmt.* **2022**, *310*, 114718.
216. Geldard, J.; Lowe, R.; Draper, S.; Ghisalberti, M.; Westera, S.; Ellwood, G.; Cuttler, M.; Smith, D.; McArdle, A. Effectiveness of coral reef restoration in wave attenuation applications. *Coast. Eng. Proc.* **2022**, *37*, 90-91.
217. Norris, B.K.; Storlazzi, C.D.; Pomeroy, A.W.; Rosenberger, K.J.; Logan, J.B.; Cheriton, O.M. Combining field observations and high-resolution numerical modeling to demonstrate the effect of coral reef roughness on turbulence and its implications for reef restoration design. *Coast. Eng.* **2023**, *184*, 104331.
218. Norris, B.K.; Reguero, B.; Beck, M.; Bartolai, J.; Yukish, M.A.; Rhode-Barbarigos, L.; Haus, B.K.; Barajas, G.; Maza, M.; Lara, J.L. Designing modular, artificial reefs for both coastal defense and coral restoration. In, *Artificial Reefs for Both Coastal Defense and Coral Restoration*, **2024**.
219. Tagliafico, A.; Baker, P.; Kelaher, B.; Ellis, S.; Harrison, D. The effects of shade and light on corals in the context of coral bleaching and shading technologies. *Front. Mar. Sci.* **2022**, *9*, 919382.

220. Butcherine, P.; Tagliafico, A.; Ellis, S.L.; Kelaher, B.P.; Hendrickson, C.; Harrison, D. Intermittent shading can moderate coral bleaching on shallow reefs. *Front. Mar. Sci.* **2023**, *10*, 1162896.
221. Razak, T.B.; Boström-Einarsson, L.; Alisa, C.A.; Vida, R.T.; Lamont, T.A. Coral reef restoration in Indonesia: A review of policies and projects. *Mar. Pol.* **2022**, *137*, 104940.
222. Morrison, T.H.; Adger, N.; Barnett, J.; Brown, K.; Possingham, H.; Hughes, T. Advancing coral reef governance into the Anthropocene. *One Earth.* **2020**, *2*(1), 64-74.
223. Westoby, R.; Becken, S.; Laria, A.P. Perspectives on the human dimensions of coral restoration. *Reg. Env. Change.* **2020**, *20*(4), 109.
224. Shumway, N.; Foster, R.; Fidelman, P. The governance of marine and coral reef restoration, lessons and paths forward for novel interventions. *Env. Sci. Pol.* **2025**, *164*, 103999.
225. Rinkevich, B. Novel tradable instruments in the conservation of coral reefs, based on the coral gardening concept for reef restoration. *J. Env. Mgmt.* **2015**, *162*, 199-205.
226. Hernández-Delgado, E.A.; Ramos-Scharrón, C.E.; Guerrero, C.; Lucking, M.A.; Laureano, R.; Méndez-Lázaro, P.A.; Meléndez-Díaz, J.O. Long-term impacts of tourism and urban development in tropical coastal habitats in a changing climate: Lessons learned from Puerto Rico. 357-398. In, M. Kasimoglu (ed.), *Visions from Global Tourism Industry-Creating and Sustaining Competitive Strategies*. **2012**, Intech Publications, Rikeja, Croatia.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.