

---

# Comparative Study of the Effects of Carvacrol and P-Cymene on the Motor Activity of Rats and Movement of *Caenorhabditis elegans*

---

[Oliver Stošić](#) , [Dragana Medić](#) , [Djordje S Marjanović](#) , [Tihomir Marić](#) , Veljko Savić ,  
[Jelena Nedeljković Trailović](#) , [Nemanja Zdravković](#) , [Saša M Trailović](#) \*

Posted Date: 28 February 2026

doi: 10.20944/preprints202602.1911.v1

Keywords: carvacrol; p-cymene; rats; *Caenorhabditis elegans*



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

# Comparative Study of the Effects of Carvacrol and P-Cymene on the Motor Activity of Rats and Movement of *Caenorhabditis elegans*

Oliver Stošić<sup>1</sup>, Dragana Medić<sup>2</sup>, Djordje S. Marjanović<sup>2</sup>, Tihomir Marić<sup>2</sup>, Veljko Savić<sup>1</sup>, Jelena Nedeljković Trailović<sup>3</sup>, Nemanja Zdravković<sup>4</sup> and Saša M. Trailović<sup>2,\*</sup>

<sup>1</sup> Department of Experimental Surgery and Laboratory Animal Breeding, Veterinary Service Center, Military Health Department, Ministry of Defense, Serbia

<sup>2</sup> Faculty of Veterinary Medicine, Department of Pharmacology and Toxicology, University of Belgrade

<sup>3</sup> Faculty of Veterinary Medicine, Department of Animal Nutrition and Botany

<sup>4</sup> Scientific Institute of Veterinary Medicine of Serbia

\* Correspondence: Saša M. Trailović sasa@vet.bg.ac.rs

## Abstract

The active constituents of essential plant oils (EOAIs), monoterpenoid carvacrol and monoterpene p-cymene, are widely distributed in many aromatic plants and their products. They differ in that carvacrol has a phenolic functional group. The numerous pharmacological effects of these two EOAs are well known. In different doses/concentrations, they exhibit analgesic, neuroprotective, vasorelaxant, anti-inflammatory, antiviral, antibacterial and antiparasitic effects. The acute toxicity of carvacrol and p-cymene in rats and the free-living nematode *Caenorhabditis elegans* was investigated. Furthermore, the impact of subacute administration of these two terpenes on general health, CNS integration, i.e. motor coordination and balance of rats, as well as their effects on the movement of adult *C. elegans*, was also examined. The aim was to compare the effects and describe in more detail the selective toxicity of carvacrol and p-cymene. The calculated LD<sub>50</sub> value of carvacrol was 790.15±1.15 mg/kg, while the LD<sub>50</sub> value of p-cymene is above 3000 mg/kg. Tested doses of carvacrol and p-cymene administered for 28 days (50, 100, and 200 mg/kg) did not exert any effect on the CNS of rats or cause any clinical disorders. LC<sub>50</sub> value of carvacrol for adult *C. elegans* was 184.13±1.51 µM and for p-cymene 1268±1.65 µM. In subacute testing, carvacrol showed negative effects on *C. elegans* reproduction, distance traveled, movement speed and rotational index at lower concentrations than p-cymene, indicating higher toxicity. On the other hand, although less toxic to *C. elegans*, p-cymene exhibited a specific effect on worm motility, with more rolling which should be further investigated, and can be a consequence of cuticle damage or loss of orientation.

**Keywords:** carvacrol; p-cymene; rats; *Caenorhabditis elegans*

## 1. Introduction

Active ingredients of essential plant oils (EOAIs), monoterpenoid carvacrol and monoterpene p-cymene are increasingly being studied and used in various therapies [1,2,3]. Both are widely distributed in several aromatic plants and their products. p-cymene is a main compound in essential oil of genus *Artemisia* (Asteraceae), *Protium* (Burseraceae), *Origanum*, *Ocimum*, *Thymus*, (Lamiaceae), and *Eucalyptus* (Myrtaceae) [4,5], while carvacrol is the major natural constituent in the essential oil of aromatic plants belonging to the family Lamiaceae, such as oregano and thyme. Carvacrol is a derivative of cymene, it is formed during the biosynthesis process in plants, and it is fundamentally categorized as a monoterpenoid due to its phenolic functional group. Numerous pharmacological effects of these two EOAs have been described. They are known to exhibit analgesic, neuroprotective, vasorelaxant, anti-inflammatory, antiviral, antibacterial, and antiparasitic

effects [2,6,7]. Until now, we were primarily interested in investigations of the mechanism of anthelmintic action of EOAs. We proved that carvacrol is a non-competitive antagonist of different types of nicotinic-acetylcholine receptors (nAChR) of parasitic nematodes and that it most likely binds to the orthosteric site on the receptor [8,9,10]. The action of carvacrol on mammalian muscle-type nAChR is somewhat different. Our studies show that carvacrol acts on the presynaptic nAChR causing tetanic fade [9]. On the other hand, a single oral administration of carvacrol to rats disrupts motor coordination on Rota-rod test with an ED<sub>50</sub> value of 421.6 mg/kg. However, no obvious signs of brain or nervous system depression were seen. The animals still had normal reflexes, could walk normally on a flat surface after being pinched on the tail, and all responded normally to external stimuli like approaching or touching [8]. Long-term antiparasitic, or any other therapeutic administration of carvacrol and p-cymene would involve the issue of their potential side effects and toxicity for the host. For this reason, our goal was to examine the impact of subacute administration of these two terpenes and the consequences for the general health condition, integration of the rat CNS, i.e. motor coordination and balance. We also examined the movement of wild-type free-living nematode *Caenorhabditis elegans* after subacute exposure to carvacrol and p-cymene. Our aim was to compare the effects and describe more closely the selective toxicity of the tested EOAs.


## 2. Results

### 2.1. Determination of the Median Lethal Dose (LD<sub>50</sub>) of Carvacrol and P-Cymene in Rats

At the very beginning of the investigation, we determined the LD<sub>50</sub> of carvacrol and p-cymene in rats and the LC<sub>50</sub> of these two monoterpenoids for *C. elegans*. The calculated LD<sub>50</sub> value of carvacrol was 790.15±1.15µg/kg. However, unlike carvacrol, it was not possible to calculate the LD<sub>50</sub> value of p-cymene because even the maximum possible volume of p-cymene that was administrated, does not cause deaths. So, we can only state that the LD<sub>50</sub> of p-cymene is above 3000mg/kg.

### 2.2. Determination of the Median Lethal Concentration (LC<sub>50</sub>) of P-Cymene and Carvacrol in *C. elegans*

The LC<sub>50</sub> value of carvacrol for adult *C. elegans* was 184.13±1.51µM (Figure 1A). On the other hand, the LC<sub>50</sub> value of p-cymene for adult *C. elegans* was 1268±1.65µM (Figure 1B).



**Figure 1.** A) Sigmoid curve of the relationship between the concentration of carvacrol and its lethal effect for adult *C. elegans*; B) Sigmoid curve of the relationship between the concentration of p-cymene and its lethal effect for adult *C. elegans*.

### 2.3. Rota-Rod Performance

In the following third part of the research, we treated 6 groups of rats with increasing oral doses of carvacrol or p-cymene for 28 days, while the 7th group was an untreated control. Every day the

rats were tested on the Rota-rod apparatus and the number of falls is recorded. During the entire observation period of 28 days, not a single fall was recorded in the rats treated with carvacrol or p-cymene. Furthermore, the righting reflex was completely preserved and movement on a flat surface was normal. No signs of CNS disorders, motor coordination problems, or other clinical symptoms were recorded.

#### 2.4. Investigation of the Influence of Carvacrol and P-Cymene on the Motility of Adult *C. elegans*

In the examination of the subacute effects of carvacrol on adult *C. elegans*, it was shown that concentrations of 1, 10, and 100 $\mu$ M do not affect the number of adults. The number increased every day and at the end of the trial it was 20 to 78% higher than at the beginning. However, the highest concentration of carvacrol tested (300 $\mu$ M), caused a reduction in the number of adults by almost 50% (Table 1). Increasing concentrations of p-cymene of 30, 100 and 300 $\mu$ M did not significantly affect the number of adults which increased by 60, 81, 3.5 and 116% after three days of exposure. However, the highest tested concentration of p-cymene of 1000 $\mu$ M reduced the number of adults by more than 50%. In the control group, an increase in the number of adults by approximately 60% was recorded (Table 2).


**Table 1.** The number of *C. elegans* adults during a 3-day exposure to increasing concentrations of carvacrol (n=3).

	Control	Carvacrol 1mM	Carvacrol 10 $\mu$ M	Carvacrol 30 $\mu$ M	Carvacrol 100 $\mu$ M	Carvacrol 300 $\mu$ M
Day 1	20.00 $\pm$ 4.16	14.33 $\pm$ 7.84	19.00 $\pm$ 11.79	20.00 $\pm$ 8.33	8.33 $\pm$ 6.33	10.00 $\pm$ 2.65
Day 2	27.67 $\pm$ 6.38	18.33 $\pm$ 10.34	21.67 $\pm$ 7.22	24.00 $\pm$ 8.00	11.67 $\pm$ 4.67	8.00 $\pm$ 1.73
Day 3	32.33 $\pm$ 2.03	25.66 $\pm$ 3.38	25.67 $\pm$ 3.38	24.67 $\pm$ 7.88	14.33 $\pm$ 5.49	5.33 $\pm$ 1.33

**Table 2.** The number of *C. elegans* adults during a 3-day exposure to increasing concentrations of p-cymene (n=3).

	Control	P-cymene 30mM	P-cymene 100 $\mu$ M	P-cymene 300 $\mu$ M	P-cymene 1000 $\mu$ M
Day 1	20.00 $\pm$ 4.16	22.00 $\pm$ 10.26	29.00 $\pm$ 10.44	12.00 $\pm$ 2.52	32.67 $\pm$ 2.40
Day 2	27.67 $\pm$ 6.38	25.33 $\pm$ 8.51	28.00 $\pm$ 4.93	11.00 $\pm$ 2.65	27.67 $\pm$ 11.15
Day 3	32.33 $\pm$ 2.03	40.00 $\pm$ 4.00	30.33 $\pm$ 2.90	26.00 $\pm$ 0.58	16.00 $\pm$ 3.21


All tested concentrations of carvacrol significantly reduced the distance of movement compared to control (Figure 2A). Similar to carvacrol, p-cymene caused a decrease in the movement distance of *C. elegans* in all applied concentrations (Figure 2B).




**Figure 2. A)** Distance traveled during the three-day activity measurement of *C. elegans* (mm) in the presence of increasing concentrations of carvacrol (n=3). Statistically significant difference compared to the control finding, \*\*\*\*-p<0.0001; \*\*\*-p=0.0008; \*\*-p=0.0005; \*-p= 0.0112. **B)** Distance traveled during the three-day activity measurement of *C. elegans* (mm) in the presence of increasing concentrations of P-cymene (n=3). Statistically significant difference compared to the control finding, \*\*\*\*-p<0.0001; \*-p=0.0468; \*\*-p=0.021; \*-p=0.0150; \*-p=0.0141; \*\*-p=0.0076; \*\*\*-p=0.0002.

All tested concentrations of carvacrol significantly reduced the average speed of *C. elegans* movement in the 1st, 2nd and 3rd day of observation (Figure 3A). At the highest concentrations of carvacrol (100 and 300μM), a decrease in movement speed was recorded from the 1st to the 3rd day of testing. It is interesting that in the presence of the p-cymene, the speed increased from the first to the third day, which was not observed in the control. This could be considered some kind of worm adaptation (Figure 3B).

The rotation index of *C. elegans* was reduced under the influence of all applied concentrations of carvacrol compared to the control, but the reduction was significant only after the first 24h of exposure to 1, 100 and 300μM of carvacrol (Figure 4A). Otherwise, in the control and all other treated *C. elegans*, the rotation index increased insignificantly from the 1st to the 3rd day of exposure to carvacrol. Unlike carvacrol, p-cymene caused a non-significant increase in the rotation index in *C. elegans*, except at the highest concentration tested (Figure 4B). The motility score of *C. elegans* was non-significantly reduced at all tested concentrations of carvacrol and p-cymene (Figure 5A and 5B).




**Figure 3.** A) Average movement speed of adult *C. elegans* measured during three days in the presence of increasing concentrations of carvacrol. Statistically significant difference compared to the control: \*-p=0.033; \*\*-p=0.0084; \*\*-p=0.0059; \*\*-p=0.0013; \*\*\*\*-p<0.0001; \*\*-p=0.0028; \*\*-p=0.0059; \*\*\*\*-p<0.0001; \*\*-p=0.0041; \*-p=0.0462; \*\*\*-p=0.0006; \*\*\*-p=0.0009; \*-p=0.0239; \*\*\*-p=0.0001; \*\*\*-p=0.0006. B) Average movement speed of adult *C. elegans* measured during three days in the presence of increasing concentrations of P-cymene. Statistically significant difference compared to the control: \*-p=0.0422; \*-p=0.0178; \*-p=0.0342; \*-p=0.0342.



**Figure 4.** A) Rotation index, measured during three-day exposure of adult *C. elegans* to increasing concentrations of carvacrol: \*-p=0.0279; \*\*\*-p=0.0008; \*\*\*-p=0.0004. B) Rotation index, measured during three-day exposure of adult *C. elegans* to increasing concentrations of p-cymene.

### 3. Discussion

Based on the results of the acute toxicity test, the LD<sub>50</sub> of carvacrol was 790 mg/kg. Only one LD<sub>50</sub> value for carvacrol has been found in the literature, at 810mg/kg [11], which is in agreement with our findings. On the other hand, we were unable to determine the LD<sub>50</sub> value of p-cymene because even the maximum volume that was applicable did not cause death in rats. This is the reason why we concluded that the LD<sub>50</sub> of p-cymene in rats is higher than 3000mg/kg. Data from the European Chemicals Agency (ECHA) show that the LD<sub>50</sub> value of p-cymene determined by the Acute Toxic Class Method is 4750mg/kg [12]. Obviously, this data is also in agreement with our results and indicate that p-cymene in comparison with carvacrol is far less acutely toxic for rats. Similar to the results obtained in rats, carvacrol exhibited higher acute toxicity than p-cymene for *C. elegans*. The LC<sub>50</sub> value of carvacrol obtained in this study corresponded to our previously published results [13]. However, the LC<sub>50</sub> value of carvacrol obtained by Fuentes et al. [14] is much higher and was 1.10±0.07mM. The reason may be the medium in which the examination was carried out. We tested *C. elegans* in NGM medium, while Fuentes et al. [14] used a liquid medium. We emphasize that *E. coli*, which *C. elegans* feeds on, was present in both media, so probably the NGM medium provides better feeding efficiency (and thus the introduction of carvacrol by pharyngeal pumping), as well transcuticular penetration of carvacrol. Data on the LC<sub>50</sub> value of p-cymene in *C. elegans* are not available in the literature. However, Lei et al. [15] state that concentration of p-cymene of 750μM causes worm mortality of 16 ± 1%. Definitely, our results indicate that p-cymene exhibits lower acute toxicity than carvacrol, both for rats and for *C. elegans*. These two models exhibit a high degree of agreement that can be exploited in toxicological studies.



**Figure 5.** A) Motility score of *C. elegans* in the presence of increasing concentrations of carvacrol and B) p-cymene.

Doses of carvacrol and p-cymene of 50, 100 and 200mg/kg administered orally for 28 days did not cause rats to fall off the rotating rod. The righting reflex was completely preserved and the rats moved normally on a flat surface. It is obvious that the tested doses applied for 28 days do not lead to disturbances in the integration of rats' nervous system, nor do they affect the peripheral nervous system function. We have previously demonstrated that carvacrol exhibited a time and dose-dependent effect on the Rota-rod performances of rats, with a high value of the  $ED_{50}=421.6$  mg/kg. Furthermore, carvacrol acts on the presynaptic nACh receptor on the neuromuscular junction and produces tetanic fades in isolated rat diaphragm contractions [8]. Nevertheless, the  $ED_{50}$  value of carvacrol now cumulatively reached after two days of administration, but without the previously described effect on the Rota-rod test. For p-cymene, there are available data on mice test. Single intraperitoneal administration of p-cymene in doses of 25, 50, and 100 mg/kg did not induce changes in motor performance, since the animals remained on the rotating rod after 3, 30, 60, and 120 min [16]. According to our results and data from the literature, it is obvious that carvacrol and p-cymene do not exert a cumulative toxic effect in rats. This point of view is supported by the fact that a single administration of high doses of carvacrol and p-cymene caused disturbances of the nervous system functions and clinical symptoms of toxicity. The absence of cumulative effects of carvacrol and p-cymene can be explained by their rapid biotransformation and elimination. Carvacrol is known to be rapidly metabolized and excreted in rats within the first 24 hours of administration. Its excretion after 24 hours is very limited, and the molecule is found unchanged [17]. After intragastric administration of p-cymene 100mg/kg to rats, urinary metabolite excretion was nearly complete within 48h, amounting to 60-80% of the dose [18].

Exposure of *C. elegans* to carvacrol in concentrations from 1 to 100  $\mu$ M did not affect its development. The development of *C. elegans* from egg to adult at a temperature of 20°C takes about 3 days [19]. So, we could expect an increase in the number of adults from the beginning to the end of our study. However, the highest tested concentration of carvacrol of 300 $\mu$ M caused a 50% reduction in the number of adults, indicating that carvacrol interferes with the development of *C. elegans*. On the other hand, p-cymene reduced the number of adults by 50% only at a concentration of 1000 $\mu$ M. Obviously, carvacrol exerts an effect on the development of *C. elegans* in a three times lower concentration than p-cymene.

The length of the traveled distance in the control worms did not differ during the three days of exposure. For 5 minutes of observation, the average distance covered on the first, second, and third day was  $12.407\pm 1.874$ ,  $8.707\pm 1.117$ ,  $10.297\pm 1.946$  mm. However, even the lowest tested concentration of carvacrol (1 $\mu$ M) significantly reduced the length of the traveled distance to  $3.750\pm 0.66$ ,  $3.610\pm 0.712$  and  $3.910\pm 0.800$  mm. The other applied concentrations of carvacrol also caused a significant reduction in the distance, but at the same time, only at the two highest concentrations tested, 100 and 300 $\mu$ M,

the distance decreased from day 1 to day 3 ( $3.900\pm 0.240$ ,  $2.520\pm 0.736$ ,  $1.853\pm 0.225$ mm and  $2.433\pm 0.175$ ,  $1.570\pm 0.180$ ,  $1.080\pm 0.132$ mm) (Figure 2A). The presence of p-cymene in all tested concentrations shortened the movement distance of *C. elegans* (Figure 2B). However, this reduction was highly significant only after the first day of exposure to all tested concentrations, and by the third day in all worms the distance increased (but without reaching the control level), which could indicate some kind of adaptation on the presence and effect of p-cymene.

A similar effect of carvacrol was recorded on the movement speed of *C. elegans*. All applied concentrations significantly decreased the speed compared to the control, but it is interesting that only the two highest concentrations also decreased the speed from the 1st to the 3rd day of observation (Figure 3A). Unlike carvacrol, p-cymene significantly reduced movement speed only at the highest tested concentration. Also, the speed was reduced on the first day after exposure to lower concentrations, but it was observed that on the 2nd and 3rd day, despite the presence of p-cymene, the speed of movement increased. This could be interpreted as some kind of adaptation of the worms to the presence of p-cymene. The observed effect of carvacrol on the length of the distance and the speed of movement can be explained by its proven inhibitory effect on the nicotinic acetylcholine receptor of nematodes [8,9,20]. This inhibition affects the speed of movement, and the distance covered. On the other hand, it is obvious that p-cymene exerts a weaker effect on these two observed parameters of *C. elegans* motility. It is important that p-cymene did not exhibit any inhibitory effect on contractions of the neuromuscular preparation of *A. suum* even at a concentration up to 1mM (Marjanović, unpublished results). We can assume that p-cymene affects *C. elegans* movement by a different mechanism compared to carvacrol.

The values of the average motility score and the rotation index are considered together. Carvacrol reduced the rotational index only on the first day of observation and significantly only in concentrations of 100 and 300  $\mu$ M. At the same time, the motility score was also insignificantly reduced at all tested concentrations. On the other hand, the rotation index in worms exposed to p-cymene increased from the 1st to the 3rd day and was even higher than the control value, except at the highest tested concentration where it decreased until the third day. The motility score was lower than the control at all tested concentrations of p-cymene, but it also increased from the 1st to the 3rd day, except for the highest tested concentration. Rotation index is particularly useful for distinguishing between worms that are inactive and those that are merely rotating in place, as observed in certain roller strains. In the context of a paralysis assay where worms exhibit a low motility index, two scenarios arise: A low motility index with a low rotation index value typically indicates that the worms are motionless, which is observed as an effect of carvacrol; A low motility index with a high rotation index suggests that the worms are engaged in rolling movements, which is noted with p-cymene. This type of specific movement that occurs in some *C. elegans* mutants is a consequence of damage to the cuticle [21] or a maneuver for reorientation.

## 4. Materials and Methods

### 4.1. Animals

#### 4.1.1. Rats

White male albino Wistar rats (Mmab:W, Strain Report - Rat Genome Database), weighing 160-230 g were housed under standard conditions for laboratory animals in groups of five with controlled 12-h light/dark cycle, temperature of 21-24 °C, and “ad libitum” access to standard diet and water. All procedures in the study conformed to EEC Directive 86/609 and were approved by the Ethic Committee of the Faculty of Veterinary Medicine University of Belgrade and by Decision of the Veterinary Administration, Ministry of Agriculture, Forestry and Water Management, number: 7492 2023 14841 002 001 323 022, from 09.01.2024. At the end of the experiments all the rats were humanely euthanized by the overdose of pentobarbitone in accordance with the Home Office Code of Practice [25] (1997).

#### 4.1.2. *C. elegans*

*C. elegans*, wild type (N2 Bristol) was obtained from the Caenorhabditis Genetics Center [22]. Worms were cultivated and adults were separated for testing as we previously explained in Stojković et al. [13].

#### 4.2. Chemicals and Method of Administration

Sigma Chemical Co. (St. Louis, MO, USA) supplied p-cymene and carvacrol. Both were dissolved in peanut oil and administered to rats orally by gavage in a volume of 0.1ml/100g of body weight.

#### 4.3. Procedures

##### 4.3.1. Determination of the Median Lethal Dose (LD<sub>50</sub>) of carvacrol and p-cymene in rats

Fifty-four male rats were split into 9 groups with 6 rats in each group. After a period of adaptation, the rats were given increasing doses of carvacrol through oral gavage: 400, 600, 800, and 1000 mg per kilogram of body weight, or p-cymene: 500, 1000, 2000, and 3000 mg per kilogram. Following the treatment, the rats were observed for 24 hours to check for any deaths, and the results were used to find out the LD<sub>50</sub> values for both carvacrol and p-cymene.

##### 4.3.2. Determination of the Median Lethal Concentration (LC<sub>50</sub>) of p-cymene and carvacrol in *C. elegans*

To find out the LC<sub>50</sub> of p-cymene and carvacrol, adult nematodes were placed in Petri dishes 3 cm in diameter. Each dish had 2.5 milliliters of NGM substrate mixed with different amounts of p-cymene (30, 100, 300, 1000, and 3000 µM) or carvacrol (30, 100, 300, and 1000 µM). The number of worms in each 20-microliter sample was between 20 and 37. Each concentration was tested on three Petri dishes. The determination of LC<sub>50</sub> is described in Stojković et al., [20].

##### 4.3.3. Rota-Rod Performance

The Rota-rod test is used to check how well rodents can keep their balance and coordination [23]. This test evaluates how long a rat can stay balanced on a rotating rod (EI Unit, Serbia). Before starting the tests, the animals were trained for three days so they could stay on the rod for 180 seconds while it spins at a steady speed of 8 rpm. The rats were divided into 7 groups with 6 rats in each group. Each group was given carvacrol or p-cymene dissolved in peanut oil at doses of 50, 100, and 200 mg/kg for 28 days. Every day, they did a 3-minute test on the Rota-rod machine, and the number of times they fell off was recorded. The righting reflex and the test of moving on a flat surface were used to check if the doses of p-cymene and carvacrol had any noticeable effects on the rats' central nervous system. All the tests were done following the methods described in Trailović et al. [24].

##### 4.3.4. Investigation of the Influence of Carvacrol and P-Cymene on the Motility of Adult *C. elegans*.

*C. elegans*, N2 wild type was obtained from the Caenorhabditis Genetics Center [22]. The effect of p-cymene and carvacrol on the motility of adult *C. elegans* was investigated by using the Microtracker SMART 8 device (PHYLUMTECH S.A., Argentina). In the Petri dishes (diameter 3cm) filled with 2.5 ml of NGM substrate and increasing concentrations of p-cymene (30, 100, 300 and 1000 µM) or carvacrol (1, 10, 30, 100 and 300 µM) the suspension (20 µL) of adult *C. elegans* was added. Data collection was performed once a day, after 24, 48 and 72 hours of exposure during a five-minute interval. The plates were subjected to "tapping" stimulation before each measurement. The following *C. elegans* motility parameters were monitored and analyzed: number of adults, movement speed, distance traveled, rotation index and motility score.

#### 4.4. Statistics

All values are expressed as mean  $\pm$  standard error of the mean (mean  $\pm$  SE). Unless stated otherwise, results were tested using the one-way ANOVA or t-test, and differences were considered significant at  $p < 0.05$ . Furthermore, nonlinear regression was used to examine dose-response relationship and determination of the Median Lethal Dose ( $LD_{50}$ ) as well as the Median Lethal Concentration ( $LC_{50}$ ) of carvacrol and p-cymene.

All data was analyzed using GraphPad Prism version 6.00 for Windows (GraphPad Software, La Jolla California USA).

## 5. Conclusions

The results obtained definitely indicate that p-cymene exhibits lower acute toxicity than carvacrol, both for rats and *C. elegans*. These two models exhibit a high degree of agreement that can be exploited in toxicological studies. Tested doses of carvacrol and p-cymene administered for 28 days did not exert any effect on the CNS of rats or cause any clinical disorders. The absence of cumulative effects is probably due to rapid biotransformation and elimination in rats. This is supported by the fact that a cumulative effect was recorded in *C. elegans*. Carvacrol exhibited negative effects on *C. elegans* reproduction, distance traveled, movement speed, and rotational index at lower concentrations than p-cymene, indicating higher toxicity. On the other hand, although less toxic to *C. elegans*, p-cymene exhibited a specific effect on worm motility, with more rolling, which is characteristic of mutants with damaged cuticle or a maneuver for reorientation. In any case, this effect of p-cymene should be further investigated because it indicates a specific action.

**Author Contributions:** “Conceptualization, O.S. and S.M.T.; methodology, S.M.T, V.S., DJ.S.M., N.Z.; software, Dj.S.M., T.M., D.M.; validation, S.M.T., O.S.; formal analysis, DJ.S.M., S.M.T.; investigation, O.S., D.M., DJ.S.M., T.M., N.Z.; resources, S.M.T., V.S.; data curation, S.M.T.; writing—original draft preparation, O.S., S.M.T.; writing—review and editing, S.M.T., J.N.T.; visualization, S.M.T., T.M., V.S.; supervision, S.M.T.; project administration, D.M.; funding acquisition, S.M.T. All authors have read and agreed to the published version of the manuscript.”

**Funding:** “The study was supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia (Contract number 451-03-136/2025-03/200143), Science Fund of the Republic of Serbia, #GRANT No, 7355 Project title—FARMASCA (<https://farmasca.vet.bg.ac.rs>).

**Institutional Review Board Statement:** “All procedures in the study conformed to EEC Directive 86/609 and were approved by the Ethic Committee of the Faculty of Veterinary Medicine University of Belgrade and by Decision of the Veterinary Administration, Ministry of Agriculture, Forestry and Water Management, number: 7492 2023 14841 002 001 323 022, from 09.01.2024.”

**Data Availability Statement:** The data that support the findings of this study are incorporated into the article.

**Conflicts of Interest:** “The authors declare no conflicts of interest.”

## References

1. Imran, M.; Aslam, M.; Alsagaby, S.A.; Saeed, F.; Ahmad, I.; Afzaal, M.; Arshad, M.U.; Abdelgawad, M.A.; El-Ghorab, A.H.; Khames, A.; Shariati, M.A.; Ahmad, A.; Hussain, M.; Imran, A.; Islam, S. Therapeutic application of carvacrol: A comprehensive review. *Food Sci Nutr.* **2022**, *10*(11), 3544-3561.
2. Balahbib, A.; El Omari, N.; Hachlafi, N.E.; Lakhdar, F.; El Menyiy, N.; Salhi, N.; Mrabtiž, H.N.; Bakrim, S.; Zengin, G.; Bouyahya, A. Health beneficial and pharmacological properties of p-cymene. *Food Chem Toxicol.* **2021**, *153*:112259.
3. Hyldgaard, M.; Mygind, T.; Meyer, R.L. Essential oils in food preservation: mode of action, synergies, and interactions with food matrix components. *Front Microbiol.* **2012**, *3*, 12.
4. Philis, J.G. The S1  $\leftarrow$  S0 spectrum of jet-cooled p-cymene. *Spectrochim Acta A Mol Biomol Spectrosc.* **2005**, *61*(6), 1239-41.

5. de Santana, M.F.; Guimarães, A.G.; Chaves, D.O.; Silva, J.C.; Bonjardim, L.R.; de Lucca Júnior, W.; Ferro, J.N.; Barreto, Ede. O.; dos Santos, F.E.; Soares, M.B., Villarreal, C.F.; Quintans, Jde, S.; Quintans-Júnior, L.J. The anti-hyperalgesic and anti-inflammatory profiles of p-cymene: Evidence for the involvement of opioid system and cytokines. *Pharm Biol.* **2015**, *53*(11),1583-90.
6. Milovanović, M.; Marjanović, Đ.; Trailović, S.; Stojanović D. Antinociceptive effect of p-cymene and cinnamaldehyde: their relationship with the L-arginine-nitric oxide pathway in rats. *Acta Veterinaria* **2025**, *75*(2), 193-211.
7. Suntres, Z.E.; Coccimiglio, J.; Alipour, M. The bioactivity and toxicological actions of carvacrol. *Crit Rev Food Sci Nutr.* **2015**, *55*(3), 304-18.
8. Marjanović, D.S.; Zdravković, N.; Milovanović, M.; Trailović, J.N.; Robertson, A.P.; Todorović, Z.; Trailović, S.M. Carvacrol acts as a potent selective antagonist of different types of nicotinic acetylcholine receptors and enhances the effect of monepantel in the parasitic nematode *Ascaris suum*. *Veterinary parasitology* **2020**, *278*, 109031
9. Trailovic, S.M.; Rajkovic, M.; Marjanovic, D.S.; Neveu, C.; Charvet, C.L. Action of Carvacrol on *Parascaris* sp. and Antagonistic Effect on Nicotinic Acetylcholine Receptors. *Pharmaceuticals (Basel)*. **2021**, *14*(6), 505.
10. Marjanović, D.S.; Trailović, S.M.; Milovanović, M.; Interaction of agonists of a different subtype of the nAChR and carvacrol with GABA in *Ascaris suum* somatic muscle contractions. *J Nematol.* **2021**, *53*:e2021-22.
11. Baser, K.H. Biological and pharmacological activities of carvacrol and carvacrol bearing essential oils. *Curr Pharm Des.* **2008**, *14*(29), 3106-19.
12. European Chemicals Agency (ECHA), p-cymene, **2014**.
13. Stojković, M.; Todorović, Z.; Protic, D.; Stevanovic, S.; Medić, D., Charvet, C.L.; Courtot, E., Marjanović, D.S.; Nedeljković Trailović, J.; Trailović, S.M. Corrigendum: Pharmacological effects of monoterpene carveol on the neuromuscular system of nematodes and mammals. *Front Pharmacol.* **2024**, *15*, 1466575.
14. Fuentes, C.; Verdú, S.; Fuentes, A.; Ruiz, M.J.; Barat, J.M. Effects of essential oil components exposure on biological parameters of *Caenorhabditis elegans*. *Food Chem Toxicol.* **2022**, *159*, 112763.
15. Lei, J.; Leser, M.; Enan, E. Nematicidal activity of two monoterpenoids and SER-2 tyramine receptor of *Caenorhabditis elegans*. *Biochem Pharmacol.* **2010**, *79*(7), 1062-71.
16. Santana, M.F.; Lucindo J.; Quintans-Júnior, S.C.H.; Cavalcanti, M.G.B.; Oliveira, A. G.; Guimarães, E.S.; Cunha, M.S.; Melo, M.R.V.; Santos, A.A.S.; Araújo, L.R.B. p-Cymene reduces orofacial nociceptive response in mice, *Revista Brasileira de Farmacognosia* **2011**, *21*(6),1138-1143.
17. Austgulen, L.T.; Solheim, E.; Scheline, R.R. Metabolism in rats of p-cymene derivatives: carvacrol and thymol. *Pharmacol Toxicol.* **1987**, *61*(2), 98-102.
18. Walde, A.; Ve, B.; Scheline, R.R.; Monge, P. p-Cymene metabolism in rats and guinea-pigs. *Xenobiotica* **1983**, *13*(8), 503-12.
19. Braendle, C.; Paaby, A. Life history in *Caenorhabditis elegans*: from molecular genetics to evolutionary ecology. *Genetics.* **2024**, *228*(3), iyae151.
20. Stojković, M.; Marjanović, D.S.; Medić, D.; Charvet, C.L.; Trailović, S.M. Neuromuscular System of Nematodes Is a Target of Synergistic Pharmacological Effects of Carvacrol and Geraniol. *Pharmaceuticals (Basel)* **2025**. *18*(8), 1232.
21. Higgins, B.J.; Hirsh, D. Roller mutants of the nematode *Caenorhabditis elegans*. *Mol Gen Genet.* **1977**, *150*(1), 63-72.
22. Charvet, C.L.; Guégnard, F.; Courtot, E.; Cortet, J.; Neveu, C. Nicotine-sensitive acetylcholine receptors are relevant pharmacological targets for the control of multidrug resistant parasitic nematodes. *Int J Parasitol Drugs Drug Resist.* **2018**, *8*(3), 540-549.
23. Jones, B.J.; Roberts, D.J. The quantitative measurement of motor inco-ordination in naive mice using an accelerating rotarod. *J Pharm Pharmacol.* **1968**, *20*(4), 302-4.

24. Trailovic, S.M.; Ivanovic, S.R.; Varagić, V.M. Ivermectin effects on motor coordination and contractions of isolated rat diaphragm. *Res Vet Sci.* **2011**, *91*(3), 426-33.
25. Home Office, Code of Practice for the Humane Killing of Animals under Schedule 1 to the Animals (Scientific Procedures) Act 1986, **1997**.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.