

Article

Not peer-reviewed version

---

# On $j$ -Fuzzy $\gamma$ $l$ -Open Sets with Some Applications

---

[Fahad Alsharari](#)\*, [Jawaher Al-Mufarrij](#), [Islam M. Taha](#)\*

Posted Date: 22 May 2025

doi: 10.20944/preprints202505.1738.v1

Keywords: fuzzy ideals; fuzzy topology;  $j$ -fuzzy  $\gamma$  $l$ -open set; fuzzy  $\gamma$  $l$ -continuity; fuzzy  $\gamma$  $l$ -irresoluteness



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

# On $j$ -Fuzzy $\gamma\mathcal{I}$ -Open Sets with Some Applications

Fahad Alsharari <sup>1</sup>, Jawaher Al-Mufarrij <sup>2</sup> and Islam M. Taha <sup>3,\*</sup>

<sup>1</sup> Department of Mathematics, College of Science, Jouf University, Sakaka 72311, Saudi Arabia

<sup>2</sup> Department of Mathematics, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

<sup>3</sup> Department of Mathematics, College of Science, Sohag University, Sohag, Egypt

\* Correspondence: f.alsharari@ju.edu.sa (F.A.); imtaha2010@yahoo.com (I.M.T.)

**Abstract:** In this article, we first explored and investigated a new class of fuzzy sets, called  $j$ -fuzzy  $\gamma\mathcal{I}$ -open ( $j$ - $F\gamma\mathcal{I}$ -open) sets on fuzzy ideal topological spaces ( $\mathcal{FIT}$  spaces). The class of  $j$ - $F\gamma\mathcal{I}$ -open sets is contained in the class of  $j$ -fuzzy strong  $\beta\mathcal{I}$ -open ( $j$ - $FS\beta\mathcal{I}$ -open) sets and contains all  $j$ -fuzzy pre- $\mathcal{I}$ -open ( $j$ - $FPL$ -open) sets and  $j$ -fuzzy semi- $\mathcal{I}$ -open ( $j$ - $FSL$ -open) sets. We also introduced and studied the closure and interior operators with respect to the classes of  $j$ - $F\gamma\mathcal{I}$ -closed sets and  $j$ - $F\gamma\mathcal{I}$ -open sets. However, we defined and discussed novel types of fuzzy  $\mathcal{I}$ -separation axioms using  $j$ - $F\gamma\mathcal{I}$ -closed sets, called  $j$ - $F\gamma\mathcal{I}$ -regular spaces and  $j$ - $F\gamma\mathcal{I}$ -normal spaces. Thereafter, we displayed and investigated the concept of fuzzy  $\gamma\mathcal{I}$ -continuity ( $F\gamma\mathcal{I}$ -continuity) using  $j$ - $F\gamma\mathcal{I}$ -open sets. Moreover, we presented and characterized the concepts of fuzzy weak  $\gamma\mathcal{I}$ -continuity ( $FW\gamma\mathcal{I}$ -continuity) and fuzzy almost  $\gamma\mathcal{I}$ -continuity ( $FA\gamma\mathcal{I}$ -continuity), which are weaker forms of  $F\gamma\mathcal{I}$ -continuity. Finally, we introduced and studied some new fuzzy  $\gamma\mathcal{I}$ -mappings via  $j$ - $F\gamma\mathcal{I}$ -open sets and  $j$ - $F\gamma\mathcal{I}$ -closed sets, called  $F\gamma\mathcal{I}$ -open mappings,  $F\gamma\mathcal{I}$ -closed mappings,  $F\gamma\mathcal{I}$ -irresolute mappings,  $F\gamma\mathcal{I}$ -irresolute open mappings, and  $F\gamma\mathcal{I}$ -irresolute closed mappings.

**Keywords:** fuzzy ideals; fuzzy topology;  $j$ -fuzzy  $\gamma\mathcal{I}$ -open set; fuzzy  $\gamma\mathcal{I}$ -continuity; fuzzy  $\gamma\mathcal{I}$ -irresoluteness

**MSC:** 54A40; 54C05; 54C08; 54D15

## 1. Introduction

The concept of a fuzzy set was first defined in 1965 by Zadeh [1] as a suitable approach to address with uncertainty cases that we cannot be efficiently managed via classical techniques. Over the last decades, the researches of fuzzy sets have a vital role in mathematics and applied sciences and garnered significant attention due to its ability to handle uncertain and vague information in various real life applications such as control systems [2,3], artificial intelligence [4,5], image processing [6,7], decision-making [8,9], etc. The integration between fuzzy sets and some uncertainty approaches such as rough sets and soft sets has been discussed in [10-12]. The notion of a fuzzy topology was introduced in 1968 by Chang [13] and this development has led to the expansion and discussion of several classical topological concepts in the context of a fuzzy topology [14-17], providing more accurate and flexible models to address problems of uncertainty in various real life ears. Overall, according to Šostak [18], the concept of a fuzzy topology being a crisp subclass of the class of fuzzy sets and fuzziness in the concept of openness of a fuzzy set have not been considered, which seems to be a drawback in the process of fuzzification of a topological space. Thereafter, Šostak [18] introduced a new notion of a fuzzy topology as the notion of openness of fuzzy sets. It is an extension of a fuzzy topology defined by Chang [13]. Furthermore, several researchers (see [19-27]) have redisplayed the same concept and investigated fuzzy topological spaces ( $\mathcal{FT}$  spaces) being unaware of Šostak's work.

The generalizations of fuzzy open sets plays an effective role in a fuzzy topology through their ability to improve on several results, or to open the door to explore and discuss many fuzzy topological concepts such as fuzzy compactness [20,21], fuzzy connectedness [20], fuzzy continuity [19,20], etc.

Furthermore, the concepts of  $j$ -fuzzy pre-open ( $j$ -FP-open) sets,  $j$ -fuzzy semi-open ( $j$ -FS-open) sets,  $j$ -fuzzy  $\beta$ -open ( $j$ -F $\beta$ -open) sets, and  $j$ -fuzzy  $\alpha$ -open ( $j$ -F $\alpha$ -open) sets were presented and investigated by the authors of [24,26] in  $\mathcal{F}T\mathcal{S}$ s based on Šostak's sense [18]. Kim et al. [24] displayed and investigated weaker forms of fuzzy continuity, called FS-continuity (resp. FP-continuity and F $\alpha$ -continuity) between  $\mathcal{F}T\mathcal{S}$ s in the sense of Šostak. Abbas [26] defined and discussed the concepts of F $\beta$ -continuous (resp. F $\beta$ -irresolute) mappings. Also, Kim and Abbas [27] explored and characterized new types of  $j$ -fuzzy compactness. Overall, the notions of  $j$ -fuzzy  $\gamma$ -open ( $j$ -F $\gamma$ -open) sets and  $j$ -fuzzy  $\gamma$ -closed ( $j$ -F $\gamma$ -closed) sets were introduced and discussed by the authors of [28].

The notion of  $j$ -fuzzy local function was presented and investigated by Taha and Abbas [29] in an  $\mathcal{F}I\mathcal{T}\mathcal{S}$   $(Z, \zeta, \mathcal{I})$  based on Šostak's sense [18]. Moreover, the notions of fuzzy lower (resp. upper) weakly and almost  $\mathcal{I}$ -continuous multifunctions were displayed and investigated by Taha and Abbas [29]. Also, Taha [30-32] introduced the notions of  $j$ -F $\mathcal{S}\mathcal{I}$ -open sets,  $j$ -F $\mathcal{P}\mathcal{I}$ -open sets,  $j$ -F $\alpha\mathcal{I}$ -open sets,  $j$ -F $\beta\mathcal{I}$ -open sets,  $j$ -F $\mathcal{S}\beta\mathcal{I}$ -open sets,  $j$ -F $\delta\mathcal{I}$ -open sets, and  $j$ -G $\mathcal{F}\mathcal{I}$ -closed sets in an  $\mathcal{F}I\mathcal{T}\mathcal{S}$   $(Z, \zeta, \mathcal{I})$  based on Šostak's sense. Overall, Taha [31-33] presented the concepts of fuzzy lower (resp. upper) generalized  $\mathcal{I}$ -continuous (resp. semi- $\mathcal{I}$ -continuous, pre- $\mathcal{I}$ -continuous,  $\delta$ - $\mathcal{I}$ -continuous,  $\alpha$ - $\mathcal{I}$ -continuous,  $\beta$ - $\mathcal{I}$ -continuous, and strong  $\beta$ - $\mathcal{I}$ -continuous) multifunctions via fuzzy ideals [34].

The arrangement of this research is as follows.

(a) Section 2 provides fundamental results and concepts which we use them in our article.

(b) In Section 3, we introduce and study a new class of fuzzy sets, called  $j$ -F $\gamma\mathcal{I}$ -open sets on  $\mathcal{F}I\mathcal{T}\mathcal{S}$ s in the sense of Šostak. We also define and discuss the interior and closure operators with respect to the classes of  $j$ -F $\gamma\mathcal{I}$ -open sets and  $j$ -F $\gamma\mathcal{I}$ -closed sets. Furthermore, we explore new types of fuzzy  $\mathcal{I}$ -separation axioms using  $j$ -F $\gamma\mathcal{I}$ -closed sets, called  $j$ -F $\gamma\mathcal{I}$ -regular spaces and  $j$ -F $\gamma\mathcal{I}$ -normal spaces.

(c) In Section 4, we display and characterize the notion of F $\gamma\mathcal{I}$ -continuous mappings using  $j$ -F $\gamma\mathcal{I}$ -open sets. However, we present and discuss the notions of F $\mathcal{A}\gamma\mathcal{I}$ -continuous and F $\mathcal{W}\gamma\mathcal{I}$ -continuous mappings, which are weaker forms of F $\gamma\mathcal{I}$ -continuous mappings.

(d) In Section 5, we explore and investigate new F $\gamma\mathcal{I}$ -mappings via  $j$ -F $\gamma\mathcal{I}$ -open sets and  $j$ -F $\gamma\mathcal{I}$ -closed sets, called F $\gamma\mathcal{I}$ -closed mappings, F $\gamma\mathcal{I}$ -open mappings, F $\gamma\mathcal{I}$ -irresolute mappings, F $\gamma\mathcal{I}$ -irresolute open mappings, and F $\gamma\mathcal{I}$ -irresolute closed mappings.

(e) In Section 6, we give some potential future studies and conclusions.

## 2. Preliminaries

In this research, non-empty sets will be denoted by  $Y, X, Z$ , etc. For any fuzzy set  $\omega \in I^Z$  (where  $I = [0, 1]$  and  $I^Z$  is the class of all fuzzy sets on  $Z$ ),  $\omega^c(z) = 1 - \omega(z)$ , for each  $z \in Z$ . Also, for  $s \in I$ ,  $\underline{s}(z) = s$ , for each  $z \in Z$ .

On  $Z$ , a fuzzy point  $z_s$  is a fuzzy set, is defined as follows:  $z_s(r) = s$  if  $r = z$ , and  $z_s(r) = 0$  for any  $r \in Z - \{z\}$ . We say that  $z_s$  belongs to  $\omega \in I^Z$  ( $z_s \in \omega$ ), if  $s \leq \omega(z)$ . Also,  $P_s(Z)$  is the class of all fuzzy points.

A fuzzy set  $\nu \in I^Z$  is a quasi coincident with  $\mu \in I^Z$  ( $\nu \mathcal{Q} \mu$ ) on  $Z$ , if there is  $z \in Z$ , with  $\nu(z) + \mu(z) > 1$ . Also,  $\nu$  is not a quasi coincident with  $\mu$  ( $\nu \overline{\mathcal{Q}} \mu$ ) otherwise.

The difference between  $\psi, \mu \in I^Z$  [29] is defined as follows:

$$\psi \bar{\wedge} \mu = \begin{cases} 0, & \text{if } \psi \leq \mu, \\ \psi \wedge \mu^c, & \text{otherwise.} \end{cases}$$

**Lemma 1.** [35] Let  $\omega, \nu \in I^Z$ . Thus,

- (a) if  $\omega \mathcal{Q} \nu$ , then  $\omega \wedge \nu \neq 0$ ,
- (b)  $\omega \mathcal{Q} \nu$  iff there is  $z_s \in \omega$  such that  $z_s \mathcal{Q} \nu$ ,
- (c)  $\omega \bar{\mathcal{Q}} \nu$  iff  $\omega \leq \nu^c$ ,
- (d)  $\omega \leq \nu$  iff  $z_s \in \omega$  implies  $z_s \in \nu$  iff  $z_s \mathcal{Q} \omega$  implies  $z_s \mathcal{Q} \nu$  iff  $z_s \bar{\mathcal{Q}} \nu$  implies  $z_s \bar{\mathcal{Q}} \omega$ .

**Definition 1.** [18, 19] A mapping  $\zeta : I^Z \rightarrow I$  is called a fuzzy topology on  $Z$  if it satisfies the following conditions:

- (a)  $\zeta(0) = \zeta(1) = 1$ .
- (b)  $\zeta(\omega \wedge \nu) \geq \zeta(\omega) \wedge \zeta(\nu)$ , for any  $\omega, \nu \in I^Z$ .
- (c)  $\zeta(\bigvee_{i \in \Gamma} \omega_i) \geq \bigwedge_{i \in \Gamma} \zeta(\omega_i)$ , for any  $\omega_i \in I^Z$ .

Thus,  $(Z, \zeta)$  is called a fuzzy topological space ( $\mathcal{F}\mathcal{T}\mathcal{S}$ ) in the sense of Šostak.

**Definition 2.** [19, 24] A fuzzy mapping  $\mathbb{P} : (Z, \zeta) \rightarrow (Y, \mathfrak{S})$  is called

- (a) fuzzy continuous if  $\zeta(\mathbb{P}^{-1}(v)) \geq \mathfrak{S}(v)$ , for any  $v \in I^Y$ ;
- (b) fuzzy open if  $\mathfrak{S}(\mathbb{P}(\omega)) \geq \zeta(\omega)$ , for any  $\omega \in I^Z$ ;
- (c) fuzzy closed if  $\mathfrak{S}((\mathbb{P}(\omega))^c) \geq \zeta(\omega^c)$ , for any  $\omega \in I^Z$ .

**Definition 3.** [20, 23] For any  $\omega \in I^Z$  and  $j \in I_0$  (where  $I_0 = (0, 1)$ ) in an  $\mathcal{F}\mathcal{T}\mathcal{S}$   $(Z, \zeta)$ , we define fuzzy operators  $C_\zeta$  and  $I_\zeta : I^Z \times I_0 \rightarrow I^Z$  as follows:


$$C_\zeta(\omega, j) = \bigwedge \{v \in I^Z : \omega \leq v, \zeta(v^c) \geq j\}.$$

$$I_\zeta(\omega, j) = \bigvee \{v \in I^Z : v \leq \omega, \zeta(v) \geq j\}.$$

**Definition 4.** [24, 26, 28] Let  $(Z, \zeta)$  be an  $\mathcal{F}\mathcal{T}\mathcal{S}$  and  $j \in I_0$ . A fuzzy set  $\omega \in I^Z$  is called

- (a)  $j$ -F-open if  $\omega = I_{\zeta}(\omega, j)$ ;
- (b)  $j$ -FP-open if  $\omega \leq I_{\zeta}(C_{\zeta}(\omega, j), j)$ ;
- (c)  $j$ -FS-open if  $\omega \leq C_{\zeta}(I_{\zeta}(\omega, j), j)$ ;
- (d)  $j$ -FR-open if  $\omega = I_{\zeta}(C_{\zeta}(\omega, j), j)$ ;
- (e)  $j$ -F $\alpha$ -open if  $\omega \leq I_{\zeta}(C_{\zeta}(I_{\zeta}(\omega, j), j), j)$ ;
- (f)  $j$ -F $\beta$ -open if  $\omega \leq C_{\zeta}(I_{\zeta}(C_{\zeta}(\omega, j), j), j)$ ;
- (g)  $j$ -F $\gamma$ -open if  $\omega \leq C_{\zeta}(I_{\zeta}(\omega, j), j) \vee I_{\zeta}(C_{\zeta}(\omega, j), j)$ .

**Remark 1.** [24, 26, 28] We have the following diagram from the previous definitions.



**Definition 5.** [24, 26, 28] A fuzzy mapping  $\mathbb{P} : (Z, \zeta) \rightarrow (Y, \mathfrak{S})$  is called FS-continuous (resp. FP-continuous, F $\alpha$ -continuous, F $\beta$ -continuous, and F $\gamma$ -continuous) if  $\mathbb{P}^{-1}(\omega)$  is a  $j$ -FS-open (resp.  $j$ -FP-open,  $j$ -F $\alpha$ -open,  $j$ -F $\beta$ -open, and  $j$ -F $\gamma$ -open) set, for any  $\omega \in I^Y$  with  $\mathfrak{S}(\omega) \geq j$  and  $j \in I_0$ .

**Definition 6.** [28] For any  $\omega \in I^Z$  and  $j \in I_0$  in an  $\mathcal{FTS} (Z, \zeta)$ , we define fuzzy operators  $\gamma C_{\zeta}$  and  $\gamma I_{\zeta} : I^Z \times I_0 \rightarrow I^Z$  as follows:

$$\gamma C_{\zeta}(\omega, j) = \bigwedge \{ \mu \in I^Z : \omega \leq \mu, \mu \text{ is } j\text{-F}\gamma\text{-closed} \}.$$

$$\gamma I_{\zeta}(\omega, j) = \bigvee \{ \mu \in I^Z : \mu \leq \omega, \mu \text{ is } j\text{-F}\gamma\text{-open} \}.$$

**Definition 7.** [34] A fuzzy ideal  $\mathcal{I}$  on  $Z$ , is a map  $\mathcal{I} : I^Z \rightarrow I$  that satisfies the following:

- (a)  $\forall \omega, \mu \in I^Z$  and  $\omega \leq \mu \Rightarrow \mathcal{I}(\mu) \leq \mathcal{I}(\omega)$ .
- (b)  $\forall \omega, \mu \in I^Z \Rightarrow \mathcal{I}(\omega \vee \mu) \geq \mathcal{I}(\omega) \wedge \mathcal{I}(\mu)$ .

Moreover,  $\mathcal{I}_0$  is the simplest fuzzy ideal on  $Z$ , and is defined as follows:

$$\mathcal{I}_0(\mu) = \begin{cases} 1, & \text{if } \mu = 0, \\ 0, & \text{otherwise.} \end{cases}$$

**Definition 8.** [29] Let  $(Z, \zeta, \mathcal{I})$  be an  $\mathcal{FITS}$ ,  $j \in I_o$ , and  $\omega \in I^Z$ . Then the  $j$ -fuzzy local function  $\omega_j^*$  of  $\omega$  is defined as follows:

$$\omega_j^* = \bigwedge \{ \rho \in I^Z : \mathcal{I}(\omega \bar{\wedge} \rho) \geq j, \zeta(\rho^c) \geq j \}.$$

**Remark 2.** [29] If we take  $\mathcal{I} = \mathcal{I}_0$ , for each  $\omega \in I^Z$  we have:

$$\omega_j^* = \bigwedge \{ \rho \in I^Z : \omega \leq \rho, \zeta(\rho^c) \geq j \} = C_\zeta(\omega, j).$$

**Definition 9.** [29] Let  $(Z, \zeta, \mathcal{I})$  be an  $\mathcal{FITS}$ ,  $j \in I_o$ , and  $\omega \in I^Z$ . Then we define fuzzy operator  $C_\zeta^* : I^Z \times I_o \rightarrow I^Z$  as follows:

$$C_\zeta^*(\omega, j) = \omega \vee \omega_j^*.$$

Now if,  $\mathcal{I} = \mathcal{I}_0$  then  $C_\zeta^*(\omega, j) = \omega \vee \omega_j^* = \omega \vee C_\zeta(\omega, j) = C_\zeta(\omega, j)$  for each  $\omega \in I^Z$ .

**Theorem 1.** [29] Let  $(Z, \zeta, \mathcal{I})$  be an  $\mathcal{FITS}$ ,  $j \in I_o$ , and  $\omega, \rho \in I^Z$ . The operator  $C_\zeta^* : I^Z \times I_o \rightarrow I^Z$  satisfies the following properties:

- (a)  $C_\zeta^*(0, j) = 0$ .
- (b)  $\omega \leq C_\zeta^*(\omega, j) \leq C_\zeta(\omega, j)$ .
- (c) If  $\omega \leq \rho$ , then  $C_\zeta^*(\omega, j) \leq C_\zeta^*(\rho, j)$ .
- (d)  $C_\zeta^*(\omega \vee \rho, j) = C_\zeta^*(\omega, j) \vee C_\zeta^*(\rho, j)$ .
- (e)  $C_\zeta^*(\omega \wedge \rho, j) \leq C_\zeta^*(\omega, j) \wedge C_\zeta^*(\rho, j)$ .
- (f)  $C_\zeta^*(C_\zeta^*(\omega, j), j) = C_\zeta^*(\omega, j)$ .

**Definition 10.** [30, 32] Let  $(Z, \zeta, \mathcal{I})$  be an  $\mathcal{FITS}$  and  $j \in I_o$ . A fuzzy set  $\omega \in I^Z$  is called

- (a)  $j$ -FS $\mathcal{I}$ -open if  $\omega \leq C_\zeta^*(I_\zeta(\omega, j), j)$ ;
- (b)  $j$ -FP $\mathcal{I}$ -open if  $\omega \leq I_\zeta(C_\zeta^*(\omega, j), j)$ ;
- (c)  $j$ -F $\alpha$  $\mathcal{I}$ -open if  $\omega \leq I_\zeta(C_\zeta^*(I_\zeta(\omega, j), j), j)$ ;
- (d)  $j$ -F $\beta$  $\mathcal{I}$ -open if  $\omega \leq C_\zeta(I_\zeta(C_\zeta^*(\omega, j), j), j)$ ;

(e)  $j$ -FS $\beta\mathcal{I}$ -open if  $\omega \leq C_{\zeta}^*(I_{\zeta}(C_{\zeta}^*(\omega, j), j), j)$ ;

(f)  $j$ -FR $\mathcal{I}$ -open if  $\omega = I_{\zeta}(C_{\zeta}^*(\omega, j), j)$ .

**Definition 11.** A fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Y, \mathfrak{S})$  is called Fa $\mathcal{I}$ -continuous (resp. FPT-continuous, FS $\mathcal{I}$ -continuous, and FS $\beta\mathcal{I}$ -continuous) if  $\mathbb{P}^{-1}(\omega)$  is a  $j$ -Fa $\mathcal{I}$ -open (resp.  $j$ -FPT-open,  $j$ -FS $\mathcal{I}$ -open, and  $j$ -FS $\beta\mathcal{I}$ -open) set, for any  $\omega \in I^Y$  and  $\mathfrak{S}(\omega) \geq j$  with  $j \in I_0$ .

Some basic results and concepts that we need in the sequel are found in [19-21,29-32].

### 3. On $j$ -fuzzy $\gamma\mathcal{I}$ -open sets

**Definition 12.** Let  $(Z, \zeta, \mathcal{I})$  be an  $\mathcal{FIT}\mathcal{S}$  and  $j \in I_0$ . A fuzzy set  $\rho \in I^Z$  is called a  $j$ -F $\gamma\mathcal{I}$ -open set if  $\rho \leq C_{\zeta}^*(I_{\zeta}(\rho, j), j) \vee I_{\zeta}(C_{\zeta}^*(\rho, j), j)$ .

**Remark 3.** The complement of  $j$ -F $\gamma\mathcal{I}$ -open sets are  $j$ -F $\gamma\mathcal{I}$ -closed sets.

**Lemma 2.** Each  $j$ -F $\gamma\mathcal{I}$ -open set is  $j$ -F $\gamma$ -open [28].

**Proof.** The proof follows by Theorem 2.1(b) and by Definitions 2.4 and 3.1.  $\square$

**Remark 4.** If we take  $\mathcal{I} = \mathcal{I}_0$ , then  $j$ -F $\gamma\mathcal{I}$ -open set and  $j$ -F $\gamma$ -open set [28] are equivalent.

**Remark 5.** The converse of Lemma 3.1 fails, as can be seen in Example 3.1.

**Example 1.** Define  $\zeta, \mathcal{I} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{\underline{1}, \underline{0}\}, \\ \frac{2}{3}, & \text{if } \psi = \underline{0.7}, \\ \frac{1}{3}, & \text{if } \psi = \underline{0.3}, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(v) = \begin{cases} 1, & \text{if } v = \underline{0}, \\ \frac{2}{3}, & \text{if } \underline{0} < v \leq \underline{0.6}, \\ 0, & \text{otherwise.} \end{cases}$$

Thus,  $\underline{0.6}$  is an  $\frac{1}{3}$ -F $\gamma$ -open set, but it is not  $\frac{1}{3}$ -F $\gamma\mathcal{I}$ -open.

**Proposition 1.** In an  $\mathcal{FIT}\mathcal{S}$   $(Z, \zeta, \mathcal{I})$ , for each  $\omega \in I^Z$  and  $j \in I_0$ . Then

- (a) each  $j$ -FPT-open set [30] is  $j$ -F $\gamma\mathcal{I}$ -open;
- (b) each  $j$ -F $\gamma\mathcal{I}$ -open set is  $j$ -FS $\beta\mathcal{I}$ -open [32];

(c) each  $j$ -FSL-open set [30] is  $j$ -F $\gamma$  $\mathcal{I}$ -open.

**Proof.** (a) If  $\omega$  is an  $j$ -FPI-open set. Then

$$\begin{aligned}\omega &\leq I_{\zeta}(C_{\zeta}^*(\omega, j), j) \\ &\leq I_{\zeta}(C_{\zeta}^*(\omega, j), j) \vee I_{\zeta}(\omega, j) \\ &\leq I_{\zeta}(C_{\zeta}^*(\omega, j), j) \vee C_{\zeta}^*(I_{\zeta}(\omega, j), j).\end{aligned}$$

Thus,  $\omega$  is  $j$ -F $\gamma$  $\mathcal{I}$ -open.

(b) If  $\omega$  is an  $j$ -F $\gamma$  $\mathcal{I}$ -open set. Then

$$\begin{aligned}\omega &\leq C_{\zeta}^*(I_{\zeta}(\omega, j), j) \vee I_{\zeta}(C_{\zeta}^*(\omega, j), j) \\ &\leq C_{\zeta}^*(I_{\zeta}(C_{\zeta}^*(\omega, j), j), j) \vee I_{\zeta}(C_{\zeta}^*(\omega, j), j) \\ &\leq C_{\zeta}^*(I_{\zeta}(C_{\zeta}^*(\omega, j), j), j).\end{aligned}$$

Thus,  $\omega$  is  $j$ -FS $\beta$  $\mathcal{I}$ -open.


(c) If  $\omega$  is an  $j$ -FSL-open set. Then

$$\begin{aligned}\omega &\leq C_{\zeta}^*(I_{\zeta}(\omega, j), j) \\ &\leq C_{\zeta}^*(I_{\zeta}(\omega, j), j) \vee I_{\zeta}(\omega, j) \\ &\leq C_{\zeta}^*(I_{\zeta}(\omega, j), j) \vee I_{\zeta}(C_{\zeta}^*(\omega, j), j).\end{aligned}$$

Thus,  $\omega$  is  $j$ -F $\gamma$  $\mathcal{I}$ -open.

□

**Remark 6.** We have the following diagram from the previous definitions and discussions.



**Remark 7.** The reverse implication of the above diagram does not hold, as demonstrated by the Examples 3.2, 3.3, and 3.4.

**Example 2.** Let  $Z = \{z_1, z_2\}$  and define  $\omega, \rho, \lambda \in I^Z$  as follows:  $\omega = \{\frac{z_1}{0.4}, \frac{z_2}{0.3}\}$ ,  $\rho = \{\frac{z_1}{0.5}, \frac{z_2}{0.4}\}$ ,  $\lambda = \{\frac{z_1}{0.4}, \frac{z_2}{0.5}\}$ . Define  $\zeta, \mathcal{I} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{4}, & \text{if } \psi = \rho, \\ \frac{1}{2}, & \text{if } \psi = \omega, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(\mu) = \begin{cases} 1, & \text{if } \mu = \underline{0}, \\ \frac{1}{2}, & \text{if } \underline{0} < \mu < \underline{0.3}, \\ 0, & \text{otherwise.} \end{cases}$$

Thus,  $\lambda$  is an  $\frac{1}{4}$ -F $\gamma\mathcal{I}$ -open set, but it is not  $\frac{1}{4}$ -FP $\mathcal{I}$ -open.

**Example 3.** Let  $Z = \{z_1, z_2\}$  and define  $\omega, \rho, \lambda \in I^Z$  as follows:  $\omega = \{\frac{z_1}{0.3}, \frac{z_2}{0.2}\}$ ,  $\rho = \{\frac{z_1}{0.7}, \frac{z_2}{0.8}\}$ ,  $\lambda = \{\frac{z_1}{0.5}, \frac{z_2}{0.4}\}$ . Define  $\zeta, \mathcal{I} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{3}, & \text{if } \psi = \omega, \\ \frac{1}{2}, & \text{if } \psi = \rho, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(\mu) = \begin{cases} 1, & \text{if } \mu = \underline{0}, \\ \frac{1}{2}, & \text{if } \underline{0} < \mu < \underline{0.5}, \\ 0, & \text{otherwise.} \end{cases}$$

Thus,  $\lambda$  is an  $\frac{1}{3}$ -F $\gamma\mathcal{I}$ -open set, but it is neither  $\frac{1}{3}$ -FS $\mathcal{I}$ -open nor  $\frac{1}{3}$ -F $\alpha\mathcal{I}$ -open.

**Example 4.** Let  $Z = \{z_1, z_2\}$  and define  $\omega, \lambda \in I^Z$  as follows:  $\omega = \{\frac{z_1}{0.5}, \frac{z_2}{0.4}\}$ ,  $\lambda = \{\frac{z_1}{0.4}, \frac{z_2}{0.5}\}$ . Define  $\zeta, \mathcal{I} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{2}, & \text{if } \psi = \omega, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(\mu) = \begin{cases} 1, & \text{if } \mu = \underline{0}, \\ \frac{1}{2}, & \text{if } \underline{0} < \mu < \underline{0.4}, \\ 0, & \text{otherwise.} \end{cases}$$

Thus,  $\lambda$  is an  $\frac{1}{3}$ -FS $\beta\mathcal{I}$ -open set, but it is not  $\frac{1}{3}$ -F $\gamma\mathcal{I}$ -open.

**Definition 13.** For each  $\omega \in I^Z$  and  $j \in I_0$  in an  $FITS (Z, \zeta, \mathcal{I})$ , we define a fuzzy  $\gamma\mathcal{I}$ -closure operator  $\gamma C_\zeta^* : I^Z \times I_0 \rightarrow I^Z$  as follows:

$$\gamma C_\zeta^*(\omega, j) = \bigwedge \{v \in I^Z : \omega \leq v, v \text{ is } j\text{-F}\gamma\mathcal{I}\text{-closed}\}.$$

**Proposition 2.** For each  $\omega \in I^Z$  and  $j \in I_0$  in an  $FITS (Z, \zeta, \mathcal{I})$ , a fuzzy set  $\omega$  is  $j$ -F $\gamma\mathcal{I}$ -closed iff  $\gamma C_\zeta^*(\omega, j) = \omega$ .

**Proof.** This follows directly from Definition 3.2.

□

**Theorem 2.** For each  $\rho, \omega \in I^Z$  and  $j \in I_0$  in an  $FITS (Z, \zeta, \mathcal{I})$ , a fuzzy  $\gamma\mathcal{I}$ -closure operator  $\gamma C_\zeta^* : I^Z \times I_0 \rightarrow I^Z$  satisfies the following properties.

- $\gamma C_\zeta^*(\underline{0}, j) = \underline{0}$ .
- $\omega \leq \gamma C_\zeta^*(\omega, j) \leq C_\zeta(\omega, j)$ .

- (c)  $\gamma C_{\zeta}^*(\omega, j) \leq \gamma C_{\zeta}^*(\rho, j)$  if  $\omega \leq \rho$ .
- (d)  $\gamma C_{\zeta}^*(\gamma C_{\zeta}^*(\omega, j), j) = \gamma C_{\zeta}^*(\omega, j)$ .
- (e)  $\gamma C_{\zeta}^*(\omega \vee \rho, j) \geq \gamma C_{\zeta}^*(\omega, j) \vee \gamma C_{\zeta}^*(\rho, j)$ .

**Proof.** (a), (b), and (c) are easily proved by Definition 3.2.

(d) From (b) and (c),  $\gamma C_{\zeta}^*(\omega, j) \leq \gamma C_{\zeta}^*(\gamma C_{\zeta}^*(\omega, j), j)$ . Now, we show  $\gamma C_{\zeta}^*(\omega, j) \geq \gamma C_{\zeta}^*(\gamma C_{\zeta}^*(\omega, j), j)$ . If  $\gamma C_{\zeta}^*(\omega, j)$  does not contain  $\gamma C_{\zeta}^*(\gamma C_{\zeta}^*(\omega, j), j)$ , there is  $z \in Z$  and  $s \in (0, 1)$  with

$$\gamma C_{\zeta}^*(\omega, j)(z) < s < \gamma C_{\zeta}^*(\gamma C_{\zeta}^*(\omega, j), j)(z). \quad (1)$$

Since  $\gamma C_{\zeta}^*(\omega, j)(z) < s$ , by Definition 3.2, there is  $\mu \in I^Z$  as a  $j$ -F $\gamma\mathcal{I}$ -closed set and  $\omega \leq \mu$  with  $\gamma C_{\zeta}^*(\omega, j)(z) \leq \mu(z) < s$ . Since  $\omega \leq \mu$ , then  $\gamma C_{\zeta}^*(\omega, j) \leq \mu$ . Again, by the definition of  $\gamma C_{\zeta}^*$ , then  $\gamma C_{\zeta}^*(\gamma C_{\zeta}^*(\omega, j), j) \leq \mu$ . Hence,  $\gamma C_{\zeta}^*(\gamma C_{\zeta}^*(\omega, j), j)(z) \leq \mu(z) < s$ , which is a contradiction for (1). Thus,  $\gamma C_{\zeta}^*(\omega, j) \geq \gamma C_{\zeta}^*(\gamma C_{\zeta}^*(\omega, j), j)$ . Therefore,  $\gamma C_{\zeta}^*(\gamma C_{\zeta}^*(\omega, j), j) = \gamma C_{\zeta}^*(\omega, j)$ .

(e) Since  $\omega \leq \omega \vee \rho$  and  $\rho \leq \omega \vee \rho$ , then by (c),  $\gamma C_{\zeta}^*(\omega, j) \leq \gamma C_{\zeta}^*(\omega \vee \rho, j)$  and  $\gamma C_{\zeta}^*(\rho, j) \leq \gamma C_{\zeta}^*(\omega \vee \rho, j)$ . Hence,  $\gamma C_{\zeta}^*(\omega \vee \rho, j) \geq \gamma C_{\zeta}^*(\omega, j) \vee \gamma C_{\zeta}^*(\rho, j)$ .

□

**Definition 14.** For each  $\omega \in I^Z$  and  $j \in I_0$  in an  $\mathcal{FITS} (Z, \zeta, \mathcal{I})$ , we define a fuzzy  $\gamma$ - $\mathcal{I}$ -interior operator  $\gamma I_{\zeta}^* : I^Z \times I_0 \rightarrow I^Z$  as follows:  $\gamma I_{\zeta}^*(\omega, j) = \bigvee \{v \in I^Z : v \leq \omega, v \text{ is } j\text{-F}\gamma\mathcal{I}\text{-open}\}$ .

**Proposition 3.** Let  $(Z, \zeta, \mathcal{I})$  be an  $\mathcal{FITS}$ ,  $\omega \in I^Z$ , and  $j \in I_0$ . Then

- (a)  $\gamma C_{\zeta}^*(\omega^c, j) = (\gamma I_{\zeta}^*(\omega, j))^c$ ;
- (b)  $\gamma I_{\zeta}^*(\omega^c, j) = (\gamma C_{\zeta}^*(\omega, j))^c$ .

**Proof.** (a) For each  $\omega \in I^Z$ , we have  $\gamma C_{\zeta}^*(\omega^c, j) = \bigwedge \{\rho \in I^Z : \omega^c \leq \rho, \rho \text{ is } j\text{-F}\gamma\mathcal{I}\text{-closed}\} = [\bigvee \{\rho^c \in I^Z : \rho^c \leq \omega, \rho^c \text{ is } j\text{-F}\gamma\mathcal{I}\text{-open}\}]^c = (\gamma I_{\zeta}^*(\omega, j))^c$ .

(b) This is similar to that of (a).

□

**Proposition 4.** For each  $\omega \in I^Z$  and  $j \in I_0$  in an  $\mathcal{FITS} (Z, \zeta, \mathcal{I})$ , a fuzzy set  $\omega$  is  $j$ -F $\gamma\mathcal{I}$ -open iff  $\gamma I_{\zeta}^*(\omega, j) = \omega$ .

**Proof.** This is immediate from Definition 3.3.

□

**Theorem 3.** For each  $\rho, \omega \in I^Z$  and  $j \in I_0$  in an  $\mathcal{FITS} (Z, \zeta, \mathcal{I})$ , a fuzzy  $\gamma$ - $\mathcal{I}$ -interior operator  $\gamma I_{\zeta}^* : I^Z \times I_0 \rightarrow I^Z$  satisfies the following properties.

- (a)  $\gamma I_{\zeta}^*(\mathbf{1}, j) = \mathbf{1}$ .
- (b)  $I_{\zeta}(\omega, j) \leq \gamma I_{\zeta}^*(\omega, j) \leq \omega$ .
- (c)  $\gamma I_{\zeta}^*(\omega, j) \leq \gamma I_{\zeta}^*(\rho, j)$  if  $\omega \leq \rho$ .
- (d)  $\gamma I_{\zeta}^*(\gamma I_{\zeta}^*(\omega, j), j) = \gamma I_{\zeta}^*(\omega, j)$ .

$$(e) \gamma I_{\zeta}^*(\omega, j) \wedge \gamma I_{\zeta}^*(\rho, j) \geq \gamma I_{\zeta}^*(\omega \wedge \rho, j).$$

**Proof.** This can be proven using the same approach as in Theorem 3.1.

□

**Definition 15.** Let  $z_s \in P_s(Z)$ ,  $\omega \in I^Z$ , and  $j \in I_o$ . An  $FITS (Z, \zeta, \mathcal{I})$  is said to be a  $j$ - $F\gamma\mathcal{I}$ -regular space if  $z_s \overline{Q} \omega$  for any  $j$ - $F\gamma\mathcal{I}$ -closed set  $\omega$ , there is  $\mu_i \in I^Z$  with  $\zeta(\mu_i) \geq j$  for  $i = 1, 2$ , such that  $z_s \in \mu_1$ ,  $\omega \leq \mu_2$ , and  $\mu_1 \overline{Q} \mu_2$ .

**Definition 16.** Let  $\omega, \rho \in I^Z$  and  $j \in I_o$ . An  $FITS (Z, \zeta, \mathcal{I})$  is said to be a  $j$ - $F\gamma\mathcal{I}$ -normal space if  $\omega \overline{Q} \rho$  for any  $j$ - $F\gamma\mathcal{I}$ -closed sets  $\omega$  and  $\rho$ , there is  $\mu_i \in I^Z$  with  $\zeta(\mu_i) \geq j$  for  $i = 1, 2$ , such that  $\omega \leq \mu_1$ ,  $\rho \leq \mu_2$ , and  $\mu_1 \overline{Q} \mu_2$ .

**Theorem 4.** Let  $(Z, \zeta, \mathcal{I})$  be an  $FITS$ ,  $z_s \in P_s(Z)$ ,  $\omega \in I^Z$ , and  $j \in I_o$ . Each of the following statements implies the others.

(a)  $(Z, \zeta, \mathcal{I})$  is an  $j$ - $F\gamma\mathcal{I}$ -regular space.

(b) If  $z_s \in \omega$  for any  $j$ - $F\gamma\mathcal{I}$ -open set  $\omega$ , there is  $\mu \in I^Z$  with  $\zeta(\mu) \geq j$ , and

$$z_s \in \mu \leq C_{\zeta}(\mu, j) \leq \omega.$$

(c) If  $z_s \overline{Q} \omega$  for any  $j$ - $F\gamma\mathcal{I}$ -closed set  $\omega$ , there is  $\mu_i \in I^Z$  with  $\zeta(\mu_i) \geq j$  for  $i = 1, 2$ , such that  $z_s \in \mu_1$ ,  $\omega \leq \mu_2$ , and  $C_{\zeta}(\mu_1, j) \overline{Q} C_{\zeta}(\mu_2, j)$ .

**Proof.** (a)  $\Rightarrow$  (b) Let  $z_s \in \omega$  for any  $j$ - $F\gamma\mathcal{I}$ -open set  $\omega$ , then  $z_s \overline{Q} \omega^c$ . Since  $(Z, \zeta, \mathcal{I})$  is  $j$ - $F\gamma\mathcal{I}$ -regular, then there is  $\mu, v \in I^Z$  with  $\zeta(\mu) \geq j$  and  $\zeta(v) \geq j$ , such that  $z_s \in \mu$ ,  $\omega^c \leq v$ , and  $\mu \overline{Q} v$ . Thus,  $z_s \in \mu \leq v^c \leq \omega$ , so  $z_s \in \mu \leq C_{\zeta}(\mu, j) \leq \omega$ .

(b)  $\Rightarrow$  (c) Let  $z_s \overline{Q} \omega$  for any  $j$ - $F\gamma\mathcal{I}$ -closed set  $\omega$ , then  $z_s \in \omega^c$ . By (b), there is  $v \in I^Z$  with  $\zeta(v) \geq j$  and  $z_s \in v \leq C_{\zeta}(v, j) \leq \omega^c$ . Since  $\zeta(v) \geq j$ , then  $v$  is an  $j$ - $F\gamma\mathcal{I}$ -open set and  $z_s \in v$ . Again, by (b), there is  $\mu \in I^Z$  such that  $\zeta(\mu) \geq j$ , and  $z_s \in \mu \leq C_{\zeta}(\mu, j) \leq v \leq C_{\zeta}(v, j) \leq \omega^c$ . Therefore,  $\omega \leq (C_{\zeta}(v, j))^c = I_{\zeta}(v^c, j) \leq v^c$ . Set  $\lambda = I_{\zeta}(v^c, j)$ , and then  $\zeta(\lambda) \geq j$ . Thus,  $C_{\zeta}(\lambda, j) \leq v^c \leq (C_{\zeta}(\mu, j))^c$ . Hence,  $C_{\zeta}(\mu, j) \overline{Q} C_{\zeta}(\lambda, j)$ .

(c)  $\Rightarrow$  (a) This is immediate from Definition 3.4.

□

**Theorem 5.** Let  $(Z, \zeta, \mathcal{I})$  be an  $FITS$ ,  $\omega, \rho \in I^Z$ , and  $j \in I_o$ . Each of the following statements implies the others.

(a)  $(Z, \zeta, \mathcal{I})$  is an  $j$ - $F\gamma\mathcal{I}$ -normal space.

(b) If  $\mu \leq \omega$  for any  $j$ - $F\gamma\mathcal{I}$ -closed set  $\mu$  and  $j$ - $F\gamma\mathcal{I}$ -open set  $\omega$ , there is  $v \in I^Z$  with  $\zeta(v) \geq j$ , and  $\mu \leq v \leq C_{\zeta}(v, j) \leq \omega$ .

(c) If  $\omega \overline{Q} \rho$  for any  $j$ - $F\gamma\mathcal{I}$ -closed sets  $\omega$  and  $\rho$ , there is  $\mu_i \in I^Z$  with  $\zeta(\mu_i) \geq j$  for  $i = 1, 2$ , such that  $\omega \leq \mu_1$ ,  $\rho \leq \mu_2$ , and  $C_{\zeta}(\mu_1, j) \overline{Q} C_{\zeta}(\mu_2, j)$ .

**Proof.** This can be proven using the same approach as in Theorem 3.3.

□

#### 4. Fuzzy $\gamma\mathcal{I}$ -continuity

**Definition 17.** A fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Y, \mathfrak{S})$  is called  $F\gamma\mathcal{I}$ -continuous if  $\mathbb{P}^{-1}(\omega)$  is a  $j$ - $F\gamma\mathcal{I}$ -open set, for any  $\omega \in I^Y$  with  $\mathfrak{S}(\omega) \geq j$  and  $j \in I_0$ .

**Lemma 3.** Every  $F\gamma\mathcal{I}$ -continuity is an  $F\gamma$ -continuity [28].

**Proof.** The proof follows by Lemma 3.1 and by Definitions 2.5 and 4.1. □

**Remark 8.** If we take  $\mathcal{I} = \mathcal{I}_0$ , then  $F\gamma\mathcal{I}$ -continuity and  $F\gamma$ -continuity [28] are equivalent.

**Remark 9.** The converse of Lemma 4.1 fails, as can be seen in Example 4.1.


**Example 5.** Define  $\zeta, \mathcal{I}, \mathfrak{S} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{2}, & \text{if } \psi = \underline{0.Z}, \\ \frac{1}{3}, & \text{if } \psi = \underline{0.3}, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(v) = \begin{cases} 1, & \text{if } v = \underline{0}, \\ \frac{1}{2}, & \text{if } \underline{0} < v \leq \underline{0.6}, \\ 0, & \text{otherwise,} \end{cases}$$

$$\mathfrak{S}(\theta) = \begin{cases} 1, & \text{if } \theta \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{3}, & \text{if } \theta = \underline{0.6}, \\ 0, & \text{otherwise.} \end{cases}$$

Then, the identity fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Z, \mathfrak{S})$  is  $F\gamma$ -continuous, but it is not  $F\gamma\mathcal{I}$ -continuous.

**Remark 10.** We have the following diagram from the previous definitions.



**Remark 11.** The reverse implication of the above diagram does not hold, as demonstrated by the Examples 4.2, 4.3, and 4.4.

**Example 6.** Let  $Z = \{z_1, z_2\}$  and define  $\omega, \rho, \lambda \in I^Z$  as follows:  $\omega = \{\frac{z_1}{0.4}, \frac{z_2}{0.3}\}$ ,  $\rho = \{\frac{z_1}{0.5}, \frac{z_2}{0.4}\}$ ,  $\lambda = \{\frac{z_1}{0.4}, \frac{z_2}{0.5}\}$ . Define  $\zeta, \mathcal{I}, \mathfrak{S} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{\underline{0}, \underline{1}\}, \\ \frac{1}{5}, & \text{if } \psi = \rho, \\ \frac{1}{2}, & \text{if } \psi = \omega, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(\mu) = \begin{cases} 1, & \text{if } \mu = \underline{0}, \\ \frac{1}{2}, & \text{if } \underline{0} < \mu < \underline{0.3}, \\ 0, & \text{otherwise,} \end{cases}$$

$$\mathfrak{S}(\theta) = \begin{cases} 1, & \text{if } \theta \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{5}, & \text{if } \theta = \lambda, \\ 0, & \text{otherwise.} \end{cases}$$

Then, the identity fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Z, \mathfrak{S})$  is  $F\gamma\mathcal{I}$ -continuous, but it is not  $F\mathcal{P}\mathcal{I}$ -continuous.

**Example 7.** Let  $Z = \{z_1, z_2\}$  and define  $\omega, \rho, \lambda \in I^Z$  as follows:  $\omega = \{\frac{z_1}{0.3}, \frac{z_2}{0.2}\}$ ,  $\rho = \{\frac{z_1}{0.7}, \frac{z_2}{0.8}\}$ ,  $\lambda = \{\frac{z_1}{0.5}, \frac{z_2}{0.4}\}$ . Define  $\zeta, \mathcal{I}, \mathfrak{S} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{3}, & \text{if } \psi = \omega, \\ \frac{1}{2}, & \text{if } \psi = \rho, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(\mu) = \begin{cases} 1, & \text{if } \mu = \underline{0}, \\ \frac{1}{2}, & \text{if } \underline{0} < \mu < \underline{0.5}, \\ 0, & \text{otherwise,} \end{cases}$$

$$\mathfrak{S}(\theta) = \begin{cases} 1, & \text{if } \theta \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{3}, & \text{if } \theta = \lambda, \\ 0, & \text{otherwise.} \end{cases}$$

Then, the identity fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Z, \mathfrak{S})$  is  $F\gamma\mathcal{I}$ -continuous, but it is neither  $F\mathcal{S}\mathcal{I}$ -continuous nor  $F\alpha\mathcal{I}$ -continuous.

**Example 8.** Let  $Z = \{z_1, z_2\}$  and define  $\omega, \lambda \in I^Z$  as follows:  $\omega = \{\frac{z_1}{0.5}, \frac{z_2}{0.4}\}$ ,  $\lambda = \{\frac{z_1}{0.4}, \frac{z_2}{0.5}\}$ . Define  $\zeta, \mathcal{I}, \mathfrak{S} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{2}, & \text{if } \psi = \omega, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(\mu) = \begin{cases} 1, & \text{if } \mu = \underline{0}, \\ \frac{1}{2}, & \text{if } \underline{0} < \mu < \underline{0.4}, \\ 0, & \text{otherwise,} \end{cases}$$

$$\mathfrak{S}(\theta) = \begin{cases} 1, & \text{if } \theta \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{3}, & \text{if } \theta = \lambda, \\ 0, & \text{otherwise.} \end{cases}$$

Then, the identity fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \longrightarrow (Z, \mathfrak{S})$  is  $\text{FS}\beta\mathcal{I}$ -continuous, but it is not  $\text{F}\gamma\mathcal{I}$ -continuous.

**Theorem 6.** A fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \longrightarrow (Y, \mathfrak{S})$  is  $\text{F}\gamma\mathcal{I}$ -continuous iff for any  $z_s \in P_s(Z)$  and any  $\psi \in I^Y$  with  $\mathfrak{S}(\psi) \geq j$  containing  $\mathbb{P}(z_s)$ , there is  $\omega \in I^Z$  that is  $j$ - $\text{F}\gamma\mathcal{I}$ -open containing  $z_s$  and  $\mathbb{P}(\omega) \leq \psi$  with  $j \in I_o$ .

**Proof.** ( $\Rightarrow$ ) Let  $z_s \in P_s(Z)$  and  $\psi \in I^Y$  with  $\mathfrak{S}(\psi) \geq j$  containing  $\mathbb{P}(z_s)$ , and hence  $\mathbb{P}^{-1}(\psi) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(\psi), j)$ . Since  $z_s \in \mathbb{P}^{-1}(\psi)$ ,  $z_s \in \gamma I_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) = \omega$  (say). Hence,  $\omega \in I^Z$  is  $j$ - $\text{F}\gamma\mathcal{I}$ -open containing  $z_s$  with  $\mathbb{P}(\omega) \leq \psi$ .

( $\Leftarrow$ ) Let  $z_s \in P_s(Z)$  and  $\psi \in I^Y$  with  $\mathfrak{S}(\psi) \geq j$  containing  $\mathbb{P}(z_s)$ . By the given assumption, there exists  $\omega \in I^Z$  that is  $j$ - $\text{F}\gamma\mathcal{I}$ -open containing  $z_s$  with  $\mathbb{P}(\omega) \leq \psi$ . Thus,  $z_s \in \omega \leq \mathbb{P}^{-1}(\psi)$  and  $z_s \in \gamma I_{\zeta}^*(\mathbb{P}^{-1}(\psi), j)$ . Then,  $\mathbb{P}^{-1}(\psi) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(\psi), j)$ , so  $\mathbb{P}^{-1}(\psi)$  is an  $j$ - $\text{F}\gamma\mathcal{I}$ -open set. Hence,  $\mathbb{P}$  is  $\text{F}\gamma\mathcal{I}$ -continuous.

□

**Theorem 7.** Let  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \longrightarrow (Y, \mathfrak{S})$  be a fuzzy mapping and  $j \in I_o$ . Each of the following statements implies the others for any  $\omega \in I^Z$  and  $\psi \in I^Y$ :

- (a)  $\mathbb{P}$  is  $\text{F}\gamma\mathcal{I}$ -continuous.
- (b)  $\mathbb{P}^{-1}(\psi)$  is  $j$ - $\text{F}\gamma\mathcal{I}$ -closed, for every  $\psi \in I^Y$  with  $\mathfrak{S}(\psi^c) \geq j$ .
- (c)  $\mathbb{P}(\gamma C_{\zeta}^*(\omega, j)) \leq C_{\mathfrak{S}}(\mathbb{P}(\omega), j)$ .
- (d)  $\gamma C_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi, j))$ .
- (e)  $\mathbb{P}^{-1}(I_{\mathfrak{S}}(\psi, j)) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(\psi), j)$ .

**Proof.** (a)  $\Leftrightarrow$  (b) The proof follows from Definition 4.1 and  $\mathbb{P}^{-1}(\psi^c) = (\mathbb{P}^{-1}(\psi))^c$ .

(b)  $\Rightarrow$  (c) Let  $\omega \in I^Z$ . By (b), we obtain  $\mathbb{P}^{-1}(C_{\mathfrak{S}}(\mathbb{P}(\omega), j))$  is  $j$ - $\text{F}\gamma\mathcal{I}$ -closed. Hence,

$$\gamma C_{\zeta}^*(\omega, j) \leq \gamma C_{\zeta}^*(\mathbb{P}^{-1}(\mathbb{P}(\omega)), j) \leq \gamma C_{\zeta}^*(\mathbb{P}^{-1}(C_{\mathfrak{S}}(\mathbb{P}(\omega), j)), j) = \mathbb{P}^{-1}(C_{\mathfrak{S}}(\mathbb{P}(\omega), j)).$$

Therefore,  $\mathbb{P}(\gamma C_{\zeta}^*(\omega, j)) \leq C_{\mathfrak{S}}(\mathbb{P}(\omega), j)$ .

(c)  $\Rightarrow$  (d) Let  $\psi \in I^Y$ . By (c), we obtain  $\mathbb{P}(\gamma C_{\zeta}^*(\mathbb{P}^{-1}(\psi), j)) \leq C_{\mathfrak{S}}(\mathbb{P}(\mathbb{P}^{-1}(\psi)), j) \leq C_{\mathfrak{S}}(\psi, j)$ . Hence,  $\gamma C_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(\mathbb{P}(\gamma C_{\zeta}^*(\mathbb{P}^{-1}(\psi), j))) \leq \mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi, j))$ .

(d)  $\Leftrightarrow$  (e) The proof follows from Proposition 3.3 and  $\mathbb{P}^{-1}(\psi^c) = (\mathbb{P}^{-1}(\psi))^c$ .

(e)  $\Rightarrow$  (a) Let  $\psi \in I^Y$  with  $\mathfrak{S}(\psi) \geq j$ . By (e), we have  $\mathbb{P}^{-1}(\psi) = \mathbb{P}^{-1}(I_{\mathfrak{S}}(\psi, j)) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(\psi)$ . Then,  $\gamma I_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) = \mathbb{P}^{-1}(\psi)$ . Hence,  $\mathbb{P}^{-1}(\psi)$  is  $j$ - $\text{F}\gamma\mathcal{I}$ -open, so  $\mathbb{P}$  is  $\text{F}\gamma\mathcal{I}$ -continuous. □

**Definition 18.** A fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \longrightarrow (Y, \mathfrak{S})$  is called  $\text{FA}\gamma\mathcal{I}$ -continuous if  $\mathbb{P}^{-1}(\omega) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(I_{\mathfrak{S}}(C_{\mathfrak{S}}(\omega, j), j)), j)$ , for any  $\omega \in I^Y$  with  $\mathfrak{S}(\omega) \geq j$  and  $j \in I_o$ .

**Lemma 4.** Every  $F\gamma\mathcal{I}$ -continuity is an  $FA\gamma\mathcal{I}$ -continuity.

**Proof.** This follows directly from Definitions 4.2 and 4.1.

□

**Remark 12.** It is clear from Example 4.5 that the converse of Lemma 4.2 does not apply.

**Example 9.** Let  $Z = \{z_1, z_2, z_3\}$  and define  $\omega, \rho, \lambda \in I^Z$  as follows:  $\omega = \{\frac{z_1}{0.4}, \frac{z_2}{0.2}, \frac{z_3}{0.4}\}$ ,  $\rho = \{\frac{z_1}{0.5}, \frac{z_2}{0.5}, \frac{z_3}{0.4}\}$ ,  $\lambda = \{\frac{z_1}{0.3}, \frac{z_2}{0.2}, \frac{z_3}{0.6}\}$ . Define  $\zeta, \mathcal{I}, \mathfrak{S} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{\underline{1}, \underline{0}\}, \\ \frac{2}{3}, & \text{if } \psi = \omega, \\ \frac{1}{2}, & \text{if } \psi = \rho, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(\mu) = \begin{cases} 1, & \text{if } \mu = \underline{0}, \\ \frac{1}{2}, & \text{if } \underline{0} < \mu \leq \underline{0.6}, \\ 0, & \text{otherwise,} \end{cases}$$

$$\mathfrak{S}(\theta) = \begin{cases} 1, & \text{if } \theta \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{2}, & \text{if } \theta = \lambda, \\ 0, & \text{otherwise.} \end{cases}$$

Then, the identity fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Z, \mathfrak{S})$  is  $FA\gamma\mathcal{I}$ -continuous, but it is not  $F\gamma\mathcal{I}$ -continuous.

**Theorem 8.** A fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Y, \mathfrak{S})$  is  $FA\gamma\mathcal{I}$ -continuous iff for any  $z_s \in P_s(Z)$  and any  $\rho \in I^Y$  with  $\mathfrak{S}(\rho) \geq j$  containing  $\mathbb{P}(z_s)$ , there is  $\omega \in I^Z$  that is  $j$ - $F\gamma\mathcal{I}$ -open containing  $z_s$  and  $\mathbb{P}(\omega) \leq I_{\mathfrak{S}}(C_{\mathfrak{S}}(\rho, j), j)$  with  $j \in I_0$ .

**Proof.** ( $\Rightarrow$ ) Let  $z_s \in P_s(Z)$  and  $\rho \in I^Y$  with  $\mathfrak{S}(\rho) \geq j$  containing  $\mathbb{P}(z_s)$ , and hence

$$\mathbb{P}^{-1}(\rho) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(I_{\mathfrak{S}}(C_{\mathfrak{S}}(\rho, j), j)), j).$$

Since  $z_s \in \mathbb{P}^{-1}(\rho)$ , then  $z_s \in \gamma I_{\zeta}^*(\mathbb{P}^{-1}(I_{\mathfrak{S}}(C_{\mathfrak{S}}(\rho, j), j)), j) = \omega$  (say). Hence,  $\omega \in I^Z$  is  $j$ - $F\gamma\mathcal{I}$ -open containing  $z_s$  and  $\mathbb{P}(\omega) \leq I_{\mathfrak{S}}(C_{\mathfrak{S}}(\rho, j), j)$ .

( $\Leftarrow$ ) Let  $z_s \in P_s(Z)$ ,  $\rho \in I^Y$ , and  $\mathfrak{S}(\rho) \geq j$  with  $z_s \in \mathbb{P}^{-1}(\rho)$ . By the given assumption, there exists  $\omega \in I^Z$  that is  $j$ - $F\gamma\mathcal{I}$ -open containing  $z_s$  and  $\mathbb{P}(\omega) \leq I_{\mathfrak{S}}(C_{\mathfrak{S}}(\rho, j), j)$ . Hence,  $z_s \in \omega \leq \mathbb{P}^{-1}(I_{\mathfrak{S}}(C_{\mathfrak{S}}(\rho, j), j))$  and

$$z_s \in \gamma I_{\zeta}^*(\mathbb{P}^{-1}(I_{\mathfrak{S}}(C_{\mathfrak{S}}(\rho, j), j)), j).$$

Then,  $\mathbb{P}^{-1}(\rho) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(I_{\mathfrak{S}}(C_{\mathfrak{S}}(\rho, j), j)), j)$ . Thus,  $\mathbb{P}$  is  $FA\gamma\mathcal{I}$ -continuous.

□

**Theorem 9.** Let  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Y, \mathfrak{S})$  be a fuzzy mapping,  $\psi \in I^Y$ , and  $j \in I_0$ . Each of the following statements implies the others.

- (a)  $\mathbb{P}$  is  $\text{FA}\gamma\mathcal{I}$ -continuous.
- (b)  $\mathbb{P}^{-1}(\psi)$  is  $j\text{-F}\gamma\mathcal{I}$ -open, for each  $j\text{-FR}$ -open set  $\psi$ .
- (c)  $\mathbb{P}^{-1}(\psi)$  is  $j\text{-F}\gamma\mathcal{I}$ -closed, for each  $j\text{-FR}$ -closed set  $\psi$ .
- (d)  $\gamma C_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi, j))$ , for each  $j\text{-F}\gamma$ -open set  $\psi$ .
- (e)  $\gamma C_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi, j))$ , for each  $j\text{-FS}$ -open set  $\psi$ .

**Proof.** (a)  $\Rightarrow$  (b) Let  $z_s \in P_s(Z)$  with  $\psi \in I^Y$  be a  $j\text{-FR}$ -open set and  $z_s \in \mathbb{P}^{-1}(\psi)$ . Thus, by (a), there is  $\omega \in I^Z$  that is  $j\text{-F}\gamma\mathcal{I}$ -open and  $z_s \in \omega$  with  $\mathbb{P}(\omega) \leq I_{\mathfrak{S}}(C_{\mathfrak{S}}(\psi, j), j)$ . Then,  $\omega \leq \mathbb{P}^{-1}(I_{\mathfrak{S}}(C_{\mathfrak{S}}(\psi, j), j)) = \mathbb{P}^{-1}(\psi)$  and  $z_s \in \gamma I_{\zeta}^*(\mathbb{P}^{-1}(\psi), j)$ . Hence,  $\mathbb{P}^{-1}(\psi) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(\psi), j)$ , so  $\mathbb{P}^{-1}(\psi)$  is  $j\text{-F}\gamma\mathcal{I}$ -open.

(b)  $\Rightarrow$  (c) If  $\psi \in I^Y$  is  $j\text{-FR}$ -closed, hence by (b),  $\mathbb{P}^{-1}(\psi^c) = (\mathbb{P}^{-1}(\psi))^c$  is  $j\text{-F}\gamma\mathcal{I}$ -open. Thus,  $\mathbb{P}^{-1}(\psi)$  is  $j\text{-F}\gamma\mathcal{I}$ -closed.

(c)  $\Rightarrow$  (d) If  $\psi \in I^Y$  is  $j\text{-F}\gamma$ -open and since  $C_{\mathfrak{S}}(\psi, j)$  is  $j\text{-FR}$ -closed, then by (c),  $\mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi, j))$  is  $j\text{-F}\gamma\mathcal{I}$ -closed. Since  $\mathbb{P}^{-1}(\psi) \leq \mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi, j))$ , thus

$$\gamma C_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi, j)).$$

(d)  $\Rightarrow$  (e) The proof follows by the fact that each  $j\text{-FS}$ -open set is  $j\text{-F}\gamma$ -open.

(e)  $\Rightarrow$  (c) If  $\psi \in I^Y$  is  $j\text{-FR}$ -closed, and then  $\psi$  is  $j\text{-FS}$ -open. By (e),

$$\gamma C_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi, j)) = \mathbb{P}^{-1}(\psi).$$

Thus,  $\mathbb{P}^{-1}(\psi)$  is  $j\text{-F}\gamma\mathcal{I}$ -closed.

(c)  $\Rightarrow$  (a) If  $z_s \in P_s(Z)$  with  $\psi \in I^Y$  and  $\mathfrak{S}(\psi) \geq j$  such that  $z_s \in \mathbb{P}^{-1}(\psi)$ , and then  $z_s \in \mathbb{P}^{-1}(I_{\mathfrak{S}}(C_{\mathfrak{S}}(\psi, j), j))$ . Since  $[I_{\mathfrak{S}}(C_{\mathfrak{S}}(\psi, j), j)]^c$  is  $j\text{-FR}$ -closed, by (c),  $\mathbb{P}^{-1}([I_{\mathfrak{S}}(C_{\mathfrak{S}}(\psi, j), j)]^c)$  is  $j\text{-F}\gamma\mathcal{I}$ -closed. Thus,  $\mathbb{P}^{-1}(I_{\mathfrak{S}}(C_{\mathfrak{S}}(\psi, j), j))$  is  $j\text{-F}\gamma\mathcal{I}$ -open and  $z_s \in \gamma I_{\zeta}^*(\mathbb{P}^{-1}(I_{\mathfrak{S}}(C_{\mathfrak{S}}(\psi, j), j)), j)$ . Then,  $\mathbb{P}^{-1}(\psi) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(I_{\mathfrak{S}}(C_{\mathfrak{S}}(\psi, j), j)), j)$ . Hence,  $\mathbb{P}$  is  $\text{FA}\gamma\mathcal{I}$ -continuous.

□

**Definition 19.** A fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Y, \mathfrak{S})$  is called  $\text{FW}\gamma\mathcal{I}$ -continuous if  $\mathbb{P}^{-1}(\omega) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(C_{\mathfrak{S}}(\omega, j)), j)$ , for any  $\omega \in I^Y$  with  $\mathfrak{S}(\omega) \geq j$  and  $j \in I_0$ .

**Lemma 5.** Every  $\text{F}\gamma\mathcal{I}$ -continuity is an  $\text{FW}\gamma\mathcal{I}$ -continuity.

**Proof.** This follows directly from Definitions 4.3 and 4.1.

□

**Remark 13.** It is clear from Example 4.6 that the converse of Lemma 4.3 does not apply.

**Example 10.** Let  $Z = \{z_1, z_2, z_3\}$  and define  $\omega, \rho, \lambda \in I^Z$  as follows:  $\omega = \{\frac{z_1}{0.4}, \frac{z_2}{0.2}, \frac{z_3}{0.4}\}$ ,  $\rho = \{\frac{z_1}{0.5}, \frac{z_2}{0.5}, \frac{z_3}{0.4}\}$ ,  $\lambda = \{\frac{z_1}{0.3}, \frac{z_2}{0.2}, \frac{z_3}{0.6}\}$ . Define  $\zeta, \mathcal{I}, \mathfrak{S} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{0, \underline{1}\}, \\ \frac{1}{3}, & \text{if } \psi = \omega, \\ \frac{1}{2}, & \text{if } \psi = \rho, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(\mu) = \begin{cases} 1, & \text{if } \mu = 0, \\ \frac{1}{2}, & \text{if } 0 < \mu \leq 0.6, \\ 0, & \text{otherwise,} \end{cases}$$

$$\mathfrak{S}(\theta) = \begin{cases} 1, & \text{if } \theta \in \{1, 0\}, \\ \frac{1}{3}, & \text{if } \theta = \lambda, \\ 0, & \text{otherwise.} \end{cases}$$

Then, the identity fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Z, \mathfrak{S})$  is  $\text{FW}\gamma\mathcal{I}$ -continuous, but it is not  $\text{F}\gamma\mathcal{I}$ -continuous.

**Theorem 10.** A fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Y, \mathfrak{S})$  is  $\text{FW}\gamma\mathcal{I}$ -continuous iff for any  $z_s \in P_s(Z)$  and any  $\rho \in I^Y$  with  $\mathfrak{S}(\rho) \geq j$  containing  $\mathbb{P}(z_s)$ , there is  $\omega \in I^Z$  that is  $j$ - $\text{F}\gamma\mathcal{I}$ -open containing  $z_s$  and  $\mathbb{P}(\omega) \leq C_{\mathfrak{S}}(\rho, j)$  with  $j \in I_0$ .

**Proof.** ( $\Rightarrow$ ) Let  $z_s \in P_s(Z)$  and  $\rho \in I^Y$  with  $\mathfrak{S}(\rho) \geq j$  containing  $\mathbb{P}(z_s)$ , and hence

$$\mathbb{P}^{-1}(\rho) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(C_{\mathfrak{S}}(\rho, j)), j).$$

Since  $z_s \in \mathbb{P}^{-1}(\rho)$ , then  $z_s \in \gamma I_{\zeta}^*(\mathbb{P}^{-1}(C_{\mathfrak{S}}(\rho, j)), j) = \omega$  (say). Thus,  $\omega \in I^Z$  is  $j$ - $\text{F}\gamma\mathcal{I}$ -open containing  $z_s$  and  $\mathbb{P}(\omega) \leq C_{\mathfrak{S}}(\rho, j)$ .

( $\Leftarrow$ ) Let  $z_s \in P_s(Z)$ ,  $\rho \in I^Y$  and  $\mathfrak{S}(\rho) \geq j$  with  $z_s \in \mathbb{P}^{-1}(\rho)$ . By the given assumption, there exists  $\omega \in I^Z$  that is  $j$ - $\text{F}\gamma\mathcal{I}$ -open containing  $z_s$  with  $\mathbb{P}(\omega) \leq C_{\mathfrak{S}}(\rho, j)$ . Hence,  $z_s \in \omega \leq \mathbb{P}^{-1}(C_{\mathfrak{S}}(\rho, j))$  and  $z_s \in \gamma I_{\zeta}^*(\mathbb{P}^{-1}(C_{\mathfrak{S}}(\rho, j)), j)$ . Thus,  $\mathbb{P}^{-1}(\rho) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(C_{\mathfrak{S}}(\rho, j)), j)$ . Therefore,  $\mathbb{P}$  is  $\text{FW}\gamma\mathcal{I}$ -continuous.  $\square$

**Theorem 11.** Let  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Y, \mathfrak{S})$  be a fuzzy mapping,  $\psi \in I^Y$ , and  $j \in I_0$ . Each of the following statements implies the others.

- (a)  $\mathbb{P}$  is  $\text{FW}\gamma\mathcal{I}$ -continuous.
- (b)  $\mathbb{P}^{-1}(\psi) \geq \gamma C_{\zeta}^*(\mathbb{P}^{-1}(I_{\mathfrak{S}}(\psi, j)), j)$ , if  $\mathfrak{S}(\psi^c) \geq j$ .
- (c)  $\gamma I_{\zeta}^*(\mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi, j)), j) \geq \mathbb{P}^{-1}(I_{\mathfrak{S}}(\psi, j))$ .
- (d)  $\gamma C_{\zeta}^*(\mathbb{P}^{-1}(I_{\mathfrak{S}}(\psi, j)), j) \leq \mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi, j))$ .

**Proof.** (a)  $\Leftrightarrow$  (b) This follows directly from Definition 4.3 and Proposition 3.3.

(b)  $\Rightarrow$  (c) Let  $\psi \in I^Y$ . Then by (b),

$$\gamma C_{\zeta}^*(\mathbb{P}^{-1}(I_{\mathfrak{S}}(C_{\mathfrak{S}}(\psi^c, j)), j)), j) \leq \mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi^c, j)).$$

Thus,  $\mathbb{P}^{-1}(I_{\mathfrak{S}}(\psi, j)) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi, j)), j)$ .

(c)  $\Leftrightarrow$  (d) This follows directly from Proposition 3.3.

(d)  $\Rightarrow$  (a) Let  $\psi \in I^Y$  with  $\mathfrak{S}(\psi) \geq j$ . Then by (d),  $\gamma C_{\zeta}^*(\mathbb{P}^{-1}(I_{\mathfrak{S}}(\psi^c, j)), j) \leq \mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi^c, j)) = \mathbb{P}^{-1}(\psi^c)$ . Hence,  $\mathbb{P}^{-1}(\psi) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(C_{\mathfrak{S}}(\psi, j)), j)$ , so  $\mathbb{P}$  is  $\text{FW}\gamma\mathcal{I}$ -continuous.  $\square$

**Lemma 6.** Every  $\text{FA}\gamma\mathcal{I}$ -continuity is an  $\text{FW}\gamma\mathcal{I}$ -continuity.

**Proof.** This follows directly from Definitions 4.3 and 4.2.  $\square$

**Remark 14.** It is clear from Example 4.7 that the converse of Lemma 4.4 does not apply.

**Example 11.** Let  $Z = \{z_1, z_2, z_3\}$  and define  $\omega, \lambda, \rho \in I^Z$  as follows:  $\omega = \{\frac{z_1}{0.6}, \frac{z_2}{0.2}, \frac{z_3}{0.4}\}$ ,  $\lambda = \{\frac{z_1}{0.3}, \frac{z_2}{0.2}, \frac{z_3}{0.5}\}$ ,  $\rho = \{\frac{z_1}{0.3}, \frac{z_2}{0.2}, \frac{z_3}{0.4}\}$ . Define  $\zeta, \mathcal{I}, \mathfrak{S} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{0, 1\}, \\ \frac{1}{4}, & \text{if } \psi = \omega, \\ \frac{1}{2}, & \text{if } \psi = \rho, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(\mu) = \begin{cases} 1, & \text{if } \mu = 0, \\ \frac{1}{2}, & \text{if } 0 < \mu \leq 0.5, \\ 0, & \text{otherwise,} \end{cases}$$

$$\mathfrak{S}(\theta) = \begin{cases} 1, & \text{if } \theta \in \{1, 0\}, \\ \frac{1}{4}, & \text{if } \theta = \lambda, \\ 0, & \text{otherwise.} \end{cases}$$

Then, the identity fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Z, \mathfrak{S})$  is  $\text{FW}\gamma\mathcal{I}$ -continuous, but it is not  $\text{FA}\gamma\mathcal{I}$ -continuous.

**Remark 15.** We have the following diagram from the previous definitions and discussions.

$$\text{Ff}\mathcal{I}\text{-continuity} \rightarrow \text{FAf}\mathcal{I}\text{-continuity} \rightarrow \text{FWf}\mathcal{I}\text{-continuity}$$

**Proposition 5.** Let  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (X, \eta)$  and  $\mathbb{Y} : (X, \eta) \rightarrow (Y, \mathfrak{S})$  be two fuzzy mappings. Then the composition  $\mathbb{Y} \circ \mathbb{P}$  is  $\text{FA}\gamma\mathcal{I}$ -continuous if  $\mathbb{P}$  is  $\text{F}\gamma\mathcal{I}$ -continuous and  $\mathbb{Y}$  is fuzzy continuous.

**Proof.** This follows directly from Definitions 2.2, 4.1, and 4.2.  $\square$

## 5. Fuzzy $\gamma\mathcal{I}$ -irresoluteness

**Definition 20.** A fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Y, \mathfrak{S})$  is called  $\text{F}\gamma\mathcal{I}$ -irresolute if  $\mathbb{P}^{-1}(\omega)$  is a  $j$ - $\text{F}\gamma\mathcal{I}$ -open set, for any  $j$ - $\text{F}\gamma$ -open set  $\omega \in I^Y$  with  $j \in I_0$ .

**Lemma 7.** Every  $F\gamma\mathcal{I}$ -irresolute mapping is  $F\gamma\mathcal{I}$ -continuous.

**Proof.** This follows directly from Definitions 4.1, 5.1, and Remark 2.1.  $\square$

**Remark 16.** It is clear from Example 5.1 that the converse of Lemma 5.1 does not apply.

**Example 12.** Let  $Z = \{z_1, z_2\}$  and define  $\lambda, \rho \in I^Z$  as follows:  $\lambda = \{\frac{z_1}{0.5}, \frac{z_2}{0.5}\}$ ,  $\rho = \{\frac{z_1}{0.5}, \frac{z_2}{0.4}\}$ . Define  $\zeta, \mathcal{I}, \mathfrak{S} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{\underline{0}, \underline{1}\}, \\ \frac{1}{2}, & \text{if } \psi = \rho, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(\mu) = \begin{cases} 1, & \text{if } \mu = \underline{0}, \\ \frac{1}{2}, & \text{if } \underline{0} < \mu < \underline{0.5}, \\ 0, & \text{otherwise,} \end{cases}$$

$$\mathfrak{S}(\theta) = \begin{cases} 1, & \text{if } \theta \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{3}, & \text{if } \theta = \lambda, \\ 0, & \text{otherwise.} \end{cases}$$

Then, the identity fuzzy mapping  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Z, \mathfrak{S})$  is  $F\gamma\mathcal{I}$ -continuous, but it is not  $F\gamma\mathcal{I}$ -irresolute.

**Theorem 12.** Let  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (Y, \mathfrak{S})$  be a fuzzy mapping and  $j \in I_0$ . Each of the following statements implies the others for any  $\omega \in I^Z$  and  $\psi \in I^Y$ :

- (a)  $\mathbb{P}$  is  $F\gamma\mathcal{I}$ -irresolute.
- (b)  $\mathbb{P}^{-1}(\psi)$  is  $j$ - $F\gamma\mathcal{I}$ -closed, for each  $j$ - $F\gamma$ -closed set  $\psi$ .
- (c)  $\mathbb{P}(\gamma C_{\zeta}^*(\omega, j)) \leq \gamma C_{\mathfrak{S}}(\mathbb{P}(\omega), j)$ .
- (d)  $\gamma C_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(\gamma C_{\mathfrak{S}}(\psi, j))$ .
- (e)  $\mathbb{P}^{-1}(\gamma I_{\mathfrak{S}}(\psi, j)) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(\psi), j)$ .

**Proof.** (a)  $\Leftrightarrow$  (b) This follows directly from Definition 5.1 and  $\mathbb{P}^{-1}(\psi^c) = (\mathbb{P}^{-1}(\psi))^c$ .

(b)  $\Rightarrow$  (c) Let  $\omega \in I^Z$ . By (b),  $\mathbb{P}^{-1}(\gamma C_{\mathfrak{S}}(\mathbb{P}(\omega), j))$  is  $j$ - $F\gamma\mathcal{I}$ -closed. Then,

$$\gamma C_{\zeta}^*(\omega, j) \leq \gamma C_{\zeta}^*(\mathbb{P}^{-1}(\mathbb{P}(\omega)), j) \leq \gamma C_{\zeta}^*(\mathbb{P}^{-1}(\gamma C_{\mathfrak{S}}(\mathbb{P}(\omega), j)), j) = \mathbb{P}^{-1}(\gamma C_{\mathfrak{S}}(\mathbb{P}(\omega), j)).$$

Thus,  $\mathbb{P}(\gamma C_{\zeta}^*(\omega, j)) \leq \gamma C_{\mathfrak{S}}(\mathbb{P}(\omega), j)$ .

(c)  $\Rightarrow$  (d) Let  $\psi \in I^Y$ . By (c),  $\mathbb{P}(\gamma C_{\zeta}^*(\mathbb{P}^{-1}(\psi), j)) \leq \gamma C_{\mathfrak{S}}(\mathbb{P}(\mathbb{P}^{-1}(\psi)), j) \leq \gamma C_{\mathfrak{S}}(\psi, j)$ . Hence,  $\gamma C_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(\mathbb{P}(\gamma C_{\zeta}^*(\mathbb{P}^{-1}(\psi), j))) \leq \mathbb{P}^{-1}(\gamma C_{\mathfrak{S}}(\psi, j))$ .

(d)  $\Leftrightarrow$  (e) This follows directly from Proposition 3.3 and  $\mathbb{P}^{-1}(\psi^c) = (\mathbb{P}^{-1}(\psi))^c$ .

(e)  $\Rightarrow$  (a) Let  $\psi \in I^Y$  be a  $j$ - $F\gamma$ -open set. By (e),

$$\mathbb{P}^{-1}(\psi) = \mathbb{P}^{-1}(\gamma I_{\mathfrak{S}}(\psi, j)) \leq \gamma I_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(\psi).$$

Then,  $\gamma I_{\zeta}^*(\mathbb{P}^{-1}(\psi), j) = \mathbb{P}^{-1}(\psi)$ . Thus,  $\mathbb{P}^{-1}(\psi)$  is  $j$ - $F\gamma\mathcal{I}$ -open, so  $\mathbb{P}$  is  $F\gamma\mathcal{I}$ -irresolute.  $\square$

**Proposition 6.** Let  $\mathbb{P} : (Z, \zeta, \mathcal{I}) \rightarrow (X, \eta)$  and  $\mathbb{Y} : (X, \eta) \rightarrow (Y, \mathfrak{S})$  be two fuzzy mappings. Then the composition  $\mathbb{Y} \circ \mathbb{P}$  is  $F\gamma\mathcal{I}$ -irresolute (resp.  $F\gamma\mathcal{I}$ -continuous) if  $\mathbb{P}$  is  $F\gamma\mathcal{I}$ -irresolute and  $\mathbb{Y}$  is  $F\gamma$ -irresolute (resp. fuzzy continuous).

**Proof.** This follows directly from Definitions 2.2, 4.1, and 5.1.  $\square$

**Definition 21.** A fuzzy mapping  $\mathbb{P} : (Z, \zeta) \rightarrow (Y, \mathfrak{S}, \mathcal{I})$  is called  $F\gamma\mathcal{I}$ -open if  $\mathbb{P}(\omega)$  is a  $j$ - $F\gamma\mathcal{I}$ -open set, for any  $\omega \in I^Z$  with  $\zeta(\omega) \geq j$  and  $j \in I_0$ .

**Definition 22.** A fuzzy mapping  $\mathbb{P} : (Z, \zeta) \rightarrow (Y, \mathfrak{S}, \mathcal{I})$  is called  $F\gamma\mathcal{I}$ -irresolute open if  $\mathbb{P}(\omega)$  is a  $j$ - $F\gamma\mathcal{I}$ -open set, for any  $j$ - $F\gamma$ -open set  $\omega \in I^Z$  with  $j \in I_0$ .

**Lemma 8.** Each  $F\gamma\mathcal{I}$ -irresolute open mapping is  $F\gamma\mathcal{I}$ -open.

**Proof.** This follows directly from Definitions 5.2, 5.3, and Remark 2.1.  $\square$

**Remark 17.** It is clear from Example 5.2 that the converse of Lemma 5.2 does not apply.

**Example 13.** Let  $Z = \{z_1, z_2\}$  and define  $\omega, \lambda \in I^Z$  as follows:  $\omega = \{\frac{z_1}{0.5}, \frac{z_2}{0.5}\}$ ,  $\lambda = \{\frac{z_1}{0.5}, \frac{z_2}{0.4}\}$ . Define  $\zeta, \mathfrak{S}, \mathcal{I} : I^Z \rightarrow I$  as follows:

$$\zeta(\psi) = \begin{cases} 1, & \text{if } \psi \in \{\underline{0}, \underline{1}\}, \\ \frac{1}{5}, & \text{if } \psi = \omega, \\ 0, & \text{otherwise,} \end{cases} \quad \mathcal{I}(\mu) = \begin{cases} 1, & \text{if } \mu = \underline{0}, \\ \frac{1}{2}, & \text{if } \underline{0} < \mu < \underline{0.5}, \\ 0, & \text{otherwise,} \end{cases}$$

$$\mathfrak{S}(\theta) = \begin{cases} 1, & \text{if } \theta \in \{\underline{1}, \underline{0}\}, \\ \frac{1}{5}, & \text{if } \theta = \lambda, \\ 0, & \text{otherwise.} \end{cases}$$

Then, the identity fuzzy mapping  $\mathbb{P} : (Z, \zeta) \rightarrow (Z, \mathfrak{S}, \mathcal{I})$  is  $F\gamma\mathcal{I}$ -open, but it is not  $F\gamma\mathcal{I}$ -irresolute open.

**Theorem 13.** Let  $\mathbb{P} : (Z, \zeta) \rightarrow (Y, \mathfrak{S}, \mathcal{I})$  be a fuzzy mapping and  $j \in I_0$ . Each of the following statements implies the others for any  $\omega \in I^Z$  and  $\psi \in I^Y$ :

(a)  $\mathbb{P}$  is  $F\gamma\mathcal{I}$ -open.

$$(b) \mathbb{P}(I_{\zeta}(\omega, j)) \leq \gamma I_{\mathfrak{S}}^*(\mathbb{P}(\omega), j).$$

$$(c) I_{\zeta}(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(\gamma I_{\mathfrak{S}}^*(\psi, j)).$$

(d) For each  $\psi$  and each  $\omega$  with  $\zeta(\omega^c) \geq j$  and  $\mathbb{P}^{-1}(\psi) \leq \omega$ , there is  $\mu \in I^Y$  is  $j$ -F $\gamma\mathcal{I}$ -closed with  $\psi \leq \mu$  and  $\mathbb{P}^{-1}(\mu) \leq \omega$ .

**Proof.** (a)  $\Rightarrow$  (b) Since  $\mathbb{P}(I_{\zeta}(\omega, j)) \leq \mathbb{P}(\omega)$ , hence by (a),  $\mathbb{P}(I_{\zeta}(\omega, j))$  is  $j$ -F $\gamma\mathcal{I}$ -open. Thus,

$$\mathbb{P}(I_{\zeta}(\omega, j)) \leq \gamma I_{\mathfrak{S}}^*(\mathbb{P}(\omega), j).$$

(b)  $\Rightarrow$  (c) Set  $\omega = \mathbb{P}^{-1}(\psi)$ , and hence by (b),  $\mathbb{P}(I_{\zeta}(\mathbb{P}^{-1}(\psi), j)) \leq \gamma I_{\mathfrak{S}}^*(\mathbb{P}(\mathbb{P}^{-1}(\psi)), j) \leq \gamma I_{\mathfrak{S}}^*(\psi, j)$ . Then,  $I_{\zeta}(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(\gamma I_{\mathfrak{S}}^*(\psi, j))$ .

(c)  $\Rightarrow$  (d) Let  $\psi \in I^Y$  and  $\omega \in I^Z$  with  $\zeta(\omega^c) \geq j$  such that  $\mathbb{P}^{-1}(\psi) \leq \omega$ . Since  $\omega^c \leq \mathbb{P}^{-1}(\psi^c)$ ,  $\omega^c = I_{\zeta}(\omega^c, j) \leq I_{\zeta}(\mathbb{P}^{-1}(\psi^c), j)$ . Hence by (c),  $\omega^c \leq I_{\zeta}(\mathbb{P}^{-1}(\psi^c), j) \leq \mathbb{P}^{-1}(\gamma I_{\mathfrak{S}}^*(\psi^c, j))$ . Then, we have

$$\omega \geq (\mathbb{P}^{-1}(\gamma I_{\mathfrak{S}}^*(\psi^c, j)))^c = \mathbb{P}^{-1}(\gamma C_{\mathfrak{S}}^*(\psi, j)).$$

Thus,  $\gamma C_{\mathfrak{S}}^*(\psi, j) \in I^Y$  is  $j$ -F $\gamma\mathcal{I}$ -closed with  $\psi \leq \gamma C_{\mathfrak{S}}^*(\psi, j)$  and  $\mathbb{P}^{-1}(\gamma C_{\mathfrak{S}}^*(\psi, j)) \leq \omega$ .

(d)  $\Rightarrow$  (a) Let  $\sigma \in I^Z$  with  $\zeta(\sigma) \geq j$ . Set  $\psi = (\mathbb{P}(\sigma))^c$  and  $\omega = \sigma^c$ ,  $\mathbb{P}^{-1}(\psi) = \mathbb{P}^{-1}((\mathbb{P}(\sigma))^c) \leq \omega$ . By (d), there exists  $\mu \in I^Y$  is  $j$ -F $\gamma\mathcal{I}$ -closed with  $\psi \leq \mu$  and  $\mathbb{P}^{-1}(\mu) \leq \omega = \sigma^c$ . Then,  $\mathbb{P}(\sigma) \leq \mathbb{P}(\mathbb{P}^{-1}(\mu^c)) \leq \mu^c$ . Since  $\psi \leq \mu$ ,  $\mathbb{P}(\sigma) = \psi^c \geq \mu^c$ . Thus,  $\mathbb{P}(\sigma) = \mu^c$ , so  $\mathbb{P}(\sigma)$  is a  $j$ -F $\gamma\mathcal{I}$ -open set. Hence,  $\mathbb{P}$  is F $\gamma\mathcal{I}$ -open.

□

**Theorem 14.** Let  $\mathbb{P} : (Z, \zeta) \longrightarrow (Y, \mathfrak{S}, \mathcal{I})$  be a fuzzy mapping and  $j \in I_{\circ}$ . Each of the following statements implies the others for any  $\omega \in I^Z$  and  $\psi \in I^Y$ :

(a)  $\mathbb{P}$  is F $\gamma\mathcal{I}$ -irresolute open.

$$(b) \mathbb{P}(\gamma I_{\zeta}(\omega, j)) \leq \gamma I_{\mathfrak{S}}^*(\mathbb{P}(\omega), j).$$

$$(c) \gamma I_{\zeta}(\mathbb{P}^{-1}(\psi), j) \leq \mathbb{P}^{-1}(\gamma I_{\mathfrak{S}}^*(\psi, j)).$$

(d) For each  $\psi$  and each  $\omega$  is an  $j$ -F $\gamma$ -closed set with  $\mathbb{P}^{-1}(\psi) \leq \omega$ , there is  $\mu \in I^Y$  is  $j$ -F $\gamma\mathcal{I}$ -closed with  $\psi \leq \mu$  and  $\mathbb{P}^{-1}(\mu) \leq \omega$ .

**Proof.** This can be proven using the same approach as in Theorem 5.2.

□

**Definition 23.** A fuzzy mapping  $\mathbb{P} : (Z, \zeta) \longrightarrow (Y, \mathfrak{S}, \mathcal{I})$  is called F $\gamma\mathcal{I}$ -closed if  $\mathbb{P}(\omega)$  is a  $j$ -F $\gamma\mathcal{I}$ -closed set, for any  $\omega \in I^Z$  with  $\zeta(\omega^c) \geq j$  and  $j \in I_{\circ}$ .

**Definition 24.** A fuzzy mapping  $\mathbb{P} : (Z, \zeta) \longrightarrow (Y, \mathfrak{S}, \mathcal{I})$  is called F $\gamma\mathcal{I}$ -irresolute closed if  $\mathbb{P}(\omega)$  is a  $j$ -F $\gamma\mathcal{I}$ -closed set, for any  $j$ -F $\gamma$ -closed set  $\omega \in I^Z$  and  $j \in I_{\circ}$ .

**Lemma 9.** Each  $F\gamma\mathcal{I}$ -irresolute closed mapping is  $F\gamma\mathcal{I}$ -closed.

**Proof.** This follows directly from Definitions 5.4 and 5.5.  $\square$

**Theorem 15.** Let  $\mathbb{P} : (Z, \zeta) \longrightarrow (Y, \mathfrak{S}, \mathcal{I})$  be a fuzzy mapping and  $j \in I_0$ . Each of the following statements implies the others for any  $\omega \in I^Z$  and  $\psi \in I^Y$ :

(a)  $\mathbb{P}$  is  $F\gamma\mathcal{I}$ -closed.

(b)  $\gamma C_{\mathfrak{S}}^*(\mathbb{P}(\omega), j) \leq \mathbb{P}(C_{\zeta}(\omega, j))$ .

(c)  $\mathbb{P}^{-1}(\gamma C_{\mathfrak{S}}^*(\psi, j)) \leq C_{\zeta}(\mathbb{P}^{-1}(\psi), j)$ .

(d) For each  $\psi$  and each  $\omega$  with  $\zeta(\omega) \geq j$  and  $\mathbb{P}^{-1}(\psi) \leq \omega$ , there is  $\mu \in I^Y$  is  $j$ - $F\gamma\mathcal{I}$ -open with  $\psi \leq \mu$  and  $\mathbb{P}^{-1}(\mu) \leq \omega$ .

**Proof.** This can be proven using the same approach as in Theorem 5.2.

$\square$

**Theorem 16.** Let  $\mathbb{P} : (Z, \zeta) \longrightarrow (Y, \mathfrak{S}, \mathcal{I})$  be a fuzzy mapping and  $j \in I_0$ . Each of the following statements implies the others for any  $\omega \in I^Z$  and  $\psi \in I^Y$ :

(a)  $\mathbb{P}$  is  $F\gamma\mathcal{I}$ -irresolute closed.

(b)  $\gamma C_{\mathfrak{S}}^*(\mathbb{P}(\omega), j) \leq \mathbb{P}(\gamma C_{\zeta}(\omega, j))$ .

(c)  $\mathbb{P}^{-1}(\gamma C_{\mathfrak{S}}^*(\psi, j)) \leq \gamma C_{\zeta}(\mathbb{P}^{-1}(\psi), j)$ .

(d) For each  $\psi$  and each  $\omega$  is an  $j$ - $F\gamma$ -open set with  $\mathbb{P}^{-1}(\psi) \leq \omega$ , there is  $\mu \in I^Y$  is  $j$ - $F\gamma\mathcal{I}$ -open with  $\psi \leq \mu$  and  $\mathbb{P}^{-1}(\mu) \leq \omega$ .

**Proof.** This can be proven using the same approach as in Theorem 5.2.

$\square$

**Proposition 7.** Let  $\mathbb{P} : (Z, \zeta) \longrightarrow (Y, \mathfrak{S}, \mathcal{I})$  be a fuzzy mapping and bijective,  $\mathbb{P}$  is  $F\gamma\mathcal{I}$ -irresolute closed iff  $\mathbb{P}$  is  $F\gamma\mathcal{I}$ -irresolute open.

**Proof.** This follows directly from:

$$\mathbb{P}^{-1}(\gamma C_{\mathfrak{S}}^*(v, j)) \leq \gamma C_{\zeta}(\mathbb{P}^{-1}(v), j) \iff \mathbb{P}^{-1}(\gamma I_{\mathfrak{S}}^*(v^c, j)) \leq \gamma I_{\zeta}(\mathbb{P}^{-1}(v^c), j).$$

$\square$

## 6. Conclusions

In this research, a novel class of fuzzy sets, called  $j$ - $F\gamma\mathcal{I}$ -open sets, has been defined and investigated on  $\mathcal{FITT}$ s in the sense of Šostak. After that, the concepts of  $F\gamma\mathcal{I}$ -interior operators and  $F\gamma\mathcal{I}$ -closure operators have been introduced and studied. We also presented and investigated some types of fuzzy  $\mathcal{I}$ -separation axioms, called  $j$ - $F\gamma\mathcal{I}$ -normal spaces and  $j$ - $F\gamma\mathcal{I}$ -regular spaces via  $j$ - $F\gamma\mathcal{I}$ -closed sets. Moreover, the notion of  $F\gamma\mathcal{I}$ -continuity has been defined and discussed. The concepts of  $FA\gamma\mathcal{I}$ -continuous and  $FW\gamma\mathcal{I}$ -continuous mappings, which are weaker forms of  $F\gamma\mathcal{I}$ -continuous

mappings, have been introduced and studied. Finally, we explored and characterized some new fuzzy  $\gamma\mathcal{I}$ -mappings via  $j$ - $F\gamma\mathcal{I}$ -open sets and  $j$ - $F\gamma\mathcal{I}$ -closed sets, called  $F\gamma\mathcal{I}$ -open mappings,  $F\gamma\mathcal{I}$ -closed mappings,  $F\gamma\mathcal{I}$ -irresolute mappings,  $F\gamma\mathcal{I}$ -irresolute open mappings, and  $F\gamma\mathcal{I}$ -irresolute closed mappings.

In upcoming paper, we intend to study the following topics: (a) extending these concepts given here to include fuzzy soft minimal (topological) spaces as introduced in [36,37]; (b) introducing fuzzy lower and upper  $\gamma\mathcal{I}$ -continuous multifunctions and  $j$ -fuzzy  $\gamma\mathcal{I}$ -connected sets; (c) finding a use for these concepts given here to include double fuzzy topological spaces as introduced in [38]; and (d) defining these concepts given here base on lattice valued fuzzy sets.

## References

1. Zadeh, L.A. Fuzzy Sets. *Inf. Control* **1965**, *8*, 338-353.
2. Kumar, R.; Singh, P.; Zhang, T. Fuzzy control systems: Advances and applications. *Control Tech.* **2021**, *19*, 210-234.
3. Ali, M.; Wang, H. Application of fuzzy logic in control systems: A comprehensive survey. *J. Control Eng. Tech.* **2022**, *18*, 23-45.
4. Smith, J.; Gupta, A. Fuzzy logic in artificial intelligence: Current trends and future directions. *Art. Intel. Rev.* **2022**, *55*, 345-367.
5. Zhang, Q.; Li, L.; Zhang, T. Fuzzy sets and their applications in artificial intelligence: A review. *IEEE Trans. Fuzzy Sys.* **2023**, *31*, 789-804.
6. Ahmed, N.; Kumar, P. Fuzzy logic in image processing: A survey of methods and applications. *Image Vis. Comput.* **2022**, *100*, 100-120.
7. Sharma, V.; Kumar, R.; Patel, M. Fuzzy image processing: Techniques and applications. *Pat. Recognit. Lett.* **2023**, *142*, 45-60.
8. Dubois, D. Fuzzy sets and systems: theory and applications. *Academic Press: Cambridge, MA, USA* **1980**, Volume 144.
9. Chen, L.; Zhao, Y. Fuzzy decision-making models: A review of trends and applications. *J. Decis. Syst.* **2023**, *32*, 110-135.
10. Ahmad, B.; Kharal, A. On fuzzy soft sets. *Adv. Fuzzy Syst.* **2009**, *2009*, 586507.
11. Cagman, N.; Enginoglu, S.; Citak, F. Fuzzy soft set theory and its application. *Iran. J. Fuzzy Syst.* **2011**, *8*, 137-147.
12. Atef, M.; Ali, M.I.; Al-shami, T.M. Fuzzy soft covering based multi-granulation fuzzy rough sets and their applications. *Comput. Appl. Math.* **2021**, *40*, 115.
13. Chang, C.L. Fuzzy topological spaces. *J. Math. Anal. Appl.* **1968**, *24*, 182-190.
14. Ameen, Z.A.; Al-shami, T.M.; Azzam, A.A.; Mhemdi, A. A novel fuzzy structure: infra-fuzzy topological spaces. *J. Fun. Spaces* **2022**, *2022*, 1-11.
15. Saleh, S.; Al-shami, T.M.; Azzam, A.A.; Hosny, M. Stronger forms of fuzzy pre-separation and regularity axioms via fuzzy topology. *Mathematics* **2023**, *11*(23), 4801.
16. Ameen, Z.A.; Mohammed, R.A.; Al-shami, T.M.; Asaad, B.A. Novel fuzzy topologies formed by fuzzy primal frameworks. *J. Intel. Fuzzy Sys.* **2024**.
17. Saleh, S.; Birkea, F.M.O.; Al-shami, T.M.; Arar, M.; Omran, M. Novel categories of spaces in the frame of generalized fuzzy topologies via fuzzy  $g\mu$ -closed sets. *Eur. J. Pure Appl. Math.* **2025**, *18*(1), 5856.
18. Šostak, A.P. On a fuzzy topological structure. In *Proceedings of the 13th Winter School on Abstract Analysis, Section of Topology, Palermo: Circolo Matematico di Palermo, Srni, Czech Republic, 27 January–3 February 1985*, pp. 89-103.
19. Ramadan, A.A. Smooth topological spaces. *Fuzzy Sets Syst.* **1992**, *48*, 371-375.
20. Chattopadhyay, K.C.; Samanta, S.K. Fuzzy topology: fuzzy closure operator, fuzzy compactness and fuzzy connectedness. *Fuzzy Sets Syst.* **1993**, *54*, 207-212.
21. El-Gayyar, M.K.; Kerre, E.E.; Ramadan, A.A. Almost compactness and near compactness in smooth topological spaces. *Fuzzy Sets Syst.* **1994**, *62*, 193-202.
22. Höhle, U.; Šostak, A.P. A general theory of fuzzy topological spaces. *Fuzzy Sets Syst.* **1995**, *73*, 131-149.
23. Ramadan, A.A.; Abbas, S.E.; Kim, Y.C. Fuzzy irresolute mappings in smooth fuzzy topological spaces. *J. Fuzzy Math.* **2001**, *9*, 865-877.

24. Kim, Y.C.; Ramadan A.A.; Abbas, S.E. Weaker forms of continuity in Šostaks fuzzy topology. *Indian J. Pure Appl. Math.* **2003**, *34*, 311-333.
25. Abbas, S.E. Fuzzy super irresolute functions. *Intert. J. Math. Math. Sci.* **2003**, *42*, 2689-2700.
26. Abbas, S.E. Fuzzy  $\beta$ -irresolute functions. *Appl. Math. Comp.* **2004**, *157*, 369-380.
27. Kim, Y.C.; Abbas, S.E. On several types of  $r$ -fuzzy compactness. *J. Fuzzy Math.* **2004**, *12*, 827-844.
28. Alsharari, F.; Saleh, H.Y.; Taha, I.M. Some characterizations of  $k$ -fuzzy  $\gamma$ -open sets and fuzzy  $\gamma$ -continuity with further selected topics. *Symmetry* **2025**, *17*, 678.
29. Taha, I.M.; Abbas, S.E. A new notion of fuzzy local function and some applications. *Adv. Fuzzy Sys.* **2022**, *8954163*, 1-10.
30. Taha, I.M. On  $r$ -fuzzy  $\ell$ -open sets and continuity of fuzzy multifunctions via fuzzy ideals. *J. Math. Comput. Sci.* **2020**, *10*, 2613-2633.
31. Taha, I.M. On  $r$ -generalized fuzzy  $\ell$ -closed sets: properties and applications. *J. Math.* **2021**, *2021*, 1-8.
32. Taha, I.M.  $r$ -fuzzy  $\delta$ - $\ell$ -open sets and fuzzy upper (lower)  $\delta$ - $\ell$ -continuity via fuzzy idealization. *J. Math. Comput. Sci.* **2022**, *25*, 1-9.
33. Taha, I.M. On fuzzy upper and lower  $\alpha$ - $\ell$ -continuity and their decomposition. *J. Math. Comput. Sci.* **2021**, *11*, 427-441.
34. Ramadan, A.A.; Kim, Y.C.; Ei-Gayyar, M.K. Smooth ideals. *J. Korean Instit. Intell. Sys.* **2002**, *12*, 90-95.
35. Kandil, A.; El-Shafei, M.E. Regularity axioms in fuzzy topological spaces and FRi-proximities. *Fuzzy Sets Syst.* **1988**, *27*, 217-231.
36. Aygünoğlu, A.; Çetkin, V.; Aygün, H. An introduction to fuzzy soft topological spaces. *Hacet. J. Math. Stat.* **2014**, *43*, 193-208.
37. Taha, I.M. Compactness on fuzzy soft  $r$ -minimal spaces. *Int. J. Fuzzy Log. Intell. Syst.* **2021**, *21*, 251-258.
38. Taha, I.M. Some properties of  $(r, s)$ -generalized fuzzy semi-closed sets and some applications. *J. Math. Comput. Sci.* **2022**, *27*, 164-175.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.