

Role of extracellular vesicles in cell death and inflammation

Rahul Sanwlani¹ and Lahiru Gangoda^{1,2,3,*}

¹Department of Biochemistry and Genetics, La Trobe Institute for Molecular Science, La Trobe University, Melbourne, VIC, Australia

²The Walter and Eliza Hall Institute of Medical Research (WEHI), 1G Royal Parade, Parkville, Melbourne, Victoria 3052, Australia

³Department of Medical Biology, University of Melbourne, Parkville, Melbourne, Victoria 3010, Australia

* **Corresponding:** Dr Lahiru Gangoda, The Walter and Eliza Hall Institute of Medical Research (WEHI), 1G Royal Parade, Parkville, Melbourne, Victoria 3052, Australia; Email: Gangoda.L@wehi.edu.au

Abstract: Extracellular vesicles (EVs) have been identified as novel mediators of intercellular communication. They work *via* delivering the sequestered cargo to cells in close vicinity as well as distant sites in the body, regulating pathophysiological processes. Cell death and inflammation are biologically crucial processes in both normal physiology and pathology. These processes are indistinguishably linked with their effectors modulating the other process. For instance, during an unresolvable infection, the upregulation of specific immune mediators leads to inflammation causing cell death and tissue damage. EVs have gained considerable interest as mediators of both cell death and inflammation during conditions such as sepsis. This review summarizes the types of extracellular vesicles known to date and their roles in mediating immune responses leading to cell death and inflammation with specific focus on sepsis and lung inflammation.

Keywords- Extracellular vesicles; exosomes; microvesicles; apoptotic bodies; apoptotic extracellular vesicles; cell death; inflammation; sepsis; lung inflammatory disorders; SARS-CoV-2

Abbreviations

ALI	Acute lung injury
ApoBDs	Apoptotic bodies
ApoEVs	Apoptotic EVs
ApoExos	Apoptotic exosomes
ApoMVs	Apoptotic microvesicles
ARDS	Acute respiratory distress syndrome
BALF	Bronchoalveolar lavage fluid
CD	Cluster of differentiation
CLP	Cecal ligation and puncture
COPD	Chronic obstructive pulmonary disorder
EVs	Extracellular vesicles
HSP	Heat shock protein
HIV	Human immunodeficiency virus
IECs	Intestinal epithelial cells
LPS	Lipopolysaccharide
MEVs	Milk-derived EVs
MHC	Major histocompatibility complex
miR	micro-RNA
MSCs	Mesenchymal stem cells
MVB	Multivesicular bodies
MVs	Microvesicles
PAMPs	Pathogen associated molecular patters
TNF	Tumor necrosis factor

Extracellular vesicles: introduction, subtypes, and cargo

Extracellular vesicles (EVs) have been defined as nanosized vesicles that are shed into extracellular space (1). First identified in differentiating reticulocytes three decades ago, EVs were thought to be the waste disposal system of cells which aided in the elimination of cellular waste (2, 3). However, extensive studies over the past two decades have been crucial for introducing a paradigm shift by ascertaining a more sophisticated role for EVs (4). Intercellular communication is no more thought to be a consequence of only direct cell-cell contact or secreted factors such as hormone signalling. EVs are now known to mediate multiple pathophysiological processes by facilitating intercellular communication (5). Their expansive role in health and disease has been investigated in multiple models (6, 7). As more studies continue to shed light on the diverse role played by EVs, their role is now becoming apparent to be crucial in not just mediating communication between distant cells in an individual but facilitate the transfer of bioactive compounds inter-individual, cross-species and even inter-kingdom (8-10).

Multiple subtypes of eukaryotic EVs exist and they are primarily categorized based on their biogenesis, subcellular origin, and size (**Figure 1**) (1, 11). Exosomes are vesicles that are endocytic in origin and shed as a consequence of multivesicular bodies (MVB) or late endosomes fusing with the plasma membrane (12). They range in size from 30-150 nm and were believed to be the smallest EV subtype until the recently (1, 13). Exomeres have been identified as secretory nanoparticles that are less than 50 nm in size with their mode of biogenesis currently unknown (14, 15). Whereas, shedding microvesicles (MVs) are larger in size and may range from 100-1000 nm. They originate due to budding or outward vesiculation of plasma membrane, thus also referred to as ectosomes (1,

16). Furthermore, there are new classes of EVs defined such as migrasomes and large oncosomes (17-19). Up until now, apoptotic cells were thought to only shed apoptotic bodies (ApoBDs) which are similar to microvesicles in origin as they emerge due to budding of plasma membrane. However, ApoBDs are larger in size and may range from 50-5000 nm (20, 21). More recently, apoptotic cells were found to partake in secretion of other EV subtypes and apoptotic cell-derived EVs (ApoEVs) now include apoptotic microvesicles (ApoMVs) and apoptotic exosomes (ApoExos) which are physiologically distinct from ApoBDs (22). Similar to MVs, ApoMVs range in size from 200-1000 nm and also called apoptotic microparticles. They have a higher membrane integrity than ApoBDs, promoting better molecular exchange (22-24). Whereas ApoExos are the most recent addition to ApoEVs and were initially reported to be released by endothelial cells following caspase 3-dependent MVB formation (25, 26). Though there is limited knowledge regarding their origin and function, they continue to be explored in-depth as final messengers of apoptotic cells (22).

EVs contain a diverse cargo containing plethora of proteins, nucleic acids and lipids (27). EV cargo is capable of mediating a variety of responses in the recipient cell including infections (28), signal transduction (29) and immune regulation (30). EV cargo is not only different for various subtypes of EVs but also varies between same EV subtype derived from two different cell types (31, 32). EV proteome is its most well studied cargo so far (27). Exosomes have been particularly found to be enriched with endosome associated proteins with documented roles in vesicle trafficking such as Rab, Annexin and Flotillin. Furthermore, Alix and TSG101 have been previously claimed as molecular markers of exosomes (5, 12). However, even though these proteins with role in MVB biogenesis are enriched in exosomes, they are not markers as previously claimed (1). Exosomes have also been found to be enriched with Heat Shock proteins (HSPs), and membrane proteins such integrins, Major Histocompatibility Complex (MHC) class I and II and tetraspanins CD63, CD9 and CD81 (12, 32, 33). MVs on the other hand, having reported roles in metastases, invasion, and inflammation, are enriched with proteins such as Annexins, selectin, CD40 and metalloproteinases (34, 35). Interestingly, even though ApoEVs have been suggested to have characteristics similar to EVs released by healthy cells, their cargo may enable more distinctive roles. With their unique cargo, ApoEVs may influence processes such as inflammation and autoimmunity in addition to aiding in clearance of apoptotic by-products and remnants. Furthermore, ApoBDs not only contain biomolecule cargo from the originating cell but can also contain nuclear fragments and cellular organelles such as mitochondria and endoplasmic reticulum (20, 22, 36, 37). Identification of nucleic acids including mRNA, miRNA (miR) and DNA in EVs has generated significant interest (38, 39). Functional transfer of nucleic acids *via* EVs is known to regulate recipient cells and control signalling pathways (40). Lastly, EVs are abundant in lipids and lipid-related enzymes including but not limited to sphingomyelin, ganglioside, ceramide, cholesterol, and prostaglandins. In fact, EV lipid composition has been observed to be different than that of the cell type of origin (41-43). Similar to proteins and nucleic acids, lipids in EVs also have functional role in controlling signalling pathways in recipient cells and thus contribute in more ways than only providing structural integrity to these vesicles (44).

EVs as regulators of cell death

Cell death is an important process that involves the removal of cells during certain developmental stages or those that are compromised due to exposure to toxins. Cell death can also occur as a result of natural processes, where old cells die and are replaced by new ones to maintain homeostasis in an individual. On the contrary, it could also be a result of unnatural factors such as pathological conditions or local injuries (45). As established above, cells undergoing apoptosis can shed multiple EV subtypes aiding in not only getting rid of apoptotic debris but also facilitating communication with other cell types. However, the role of EVs in cell death is not only limited to the functions of EVs shed by apoptotic cells. As more studies continue to understand the regulatory potential of EVs, it is becoming increasingly evident that EVs from living cells are also capable of moderating cell death responses in recipient cells (**Figure 2**). The cargo embodied in EVs is not just dependent on the EV subtype but also the cell type of origin, such as diseased versus healthy cells (13, 46). Thus, in a context dependent manner, these signalling moieties may be capable of promoting or attenuating cell death in a particular recipient cell type. The subsequent sections are aimed at elaborating the role of EVs in cell death in various pathophysiological conditions. As this review specifically addresses the role of EVs in inflammatory disorders, this section would emphasize the role of EVs in mediating immune and inflammatory cell death.

Role of EVs in attenuating cell death

Attenuation of cell death *via* extracellular vesicles has been studied in multiple models. It has recently been discovered that milk-derived EVs (MEVs) have a role in promoting gut health and development upon uptake by intestinal epithelial cells (IECs) (9). MEVs from multiple sources such as porcine, bovine and humans have been demonstrated to prevent cell death in IECs during oxidative stress and hypoxia and promote survival and proliferation (47-49). This ability of MEVs has been attributed to its cargo inhibiting p53 expression in IECs (50). However, the role of MEVs in attenuation of cell death may not always be beneficial for the consumer. miR-148a and miR-21 which are a part of MEV cargo have been designated as oncomiRs due to their role in promoting vital cellular processes such as proliferation and survival by inhibition of tumor suppressor genes (51-53). Thus, the ability of MEVs to attenuate cell death could be detrimental too. Multiple studies have now demonstrated the ability of tumor cell derived EVs in promoting survival and preventing cell death in recipient tumor cells (**Figure 2**). For instance, exosomes from N-myc amplified neuroblastoma cells were found to enhance survival by inducing chemoresistance in non-N-myc amplified cells, preventing doxorubicin induced apoptosis (54). Furthermore, in other studies using glioblastoma models, it has been shown that exosomes shed by apoptotic tumor cells were able to confer survival advantage to the surviving glioblastoma cells by transferring spliceosome components (55). Another model where EVs attenuated cell death, that continues to be increasingly defined with studies, is that of stem cells. The therapeutic nature of mesenchymal stem cells (MSCs) due to paracrine signalling is well established (56). Recently, it was observed that the MSC derived therapeutic benefits were mediated largely due to EVs. MSC-derived EVs were reported to enhance proliferation and migration of keratinocytes and suppress apoptosis, thus promoting

enhanced healing at wound sites (57). Similarly, MSC-derived EVs are now well known for their cardioprotective function. They have been found to inhibit apoptosis, autophagy and fibrosis by modulating cardiomyocytes and endothelial cells. Further, MSCs were also found to stimulate cell proliferation and angiogenesis, thus leading to enhanced cardiac survival (**Figure 2**) (58-60).

Attenuation of cell death is not necessarily a feature of EVs shed by live cells. ApoEVs which serve as the last messages from dying cells have been thoroughly characterised for the complex role they play in tissue development, regeneration and apoptosis (**Figure 2**) (61). Similar to MSC-derived exosomes, MSC-derived ApoEVs have been shown to promote angiogenesis and the recovery of cardiac function *via* the regulation of autophagy (62). Further, apoptotic MSC-derived EVs have also been found to have a role in nephron repair following injury. These MSC-derived ApoEV associated benefits could be attributed to compensatory proliferation signalling which has been implied to be consequence of apoptosis (63). Lastly, ApoEVs have been found to indirectly attenuate cell death in surrounding cells by aiding in the regulation of immune system in disease settings. For instance, ApoEVs from *Mycobacterium tuberculosis* infected macrophages engulfed by dendritic cells induced antimicrobial immunity. This was shown to aid in clearance of pathogen and prevented further infection of macrophages (64). A similar phenomenon has also been discovered in the clearance of prion infection by microglia cells upon engulfment of ApoEVs from prion infected neurons (65). Overall, the ability of EVs to prevent cell death directly or indirectly due to its signalling cargo is evident. These findings highlight the broad untapped therapeutic potential of EVs in controlling diseases and in treating pathological conditions that have so far been extremely challenging to tackle.

Role of EVs in promoting cell death

The role of EVs in pathological diseases has been extensively studied in host-pathogen interaction model. EVs have evolved as means of communication between host and pathogen by mediating inter-kingdom transfer of cargo. An increasing number of studies over the last decade have highlighted the ability of EVs in facilitating the establishment and accelerated spread of infections in the host. The pathogens have been found to work by exploiting host's machinery or even deploying their own EVs to suppress the host's immune system (**Figure 2**) (66-69). For instance, astrocytes infected with Human immunodeficiency virus (HIV) have also been shown to shed EVs with the viral Tat protein leading to neuronal cell death (70). Similarly, Epstein-Barr virus has been shown to deploy viral components such as latent membrane protein-1 in EVs shed by infected cells, ultimately leading to the alteration of signalling pathways and cell death in T-lymphocytes (71). Bacterial pathogen *Mycobacterium tuberculosis* too has been shown to load virulent cargo in EVs shed by infected macrophages. These EVs were capable of inducing apoptosis in Jurkat T cells (72). Macrophages infected with bacterial pathogens including *Mycobacterium tuberculosis*, *Salmonella typhimurium* and *Toxoplasma gondii* have been shown to shed small EVs which contain pathogen associated molecular patterns (PAMPs). Upon uptake by uninfected macrophages, these EVs were responsible for elevated cytokine release and resulting pro-inflammatory response (66). Additionally, it has been shown that EVs shed by a variety of bacterial

pathogens themselves harbour virulent cytotoxic cargo which enables them to establish infection in healthy cells and suppress host's defences (67). For instance, Gram negative bacterial EVs have been shown to aid in the delivery of lipopolysaccharide (LPS) to the recipient cell's cytosol. This event induces an inflammatory form of cell death, known as pyroptosis, by triggering casapase-11 dependent effector responses (73). Furthermore, *Neisseria gonorrhoeae* derived EVs have been shown to contain the Porin protein PorB which induce apoptosis in host macrophages (74). Similarly, the presence of *Shigella* toxins has been observed in pathogen derived EVs and found have cytotoxic implications via caspase-9 and caspase-3 activation in human IECs (75).

The role of EVs as promoters of cell death has been elucidated in other pathological conditions too such as cancer, autoimmunity, and pulmonary disorders (**Figure 2**). For instance, in a recent study, the role of T-lymphocyte derived exosomes in Type-1 diabetes was explored. Specific miR (miR-142-3p, miR-142-5p and miR-155) have been found to be enriched in T-lymphocyte derived EVs which led to the elevated incidence of Type-1 diabetes as they selectively targeted and induced apoptosis in pancreatic β cells along with chemokine signalling including *Ccl2*, *Ccl7* and *Cxcl10* (76). Tumor cell derived EVs have also been found to use this mechanism to suppress immune response. A recent study showed that exosomal PD-L1 was upregulated and carried in EVs released by metastatic melanomas. These EVs were found to have role in exhaustion of CD8⁺ T cells which aided the tumor cells in evading immune response (30). Similarly, sera derived from individuals with oral squamous cell carcinoma had elevated expression of Fas ligand which led to T cell apoptosis (77). In context of lung disorders such as asthma, airway inflammatory potential of eosinophil derived EVs due to pro-inflammatory cargo has been observed (78). Further, these EVs were demonstrated to lead to apoptosis in primary alveolar epithelial cells by interfering with JAK/STAT signalling (79). Similarly, monocyte derived EVs encapsulated caspase-1 and their ability to induce apoptosis of pulmonary microvascular endothelial cells was observed in acute lung injury (ALI) and acute respiratory distress syndrome (ARDS) (80, 81). Another intriguing observation of EV mediated lung pathology and cell death was made in chronic obstructive pulmonary disease (COPD). The apoptotic endothelial cells during COPD shed EVs loaded with inflammatory cargo that promote progression of the disease by inducing apoptosis in otherwise healthy endothelial cells (82). The subsequent sections further discuss the role of EVs in promoting lung inflammatory disorders in more detail.

The immense potential of EVs as mediators that promote cell death in various pathological conditions has been concretely established and continues to be explored in depth. Collectively, these studies present opportunities to exploit EVs at a therapeutic front for multiple purposes. These findings are key in devising novel therapeutic paradigms and alternatives in treating and preventing the spread of infections as well as diseases such as cancer by targeting EV mediated signalling. Although promising, these studies need further validation with better *in vivo* models and improved understanding of the molecular mechanisms and cargo components of interest before extrapolating the use of EVs to clinical setups.

EVs in inflammation

Inflammation is the immune system's response to harmful stimuli, such as pathogens, damaged cells, toxic compounds, or irradiation (83). Some forms of cell death can provoke inflammatory responses. While apoptosis is an immunologically silent process and generally does not trigger inflammation, other forms of lytic cell death such as necrosis and pyroptosis are pro-inflammatory (84). For instance, cells infected with pathogenic bacteria or viruses undergo pyroptosis induced by activation of inflammasome sensors resulting in the loss of plasma membrane integrity, leakage of cellular contents and inflammation (85). Inflammation can be further classified as either acute inflammation which occurs immediately after injury and lasts for few days or chronic inflammation which may last for several months or sometimes even years. During acute inflammatory responses, the body efficiently minimises impending damage by removing injurious stimuli and initiating the healing process. This leads to the restoration of tissue homeostasis and the resolution of inflammation (86). Another form of cell death known as pyroptosis has been long studied for its role in chronic inflammatory diseases such as Alzheimer's disease (87, 88) and atherosclerosis (89). Under chronic conditions, the inflammation fails to resolve due to excessive and uncontrollable activity of the immune system which can lead to extensive tissue damage (90). Hence new therapies aimed at preventing this hyperactivity of the immune system could have major clinical benefits.

Recently EVs have gained attention as mediators and biomarkers of inflammation. The discovery of MHC harbouring EVs shed by multiple lineages of immune cells suggested a complex functional role for these vesicles as they mediated antigen specific immune response (34, 91, 92). Further, secretion of Tumor Growth Factor $\beta 1$ via EVs was found to modulate anti-inflammatory effects (93). ApoExos containing Sphingosine 1-Phosphate Receptors 1 and SPR1/3 have been identified to induce proinflammatory cytokine production in macrophages leading to activation of NF- κ B and p38 MAPK (37). Non-coding RNAs delivered *via* ApoExos have been found to be pro-inflammatory in nature as they stimulate toll-like receptors (94). Additionally, EVs have been found to contain wide variety of proteins, including chemokines and inflammatory cytokines, such as tumor necrosis factor (TNF), IL-1 β , CXCL2 and CXCL8 suggesting an important immune modulatory functions of EVs (95-97). In the following section, we aim to discuss existing literature with the focus on the role of EVs during acute inflammation in diseases such as sepsis and ARDS, and chronic inflammation associated with diseases such as COPD and asthma.

EVs in sepsis associated inflammation

Sepsis is a dysregulated systemic inflammatory disorder that can lead to life threatening organ injury (98). Since their discovery over two decades ago, EVs originating from different type of cells have been shown to play diverse roles in sepsis. The source of EVs during sepsis can be either from the pathogen responsible for the infection or the host's own cells (99).

Cell death is the most common outcome during infections. Pathogen can trigger both non-inflammatory modes of cell death such as apoptosis and autophagy or inflammatory modes of cell death such as pyroptosis and oncosis (100). Gram-positive and Gram-negative bacteria, as the most frequent infectious agent of sepsis, can produce EVs (101) that carry

bacterial virulence factors that contribute towards the systemic inflammation (102). During sepsis, lymphocytes are depleted by apoptosis, which leads to immunosuppression (103). Exosomes from HIV type 1 infected cells has been shown to contain the accessory extracellular viral protein Nef, which was shown to induce apoptosis of CD4⁺ T-cells (104). Given that pathogen-derived EVs carry cargo that are representative of their cell of origin, bacterial protoplast-derived nanovesicles have been used to vaccinate mice against these pathogens in models of pneumonia and peritonitis (105).

On the other hand, host-derived EVs during sepsis are produced mainly from immune cells such as platelets and innate immune cells (99). Increased levels of platelet MVs has been observed in patients with Gram-negative bacterial sepsis (106). MVs originating from platelets or granulocytes were also elevated in meningococcal sepsis (107). These MVs were associated with increased coagulation activity. Consequently, MVs have also been linked with thromboembolic events and in patients with transient ischemic attacks, lacunar infarcts and multi-infarct dementias have elevated levels of MVs (108-110). Hence, host-derived MVs have been suggested as a novel target for therapeutic intervention in clinical conditions with enhanced coagulation activation (107). In sepsis neutrophils are one of the vital leukocytes to play first line of defence (111-113). Neutrophil-derived EVs dramatically increased in the blood of mice subjected to cecal ligation and puncture (CLP), a commonly used animal model of sepsis (114). Treatment with neutral sphingomyelinase inhibitor, GW4869 which blocks exosome production, significantly improved the survival of mice subjected to LPS injection or CLP, suggesting that exosomes might play an important role in sepsis (115). Studies have also shown altered contents and function of host derived EVs during sepsis. In sepsis, exosomes carry increased levels of cytokines and damage-associated molecular patterns (DAMPs) to induce inflammation. It is well documented that patients with sepsis have increased levels of serum cytokines which are important for initiation of the "cytokine storm". Exosomes have been shown to carry biologically active pro-inflammatory cytokines IL-12, IL-15, IL-17, and IFN- γ at early phase and anti-inflammatory cytokines, IL-4 and IL-10 at late phase of the cytokine storm (116, 117).

EVs play major roles in organ damage during sepsis. Activated polymorphonuclear leukocytes from septic patients have been shown to produce microparticles with increased adhesion molecules that can activate the vascular endothelium leading to endothelial injury and resultant organ dysfunction (118, 119). Moreover, platelet-derived exosomes from sepsis patients induced apoptosis of endothelial cells and vascular smooth muscle cells due to increased NADPH activity and by generating reactive oxygen species such as superoxide, nitric oxide and peroxynitrite (120). Thereby, exosomes have been implicated in the induction of oxidative stress and myocardial dysfunction and vascular cell apoptosis. Additionally, exosomes have also been shown to contribute to septic encephalopathy directly and by causing systemic immune dysfunction due to the collapse of neuroendocrine immune networks (121). Lastly, exosomes may contribute to chronic liver dysfunction after sepsis, as exosomal miR-103-3p from LPS-activated macrophages targeted KLF4 to increase liver fibrosis (122).

The role of EVs in sepsis may not be limited to causing damage to host cells and organs. Interestingly, several protective host EV-related

mechanisms were recently discovered in sepsis models. In several studies, granulocyte derived EVs have shown antibacterial effects (123) in addition to supporting the production of inflammatory mediators and protecting from vascular dysfunction (124). Likewise, EVs containing alpha-2-macroglobulin excreted by neutrophilic granulocytes were shown to help in bacterial clearance and reduce inflammation in a mouse model of sepsis (125). Immature dendritic cell-derived exosomes attenuated the acute systemic inflammatory response in sepsis by enhancing apoptotic cell clearance in septic rats (126). Overall, EVs have both protective and detrimental roles during sepsis in a contextual manner, depending on their cell type of origin and the cargo contained within (99). In this context, the protein cargo from EVs isolated from the serum of patients following sepsis diagnosis has been analysed in a clinical study to identify biomarkers for better disease management and monitoring disease progress and prognosis (127). Further knowledge from similar studies will be instrumental in developing novel therapeutic interventions and even biomarker discovery for improved management of sepsis patients.

Lung is one of the most susceptible organs to systemic inflammation. Septic patients are often accompanied by ALI or ARDS, which further add to sepsis-associated mortality (128). Thus, in the following section we discuss the role of EVs in lung inflammation.

EVs in lung inflammatory disorders

Lung is the organ with highest vascular density in the human body. Hence the endothelium of the lung contributes substantially to the circulation of EVs which in-turn play critical roles in normal lung physiology (129). Lung epithelial cell-derived exosomes contain membrane-tethered mucins that contribute to the innate defence of the airway (130). EVs have also gained attention as novel communicators in lung diseases such as COPD, asthma, ALI, and ARDS (131). Alveolar macrophages can secrete SOCS1 and 3 (suppressor of cytokine signalling 1, 3) in EVs which can further control inflammatory signalling. This process has been observed to be dysregulated in association with cigarette smoke exposure (132). EV secretion from lung endothelial cells is enhanced during adverse conditions such as infection and smoke exposure (133, 134). These vesicles are primarily generated by an apoptosis-dependent mechanism that is triggered upon pulmonary endothelial damage. Circulating lung endothelial cell-derived EVs can therefore be potential biomarkers for lung injury and inflammation. In one such study, cigarette smoke extract induced CCN1-enriched exosomes. CCN1 is generally known to play crucial roles in tissue remodelling and repair as an early response gene product and as an extracellular matrix protein (135, 136). However, these exosomes were found to induce the activation of IL-8 secretion which recruits inflammatory cells, particularly neutrophils, into the lung parenchyma, further promoting lung inflammation in COPD (137).

Enhanced secretion of exosomes by lung epithelial cells has also been observed in asthma which is a chronic, recurrent, and incurable allergy-related respiratory disease. These epithelial cell-derived exosomes induced proliferation and chemotaxis of undifferentiated macrophages and inhibition of exosome production, which resulted in a reduced population of proliferating monocytes and alleviation of various asthmatic features (138). Asthma is characterized by the infiltration of

airway with T lymphocytes, mast cells, basophils, macrophages, and eosinophils (139). Mast cells secrete exosomes that carry immunologically relevant molecules such as MHC class II, CD86, LFA-1 and ICAM-1. These exosomes can act as messengers to induce splenic B and T cell blast activation and have been suggested to play a role in the recruitment of these cells to the lungs, eventually contributing to lung inflammation (140). On the contrary, mast cell exosomes containing high-affinity IgE receptors (FcεRI) have been shown to suppress allergic reactions by binding to free IgE (141). Eosinophils are the main effector cells in asthma. Eosinophils of asthmatic patients were shown to produce more exosomes than those of healthy individuals and these exosomes promoted inflammatory behaviour of eosinophils that is related to asthma pathogenesis (79, 142). Similarly, B cell-derived exosomes have been shown to play a role in stimulating the allergic immune response via stimulation of T cell proliferation and induction of cytokine production of allergen-specific T cells (IL-4, IL-5 and IL-13 and lower levels of IFN- γ and TNF- α) (143, 144).

ALI and ARDS remain as life-threatening diseases in critically ill patients. ALI and ARDS can be triggered by both non-infectious (sterile) and infectious stimuli underlying clinical disorders, such as sepsis, pneumonia, trauma, and pancreatitis (145-147). Studies have shown that proceeding sterile stimuli, alveolar type-I epithelial cells were the main source of the EVs in the bronchoalveolar lavage fluid (BALF) whereas infectious stimuli-induced BALF EVs were mainly derived from alveolar macrophages. Both sterile stimuli-induced EVs and infection-induced EVs facilitated classic macrophage activation, and subsequently promoted inflammatory lung responses *via* different signalling pathways (148). Caspase-1 encapsulated in exosomes released from LPS-challenged macrophages induced lung endothelial cell apoptosis, indicating their contribution to the disruption of the alveolar-capillary barrier (81). Given the ability of EVs to induce an inflammatory response and cell death of lung tissue, they have the potential to be targeted for more effective therapeutic approaches to treat ALI and ARDS (149).

The novel coronavirus disease (COVID-19) is responsible for millions of deaths worldwide with the respiratory failure been the most common clinical presentation (150). Recent evidence points to a correlation between COVID-19 severity and pyroptotic inflammasome activation by SARS-CoV-2 infection (151). COVID-19 can cause lung complications such as pneumonia and in the most severe cases ARDS and sepsis (152). In a recent proteomic analysis of COVID-19 patient-derived exosomes, several molecules were identified which are involved in several biological processes. These processes include immune response, inflammation, and activation of the coagulation and complement pathways, the main mechanisms of COVID-19-associated tissue damage and multiple organ dysfunctions. Several potential biomarkers such as fibrinogen, fibronectin, complement C1r subcomponent and serum amyloid P-component that are well correlated with the COVID-19 disease's severity were also encapsulated within these exosomes (153). The presence of viral material amongst the exosomal cargo showed that SARS-CoV-2 could use the cell-to-cell communication system to spread infection in the host (153, 154). Cells that express ACE2 and CD9 can transfer these viral receptors to other cells via EVs, making recipient cells more susceptible for SARS-CoV-2 infection (154). GM3-enriched exosomes were shown to be positively correlated with COVID-19 disease

severity, suggesting they may partake in pathological processes related to COVID-19 pathogenesis (155). In SARS-CoV-2 infection, the systemic inflammatory response was shown to result in the release of substantial amounts of procoagulant EVs that may act as clotting initiation agents, contributing to disease severity (156, 157). On the other hand, exosome-based strategies were also proposed to treat COVID-19 or prevent SARS-CoV-2 infection (154, 158, 159). Especially, MSCs-derived EVs from a broad range of sources, including bone marrow, adipose tissue, umbilical cord are currently under investigation for many conditions and areas of regenerative medicine (160). They have been shown to alleviate severe inflammation (cytokine storm) and repair damaged lung cells in COVID-19 by delivery of anti-inflammatory molecules. There are several ongoing clinical trials that apply these therapeutically beneficial EVs for treatment of COVID-19 patients as drug delivery platforms or vaccines (150, 161, 162). Therefore, like sepsis, EVs from diverse sources exhibit distinct effects during COVID-19 infections (163).

Conclusion

Overall, there is strong evidence to support the significance of EVs in the modulation of inflammation and cell death and the diseases related to these processes. EVs have been shown to play critical roles in both promoting and suppressing cell death and inflammation leading to beneficial as well as detrimental health outcomes. Over the last couple of decades, studies on EVs have shown their great potential to improve diagnosis and therapy in human diseases. As has been occasionally discussed in this article as well, EVs present as a valuable target for disease treatment, targeted drug-delivery, vaccine development and biomarker discovery. Furthermore, there is a need for developing better methods for isolation of pure EV subtypes and address the challenge of EV heterogeneity in preparations. The establishment of precise isolation methods is crucial before a particular function can be attributed to any specific EV subtype. Further, MSC-derived EVs are widely explored in preclinical and clinical settings due to their therapeutic potential. However, translation of such findings regarding EVs to clinical practices would require further exploration of their exact mode of biogenesis and cargo components, which are responsible for the desired phenotype to clearly understand the underlying mode of action. Despite these knowledge gaps and shortcomings, with further exploration employing improved methodologies and techniques, EVs remain as putative therapeutic targets for the future.

Author Contributions: LG planned the review. LG and RS contributed to the writing and editing of the review. RS prepared the figures.

Acknowledgments: The authors would like to thank Dr Christina Nedeva for valuable feedbacks and help in formatting the manuscript.

Conflicts of Interest: The authors declare no conflict of interest with respect to this manuscript.

Figure legends

Figure 1- Schematic representation of EV subtypes identified in eukaryotic cells. **A.** While live cells were initially thought to shed exosomes and microvesicles, recent developments have identified novel EV subtypes including exomeres and migrasomes. Whereas cancer cells also partake in secretion of large oncosomes. On the contrary, apoptotic cells which were long thought to shed only ApoBDs are now known to secrete other EV subtypes including ApoMVs and ApoExos. **B.** Various EV subtypes shed by live or apoptotic cells are distinct in their biogenesis, cargo, origin and size.

Figure 1





Figure 2- Context dependent roles of EVs in mediating cell death and inflammation. **A.** ApoEVs have the ability to either attenuate or promote cell death. MSC derived ApoEVs have regenerative function whereas ApoEVs from pathogen infected cells induce activation of immune cells, leading to enhanced targeting and clearance of pathogen cells. **B.** Similarly, EVs from healthy MSCs have regenerative function in preventing cell death and inflammation in various cell types. **C.** Contrastingly, EVs from diseased host cells may function in different ways depending on the recipient cell. For instance, cancer cell derived EVs promote cell death and inflammation in immune cells but prevent cell death and inflammation in other cancer cells. **D.** Infected cells can shed pathogen derived EVs or host EVs loaded with pathogen’s cargo. These EVs can promote cell death and inflammation in recipient host cells.



References

1. Kalra H, Drummen GPC, Mathivanan S. Focus on Extracellular Vesicles: Introducing the Next Small Big Thing. *International journal of molecular sciences*. 2016;17(2):170-.
2. Harding C, Heuser J, Stahl P. Endocytosis and intracellular processing of transferrin and colloidal gold-transferrin in rat reticulocytes: demonstration of a pathway for receptor shedding. *Eur J Cell Biol*. 1984;35(2):256-63.
3. Pan BT, Teng K, Wu C, Adam M, Johnstone RM. Electron microscopic evidence for externalization of the transferrin receptor in vesicular form in sheep reticulocytes. *J Cell Biol*. 1985;101(3):942-8.
4. Gangoda L, Boukouris S, Liem M, Kalra H, Mathivanan S. Extracellular vesicles including exosomes are mediators of signal transduction: are they protective or pathogenic? *Proteomics*. 2015;15(2-3):260-71.
5. Raposo G, Stoorvogel W. Extracellular vesicles: Exosomes, microvesicles, and friends. *Journal of Cell Biology*. 2013;200(4):373-83.
6. Kalluri R. The biology and function of exosomes in cancer. *J Clin Invest*. 2016;126(4):1208-15.
7. Howitt J, Hill AF. Exosomes in the Pathology of Neurodegenerative Diseases. *J Biol Chem*. 2016;291(52):26589-97.
8. Samuel M, Fonseka P, Sanwlan R, Gangoda L, Chee SH, Keerthikumar S, et al. Oral administration of bovine milk-derived extracellular vesicles induces senescence in the primary tumor but accelerates cancer metastasis. *Nature Communications*. 2021;12(1):3950.
9. Sanwlan R, Fonseka P, Chitti SV, Mathivanan S. Milk-Derived Extracellular Vesicles in Inter-Organism, Cross-Species Communication and Drug Delivery. *Proteomes*. 2020;8(2):11.
10. Sanwlan R, Fonseka P, Mathivanan S. Are Dietary Extracellular Vesicles Bioavailable and Functional in Consuming Organisms? *Subcell Biochem*. 2021;97:509-21.
11. Kang T, Atukorala I, Mathivanan S. Biogenesis of Extracellular Vesicles. *Subcell Biochem*. 2021;97:19-43.
12. Théry C, Zitvogel L, Amigorena S. Exosomes: composition, biogenesis and function. *Nature Reviews Immunology*. 2002;2(8):569-79.
13. Keerthikumar S, Gangoda L, Liem M, Fonseka P, Atukorala I, Ozcitti C, et al. Proteogenomic analysis reveals exosomes are more oncogenic than ectosomes. *Oncotarget*. 2015;6(17):15375-96.
14. Zhang H, Freitas D, Kim HS, Fabijanic K, Li Z, Chen H, et al. Identification of distinct nanoparticles and subsets of extracellular vesicles by asymmetric flow field-flow fractionation. *Nature Cell Biology*. 2018;20(3):332-43.
15. Anand S, Samuel M, Mathivanan S. Exomeres: A New Member of Extracellular Vesicles Family. In: Mathivanan S, Fonseka P, Nedeva C, Atukorala I, editors. *New Frontiers: Extracellular Vesicles*. Cham: Springer International Publishing; 2021. p. 89-97.
16. György B, Szabó TG, Pásztói M, Pál Z, Misják P, Aradi B, et al. Membrane vesicles, current state-of-the-art: emerging role of extracellular vesicles. *Cellular and molecular life sciences : CMLS*. 2011;68(16):2667-88.
17. Minciacchi VR, You S, Spinelli C, Morley S, Zandian M, Aspuria P-J, et al. Large oncosomes contain distinct protein cargo and represent a separate functional class of tumor-derived extracellular vesicles. *Oncotarget*. 2015;6(13):11327-41.
18. Zijlstra A, Di Vizio D. Size matters in nanoscale communication. *Nature cell biology*. 2018;20(3):228-30.
19. Tavano S, Heisenberg C-P. Migrasomes take center stage. *Nature Cell Biology*. 2019;21(8):918-20.
20. Atkin-Smith GK, Tixeira R, Paone S, Mathivanan S, Collins C, Liem M, et al. A novel mechanism of generating extracellular vesicles during apoptosis via a beads-on-a-string membrane structure. *Nature Communications*. 2015;6(1):7439.
21. Kerr JFR, Wyllie AH, Currie AR. Apoptosis: A Basic Biological Phenomenon with Wideranging Implications in Tissue Kinetics. *British Journal of Cancer*. 1972;26(4):239-57.
22. Kakarla R, Hur J, Kim YJ, Kim J, Chwae Y-J. Apoptotic cell-derived exosomes: messages from dying cells. *Experimental & Molecular Medicine*. 2020;52(1):1-6.
23. Schiller M, Parcina M, Heyder P, Foermer S, Ostrop J, Leo A, et al. Induction of Type I IFN Is a Physiological Immune Reaction to Apoptotic Cell-Derived Membrane Microparticles. 2012;189(4):1747-56.
24. Tucher C, Bode K, Schiller P, Claßen L, Birr C, Souto-Carneiro MM, et al. Extracellular Vesicle Subtypes Released From Activated or Apoptotic T-Lymphocytes Carry a Specific and Stimulus-Dependent Protein Cargo. 2018;9(534).
25. Sirois I, Raymond MA, Brassard N, Cailhier JF, Fedjaev M, Hamelin K, et al. Caspase-3-dependent export of TCTP: a novel pathway for antiapoptotic intercellular communication. *Cell death and differentiation*. 2011;18(3):549-62.
26. Sirois I, Groleau J, Pallet N, Brassard N, Hamelin K, Londono I, et al. Caspase activation regulates the extracellular export of autophagic vacuoles. *Autophagy*. 2012;8(6):927-37.
27. Pathan M, Fonseka P, Chitti SV, Kang T, Sanwlan R, Van Deun J, et al. Vesiclepedia 2019: a compendium of RNA, proteins, lipids and metabolites in extracellular vesicles. *Nucleic Acids Res*. 2019;47(D1):D516-d9.

28. Mori Y, Koike M, Moriishi E, Kawabata A, Tang H, Oyaizu H, et al. Human Herpesvirus-6 Induces MVB Formation, and Virus Egress Occurs by an Exosomal Release Pathway. *Traffic*. 2008;9(10):1728-42.
29. Kalra H, Gangoda L, Fonseka P, Chitti SV, Liem M, Keerthikumar S, et al. Extracellular vesicles containing oncogenic mutant β -catenin activate Wnt signalling pathway in the recipient cells. *Journal of Extracellular Vesicles*. 2019;8(1):1690217.
30. Chen G, Huang AC, Zhang W, Zhang G, Wu M, Xu W, et al. Exosomal PD-L1 contributes to immunosuppression and is associated with anti-PD-1 response. *Nature*. 2018;560(7718):382-6.
31. Boukouris S, Mathivanan S. Exosomes in bodily fluids are a highly stable resource of disease biomarkers. *Proteomics Clinical applications*. 2015;9(3-4):358-67.
32. Mathivanan S, Lim JWE, Tauro BJ, Ji H, Moritz RL, Simpson RJ. Proteomics Analysis of A33 Immunoaffinity-purified Exosomes Released from the Human Colon Tumor Cell Line LIM1215 Reveals a Tissue-specific Protein Signature*. *Molecular & Cellular Proteomics*. 2010;9(2):197-208.
33. Srivastava P. Interaction of Heat Shock Proteins with Peptides and Antigen Presenting Cells: Chaperoning of the Innate and Adaptive Immune Responses. *Annual Review of Immunology*. 2002;20(1):395-425.
34. Théry C, Ostrowski M, Segura E. Membrane vesicles as conveyors of immune responses. *Nat Rev Immunol*. 2009;9(8):581-93.
35. Borges FT, Reis LA, Schor N. Extracellular vesicles: structure, function, and potential clinical uses in renal diseases. *Brazilian journal of medical and biological research = Revista brasileira de pesquisas medicas e biologicas*. 2013;46(10):824-30.
36. Poon IK, Lucas CD, Rossi AG, Ravichandran KS. Apoptotic cell clearance: basic biology and therapeutic potential. *Nat Rev Immunol*. 2014;14(3):166-80.
37. Park SJ, Kim JM, Kim J, Hur J, Park S, Kim K, et al. Molecular mechanisms of biogenesis of apoptotic exosome-like vesicles and their roles as damage-associated molecular patterns. *Proceedings of the National Academy of Sciences of the United States of America*. 2018;115(50):E11721-e30.
38. Valadi H, Ekström K, Bossios A, Sjöstrand M, Lee JJ, Lötvall JO. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nature Cell Biology*. 2007;9(6):654-9.
39. Kalluri R, LeBleu VS. Discovery of Double-Stranded Genomic DNA in Circulating Exosomes. *Cold Spring Harbor symposia on quantitative biology*. 2016;81:275-80.
40. Kurywachak P, Kalluri R. An evolving function of DNA-containing exosomes in chemotherapy-induced immune response. *Cell Research*. 2017;27(6):722-3.
41. Llorente A, Skotland T, Sylvänne T, Kauhanen D, Róg T, Orłowski A, et al. Molecular lipidomics of exosomes released by PC-3 prostate cancer cells. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*. 2013;1831(7):1302-9.
42. Lydic TA, Townsend S, Adda CG, Collins C, Mathivanan S, Reid GE. Rapid and comprehensive 'shotgun' lipidome profiling of colorectal cancer cell derived exosomes. *Methods*. 2015;87:83-95.
43. Skotland T, Sandvig K, Llorente A. Lipids in exosomes: Current knowledge and the way forward. *Progress in Lipid Research*. 2017;66:30-41.
44. Kim CW, Lee HM, Lee TH, Kang C, Kleinman HK, Gho YS. Extracellular Membrane Vesicles from Tumor Cells Promote Angiogenesis via Sphingomyelin. 2002;62(21):6312-7.
45. Galluzzi L, Vitale I, Aaronson SA, Abrams JM, Adam D, Agostinis P, et al. Molecular mechanisms of cell death: recommendations of the Nomenclature Committee on Cell Death 2018. *Cell Death & Differentiation*. 2018;25(3):486-541.
46. Reinhardt TA, Sacco RE, Nonnecke BJ, Lippolis JD. Bovine milk proteome: Quantitative changes in normal milk exosomes, milk fat globule membranes and whey proteomes resulting from *Staphylococcus aureus* mastitis. *Journal of Proteomics*. 2013;82:141-54.
47. Miyake H, Lee C, Chusilp S, Bhalla M, Li B, Pitino M, et al. Human breast milk exosomes attenuate intestinal damage. *Pediatric surgery international*. 2020;36(2):155-63.
48. Chen T, Xie M-Y, Sun J-J, Ye R-S, Cheng X, Sun R-P, et al. Porcine milk-derived exosomes promote proliferation of intestinal epithelial cells. *Scientific Reports*. 2016;6(1):33862.
49. Gao HN, Guo HY, Zhang H, Xie XL, Wen PC, Ren FZ. Yak-milk-derived exosomes promote proliferation of intestinal epithelial cells in an hypoxic environment. *Journal of dairy science*. 2019;102(2):985-96.
50. Martin C, Patel M, Williams S, Arora H, Brawner K, Sims B. Human breast milk-derived exosomes attenuate cell death in intestinal epithelial cells. *Innate immunity*. 2018;24(5):278-84.
51. Melnik BC, Schmitz G. Milk's Role as an Epigenetic Regulator in Health and Disease. *Diseases (Basel, Switzerland)*. 2017;5(1).
52. Meng F, Henson R, Wehbe-Janek H, Ghoshal K, Jacob ST, Patel T. MicroRNA-21 Regulates Expression of the PTEN Tumor Suppressor Gene in Human Hepatocellular Cancer. *Gastroenterology*. 2007;133(2):647-58.
53. Murata T, Takayama K, Katayama S, Urano T, Horie-Inoue K, Ikeda K, et al. miR-148a is an androgen-responsive microRNA that promotes LNCaP prostate cell growth by repressing its target CAND1 expression. *Prostate Cancer and Prostatic Diseases*. 2010;13(4):356-61.

54. Fonseka P, Liem M, Ozcitti C, Adda CG, Ang C-S, Mathivanan S. Exosomes from N-Myc amplified neuroblastoma cells induce migration and confer chemoresistance to non-N-Myc amplified cells: implications of intra-tumour heterogeneity. *Journal of Extracellular Vesicles*. 2019;8(1):1597614.
55. Pavlyukov MS, Yu H, Bastola S, Minata M, Shender VO, Lee Y, et al. Apoptotic Cell-Derived Extracellular Vesicles Promote Malignancy of Glioblastoma Via Intercellular Transfer of Splicing Factors. *Cancer Cell*. 2018;34(1):119-35.e10.
56. Gallet R, Dawkins J, Valle J, Simsolo E, de Couto G, Middleton R, et al. Exosomes secreted by cardiosphere-derived cells reduce scarring, attenuate adverse remodelling, and improve function in acute and chronic porcine myocardial infarction. *European heart journal*. 2017;38(3):201-11.
57. Zhao G, Liu F, Liu Z, Zuo K, Wang B, Zhang Y, et al. MSC-derived exosomes attenuate cell death through suppressing AIF nucleus translocation and enhance cutaneous wound healing. *Stem Cell Research & Therapy*. 2020;11(1):174.
58. Wang Y, Zhang L, Li Y, Chen L, Wang X, Guo W, et al. Exosomes/microvesicles from induced pluripotent stem cells deliver cardioprotective miRNAs and prevent cardiomyocyte apoptosis in the ischemic myocardium. *Int J Cardiol*. 2015;192:61-9.
59. Liu L, Jin X, Hu CF, Li R, Zhou Z, Shen CX. Exosomes Derived from Mesenchymal Stem Cells Rescue Myocardial Ischaemia/Reperfusion Injury by Inducing Cardiomyocyte Autophagy Via AMPK and Akt Pathways. *Cellular physiology and biochemistry : international journal of experimental cellular physiology, biochemistry, and pharmacology*. 2017;43(1):52-68.
60. Yu B, Kim HW, Gong M, Wang J, Millard RW, Wang Y, et al. Exosomes secreted from GATA-4 overexpressing mesenchymal stem cells serve as a reservoir of anti-apoptotic microRNAs for cardioprotection. *International Journal of Cardiology*. 2015;182:349-60.
61. Li M, Liao L, Tian W. Extracellular Vesicles Derived From Apoptotic Cells: An Essential Link Between Death and Regeneration. 2020;8(1063).
62. Liu H, Liu S, Qiu X, Yang X, Bao L, Pu F, et al. Donor MSCs release apoptotic bodies to improve myocardial infarction via autophagy regulation in recipient cells. *Autophagy*. 2020;16(12):2140-55.
63. Gupta KH, Goldufsky JW, Wood SJ, Tardi NJ, Moorthy GS, Gilbert DZ, et al. Apoptosis and Compensatory Proliferation Signaling Are Coupled by Crkl-Containing Microvesicles. *Developmental cell*. 2017;41(6):674-84.e5.
64. Winau F, Weber S, Sad S, de Diego J, Hoops SL, Breiden B, et al. Apoptotic vesicles crossprime CD8 T cells and protect against tuberculosis. *Immunity*. 2006;24(1):105-17.
65. Kranich J, Krautler NJ, Falsig J, Ballmer B, Li S, Hutter G, et al. Engulfment of cerebral apoptotic bodies controls the course of prion disease in a mouse strain-dependent manner. *J Exp Med*. 2010;207(10):2271-81.
66. Schorey JS, Cheng Y, Singh PP, Smith VL. Exosomes and other extracellular vesicles in host-pathogen interactions. *EMBO reports*. 2015;16(1):24-43.
67. Kuipers ME, Hokke CH, Smits HH, Nolte-'t Hoen ENM. Pathogen-Derived Extracellular Vesicle-Associated Molecules That Affect the Host Immune System: An Overview. 2018;9(2182).
68. Munhoz da Rocha IF, Amatuzzi RF, Lucena ACR, Faoro H, Alves LR. Cross-Kingdom Extracellular Vesicles EV-RNA Communication as a Mechanism for Host-Pathogen Interaction. *Frontiers in Cellular and Infection Microbiology*. 2020;10(684).
69. Martins SdT, Alves LR. Extracellular Vesicles in Viral Infections: Two Sides of the Same Coin? *Frontiers in Cellular and Infection Microbiology*. 2020;10(737).
70. Rahimian P, He JJ. Exosome-associated release, uptake, and neurotoxicity of HIV-1 Tat protein. *Journal of NeuroVirology*. 2016;22(6):774-88.
71. Keryer-Bibens C, Pioche-Durieu C, Villemant C, Souquère S, Nishi N, Hirashima M, et al. Exosomes released by EBV-infected nasopharyngeal carcinoma cells convey the viral Latent Membrane Protein 1 and the immunomodulatory protein galectin 9. *BMC Cancer*. 2006;6(1):283.
72. Balaji KN, Goyal G, Narayana Y, Srinivas M, Chaturvedi R, Mohammad S. Apoptosis triggered by Rv1818c, a PE family gene from Mycobacterium tuberculosis is regulated by mitochondrial intermediates in T cells. *Microbes and infection*. 2007;9(3):271-81.
73. Vanaja Sivapriya K, Russo Ashley J, Behl B, Banerjee I, Yankova M, Deshmukh Sachin D, et al. Bacterial Outer Membrane Vesicles Mediate Cytosolic Localization of LPS and Caspase-11 Activation. *Cell*. 2016;165(5):1106-19.
74. Deo P, Chow SH, Hay ID, Kleifeld O, Costin A, Elgass KD, et al. Outer membrane vesicles from Neisseria gonorrhoeae target PorB to mitochondria and induce apoptosis. *PLoS pathogens*. 2018;14(3):e1006945.
75. Kunsman L, Rüter C, Bauwens A, Greune L, Glüder M, Kemper B, et al. Virulence from vesicles: Novel mechanisms of host cell injury by Escherichia coli O104:H4 outbreak strain. *Scientific Reports*. 2015;5(1):13252.

76. Guay C, Kruit JK, Rome S, Menoud V, Mulder NL, Jurdzinski A, et al. Lymphocyte-Derived Exosomal MicroRNAs Promote Pancreatic β Cell Death and May Contribute to Type 1 Diabetes Development. *Cell Metab.* 2019;29(2):348-61.e6.
77. Wieckowski EU, Visus C, Szajnik M, Szczepanski MJ, Storkus WJ, Whiteside TL. Tumor-derived microvesicles promote regulatory T cell expansion and induce apoptosis in tumor-reactive activated CD8⁺ T lymphocytes. *J Immunol.* 2009;183(6):3720-30.
78. Mazzeo C, Cañas JA, Zafra MP, Rojas Marco A, Fernández-Nieto M, Sanz V, et al. Exosome secretion by eosinophils: A possible role in asthma pathogenesis. *J Allergy Clin Immunol.* 2015;135(6):1603-13.
79. Cañas JA, Sastre B, Rodrigo-Muñoz JM, Fernández-Nieto M, Barranco P, Quirce S, et al. Eosinophil-derived exosomes contribute to asthma remodelling by activating structural lung cells. *Clin Exp Allergy.* 2018;48(9):1173-85.
80. Mitra S, Exline M, Habyarimana F, Gavrilin MA, Baker PJ, Masters SL, et al. Microparticulate Caspase 1 Regulates Gasdermin D and Pulmonary Vascular Endothelial Cell Injury. *American journal of respiratory cell and molecular biology.* 2018;59(1):56-64.
81. Mitra S, Wewers MD, Sarkar A. Mononuclear Phagocyte-Derived Microparticulate Caspase-1 Induces Pulmonary Vascular Endothelial Cell Injury. *PLoS One.* 2015;10(12):e0145607.
82. Kubo H. Extracellular Vesicles in Lung Disease. *Chest.* 2018;153(1):210-6.
83. Medzhitov R. Inflammation 2010: new adventures of an old flame. *Cell.* 2010;140(6):771-6.
84. Davidovich P, Kearney CJ, Martin SJ. Inflammatory outcomes of apoptosis, necrosis and necroptosis. *Biol Chem.* 2014;395(10):1163-71.
85. Jorgensen I, Miao EA. Pyroptotic cell death defends against intracellular pathogens. *Immunol Rev.* 2015;265(1):130-42.
86. Hannood S, Nasuruddin DN. Acute Inflammatory Response. *StatPearls.* Treasure Island (FL)2021.
87. Heneka MT, Kummer MP, Stutz A, Delekate A, Schwartz S, Vieira-Saecker A, et al. NLRP3 is activated in Alzheimer's disease and contributes to pathology in APP/PS1 mice. *Nature.* 2013;493(7434):674-8.
88. Ising C, Venegas C, Zhang S, Scheiblich H, Schmidt SV, Vieira-Saecker A, et al. NLRP3 inflammasome activation drives tau pathology. *Nature.* 2019;575(7784):669-73.
89. Duewell P, Kono H, Rayner KJ, Sirois CM, Vladimer G, Bauernfeind FG, et al. NLRP3 inflammasomes are required for atherogenesis and activated by cholesterol crystals. *Nature.* 2010;464(7293):1357-61.
90. Chen L, Deng H, Cui H, Fang J, Zuo Z, Deng J, et al. Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget.* 2018;9(6):7204-18.
91. Raposo G, Nijman HW, Stoorvogel W, Liejendekker R, Harding CV, Melief CJ, et al. B lymphocytes secrete antigen-presenting vesicles. *J Exp Med.* 1996;183(3):1161-72.
92. Zitvogel L, Regnault A, Lozier A, Wolfers J, Flament C, Tenza D, et al. Eradication of established murine tumors using a novel cell-free vaccine: dendritic cell-derived exosomes. *Nat Med.* 1998;4(5):594-600.
93. Gasser O, Schifferli JA. Activated polymorphonuclear neutrophils disseminate anti-inflammatory microparticles by ectocytosis. *Blood.* 2004;104(8):2543-8.
94. Hardy MP, Audemard É, Migneault F, Feghaly A, Brochu S, Gendron P, et al. Apoptotic endothelial cells release small extracellular vesicles loaded with immunostimulatory viral-like RNAs. *Sci Rep.* 2019;9(1):7203.
95. Mardpour S, Hamidieh AA, Taleahmad S, Sharifzad F, Taghikhani A, Baharvand H. Interaction between mesenchymal stromal cell-derived extracellular vesicles and immune cells by distinct protein content. *J Cell Physiol.* 2019;234(6):8249-58.
96. Obregon C, Rothen-Rutishauser B, Gerber P, Gehr P, Nicod LP. Active uptake of dendritic cell-derived exovesicles by epithelial cells induces the release of inflammatory mediators through a TNF-alpha-mediated pathway. *Am J Pathol.* 2009;175(2):696-705.
97. Keerthikumar S, Chisanga D, Ariyaratne D, Al Saffar H, Anand S, Zhao K, et al. ExoCarta: A Web-Based Compendium of Exosomal Cargo. *J Mol Biol.* 2016;428(4):688-92.
98. Hotchkiss RS, Moldawer LL, Opal SM, Reinhart K, Turnbull IR, Vincent JL. Sepsis and septic shock. *Nat Rev Dis Primers.* 2016;2:16045.
99. Raeven P, Zipperle J, Drechsler S. Extracellular Vesicles as Markers and Mediators in Sepsis. *Theranostics.* 2018;8(12):3348-65.
100. Labbe K, Saleh M. Cell death in the host response to infection. *Cell death and differentiation.* 2008;15(9):1339-49.
101. Vincent JL, Sakr Y, Sprung CL, Ranieri VM, Reinhart K, Gerlach H, et al. Sepsis in European intensive care units: results of the SOAP study. *Crit Care Med.* 2006;34(2):344-53.
102. Park KS, Choi KH, Kim YS, Hong BS, Kim OY, Kim JH, et al. Outer membrane vesicles derived from *Escherichia coli* induce systemic inflammatory response syndrome. *PLoS One.* 2010;5(6):e11334.
103. Hotchkiss RS, Nicholson DW. Apoptosis and caspases regulate death and inflammation in sepsis. *Nat Rev Immunol.* 2006;6(11):813-22.

104. Lenassi M, Cagney G, Liao M, Vaupotic T, Bartholomeeusen K, Cheng Y, et al. HIV Nef is secreted in exosomes and triggers apoptosis in bystander CD4⁺ T cells. *Traffic*. 2010;11(1):110-22.
105. Kim OY, Choi SJ, Jang SC, Park KS, Kim SR, Choi JP, et al. Bacterial protoplast-derived nanovesicles as vaccine delivery system against bacterial infection. *Nano Lett*. 2015;15(1):266-74.
106. Larsson A, Lundahl T, Eriksson M, Lundkvist K, Lindahl T. Endotoxin induced platelet microvesicle formation measured by flow cytometry. *Platelets*. 1996;7(3):153-8.
107. Nieuwland R, Berckmans RJ, McGregor S, Boing AN, Romijn FP, Westendorp RG, et al. Cellular origin and procoagulant properties of microparticles in meningococcal sepsis. *Blood*. 2000;95(3):930-5.
108. Lee YJ, Jy W, Horstman LL, Janania J, Reyes Y, Kelley RE, et al. Elevated platelet microparticles in transient ischemic attacks, lacunar infarcts, and multiinfarct dementias. *Thromb Res*. 1993;72(4):295-304.
109. Holme PA, Solum NO, Brosstad F, Roger M, Abdelnoor M. Demonstration of platelet-derived microvesicles in blood from patients with activated coagulation and fibrinolysis using a filtration technique and western blotting. *Thromb Haemost*. 1994;72(5):666-71.
110. Lundahl TH, Lindahl TL, Fagerberg IH, Egberg N, Bunescu A, Larsson A. Activated platelets and impaired platelet function in intensive care patients analyzed by flow cytometry. *Blood Coagul Fibrinolysis*. 1996;7(2):218-20.
111. Denning NL, Aziz M, Gurien SD, Wang P. DAMPs and NETs in Sepsis. *Front Immunol*. 2019;10:2536.
112. Gangoda L, Schenk RL, Best SA, Nedeva C, Louis C, D'Silva DB, et al. Absence of pro-survival A1 has no impact on inflammatory cell survival in vivo during acute lung inflammation and peritonitis. *Cell death and differentiation*. 2021.
113. Nedeva C, Menassa J, Duan M, Liu C, Doerflinger M, Kueh AJ, et al. TREML4 receptor regulates inflammation and innate immune cell death during polymicrobial sepsis. *Nat Immunol*. 2020;21(12):1585-96.
114. Herrmann IK, Bertazzo S, O'Callaghan DJ, Schlegel AA, Kallepitis C, Antcliffe DB, et al. Differentiating sepsis from non-infectious systemic inflammation based on microvesicle-bacteria aggregation. *Nanoscale*. 2015;7(32):13511-20.
115. Essandoh K, Yang L, Wang X, Huang W, Qin D, Hao J, et al. Blockade of exosome generation with GW4869 dampens the sepsis-induced inflammation and cardiac dysfunction. *Biochim Biophys Acta*. 2015;1852(11):2362-71.
116. Gao K, Jin J, Huang C, Li J, Luo H, Li L, et al. Exosomes Derived From Septic Mouse Serum Modulate Immune Responses via Exosome-Associated Cytokines. *Front Immunol*. 2019;10:1560.
117. Fitzgerald W, Freeman ML, Lederman MM, Vasilieva E, Romero R, Margolis L. A System of Cytokines Encapsulated in ExtraCellular Vesicles. *Sci Rep*. 2018;8(1):8973.
118. Fujimi S, Ogura H, Tanaka H, Koh T, Hosotsubo H, Nakamori Y, et al. Activated polymorphonuclear leukocytes enhance production of leukocyte microparticles with increased adhesion molecules in patients with sepsis. *J Trauma*. 2002;52(3):443-8.
119. Shah B, Sullivan CJ, Lonergan NE, Stanley S, Soult MC, Britt LD. Circulating bacterial membrane vesicles cause sepsis in rats. *Shock*. 2012;37(6):621-8.
120. Gambim MH, do Carmo Ade O, Marti L, Verissimo-Filho S, Lopes LR, Janiszewski M. Platelet-derived exosomes induce endothelial cell apoptosis through peroxynitrite generation: experimental evidence for a novel mechanism of septic vascular dysfunction. *Crit Care*. 2007;11(5):R107.
121. Ren C, Yao RQ, Zhang H, Feng YW, Yao YM. Sepsis-associated encephalopathy: a vicious cycle of immunosuppression. *J Neuroinflammation*. 2020;17(1):14.
122. Chen L, Yao X, Yao H, Ji Q, Ding G, Liu X. Exosomal miR-103-3p from LPS-activated THP-1 macrophage contributes to the activation of hepatic stellate cells. *FASEB J*. 2020;34(4):5178-92.
123. Timar CI, Lorincz AM, Csepanyi-Komi R, Valyi-Nagy A, Nagy G, Buzas EI, et al. Antibacterial effect of microvesicles released from human neutrophilic granulocytes. *Blood*. 2013;121(3):510-8.
124. Mostefai HA, Meziani F, Mastronardi ML, Agouni A, Heymes C, Sargentini C, et al. Circulating microparticles from patients with septic shock exert protective role in vascular function. *Am J Respir Crit Care Med*. 2008;178(11):1148-55.
125. Dalli J, Norling LV, Montero-Melendez T, Federici Canova D, Lashin H, Pavlov AM, et al. Microparticle alpha-2-macroglobulin enhances pro-resolving responses and promotes survival in sepsis. *EMBO Mol Med*. 2014;6(1):27-42.
126. Miksa M, Wu R, Dong W, Komura H, Amin D, Ji Y, et al. Immature dendritic cell-derived exosomes rescue septic animals via milk fat globule epidermal growth factor-factor VIII [corrected]. *J Immunol*. 2009;183(9):5983-90.
127. Xu Y, Ku X, Wu C, Cai C, Tang J, Yan W. Exosomal proteome analysis of human plasma to monitor sepsis progression. *Biochem Biophys Res Commun*. 2018;499(4):856-61.
128. Englert JA, Bobba C, Baron RM. Integrating molecular pathogenesis and clinical translation in sepsis-induced acute respiratory distress syndrome. *JCI Insight*. 2019;4(2).

129. Fujita Y, Kosaka N, Araya J, Kuwano K, Ochiya T. Extracellular vesicles in lung microenvironment and pathogenesis. *Trends Mol Med.* 2015;21(9):533-42.
130. Kesimer M, Scull M, Brighton B, DeMaria G, Burns K, O'Neal W, et al. Characterization of exosome-like vesicles released from human tracheobronchial ciliated epithelium: a possible role in innate defense. *FASEB J.* 2009;23(6):1858-68.
131. Mohan A, Agarwal S, Clauss M, Britt NS, Dhillon NK. Extracellular vesicles: novel communicators in lung diseases. *Respir Res.* 2020;21(1):175.
132. Bourdonnay E, Zaslona Z, Penke LR, Speth JM, Schneider DJ, Przybranowski S, et al. Transcellular delivery of vesicular SOCS proteins from macrophages to epithelial cells blunts inflammatory signaling. *J Exp Med.* 2015;212(5):729-42.
133. Letsiou E, Sammani S, Zhang W, Zhou T, Quijada H, Moreno-Vinasco L, et al. Pathologic mechanical stress and endotoxin exposure increases lung endothelial microparticle shedding. *Am J Respir Cell Mol Biol.* 2015;52(2):193-204.
134. Li CJ, Liu Y, Chen Y, Yu D, Williams KJ, Liu ML. Novel proteolytic microvesicles released from human macrophages after exposure to tobacco smoke. *Am J Pathol.* 2013;182(5):1552-62.
135. Moon HG, Zheng Y, An CH, Kim YK, Jin Y. CCN1 secretion induced by cigarette smoking extracts augments IL-8 release from bronchial epithelial cells. *PLoS One.* 2013;8(7):e68199.
136. Schutze N, Rucker N, Muller J, Adamski J, Jakob F. 5' flanking sequence of the human immediate early responsive gene *ccn1* (*cyr61*) and mapping of polymorphic CA repeat sequence motifs in the human *ccn1* (*cyr61*) locus. *Mol Pathol.* 2001;54(3):170-5.
137. Moon HG, Kim SH, Gao J, Quan T, Qin Z, Osorio JC, et al. CCN1 secretion and cleavage regulate the lung epithelial cell functions after cigarette smoke. *Am J Physiol Lung Cell Mol Physiol.* 2014;307(4):L326-37.
138. Kulshreshtha A, Ahmad T, Agrawal A, Ghosh B. Proinflammatory role of epithelial cell-derived exosomes in allergic airway inflammation. *J Allergy Clin Immunol.* 2013;131(4):1194-203, 203 e1-14.
139. Hamid Q, Tulic MK, Liu MC, Moqbel R. Inflammatory cells in asthma: mechanisms and implications for therapy. *J Allergy Clin Immunol.* 2003;111(1 Suppl):S5-S12; discussion S-7.
140. Skokos D, Le Panse S, Villa I, Rousselle JC, Peronet R, Namane A, et al. Nonspecific B and T cell-stimulatory activity mediated by mast cells is associated with exosomes. *Int Arch Allergy Immunol.* 2001;124(1-3):133-6.
141. Xie G, Yang H, Peng X, Lin L, Wang J, Lin K, et al. Mast cell exosomes can suppress allergic reactions by binding to IgE. *J Allergy Clin Immunol.* 2018;141(2):788-91.
142. Canas JA, Sastre B, Mazzeo C, Fernandez-Nieto M, Rodrigo-Munoz JM, Gonzalez-Guerra A, et al. Exosomes from eosinophils autoregulate and promote eosinophil functions. *J Leukoc Biol.* 2017;101(5):1191-9.
143. Admyre C, Johansson SM, Paulie S, Gabrielsson S. Direct exosome stimulation of peripheral human T cells detected by ELISPOT. *Eur J Immunol.* 2006;36(7):1772-81.
144. Admyre C, Bohle B, Johansson SM, Focke-Tejkl M, Valenta R, Scheynius A, et al. B cell-derived exosomes can present allergen peptides and activate allergen-specific T cells to proliferate and produce TH2-like cytokines. *J Allergy Clin Immunol.* 2007;120(6):1418-24.
145. Lee KY. Pneumonia, Acute Respiratory Distress Syndrome, and Early Immune-Modulator Therapy. *Int J Mol Sci.* 2017;18(2).
146. Tsushima K, King LS, Aggarwal NR, De Gorordo A, D'Alessio FR, Kubo K. Acute lung injury review. *Intern Med.* 2009;48(9):621-30.
147. Matthay MA, Ware LB, Zimmerman GA. The acute respiratory distress syndrome. *J Clin Invest.* 2012;122(8):2731-40.
148. Lee H, Zhang D, Laskin DL, Jin Y. Functional Evidence of Pulmonary Extracellular Vesicles in Infectious and Noninfectious Lung Inflammation. *J Immunol.* 2018;201(5):1500-9.
149. Jiang K, Yang J, Guo S, Zhao G, Wu H, Deng G. Peripheral Circulating Exosome-Mediated Delivery of miR-155 as a Novel Mechanism for Acute Lung Inflammation. *Mol Ther.* 2019;27(10):1758-71.
150. Pocsfalvi G, Mammadova R, Ramos Juarez AP, Bokka R, Trepiccione F, Capasso G. COVID-19 and Extracellular Vesicles: An Intriguing Interplay. *Kidney Blood Press Res.* 2020;45(5):661-70.
151. Rodrigues TS, de Sa KSG, Ishimoto AY, Becerra A, Oliveira S, Almeida L, et al. Inflammasomes are activated in response to SARS-CoV-2 infection and are associated with COVID-19 severity in patients. *J Exp Med.* 2021;218(3).
152. Xiao K, Hou F, Huang X, Li B, Qian ZR, Xie L. Mesenchymal stem cells: current clinical progress in ARDS and COVID-19. *Stem Cell Res Ther.* 2020;11(1):305.
153. Barberis E, Vanella VV, Falasca M, Caneapero V, Cappellano G, Raineri D, et al. Circulating Exosomes Are Strongly Involved in SARS-CoV-2 Infection. *Front Mol Biosci.* 2021;8:632290.
154. Hassanpour M, Rezaie J, Nouri M, Panahi Y. The role of extracellular vesicles in COVID-19 virus infection. *Infect Genet Evol.* 2020;85:104422.
155. Song JW, Lam SM, Fan X, Cao WJ, Wang SY, Tian H, et al. Omics-Driven Systems Interrogation of Metabolic Dysregulation in COVID-19 Pathogenesis. *Cell Metab.* 2020;32(2):188-202 e5.

156. Balbi C, Burrello J, Bolis S, Lazzarini E, Biemmi V, Pianezzi E, et al. Circulating extracellular vesicles are endowed with enhanced procoagulant activity in SARS-CoV-2 infection. *EBioMedicine*. 2021;67:103369.
157. Guervilly C, Bonifay A, Burtey S, Sabatier F, Cauchois R, Abdili E, et al. Dissemination of extreme levels of extracellular vesicles: tissue factor activity in patients with severe COVID-19. *Blood Adv*. 2021;5(3):628-34.
158. Cocozza F, Nevo N, Piovesana E, Lahaye X, Buchrieser J, Schwartz O, et al. Extracellular vesicles containing ACE2 efficiently prevent infection by SARS-CoV-2 Spike protein-containing virus. *J Extracell Vesicles*. 2020;10(2):e12050.
159. Sengupta V, Sengupta S, Lazo A, Woods P, Nolan A, Bremer N. Exosomes Derived from Bone Marrow Mesenchymal Stem Cells as Treatment for Severe COVID-19. *Stem Cells Dev*. 2020;29(12):747-54.
160. Tsiapalis D, O'Driscoll L. Mesenchymal Stem Cell Derived Extracellular Vesicles for Tissue Engineering and Regenerative Medicine Applications. *Cells*. 2020;9(4).
161. O'Driscoll L. Extracellular vesicles from mesenchymal stem cells as a Covid-19 treatment. *Drug Discov Today*. 2020;25(7):1124-5.
162. Khalaj K, Figueira RL, Antounians L, Lauriti G, Zani A. Systematic review of extracellular vesicle-based treatments for lung injury: are EVs a potential therapy for COVID-19? *J Extracell Vesicles*. 2020;9(1):1795365.
163. Xia X, Yuan P, Liu Y, Wang Y, Cao W, Zheng JC. Emerging roles of extracellular vesicles in COVID-19, a double-edged sword? *Immunology*. 2021;163(4):416-30.