

Article

Not peer-reviewed version

Level Weights for Modeling with Complex Survey Data

[Charles Natuhanya](#) *

Posted Date: 20 March 2023

doi: 10.20944/preprints202303.0339.v1

Keywords: complex survey design; multilevel models; nonresponse; sampling probability; survey data; Uganda; weighting



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Level Weights for Modeling with Complex Survey Data

Charles Natuhamya

Makerere University School of Public Health; natuhamyac@gmail.com

Abstract: Introduction: Weighting is widely used in applied statistics especially while dealing with survey data. In recent years, multilevel modeling under complex survey designs has increased, resulting into demand for level weights. However, survey data that are accessible to the public for use usually do not contain level weights that are useful in multilevel modeling, but final survey weights that are only appropriate for single level analyses. In this paper, we demonstrate how the final survey weights can be used to estimate level weights for multilevel data analysis, and compare a model that applied level weights with one that applied the final survey weights. **Methods:** A framework for approximating level weights proposed by the Demographic Health Survey (DHS) program was used while estimating level weights. Models were fitted using a multilevel mixed effects logistic regression method. Estimates from a model that applied survey weights was compared to those from a model that applied level weights. **Results:** Application of final survey weights instead of level weights underestimated standard errors and led to loss of precision of model estimates. **Conclusions:** Use of level weights produces estimates with high precision and yields correct values of standard errors hence appropriately informing inference.

Keywords: complex survey design; multilevel models; nonresponse; sampling probability; survey data; Uganda; weighting

1. Introduction

Weighting is a correction technique applied to survey data to improve the accuracy of the survey estimates (Lavrakas, 2008) especially for health surveys that are typically based on complex survey designs like stratified multi-stage clustered designs (Graubard & Korn, 1996). Survey data is usually weighted for two reasons; to correct for unequal probabilities of selection that often have occurred during sampling and to compensate for survey nonresponse during data collection (Gelman, 2007; Kalton & Flores-Cervantes, 2003; Lavrakas, 2008; Lepkowski, Mosher, Groves, West, & Wagner, 2013; Little & Vartivarian, 2003; Pfeffermann-a, 1993; Seaman & White, 2013; Wagemaker, 2020).

Though widely accepted in descriptive analyses, weights have been criticized by some modelers in analytical inference (Pfeffermann-a, 1993). However, weighting sample observations produces consistent estimators of the model parameters (Bethlehem & Keller, 1987; Skinner & Mason), averts model misspecification (Pfeffermann-b, 1996) and reduces bias in the survey estimates (Chen, Gelman, Tracy, Norris, & Galea, 2015; Kim & Skinner, 2013) while analyzing data from unequal probability designs (Nahorniak, Larsen, Volk, & Jordan, 2015). Despite efforts to maximize response rates, virtually all sample surveys are prone to nonresponses (Wun, Ezzati-Rice, Diaz-Tena, & Greenblatt, 2007).

One of the critical challenges faced by analysts using Demographic and Health Survey (DHS) program datasets for multilevel modeling and other survey datasets is that sampling weights related to the specific levels of the multistage design (level weights) are not incorporated in these datasets. Providing the exact level weights may create disclosure risks where dataset users may recognize specific clusters, households, or individuals within the clusters if they have access to sampling frames. Hence, instead of level-specific weights, the final survey weights are made available in the public-used datasets. Yet, multilevel modeling requires level-specific sampling weights instead of the final survey weights, the product of the level weights. The final survey weights are only adequate for most analytical purposes besides multilevel modeling (Elkasabi, Ren, & Pullum, 2020). Ignoring these level weights however, may lead to erroneous inferences with respect to the sample design (Hakim, Bhuiyan, Akter, & Zaman, 2020; West, et al., 2015).

Notably, few guidelines for integrating weights into multilevel models are in place (Carle, 2009). Hence, this paper aimed at demonstrating the estimation of level weights and comparing weighted models; one that applied the final survey weights and another that applied the estimated level weights, using the Uganda Malaria Indicator Survey (UMIS) data of 2018/19.

2. Methods

Source of Data and Study Population

The study applied secondary data from the latest UMIS of 2018/19. The data are based on a two-stage cluster and stratified sampling procedure, meeting the requirements of a complex survey design. A total of 7,632 children below 5 years of age that were tested for anaemia and malaria infection formed the study population. Blood samples from the children to be tested were collected by finger or heel prick. Anaemia and malaria testing was carried out by health technicians (NMCD, UBOS, & ICF, 2020).

Weighting

Sampling weights are required for analysis to ensure that the survey results are representative at the national level as well as the domain level due to the non-proportional allocation of the sample to different regions and to their urban and rural areas, and the possible differences in response rates (UBOS & ICF, 2018). Since the 2018/19 UMIS sample is a two-stage stratified cluster sample, sampling weights were calculated separately, based on sampling probabilities for each sampling stage and for each cluster (NMCD, UBOS, & ICF, 2020).

In this study, we used a procedure for estimating level weights in Malaria Indicator Surveys (MIS) proposed by the Demographic Health Survey (DHS) program (Elkasabi, Ren, & Pullum, 2020). The procedure required information available in the UMIS datasets and the final report. The following steps were followed to approximate level-2 (household) and level-3 (enumeration area) as outlined in Table 1.

Table 1. Steps to Approximate Household (HH) and Enumeration Area (EA) Level Weights from Final Household Survey Weights.

Steps	HH and EA weights
1 Apply the estimated normalization factor to de-normalize the final survey weight	$d_{hi}^{HH} = HV005_{hi} \frac{\hat{M}}{m^c}$
2 Approximate the level-3 weight (w_3^{HH})	$w_3^{HH} = w_{3hi} = \frac{A_h}{a_h^c} f^\alpha,$ $f = d_{hi}^{HH} / \left(\frac{A_h}{a_h^c} \times \frac{\bar{M}_h}{s_h} \right)$ and $0 \leq \alpha \leq 1$
3 Estimate the level-2 weight ($w_{2\setminus 3}^{HH}$)	$w_{2\setminus 3}^{HH} = w_{2hi} = \frac{d_{hi}^{HH}}{w_{3hi}}$

- Where;
- HV005 is the final household survey weight variable, from the household recode (HR) dataset.
 - a_h^c is the number of finalized EAs in stratum or region h for the strata. The number of interviewed EAs, a_h^c was calculated from the household (HR) dataset.
 - M_h is the number of households in stratum h for all strata.
 - M is the number of households in the whole of Uganda according to the Uganda Population and Housing Census of 2014.
 - M_{hi} is the number of households in EA i per EA. These numbers were estimated using the average number of households in each EA in strata \bar{M}_h according to the most recent Uganda Population and Housing Census data of 2014.
 - m^c is the number of complete households in the survey.
 - \hat{M} is the approximated household number at the time of the survey in the whole country. This was approximated by the number of households in Uganda according to the Uganda Population and Housing Census of 2014.

The applied variation factor f depends on the exponent specification α , $0 \leq \alpha \leq 1$ thus contributing different degrees to $w_{2\setminus 3}^{HH}$ and w_3^{HH} . The factor is fully assigned to w_3^{HH} and also fully assigned to $w_{2\setminus 3}^{HH}$ when $\alpha = 1$ and $\alpha = 0$, respectively. In the case of this study, it is equally assigned to the two weights as $\alpha = 0.5$. In case extreme values are applied, all the weight variation is attributed to either the level-2 or level-3 weight.

De-normalization of the final survey weight was done to restore the original scale of the survey weight using the available information in the UMIS dataset and the UMIS final report. The de-normalized final survey weights in the first step were estimated as:

$$d_{hi}^{HH} = HV005_{hi} \frac{\hat{M}}{m^c}$$

where M is the number of households in the entire country.

Multilevel mixed effects models

Weighted multilevel mixed effects logistic regression models were specified to explain the contextual area associations. The model is represented in the equation below:

$$\ln\left(\frac{p_{ijk}}{1 - p_{ijk}}\right) = \beta_0 + \beta_1 X_{ijk} + \eta_k + \xi_{jk}$$

- \ln is the natural logarithm.
- p_{ijk} is the probability of testing positive for malaria for the i^{th} child in the j^{th} household and k^{th} EA.
- β_0 is the average log-odds of malaria infection.
- X_{ijk} is a covariate at level-1 for the i^{th} child in the j^{th} household and k^{th} EA.
- β_1 denotes the slope related with X_{ijk} representing the association between the individual child covariates and the log-odds of malaria infection.
- η_k is the k^{th} EA random effect.
- ξ_{jk} is the j^{th} household random effect.

Note that the random effects are assumed to be independently and identically distributed to one another with zero mean and σ_η and σ_ξ variances, for EA and household respectively.

Model comparison

Design factor (deft) was used to compare model estimates since nonresponse results into loss in the precision of survey estimates, primarily due to reduced sample size (Brick, 2013). A model was therefore considered better if it produced estimates with lower deft values in general. Lower deft values are associated with lower loss of precision of model estimates (Sturgis, 2004). The deft was calculated as follows:

$$deft = \sqrt{deff} = \sqrt{1 + rho(n - 1)}$$

Where;

- $deff$ is the design effect.
- rho is the intra-class correlation for the variable in question.
- n is the size of the cluster.

3. Results

Comparison of Models Using Standard Errors of Model Estimates

Findings from this study indicated that the model weighted with the final survey weights underestimated standard errors compared to one that was weighted with the estimated level weights (Figure 1). Underestimated standard errors in turn resulted into narrow confidence intervals (Table 2 and Figure 3), presenting seemingly high accuracy of model estimates and likely overconfidence in the model estimates. This is worth paying attention to because overconfidence in results leads to rejecting the null hypothesis when it should not be the case (Type I error) which most likely misleads inference.




Figure 1. Comparison of standard errors of model estimates (survey weights versus level weights).

Table 2. Comparison of multilevel mixed effects logistic regression model estimates.

Model estimates	Survey weighted model					Level weighted model				
	OR	SE	P	(95% CI)		OR	SE	P	(95% CI)	
β1	0.98	0.10	0.85	0.80	1.20	0.98	0.10	0.88	0.80	1.21
β2	1.40	0.05	0.00	1.31	1.50	1.42	0.05	0.00	1.33	1.52
β3	0.92	0.16	0.63	0.66	1.29	0.94	0.17	0.72	0.66	1.33
β4	0.50	0.14	0.02	0.29	0.88	0.53	0.16	0.03	0.30	0.95
β5	0.78	0.12	0.09	0.58	1.04	0.77	0.12	0.11	0.56	1.06
β6	0.99	0.18	0.96	0.69	1.43	1.03	0.20	0.87	0.70	1.52
β7	0.36	0.08	0.00	0.23	0.55	0.42	0.09	0.00	0.27	0.64
β8	0.68	0.22	0.23	0.36	1.29	1.05	0.36	0.90	0.53	2.08
β9	1.35	0.68	0.55	0.50	3.62	1.37	0.82	0.60	0.42	4.42
β10	1.19	0.17	0.23	0.90	1.57	1.12	0.16	0.44	0.84	1.49
β11	0.98	0.00	0.00	0.99	1.00	0.99	0.00	0.00	0.97	1.00

CI: Confidence interval, OR: Odds ratio, SE: Standard error, P: p-value.

Comparison of Models Using Design Factor Values of Model Estimates

Design factor (deft) values for the model weighted with final survey weights were generally higher than those for the model weighted with level weights (Figure 2 and Table 3). This indicates that the use of final survey weights resulted into a considerable loss of precision for model estimates compared to the use of level weights. Almost half (5 of the 11) estimates for the model weighted with final survey weights produced deft values greater than 1.2, a value commonly taken to indicate sizeable variance inflation compared to only 1 estimate for the model weighted with level weights.

Table 3. Deff and Deft values for survey weighted and level weighted model estimates.

Model estimates	Survey weighted model		Level weighted model	
	Deff	Deft	Deff	Deft
β_1	1.52	1.23	1.13	1.07
β_2	1.31	1.14	1.01	1.00
β_3	1.32	1.15	0.95	0.97
β_4	1.79	1.34	1.02	1.01
β_5	1.18	1.09	1.15	1.07
β_6	1.58	1.26	1.42	1.19
β_7	1.75	1.32	1.48	1.22
β_8	1.01	1.01	1.32	1.15
β_9	0.91	0.95	1.14	1.07
β_{10}	1.51	1.23	1.41	1.19
β_{11}	1.43	1.19	1.10	1.05

Deff: Design effect, Deft: Design factor.




Figure 2. Comparison of design factor values of survey-weighted and level-weighted models.




Figure 3. Confidence intervals for estimates when using survey weights versus level weights.

4. Discussions

Recently, the application of multilevel analysis on complex survey data has increased as well as the accessibility of public available survey datasets collected from large national samples. However, the challenge with modeling in these data is weighting, as researchers have continuously applied the final survey weights that are inappropriate for multi-level analysis (Elkasabi, Ren, & Pullum, 2020) instead of level weights at the sampled hierarchies (Rabe-Hesketh & Skrondal, 2006). The use of the former weights has a number of implications on model estimates among which is mis-estimating standard errors and model parameters (Carle, 2009). Ignoring specific level weights has consequences; at the first stage, it results into biased estimates on the intercept and variance of random effect, whereas at the second stage, it leads to slightly underestimated fixed effects and residual variance, in addition to the biased estimates on the intercept and variance of the random effect (Cai, 2013).

A model that applied final survey weights produced more estimates with design factor values greater than 1.2 a value commonly taken to indicate sizeable variance inflation (Sturgis, 2004) hence increasing the variance of the estimated coefficients (Skinner & Mason), an indication that level weights are useful in reducing variance of model estimate (Dargatz & Hill, 1996; Liao & Valliant, 2012).

Computation of appropriate standard errors with complex survey data has important implications for policy research since standard errors are the foundation upon which statistical significance testing is based (Davern, Jones, Lepkowski, Davidson, & Blewett, 2007) which informs inference. Besides, the DHS public available datasets like other complex survey data are used extensively for health policy research, with significant implications for national health policy formulation decisions. A model weighted by final survey weights underestimated standard errors for the model estimates. This has an effect on model estimates (Daniels, Dominici, & Zeger, 2004). Underestimation of standard errors leads to narrow confidence interval, affecting test of statistical significance and possibly misleading inference.

5. Conclusions

Use of level weights instead of the final survey weights produces correct values of standard errors hence correctly informing inference. Models that apply level weights also produce estimates with high precision. Researchers should

always estimate and apply level weights while using survey data for multilevel analysis with final survey weights especially datasets under the DHS program, for better model estimates and inference.

Funding: This work was supported through the DELTAS Africa Initiative Grant No. 107754/Z/15/Z-DELTAS Africa SSACAB. The DELTAS Africa Initiative is an independent funding scheme of the African Academy of Sciences (AAS)'s Alliance for Accelerating Excellence in Science in Africa (AESA) and supported by the New Partnership for Africa's Development Planning and Coordinating Agency (NEPAD Agency) with funding from the Wellcome Trust (Grant No. 107754/Z/15/Z) and the UK government. The views expressed in this publication are those of the author(s) and not necessarily those of AAS, NEPAD Agency, Wellcome Trust or the UK government.

References

- Bethlehem, J. G., & Keller, W. J. (1987). Linear weighting of sample survey data. *Journal of official statistics*, 3(4), 141-153. doi:<https://www.scb.se/contentassets/ca21efb41fee47d293bbee5bf7be7fb3/linear-weighting-of-sample-survey-data.pdf>
- Brick, J. M. (2013). Unit Nonresponse and Weighting Adjustments: A Critical Review. *Journal of Official Statistics*, 29(3), 329-353. Retrieved from <http://dx.doi.org/10.2478/jos-2013-0026>
- Cai, T. (2013). Investigation of Ways to Handle Sampling Weights for Multilevel Model Analyses. *Sociological Methodology*, 43(1), 178-219. Retrieved from <https://doi.org/10.1177/0081175012460221>
- Carle, A. C. (2009). Fitting multilevel models in complex survey data with design weights: Recommendations. *BMC Med Res Methodol*, 9(49). Retrieved from <https://doi.org/10.1186/1471-2288-9-49>
- Chen, Q., Gelman, A., Tracy, M., Norris, F. H., & Galea, S. (2015). Incorporating the sampling design in weighting adjustments for panel attrition. *Stat Med*, 34(28), 3637-47. doi:10.1002/sim.6618
- Daniels, M. J., Dominici, F., & Zeger, S. (2004). Underestimation of Standard Errors in Multi-Site Time Series Studies. *Epidemiology*, 15(1), 57-62. Retrieved from <https://www.jstor.org/stable/20485840>
- Dargatz, D. A., & Hill, G. W. (1996). Analysis of survey data. *Preventive Veterinary Medicine*, 28(4), 225-237. Retrieved from [https://doi.org/10.1016/0167-5877\(96\)01052-5](https://doi.org/10.1016/0167-5877(96)01052-5)
- Davern, M., Jones, A., Lepkowski, J., Davidson, G., & Blewett, L. A. (2007). Estimating Regression Standard Errors with Data from the Current Population Survey's Public Use File. *Inquiry*, 44, 211-224. Retrieved from https://journals.sagepub.com/doi/pdf/10.5034/inquiryjrn1_44.2.211
- Elkasabi, M., Ren, R., & Pullum, T. W. (2020). *Multilevel Modeling Using DHS Surveys: A Framework to Approximate Level-Weights*. Rockville, Maryland, USA: ICF.
- Gelman, A. (2007). Struggles with Survey Weighting and Regression Modeling. *Statist. Sci*, 22(2), 153-164. Retrieved from <https://doi.org/10.1214/088342306000000691>
- Graubard, B., & Korn, E. L. (1996). Modelling the sampling design in the analysis of health surveys. *Stat Methods Med Res*, 5(3), 263-81. doi:10.1177/096228029600500304
- Hakim, F., Bhuiyan, R., Akter, K., & Zaman, M. (2020). Weighting National Survey Data for Bangladeshi Population. *Research Methods Cases*. Retrieved from <https://dx.doi.org/10.4135/9781529743623>
- Kalton, G., & Flores-Cervantes, I. (2003). Weighting Methods. *Journal of Official Statistics*, 19(2), 81-97. Retrieved from <http://www.sverigeisiffror.scb.se/contentassets/ca21efb41fee47d293bbee5bf7be7fb3/weighting-methods.pdf>
- Kim, J. K., & Skinner, C. J. (2013). Weighting in survey analysis under informative sampling. *Biometrika*, 100(2), 385-398. Retrieved from <https://doi.org/10.1093/biomet/ass085>
- Lavrakas, P. J. (2008). *Encyclopedia of survey research methods* (Vols. 1-0). Thousand Oaks, Carifonia: Sage Publication, Inc. doi:10.4135/9781412963947
- Lepkowski, J. M., Mosher, W. D., Groves, R. M., West, B. T., & Wagner, J. (2013). *Responsive design, weighting, and variance estimation in the 2006-2010 National Survey of Family Growth*. Hyattsville, Maryland: National Center for Health Statistics. Retrieved from <https://stacks.cdc.gov/view/cdc/22069>
- Liao, D., & Valliant, R. (2012). Variance inflation factors in the analysis of complex survey data. *Survey Methodology*, 38(1), 53-62. Retrieved from <https://www.rti.org/publication/variance-inflation-factors-analysis-complex-survey-data/fulltext.pdf>
- Little, R. J., & Vartivarian, S. (2003). On weighting the rates in non-response weights. *Stat Med*, 22(9), 1589-99. doi:10.1002/sim.1513
- Nahorniak, M., Larsen, D. P., Volk, C., & Jordan, C. E. (2015). Using Inverse Probability Bootstrap Sampling to Eliminate Sample Induced Bias in Model Based Analysis of Unequal Probability Samples. *PLoS One*, 10(6), e0131765. doi:10.1371/journal.pone.0131765
- NMCD, UBOS, & ICF. (2020). *Uganda Malaria Indicator Survey 2018-19*. Kampala, Uganda, and Rockville, Maryland, USA: Uganda National Malaria Control Division (NMCD), Uganda Bureau of Statistics (UBOS), and ICF. Retrieved from <https://www.dhsprogram.com/pubs/pdf/MIS34/MIS34.pdf>
- Pfeffermann-a, D. (1993). The Role of Sampling Weights When Modeling Survey Data. *International Statistical Review*, 61(2), 317-337. Retrieved from <https://doi.org/10.2307/1403631>

-
- Pfeffermann-b, D. (1996). The use of sampling weights for survey data analysis. *Stat Methods Med Res*, 239-261. doi:10.1177/096228029600500303
- Rabe-Hesketh, S., & Skrondal, A. (2006). Multilevel Modelling of Complex Survey Data. *Journal of the Royal Statistical Society*, 169(4), 805–827. Retrieved from <http://www.jstor.org/stable/3877401>
- Seaman, S. R., & White, I. R. (2013). Review of inverse probability weighting for dealing with missing data. *Statistical Methods in Medical Research*, 22(3), 278–295. Retrieved from <https://doi.org/10.1177/0962280210395740>
- Skinner, C., & Mason, B. (n.d.). Weighting in the regression analysis of survey data with a cross-national application. *The Canadian Journal of Statistics*, 40(4), 697-711. Retrieved from <https://doi.org/10.1002/cjs.11155>
- Sturgis, P. (2004). Analysing Complex Survey Data: Clustering, Stratification and Weights. *social research UPDATE*(43). Retrieved from <https://sru.soc.surrey.ac.uk/SRU43.PDF>
- UBOS, & ICF. (2018). *Uganda Demographic and Health Survey 2016*. Kampala, Uganda and Rockville, Maryland, USA: Uganda Bureau of Statistics (UBOS) and ICF. Retrieved from <https://dhsprogram.com/pubs/pdf/FR333/FR333.pdf>
- Wagemaker, H. (Ed.). (2020). *Reliability and Validity of International Large-Scale Assessment* (Vol. 10). Cham, Switzerland: Springer. Retrieved from <https://doi.org/10.1007/978-3-030-53081-5>
- West, B. T., Beer, L., Gremel, G. W., Weiser, J., Johnson, C. H., Garg, S., & Skarbinski, J. (2015). Weighted Multilevel Models: A Case Study. *Am J Public Health*, 105(11), 2214–2215. doi:10.2105/AJPH.2015.302842
- Wun, L.-M., Ezzati-Rice, T. M., Diaz-Tena, N., & Greenblatt, J. (2007). On modelling response propensity for dwelling unit (DU) level non-response adjustment in the Medical Expenditure Panel Survey (MEPS). *Stat Med*, 26(8), 1875-84. doi:10.1002/sim.2809

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.