Preprints (www.preprints.org) | NOT PEER-REVIEWED | Posted: 27 March 2017 doi:10.20944/preprints201703.0200.v1

Peer-reviewed version available at AIMS Mathematics 2020, 5, 1333-1345; doi:10.3934/math.2020091

SEVERAL EXPLICIT AND RECURSIVE FORMULAS FOR THE GENERALIZED MOTZKIN NUMBERS

FENG QI

Institute of Mathematics, Henan Polytechnic University, Jiaozuo City, Henan Province, 454010, China; College of Mathematics, Inner Mongolia University for Nationalities, Tongliao City, Inner Mongolia Autonomous Region, 028043, China; Department of Mathematics, College of Science, Tianjin Polytechnic University, Tianjin City, 300387, China

BAI-NI GUO

School of Mathematics and Informatics, Henan Polytechnic University, Jiaozuo City, Henan Province, 454010, China

Abstract. In the paper, the authors find two explicit formulas and recover a recursive formula for the generalized Motzkin numbers. Consequently, the authors deduce two explicit formulas and a recursive formula for the Motzkin numbers, the Catalan numbers, and the restricted hexagonal numbers respectively.

1. Introduction

The Motzkin numbers M_n enumerate various combinatorial objects. In 1977, Donaghey and Shapiro [3] gave fourteen different manifestations of the Motzkin numbers M_n. In particular, the Motzkin numbers M_n give the numbers of paths from $(0,0)$ to $(n,0)$ which never dip below the x-axis $y=0$ and are made up only of the steps $(1,0)$, $(1,1)$, and $(1,-1)$.

The first seven Motzkin numbers M_n for $0 \le n \le 6$ are $1, 1, 2, 4, 9, 21, 51$. All the Motzkin numbers M_n can be generated by

$$M(x)=\frac{1-x-\sqrt{1-2x-3x^2}}{2x^2}=\frac{1}{1-x+\sqrt{1-2x-3x^2}}=\sum_{k=0}^{\infty}M_kx^k.$$

E-mail addresses: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com, bai.ni.guo@gmail.com, bai.ni.guo@hotmail.com.

2010 *Mathematics Subject Classification.* Primary 05A15; Secondary 05A19, 05A20, 05A15, 11B37, 11B83, 34A05.

Key words and phrases. explicit formula; recursive formula; generalized Motzkin number; Motzkin number; restricted hexagonal number; Catalan number; generating function.

This paper was typeset using $\mathcal{A}_{\mathcal{M}}\mathcal{S}$-LaTeX.

Preprints (www.preprints.org) | NOT PEER-REVIEWED | Posted: 27 March 2017 doi:10.20944/preprints201703.0200.v1

Peer-reviewed version available at AIMS Mathematics 2020, 5, 1333-1345; doi:10.3934/math.2020091

In 2007, Mansour *et al* [10] introduced the (u,l,d)-Motzkin numbers $m_n^{(u,l,d)}$ and obtained [10, Theorem 2.1] that $m_n^{(u,l,d)}=m_n^{(1,l,ud)}$,

$$M_{u,l,d}(x)=\frac{1-lx-\sqrt{(1-lx)^2-4udx^2}}{2udx^2}=\sum_{n=0}^{\infty}m_n^{(u,l,d)}x^n, \tag{1}$$

and

$$m_n^{(u,l,d)}=l^n\sum_{j=0}^{n/2}\frac{1}{j+1}\binom{2j}{j}\binom{n}{2j}\left(\frac{ud}{l^2}\right)^j. \tag{2}$$

From (1) and (2), it is easy to see that $m_n^{(u,l,d)}=m_n^{(d,l,u)}$.

In 2014, Sun [21] generalized the Motzkin numbers M_n to

$$M_n(a,b)=\sum_{k=0}^{\lfloor n/2\rfloor}\binom{n}{2k}C_k a^{n-2k}b^k \tag{3}$$

for $a,b\in\mathbb{N}$ in terms of the Catalan numbers

$$C_n=\frac{1}{n+1}\binom{2n}{n} \tag{4}$$

and established the generating function

$$\begin{aligned}M_{a,b}(x)&=\frac{1-ax-\sqrt{(1-ax)^2-4bx^2}}{2bx^2}\\&=\frac{1}{1-ax+\sqrt{(1-ax)^2-4bx^2}}=\sum_{k=0}^{\infty}M_k(a,b)x^k,\end{aligned} \tag{5}$$

where $\lfloor\lambda\rfloor$ denotes the floor function defined by the largest integer less than or equal to $\lambda\in\mathbb{R}$. Wang and Zhang pointed out [22] that

$$M_n(1,1)=M_n,\quad M_n(2,1)=C_{n+1},\quad\text{and}\quad M_n(3,1)=H_n, \tag{6}$$

where H_n denote the restricted hexagonal numbers described by Harary and Read [5].

For more information on many results, applications, and generalizations of the Motzkin numbers M_n, please refer to the papers [3, 7, 8, 21, 22] and closely related references therein. For more information on many results, applications, and generalizations of the Catalan numbers C_n, please refer to the monograph [6], the papers [9, 14, 15, 20], the survey article [12], and closely related references therein.

Comparing (1) with (5) reveals that $M_k(a,b)$ and $m_k^{(u,l,d)}$ are equivalent to each other and satisfy

$$M_k(a,b)=m_n^{(1,a,b)}=m_k^{(b,a,1)}\quad\text{and}\quad m_k^{(u,l,d)}=M_k(l,ud). \tag{7}$$

Therefore, it suffices to consider the generalized Motzkin numbers $M_k(a,b)$, rather than the (u,l,d)-Motzkin numbers $m_n^{(u,l,d)}$, in this paper.

By the second relation in (7), one can reformulated the formula (2) as

$$M_n(a,b)=a^n\sum_{j=0}^{\lfloor n/2\rfloor}\frac{1}{j+1}\binom{2j}{j}\binom{n}{2j}\left(\frac{b}{a^2}\right)^j. \tag{8}$$

Substituting (4) into (3) recovers (8) once again.

Preprints (www.preprints.org) | NOT PEER-REVIEWED | Posted: 27 March 2017 doi:10.20944/preprints201703.0200.v1

Peer-reviewed version available at AIMS Mathematics 2020, 5, 1333-1345; doi:10.3934/math.2020091

In 2015, Wang and Zhang [22, Theorem 1] combinatorially obtained, among other things, the recursive formula

$$M_{n+2}(a,b) = aM_{n+1}(a,b) + b\sum_{\ell=0}^{n} M_\ell(a,b)M_{n-\ell}(a,b), \quad n \geq 0. \tag{9}$$

In this paper, we will find two explicit formulas, different from (8), and recover the recursive formula (9) for the generalized Motzkin numbers $M_n(a,b)$. Consequently, we will derive two explicit formula and a recursive formula for the Motzkin numbers M_n, the Catalan numbers C_n, and the restricted hexagonal numbers H_n respectively.

We can state our main results as the following three theorems.

Theorem 1. *For $n \geq 0$, we can compute the generalized Motzkin numbers $M_n(a,b)$ by*

$$M_n(a,b) = \frac{1}{2b}\left(\frac{4b-a^2}{2a}\right)^{n+2}\sum_{\ell=0}^{n+2}\left(\frac{2a^2}{4b-a^2}\right)^{\ell}\frac{(2\ell-3)!!}{\ell!}\binom{\ell}{n-\ell+2}, \tag{10}$$

where $\binom{p}{q} = 0$ for $q > p \geq 0$ and the double factorial of negative odd integers $-(2n+1)$ is

$$[-(2n+1)]!! = \frac{(-1)^n}{(2n-1)!!} = (-1)^n\frac{2^n n!}{(2n)!}, \quad n = 0, 1, \ldots .$$

Consequently, we can compute the Motzkin numbers M_n and the restricted hexagonal numbers H_n respectively by

$$M_n = \frac{9}{8}\left(\frac{3}{2}\right)^{n}\sum_{\ell=0}^{n+2}\left(\frac{2}{3}\right)^{\ell}\frac{(2\ell-3)!!}{\ell!}\binom{\ell}{n-\ell+2} \tag{11}$$

and

$$H_n = (-1)^n\frac{25}{72}\left(\frac{5}{6}\right)^{n}\sum_{\ell=0}^{n+2}(-1)^{\ell}\left(\frac{18}{5}\right)^{\ell}\frac{(2\ell-3)!!}{\ell!}\binom{\ell}{n-\ell+2}. \tag{12}$$

Theorem 2. *For $n \geq 0$, we can compute the generalized Motzkin numbers $M_n(a,b)$ by*

$$M_n(a,b) = -\frac{\left(a-2\sqrt{b}\right)^{n+2}}{2b}\sum_{\ell=0}^{n+2}\frac{(2\ell-3)!!}{(2\ell)!!}\frac{[2(n-\ell+2)-3]!!}{[2(n-\ell+2)]!!}\left(\frac{a+2\sqrt{b}}{a-2\sqrt{b}}\right)^{\ell}. \tag{13}$$

Consequently, we can compute the Motzkin numbers M_n and the restricted hexagonal numbers H_n respectively by

$$M_n = \frac{(-1)^{n+1}}{2}\sum_{\ell=0}^{n+2}(-1)^{\ell}3^{\ell}\frac{(2\ell-3)!!}{(2\ell)!!}\frac{[2(n-\ell+2)-3]!!}{[2(n-\ell+2)]!!}$$

and

$$H_n = -\frac{1}{2}\sum_{\ell=0}^{n+2}5^{\ell}\frac{(2\ell-3)!!}{(2\ell)!!}\frac{[2(n-\ell+2)-3]!!}{[2(n-\ell+2)]!!}.$$

Theorem 3. *For $n \geq 0$, the generalized Motzkin numbers $M_n(a,b)$ satisfy*

$$M_0(a,b) = 1, \quad M_1(a,b) = a, \tag{14}$$

Preprints (www.preprints.org) | NOT PEER-REVIEWED | Posted: 27 March 2017 doi:10.20944/preprints201703.0200.v1

Peer-reviewed version available at AIMS Mathematics 2020, 5, 1333-1345; doi:10.3934/math.2020091

and the recursive formula (9). *Consequently, for* $n \ge 0$, *the Motzkin numbers* M_n, *the Catalan numbers* C_n, *and the restricted hexagonal numbers* H_n *meet the recursive formulas*

$$M_{n+2} = M_{n+1} + \sum_{\ell=0}^{n} M_\ell M_{n-\ell}, \tag{15}$$

$$C_{n+2} = 2C_{n+1} + \sum_{\ell=0}^{n} C_\ell C_{n-\ell}, \tag{16}$$

and

$$H_{n+2} = 3H_{n+1} + \sum_{\ell=0}^{n} H_\ell H_{n-\ell} \tag{17}$$

respectively.

2. Lemmas

In order to prove the explicit formula (10), we need the following lemmas.

Lemma 1 ([1, p. 40, Exercise 5)], [11, Section 2.2, p. 849], [13, p. 94], [17, Lemma 3], and [23, Lemma 2.1]). *Let* $u(x)$ *and* $v(x) \ne 0$ *be two differentiable functions. Let* $U_{(n+1)\times1}(x)$ *be an* $(n+1)\times1$ *matrix whose elements* $u_{k,1}(x) = u^{(k-1)}(x)$ *for* $1 \le k \le n+1$, *let* $V_{(n+1)\times n}(x)$ *be an* $(n+1)\times n$ *matrix whose elements*

$$v_{i,j}(x) = \begin{cases} \binom{i-1}{j-1} v^{(i-j)}(x), & i-j \ge 0 \\ 0, & i-j < 0 \end{cases}$$

for $1 \le i \le n+1$ *and* $1 \le j \le n$, *and let* $|W_{(n+1)\times(n+1)}(x)|$ *denote the determinant of the* $(n+1)\times(n+1)$ *matrix*

$$W_{(n+1)\times(n+1)}(x) = \begin{pmatrix} U_{(n+1)\times1}(x) & V_{(n+1)\times n}(x) \end{pmatrix}.$$

Then the n*th derivative of the ratio* $\frac{u(x)}{v(x)}$ *can be computed by*

$$\frac{\mathrm{d}^n}{\mathrm{d}x^n}\left[\frac{u(x)}{v(x)}\right] = (-1)^n \frac{\left|W_{(n+1)\times(n+1)}(x)\right|}{v^{n+1}(x)}.$$

Lemma 2 ([2, p. 134, Theorem A and p. 139, Theorem C]). *The Faà di Bruno formula can be described in terms of the Bell polynomials of the second kind*

$$\mathrm{B}_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{1\le i\le n,\ell_i\in\{0\}\cup\mathbb{N} \\ \sum_{i=1}^{n} i\ell_i = n \\ \sum_{i=1}^{n} \ell_i = k}} \frac{n!}{\prod_{i=1}^{n-k+1} \ell_i!} \prod_{i=1}^{n-k+1} \left(\frac{x_i}{i!}\right)^{\ell_i}$$

for $n \ge k \ge 0$ *by*

$$\frac{\mathrm{d}^n}{\mathrm{d}t^n}[f\circ h(t)] = \sum_{k=0}^{n} f^{(k)}(h(t))\mathrm{B}_{n,k}\left(h'(t), h''(t), \dots, h^{(n-k+1)}(t)\right) \tag{18}$$

for $n \ge 0$.

Preprints (www.preprints.org) | NOT PEER-REVIEWED | Posted: 27 March 2017 doi:10.20944/preprints201703.0200.v1

Peer-reviewed version available at AIMS Mathematics 2020, 5, 1333-1345; doi:10.3934/math.2020091

Lemma 3 ([2, p. 135]). *The Bell polynomials of the second kind* $\mathrm{B}_{n,k}$ *satisfy*

$$\mathrm{B}_{n,k}\bigl(abx_1, ab^2x_2, \dots, ab^{n-k+1}x_{n-k+1}\bigr) = a^k b^n \mathrm{B}_{n,k}(x_1, x_2, \dots, x_{n-k+1}) \tag{19}$$

for $n \ge k \ge 0$.

Lemma 4 ([4, Theorem 4.1], [16, Remark 1], [18, p. 7, (19)], [19, Section 3], and [23, Lemma 2.5]). *For* $n \ge k \ge 0$, *we have*

$$\mathrm{B}_{n,k}(x, 1, 0, \dots, 0) = \frac{(n-k)!}{2^{n-k}} \binom{n}{k} \binom{k}{n-k} x^{2k-n}. \tag{20}$$

More generally, for $n \ge k \ge 0$ *and* $\lambda \in \mathbb{R}$, *we have*

$$\mathrm{B}_{n,k}\left(1, 1-\lambda, (1-\lambda)(1-2\lambda), \dots, \prod_{\ell=0}^{n-k}(1-\ell\lambda)\right) = \frac{(-1)^k}{k!} \sum_{\ell=0}^{k} (-1)^\ell \binom{k}{\ell} \prod_{q=0}^{n-1} (\ell - q\lambda). \tag{21}$$

3. Proofs of Theorems 1 and 3

We are now in a position to prove our main results.

Proof of Theorem 1. By virtue of (18), (19), and (20), we obtain for $k \ge 0$ that

$$\begin{aligned}
\left[\sqrt{(1-ax)^2 - 4bx^2}\right]^{(k+2)} &= \sum_{\ell=0}^{k+2} \left\langle \frac{1}{2} \right\rangle_\ell \left[(1-ax)^2 - 4bx^2\right]^{1/2-\ell} \\
&\quad \times \mathrm{B}_{k+2,\ell}\bigl(-2\bigl[a + \bigl(4b - a^2\bigr)x\bigr], 2\bigl(a^2 - 4b\bigr), 0, \dots, 0\bigr) \\
&\to \sum_{\ell=0}^{k+2} \left\langle \frac{1}{2} \right\rangle_\ell \mathrm{B}_{k+2,\ell}\bigl(-2a, 2\bigl(a^2 - 4b\bigr), 0, \dots, 0\bigr) \\
&= \sum_{\ell=0}^{k+2} \left\langle \frac{1}{2} \right\rangle_\ell \bigl[2\bigl(a^2 - 4b\bigr)\bigr]^\ell \mathrm{B}_{k+2,\ell}\left(\frac{a}{4b-a^2}, 1, 0, \dots, 0\right) \\
&= \sum_{\ell=0}^{k+2} \left\langle \frac{1}{2} \right\rangle_\ell \bigl[2\bigl(a^2 - 4b\bigr)\bigr]^\ell \frac{(k-\ell+2)!}{2^{k-\ell+2}} \binom{k+2}{\ell} \binom{\ell}{k-\ell+2} \left(\frac{a}{4b-a^2}\right)^{2\ell-k-2}
\end{aligned} \tag{22}$$

as $x \to 0$, where

$$\langle x \rangle_n = \begin{cases} x(x-1)\cdots(x-n+1), & n \ge 1 \\ 1, & n = 0 \end{cases}$$

denotes the falling factorial of $x \in \mathbb{R}$.

Preprints (www.preprints.org) | NOT PEER-REVIEWED | Posted: 27 March 2017 doi:10.20944/preprints201703.0200.v1

Peer-reviewed version available at AIMS Mathematics 2020, 5, 1333-1345; doi:10.3934/math.2020091

Letting $u(x)=1-ax-\sqrt{(1-ax)^2-4bx^2}$ and $v(x)=x^2$ in Lemma 1 gives

$$\frac{\mathrm{d}^n M_{a,b}(x)}{\mathrm{d}x^n}=\frac{1}{2b}\frac{(-1)^n}{x^{2(n+1)}}\begin{vmatrix} u(x) & \binom{0}{0}x^2 & 0 & \cdots & 0 & 0 & 0\\ u'(x) & 2\binom{1}{0}x & \binom{1}{1}x^2 & \cdots & 0 & 0 & 0\\ u''(x) & 2\binom{2}{0} & 2\binom{2}{1}x & \cdots & 0 & 0 & 0\\ u^{(3)}(x) & 0 & 2\binom{3}{1} & \cdots & 0 & 0 & 0\\ u^{(4)}(x) & 0 & 0 & \cdots & 0 & 0 & 0\\ \cdots & \cdots & \cdots & \ddots & \cdots & \cdots & \cdots\\ u^{(n-2)}(x) & 0 & 0 & \cdots & 2\binom{n-2}{n-3}x & \binom{n-2}{n-2}x^2 & 0\\ u^{(n-1)}(x) & 0 & 0 & \cdots & 2\binom{n-1}{n-3} & 2\binom{n-1}{n-2}x & \binom{n-1}{n-1}x^2\\ u^{(n)}(x) & 0 & 0 & \cdots & 0 & 2\binom{n}{n-2} & 2\binom{n}{n-1}x \end{vmatrix}$$

$$=\frac{1}{2b}\frac{(-1)^n}{x^{2(n+1)}}\left[(-1)^n u^{(n)}(x)\begin{vmatrix} \binom{0}{0}x^2 & 0 & 0 & \cdots & 0 & 0 & 0\\ 2\binom{1}{0}x & \binom{1}{1}x^2 & 0 & \cdots & 0 & 0 & 0\\ 2\binom{2}{0} & 2\binom{2}{1}x & \binom{2}{2}x^2 & \cdots & 0 & 0 & 0\\ 0 & 2\binom{3}{1} & 2\binom{3}{2}x & \cdots & 0 & 0 & 0\\ 0 & 0 & 2\binom{4}{2} & \cdots & 0 & 0 & 0\\ \cdots & \cdots & \cdots & \ddots & \cdots & \cdots & \cdots\\ 0 & 0 & 0 & \cdots & 2\binom{n-2}{n-3}x & \binom{n-2}{n-2}x^2 & 0\\ 0 & 0 & 0 & \cdots & 2\binom{n-1}{n-3} & 2\binom{n-1}{n-2}x & \binom{n-1}{n-1}x^2 \end{vmatrix}\right.$$

$$+2\binom{n}{n-1}x\begin{vmatrix} u(x) & \binom{0}{0}x^2 & 0 & 0 & 0 & \cdots & 0 & 0\\ u'(x) & 2\binom{1}{0}x & \binom{1}{1}x^2 & 0 & 0 & \cdots & 0 & 0\\ u''(x) & 2\binom{2}{0} & 2\binom{2}{1}x & \binom{2}{2}x^2 & 0 & \cdots & 0 & 0\\ u^{(3)}(x) & 0 & 2\binom{3}{1} & 2\binom{3}{2}x & \binom{3}{3}x^2 & \cdots & 0 & 0\\ u^{(4)}(x) & 0 & 0 & 2\binom{4}{2} & 2\binom{4}{3}x & \cdots & 0 & 0\\ \cdots & \cdots & \cdots & \cdots & \cdots & \ddots & \cdots & \cdots\\ u^{(n-2)}(x) & 0 & 0 & 0 & 0 & \cdots & 2\binom{n-2}{n-3}x & \binom{n-2}{n-2}x^2\\ u^{(n-1)}(x) & 0 & 0 & 0 & 0 & \cdots & 2\binom{n-1}{n-3} & 2\binom{n-1}{n-2}x \end{vmatrix}$$

$$\left.-2\binom{n}{n-2}\binom{n-1}{n-1}x^2\begin{vmatrix} u(x) & \binom{0}{0}x^2 & 0 & 0 & \cdots & 0 & 0\\ u'(x) & 2\binom{1}{0}x & \binom{1}{1}x^2 & 0 & \cdots & 0 & 0\\ u''(x) & 2\binom{2}{0} & 2\binom{2}{1}x & \binom{2}{2}x^2 & \cdots & 0 & 0\\ u^{(3)}(x) & 0 & 2\binom{3}{1} & 2\binom{3}{2}x & \cdots & 0 & 0\\ u^{(4)}(x) & 0 & 0 & 2\binom{4}{2} & \cdots & 0 & 0\\ \cdots & \cdots & \cdots & \cdots & \ddots & \cdots & \cdots\\ u^{(n-3)}(x) & 0 & 0 & 0 & \cdots & 2\binom{n-3}{n-4}x & \binom{n-3}{n-3}x^2\\ u^{(n-2)}(x) & 0 & 0 & 0 & \cdots & 2\binom{n-2}{n-4} & 2\binom{n-2}{n-3}x \end{vmatrix}\right]$$

$$=\frac{1}{2b}\frac{u^{(n)}(x)}{x^2}-\frac{2n}{x}\frac{1}{2b}\frac{(-1)^{n-1}}{x^{2n}}\begin{vmatrix} u(x) & \binom{0}{0}x^2 & 0 & \cdots & 0 & 0\\ u'(x) & 2\binom{1}{0}x & \binom{1}{1}x^2 & \cdots & 0 & 0\\ u''(x) & 2\binom{2}{0} & 2\binom{2}{1}x & \cdots & 0 & 0\\ u^{(3)}(x) & 0 & 2\binom{3}{1} & \cdots & 0 & 0\\ u^{(4)}(x) & 0 & 0 & \cdots & 0 & 0\\ \cdots & \cdots & \cdots & \ddots & \cdots & \cdots\\ u^{(n-2)}(x) & 0 & 0 & \cdots & 2\binom{n-2}{n-3}x & \binom{n-2}{n-2}x^2\\ u^{(n-1)}(x) & 0 & 0 & \cdots & 2\binom{n-1}{n-3} & 2\binom{n-1}{n-2}x \end{vmatrix}$$

Preprints (www.preprints.org) | NOT PEER-REVIEWED | Posted: 27 March 2017 doi:10.20944/preprints201703.0200.v1

Peer-reviewed version available at AIMS Mathematics 2020, 5, 1333-1345; doi:10.3934/math.2020091

$$-n(n-1)\frac{1}{x^2}\frac{1}{2b}\frac{(-1)^{n-2}}{x^{2(n-1)}}\begin{vmatrix} u(x) & \binom{0}{0}x^2 & 0 & 0 & \cdots & 0 & 0 \\ u'(x) & 2\binom{1}{0}x & \binom{1}{1}x^2 & 0 & \cdots & 0 & 0 \\ u''(x) & 2\binom{2}{0} & 2\binom{2}{1}x & \binom{2}{2}x^2 & \cdots & 0 & 0 \\ u^{(3)}(x) & 0 & 2\binom{3}{1} & 2\binom{3}{2}x & \cdots & 0 & 0 \\ u^{(4)}(x) & 0 & 0 & 2\binom{4}{2} & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \ddots & \cdots & \cdots \\ u^{(n-3)}(x) & 0 & 0 & 0 & \cdots & 2\binom{n-3}{n-4}x & \binom{n-3}{n-3}x^2 \\ u^{(n-2)}(x) & 0 & 0 & 0 & \cdots & 2\binom{n-2}{n-4} & 2\binom{n-2}{n-3}x \end{vmatrix}$$

$$= \frac{1}{2b}\frac{u^{(n)}(x)}{x^2} - \frac{2n}{x}\frac{\mathrm{d}^{n-1}M_{a,b}(x)}{\mathrm{d}x^{n-1}} - \frac{n(n-1)}{x^2}\frac{\mathrm{d}^{n-2}M_{a,b}(x)}{\mathrm{d}x^{n-2}}$$

$$= \frac{1}{x^2}\left[\frac{u^{(n)}(x)}{2b} - 2nx\frac{\mathrm{d}^{n-1}M_{a,b}(x)}{\mathrm{d}x^{n-1}} - n(n-1)\frac{\mathrm{d}^{n-2}M_{a,b}(x)}{\mathrm{d}x^{n-2}}\right].$$

Therefore, by L'Hôspital's rule, we have

$$\begin{aligned}\lim_{x\to 0}\frac{\mathrm{d}^n M_{a,b}(x)}{\mathrm{d}x^n} &= \lim_{x\to 0}\left\{\frac{1}{x^2}\left[\frac{u^{(n)}(x)}{2b} - 2nx\frac{\mathrm{d}^{n-1}M_{a,b}(x)}{\mathrm{d}x^{n-1}} - n(n-1)\frac{\mathrm{d}^{n-2}M_{a,b}(x)}{\mathrm{d}x^{n-2}}\right]\right\} \\ &= \lim_{x\to 0}\left\{\frac{1}{2x}\left[\frac{u^{(n+1)}(x)}{2b} - 2nx\frac{\mathrm{d}^{n}M_{a,b}(x)}{\mathrm{d}x^{n}} - n(n+1)\frac{\mathrm{d}^{n-1}M_{a,b}(x)}{\mathrm{d}x^{n-1}}\right]\right\} \\ &= \frac{1}{2}\lim_{x\to 0}\left[\frac{u^{(n+2)}(x)}{2b} - 2nx\frac{\mathrm{d}^{n+1}M_{a,b}(x)}{\mathrm{d}x^{n+1}} - n(n+3)\frac{\mathrm{d}^{n}M_{a,b}(x)}{\mathrm{d}x^{n}}\right] \\ &= \frac{1}{2}\left[\lim_{x\to 0}\frac{u^{(n+2)}(x)}{2b} - n(n+3)\lim_{x\to 0}\frac{\mathrm{d}^{n}M_{a,b}(x)}{\mathrm{d}x^{n}}\right]\end{aligned}$$

which is equivalent to

$$\lim_{x\to 0}\frac{\mathrm{d}^n M_{a,b}(x)}{\mathrm{d}x^n} = \frac{1}{(n+1)(n+2)}\lim_{x\to 0}\frac{u^{(n+2)}(x)}{2b} = \frac{1}{2b(n+1)(n+2)}\lim_{x\to 0}u^{(n+2)}(x).$$

Considering

$$\lim_{x\to 0}\frac{\mathrm{d}^n M_{a,b}(x)}{\mathrm{d}x^n} = n!M_n(a,b),$$

making use of (22), and simplifying lead to the explicit formula (10).

Letting $(a,b)=(1,1)$ and $(a,b)=(3,1)$ respectively in (10) and considering the three relations in (6) derive (11) and (12) immediately. The proof of Theorem 1 is complete. □

Proof of Theorem 2. From (5), it is derived that

$$\sqrt{(1-ax)^2-4bx^2} = 1-ax-2b\sum_{k=0}^{\infty}M_k(a,b)x^{k+2}.$$

This implies that

$$M_k(a,b) = -\frac{1}{2b}\frac{1}{(k+2)!}\lim_{x\to 0}\left[\sqrt{(1-ax)^2-4bx^2}\right]^{(k+2)}, \quad k\geq 0. \tag{23}$$

It is easy to see that

Preprints (www.preprints.org) | NOT PEER-REVIEWED | Posted: 27 March 2017 doi:10.20944/preprints201703.0200.v1

Peer-reviewed version available at AIMS Mathematics 2020, 5, 1333-1345; doi:10.3934/math.2020091

(1) when $a^2-4b\ge0$ and $x\le\min\left\{\frac{1}{a+2\sqrt{b}},\frac{1}{a-2\sqrt{b}}\right\}=\frac{1}{a+2\sqrt{b}}$, we have

$$\begin{aligned}
\left[\sqrt{(1-ax)^2-4bx^2}\right]^{(k+2)}&=\left[\sqrt{(a^2-4b)\left(x-\frac{1}{a+2\sqrt{b}}\right)\left(x-\frac{1}{a-2\sqrt{b}}\right)}\right]^{(k+2)}\\
&=\sqrt{a^2-4b}\left(\sqrt{\frac{1}{a+2\sqrt{b}}-x}\sqrt{\frac{1}{a-2\sqrt{b}}-x}\right)^{(k+2)}\\
&=\sqrt{a^2-4b}\sum_{\ell=0}^{k+2}\binom{k+2}{\ell}\left(\sqrt{\frac{1}{a+2\sqrt{b}}-x}\right)^{(\ell)}\left(\sqrt{\frac{1}{a-2\sqrt{b}}-x}\right)^{(k-\ell+2)}\\
&=(-1)^k\sqrt{a^2-4b}\sum_{\ell=0}^{k+2}\binom{k+2}{\ell}\left\langle\frac{1}{2}\right\rangle_\ell\left(\frac{1}{a+2\sqrt{b}}-x\right)^{1/2-\ell}\left\langle\frac{1}{2}\right\rangle_{k-\ell+2}\left(\frac{1}{a-2\sqrt{b}}-x\right)^{\ell-k-3/2}\\
&\to(-1)^k\sqrt{a^2-4b}\sum_{\ell=0}^{k+2}\binom{k+2}{\ell}\left\langle\frac{1}{2}\right\rangle_\ell\left(\frac{1}{a+2\sqrt{b}}\right)^{1/2-\ell}\left\langle\frac{1}{2}\right\rangle_{k-\ell+2}\left(\frac{1}{a-2\sqrt{b}}\right)^{\ell-k-3/2}\\
&=(k+2)!\left(a-2\sqrt{b}\right)^{k+2}\sum_{\ell=0}^{k+2}\frac{(2\ell-3)!!}{(2\ell)!!}\frac{[2(k-\ell+2)-3]!!}{[2(k-\ell+2)]!!}\left(\frac{a+2\sqrt{b}}{a-2\sqrt{b}}\right)^{\ell}
\end{aligned}$$

as $x\to0$;

(2) when $a^2-4b<0$ and

$$\begin{aligned}
\frac{1}{a+2\sqrt{b}}&=\max\left\{\frac{1}{a+2\sqrt{b}},\frac{1}{a-2\sqrt{b}}\right\}>x\\
&>\min\left\{\frac{1}{a+2\sqrt{b}},\frac{1}{a-2\sqrt{b}}\right\}=\frac{1}{a-2\sqrt{b}},
\end{aligned}$$

we have

$$\begin{aligned}
\left[\sqrt{(1-ax)^2-4bx^2}\right]^{(k+2)}&=\left[\sqrt{(4b-a^2)\left(\frac{1}{a+2\sqrt{b}}-x\right)\left(x-\frac{1}{a-2\sqrt{b}}\right)}\right]^{(k+2)}\\
&=\sqrt{4b-a^2}\left(\sqrt{\frac{1}{a+2\sqrt{b}}-x}\sqrt{x-\frac{1}{a-2\sqrt{b}}}\right)^{(k+2)}\\
&=\sqrt{4b-a^2}\sum_{\ell=0}^{k+2}\binom{k+2}{\ell}\left(\sqrt{\frac{1}{a+2\sqrt{b}}-x}\right)^{(\ell)}\left(\sqrt{x-\frac{1}{a-2\sqrt{b}}}\right)^{(k-\ell+2)}\\
&=\sqrt{4b-a^2}\sum_{\ell=0}^{k+2}\binom{k+2}{\ell}(-1)^\ell\left\langle\frac{1}{2}\right\rangle_\ell\left(\frac{1}{a+2\sqrt{b}}-x\right)^{1/2-\ell}\left\langle\frac{1}{2}\right\rangle_{k-\ell+2}\left(x-\frac{1}{a-2\sqrt{b}}\right)^{1/2-(k-\ell+2)}\\
&\to\sqrt{4b-a^2}\sum_{\ell=0}^{k+2}\binom{k+2}{\ell}(-1)^\ell\left\langle\frac{1}{2}\right\rangle_\ell\left(\frac{1}{a+2\sqrt{b}}\right)^{1/2-\ell}\left\langle\frac{1}{2}\right\rangle_{k-\ell+2}\left(\frac{1}{2\sqrt{b}-a}\right)^{1/2-(k-\ell+2)}\\
&=\left(2\sqrt{b}-a\right)^{k+1}\sqrt{4b-a^2}\sum_{\ell=0}^{k+2}\binom{k+2}{\ell}\frac{(2\ell-3)!!}{2^\ell}\left(\frac{a+2\sqrt{b}}{2\sqrt{b}-a}\right)^{\ell-1/2}(-1)^{k-\ell}\frac{[2(k-\ell+2)-3]!!}{2^{k-\ell+2}}\\
&=\left(2\sqrt{b}-a\right)^{k+2}\sum_{\ell=0}^{k+2}\binom{k+2}{\ell}\frac{(2\ell-3)!!}{2^\ell}\left(\frac{a+2\sqrt{b}}{2\sqrt{b}-a}\right)^{\ell}(-1)^{k-\ell}\frac{[2(k-\ell+2)-3]!!}{2^{k-\ell+2}}
\end{aligned}$$

Preprints (www.preprints.org) | NOT PEER-REVIEWED | Posted: 27 March 2017 doi:10.20944/preprints201703.0200.v1

Peer-reviewed version available at AIMS Mathematics 2020, 5, 1333-1345; doi:10.3934/math.2020091

$$= \big(a-2\sqrt{b}\big)^{k+2}\sum_{\ell=0}^{k+2}\binom{k+2}{\ell}\frac{(2\ell-3)!!}{2^\ell}\frac{[2(k-\ell+2)-3]!!}{2^{k-\ell+2}}\left(\frac{a+2\sqrt{b}}{a-2\sqrt{b}}\right)^\ell$$

$$= (k+2)!\big(a-2\sqrt{b}\big)^{k+2}\sum_{\ell=0}^{k+2}\frac{(2\ell-3)!!}{(2\ell)!!}\frac{[2(k-\ell+2)-3]!!}{[2(k-\ell+2)]!!}\left(\frac{a+2\sqrt{b}}{a-2\sqrt{b}}\right)^\ell$$

as $x\to 0$.

By virtue of (23), we obtain the formula (13) readily.

Letting $(a,b)=(1,1)$ and $(a,b)=(3,1)$ respectively in (13) and making use of the first and third relations in (6) lead to (11) and (12) immediately. The proof of Theorem 2 is complete. □

Proof of Theorem 3. From (5), it is derived that

$$\sqrt{(1-ax)^2-4bx^2} = 1-ax-2b\sum_{k=0}^{\infty}M_k(a,b)x^{k+2}.$$

Squaring on both sides of the above equation gives

$$\begin{aligned}
(1-ax)^2-4bx^2 &= 1-2ax+\big(a^2-4b\big)x^2\\
&= \left[1-ax-2b\sum_{k=0}^{\infty}M_k(a,b)x^{k+2}\right]^2\\
&= 1+a^2x^2+4b^2\left[\sum_{k=0}^{\infty}M_k(a,b)x^{k+2}\right]^2\\
&\quad -2ax-4b\sum_{k=0}^{\infty}M_k(a,b)x^{k+2}+4abx\sum_{k=0}^{\infty}M_k(a,b)x^{k+2}\\
&= 1-2ax+a^2x^2+4b^2x^4\sum_{k=0}^{\infty}\left[\sum_{\ell=0}^{k}M_\ell(a,b)M_{k-\ell}\right]x^k\\
&\quad -4b\sum_{k=2}^{\infty}M_{k-2}(a,b)x^k+4ab\sum_{k=3}^{\infty}M_{k-3}(a,b)x^k\\
&= 1-2ax+a^2x^2-4b\sum_{k=2}^{\infty}M_{k-2}(a,b)x^k\\
&\quad +4ab\sum_{k=3}^{\infty}M_{k-3}(a,b)x^k+4b^2\sum_{k=4}^{\infty}\left[\sum_{\ell=0}^{k-4}M_\ell(a,b)M_{k-\ell-4}(a,b)\right]x^k\\
&= 1-2ax+a^2x^2-4b\big[M_0(a,b)x^2+M_1(a,b)x^3\big]+4abM_0(a,b)x^3\\
&\quad -4b\sum_{k=4}^{\infty}M_{k-2}(a,b)x^k+4ab\sum_{k=4}^{\infty}M_{k-3}(a,b)x^k\\
&\quad +4b^2\sum_{k=4}^{\infty}\left[\sum_{\ell=0}^{k-4}M_\ell(a,b)M_{k-\ell-4}(a,b)\right]x^k
\end{aligned}$$

Preprints (www.preprints.org) | NOT PEER-REVIEWED | Posted: 27 March 2017 doi:10.20944/preprints201703.0200.v1

Peer-reviewed version available at AIMS Mathematics 2020, 5, 1333-1345; doi:10.3934/math.2020091

$$= 1 - 2ax + \left[a^2 - 4bM_0(a,b)\right]x^2 + 4b[aM_0(a,b) - M_1(a,b)]x^3$$
$$- 4b\sum_{k=4}^{\infty}\left[M_{k-2}(a,b) - aM_{k-3}(a,b) - b\sum_{\ell=0}^{k-4}M_\ell(a,b)M_{k-\ell-4}(a,b)\right]x^k$$

which means that

$$a^2 - 4b = a^2 - 4bM_0(a,b), \quad 4b[aM_0(a,b) - M_1(a,b)] = 0,$$

and

$$M_{k-2}(a,b) - aM_{k-3}(a,b) - b\sum_{\ell=0}^{k-4}M_\ell(a,b)M_{k-\ell-4}(a,b) = 0, \quad k \ge 4.$$

Consequently, the identities in (14) and the recursive formula (9) follow.

Taking $(a,b) = (1,1)$, $(a,b) = (2,1)$, and $(a,b) = (3,1)$ respectively in (9) and considering the three relations in (6) lead to (15), (16), and (17) immediately. The proof of Theorem 3 is complete. □

4. Two remarks

Remark 1. From the proof of Theorem 1, we can conclude that

$$x^2\frac{\mathrm{d}^n M_{a,b}(x)}{\mathrm{d}x^n} + 2nx\frac{\mathrm{d}^{n-1}M_{a,b}(x)}{\mathrm{d}x^{n-1}} + n(n-1)\frac{\mathrm{d}^{n-2}M_{a,b}(x)}{\mathrm{d}x^{n-2}} = \frac{u^{(n)}(x)}{2b}, \quad n \ge 2.$$

This implies that the generating function $M_{a,b}(x)$ expressed in (5) is an explicit solution of the linear ordinary differential equations

$$x^2 f^{(n)}(x) + 2nxf^{(n-1)}(x) + n(n-1)f^{(n-2)}(x) = F_{n;a,b}(x)$$

for all $n \ge 2$, where, by (19) and (20) or (21),

$$F_{n;a,b}(x) = \frac{n!\left(4b-a^2\right)^n}{2^{n+1}b}\frac{\sqrt{(1-ax)^2-4bx^2}}{[a+(4b-a^2)x]^n}$$
$$\times\sum_{\ell=1}^{n}\frac{2^\ell(2\ell-3)!!}{\ell!(4b-a^2)^\ell}\binom{\ell}{n-\ell}\frac{\left[a+\left(4b-a^2\right)x\right]^{2\ell}}{[(1-ax)^2-4bx^2]^\ell}.$$

Remark 2. This paper is a company and continuation of the article [24].

References

[1] N. Bourbaki, *Elements of Mathematics: Functions of a Real Variable: Elementary Theory*, Translated from the 1976 French original by Philip Spain. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2004; Available online at `http://dx.doi.org/10.1007/978-3-642-59315-4`.

[2] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Revised and Enlarged Edition, D. Reidel Publishing Co., Dordrecht and Boston, 1974.

[3] R. Donaghey and L. W. Shapiro, *Motzkin numbers*, J. Combinatorial Theory Ser. A **23** (1977), no. 3, 291–301.

[4] B.-N. Guo and F. Qi, *Explicit formulas for special values of the Bell polynomials of the second kind and the Euler numbers*, ResearchGate Technical Report (2015), available online at `http://dx.doi.org/10.13140/2.1.3794.8808`.

[5] F. Harary and R. C. Read, *The enumeration of tree-like polyhexes*, Proc. Edinburgh Math. Soc. (2) **17** (1970), 1–13.

[6] T. Koshy, *Catalan Numbers with Applications*, Oxford University Press, Oxford, 2009.

[7] T. Lengyel, *Exact p-adic orders for differences of Motzkin numbers*, Int. J. Number Theory **10** (2014), no. 3, 653–667; Available online at `http://dx.doi.org/10.1142/S1793042113501157`.

Preprints (www.preprints.org) | NOT PEER-REVIEWED | Posted: 27 March 2017 doi:10.20944/preprints201703.0200.v1

Peer-reviewed version available at AIMS Mathematics 2020, 5, 1333-1345; doi:10.3934/math.2020091

[8] T. Lengyel, *On divisibility properties of some differences of Motzkin numbers*, Ann. Math. Inform. **41** (2013), 121–136.

[9] F.-F. Liu, X.-T. Shi, and F. Qi, *A logarithmically completely monotonic function involving the gamma function and originating from the Catalan numbers and function*, Glob. J. Math. Anal. **3** (2015), no. 4, 140–144; Available online at `http://dx.doi.org/10.14419/gjma.v3i4.5187`.

[10] T. Mansour, M. Schork, and Y. Sun, *Motzkin numbers of higher rank: Generating function and explicit expression*, J. Int. Seq. **10** (2007), Article 07.7.4, 11 pages.

[11] F. Qi, *Derivatives of tangent function and tangent numbers*, Appl. Math. Comput. **268** (2015), 844–858; Available online at `http://dx.doi.org/10.1016/j.amc.2015.06.123`.

[12] F. Qi, *Some properties and generalizations of the Catalan, Fuss, and Fuss–Catalan numbers*, ResearchGate Research (2015), available online at `http://dx.doi.org/10.13140/RG.2.1.1778.3128`. Mathematical Analysis and Applications: Selected Topics, 35 pages, edited by H. Dutta, M. Ruzhansky, and R. P. Agarwal, Wiley, October 2017. (ISBN 9781119414346)

[13] F. Qi and R. J. Chapman, *Two closed forms for the Bernoulli polynomials*, J. Number Theory **159** (2016), 89–100; Available online at `http://dx.doi.org/10.1016/j.jnt.2015.07.021`.

[14] F. Qi and B.-N. Guo, *Logarithmically complete monotonicity of a function related to the Catalan–Qi function*, Acta Univ. Sapientiae Math. **8** (2016), no. 1, 93–102; Available online at `http://dx.doi.org/10.1515/ausm-2016-0006`.

[15] F. Qi and B.-N. Guo, *Logarithmically complete monotonicity of Catalan–Qi function related to Catalan numbers*, Cogent Math. (2016), **3**:1179379, 6 pages; Available online at `http://dx.doi.org/10.1080/23311835.2016.1179379`.

[16] F. Qi and B.-N. Guo, *Viewing some ordinary differential equations from the angle of derivative polynomials*, Preprints **2016**, 2016100043, 12 pages; Available online at `http://dx.doi.org/10.20944/preprints201610.0043.v1`.

[17] F. Qi, X.-T. Shi, and B.-N. Guo, *Two explicit formulas of the Schröder numbers*, Integers **16** (2016), Paper No. A23, 15 pages.

[18] F. Qi, X.-T. Shi, F.-F. Liu, and D. V. Kruchinin *Several formulas for special values of the Bell polynomials of the second kind and applications*, J. Appl. Anal. Comput. **7** (2017), in press. ResearchGate Technical Report (2015), available online at `http://dx.doi.org/10.13140/RG.2.1.3230.1927`.

[19] F. Qi and M.-M. Zheng, *Explicit expressions for a family of the Bell polynomials and applications*, Appl. Math. Comput. **258** (2015), 597–607; Available online at `http://dx.doi.org/10.1016/j.amc.2015.02.027`.

[20] X.-T. Shi, F.-F. Liu, and F. Qi, *An integral representation of the Catalan numbers*, Glob. J. Math. Anal. **3** (2015), no. 3, 130–133; Available online at `http://dx.doi.org/10.14419/gjma.v3i3.5055`.

[21] Z.-W. Sun, *Congruences involving generalized central trinomial coefficients*, Sci. China Math. **57** (2014), no. 7, 1375–1400; Available online at `http://dx.doi.org/10.1007/s11425-014-4809-z`.

[22] Y. Wang and Z.-H. Zhang, *Combinatorics of generalized Motzkin numbers*, J. Integer Seq. **18** (2015), no. 2, Article 15.2.4, 9 pages.

[23] C.-F. Wei and F. Qi, *Several closed expressions for the Euler numbers*, J. Inequal. Appl. 2015, **2015**:219, 8 pages; Available online at `http://dx.doi.org/10.1186/s13660-015-0738-9`.

[24] J.-L. Zhao and F. Qi, *Two explicit formulas for the generalized Motzkin numbers*, J. Inequal. Appl. **2017**, 2017:44, 8 pages; Available online at `http://dx.doi.org/10.1186/s13660-017-1313-3`.

© 2017 by the authors. Licensee *Preprints,* Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons by Attribution (CC-BY) license (http://creativecommons.org/licenses/by/4.0/).