

Article

Not peer-reviewed version

At-Sea Measurement of the Effect of Ship Noise on Mussel Behaviour

[Soledad Torres-Guijarro](#)*, [David Santos-Domínguez](#), [Jose MF Babarro](#), [Laura García Peteiro](#), [Miguel Gilcoto](#)

Posted Date: 27 May 2025

doi: 10.20944/preprints202505.2027.v1

Keywords: *Mytilus galloprovincialis*; mussel; underwater noise; particle motion; behaviour



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Article

At-Sea Measurement of the Effect of Ship Noise on Mussel Behaviour

Soledad Torres-Guijarro ^{1,*}, David Santos-Domínguez ¹, Jose MF Babarro ²
and Laura García Peteiro ² and Miguel Gilcoto ²

¹atlanTTic Research Center, Universidade de Vigo, Spain; dsantos@gts.uvigo.es

² Instituto de Investigaciones Marinas de Vigo, IIM-CSIC, Spain

* Correspondence: soledadtorres@uvigo.es; Tel.: +34986818674

Abstract: Anthropogenic underwater noise is an increasing form of pollution that negatively affects biota. The effect of this pollutant on many marine species is still largely unknown, especially those that are more sensitive to particle motion than to sound pressure. In these cases, experiments at sea are necessary, due to the difficulty of recreating the particle movement of a real acoustic field under laboratory conditions. This work aims to contribute to the knowledge of the effect of ship noise on the behaviour of mussels (*Mytilus galloprovincialis*), performing measurements at sea on a real mussel cultivation raft for the first time. The study is carried out on cluster-forming individuals living in the rafts where they are cultivated. Their behaviour is monitored by means of valvometry systems, which measure the magnitude of shell opening using a High-Frequency Non-Invasive (HFNI) system. Simultaneously, the acoustic field generated by the abundant traffic in the area is measured. The results show cause-effect relationships between ship noise and valve closure events.

Keywords: *Mytilus galloprovincialis*; mussel; underwater noise; particle motion; behaviour

1. Introduction

Underwater noise of anthropogenic origin is an increasing form of pollution that negatively affects biota. The effect of this pollutant on many marine species remains largely unknown and depends on both the characteristics of the sound (level, frequency, power distribution, distance to the source, duration, etc.) and the sensitivity of the species under study.

Although the effects of underwater noise on marine fauna are complex and not yet fully understood, its potential for harm is well recognized, and policy measures are already being developed. The Marine Strategy Framework Directive (MSFD) includes a specific descriptor—Descriptor 11—focused on underwater noise as part of its criteria for achieving ‘Good Environmental Status’ (GES). These descriptors aim to assess the overall health of the marine environment, encompassing both ecological aspects, such as biodiversity and food webs, and human-induced pressures, including commercial fisheries, marine litter, chemical contaminants, and the input of energy—such as underwater noise [1,2].

Over the past decades, significant progress has been made in understanding the effects of noise, particularly in relation to behavioural responses in marine mammals [3,4]. This advancement has been driven by innovative technologies, large-scale coordinated field studies, and dedicated funding. However, our knowledge of how noise impacts fish, and especially invertebrates, remains limited and continues to lag behind [5].

Marine invertebrates represent the largest proportion of marine biomass, are indicators of environmental health, and play a key role in marine ecosystems [6]. They perform essential ecosystem services, such as bioturbation (e.g., infaunal bivalves buried in the sediment), the removal of a large part of the nutrients from the water column through the filtration of large volumes of water (bioextraction) that could otherwise generate processes of eutrophication and/or anoxia, the stabilisation of the coastline against the impact of waves and coastal erosion processes, aquaculture

as a source of high nutritional quality protein, and low-carbon footprint production systems representing socio-economic sectors of great relevance. In addition, many of these organisms are sessile or have relatively little capacity for movement, which is why they are used as environmental sentinels (biosensors) to report the magnitude and effects of abiotic and/or biological environmental changes.

In the last decades numerous studies have tried to assess in different ways the effects of different stressors on bivalves [7,8]. But only recently experiments have been conducted in relation to the effect of noise as a stressor, and how anthropogenic or low frequency sounds can interfere with the behaviour, biochemistry and ecophysiology of bivalve species such as *Mytilus edulis* [9–13], *Mytilus galloprovincialis* [10,14,15], *Sinonovacula constricta* [16], *Perna perna* [17], *Crassostrea virginica* [18] or *Pinctada radiata* [19], among others.

Valve gaping behavioural impairments related to underwater noise are the most frequently reported effects on bivalves [9–19]. Bivalve gaping behaviour can be monitored remotely and continuously in natural environments over extended periods, using HFNI (High-Frequency Non-Invasive) valvometers, allowing for the use of bivalves as biosensors to detect and provide early-warning signals of changes in GES [20]. Specifically, this technique could be used to monitor specific indicators for underwater noise, with potential applications at harbours or aquaculture sites. Thresholds and restrictions should be set according to different species, environment, and equipment before establishing valvometry as a standard monitoring technique for detecting noise-induced stress in bivalves.

All the dose-response studies carried out to date have been conducted in a laboratory setting. In these experiments, specimens are maintained within a water tank in controlled conditions, with the use of a loudspeaker to reproduce acoustic stimuli and, on certain occasions, a sensor or camera to analyse their behaviour. This experimental approach has intrinsic limitations related to the acoustic field that can be generated under these conditions and how it is perceived by the animals under study, making it more challenging to interpret the results obtained as well as extrapolating them to environmental conditions in nature. In a sound wave, particles in the medium (e.g. water molecules and suspended particulate matter) oscillate about their resting position (i.e. particle motion), causing local compressions and expansions (i.e. sound pressure) [21]. All sound causes fluctuations in both particle motion and sound pressure. Sound pressure is measured using hydrophones, while particle motion, which can be described by displacement, velocity or acceleration, can be measured by accelerometers, geophones or hydrophone pairs, or calculated (except for directional information) from sound pressure measurements under certain conditions detailed below.

Invertebrates lack an ear, and therefore do not detect sound through sound pressure, but they do possess statocysts that sense particle motion, allow them to maintain equilibrium in the water column and to perceive gravity [22]. Besides, in bivalve mollusc species such as scallops, the abdominal sensory organ (ASO) has been shown to be very sensitive to water-borne mechanical vibrations [23] and thus to particle motion.

Therefore, experiments on the effect of noise on invertebrates should be carried out with the particle motion as the acoustic variable of reference, i.e. the acoustic field to which individuals are subjected should be generated in such a way that the particle motion can be controlled. The main source of anthropogenic noise in bivalve culture areas usually comes from maritime traffic (goods and passenger transport, sea farming, fishing, etc.). For example, in the Ría de Vigo, an area of high aquaculture production in NW Spain, it has been shown that the main noise stressor in some of the mussel rafts parks where *Mytilus galloprovincialis* is cultivated is the noise of passenger ships that continuously cross the ria [24]. The difficulty in studying the effect of ship noise is due to the lack of the necessary equipment to do so in a laboratory: a loudspeaker submerged in a small tank playing a recording of boat noise is unable to generate a particle motion similar to that actually caused by a ship in the coastal zone [25]. The scientific community is aware of this limitation, and there is at least one prototype in development of a laboratory tank for invertebrates where particle velocity can be controlled [26]. But until such equipment is fully tested and widely available, experiments at sea are

essential, where the sound field that actually affects the species, with its associated particle motion, is already present.

For this reason, the study presented here has been carried out at sea, and to our knowledge it is the first time that the effect of anthropogenic noise on the behaviour of mussels in their cultivation arrangement in rafts has been studied. The species studied is *Mytilus galloprovincialis*, during its cultivation on floating platforms (rafts) in the Ría de Arousa (Spain). Its behaviour is monitored by measuring the valve opening rhythms in real time, non-invasively and at high frequency. Given that mussels in culture are open practically all the time [27], valve closure either as persistent closure or as micro-closures (high frequency micro-movements of the valves, not detectable to the human eye) respond to a certain degree of environmental stress, which has been observed in the presence of toxic phytoplankton in other studies [28]. In our case, the stressor under study is particle motion produced by the acoustic field and is estimated from in-situ pressure measurements. Its correlation with valve gaping allows us to analyse the effect of ship traffic noise on individuals of this species and to consider the potential use of *M. galloprovincialis* as biosensor for this environmental stressor.

2. Materials and Methods

2.1. Experimental Sites

The experiment was carried out on two mussel rafts between September 6th and November 1st, 2023, following the standard cultivation practices commonly used in the region for *Mytilus galloprovincialis*. The rafts were located in the Ría de Arousa (Figure 1; Table 1), a flooded valley on the northwest coast of Spain (the world’s third-largest mussel producer), which hosts the majority of the country’s mussel production, with over 3,300 cultivation rafts and significant maritime traffic in the area.




Figure 1. Location of the rafts with the mussels studied, in the Ría de Arousa (NW Spain).

Table 1. Location and depth of the water channel of the two rafts, identified by the first 3 letters of the nearest locality.

Raft ID (nearest locality)	Location	Channel depth (m)
VIL (Vilagarcía)	42.6004N, 8.8270W	20
AGU (Aguiño)	42.5184N, 8.9928W	45

The two rafts chosen for the study have very different habitats [29]: the VIL raft is in sheltered, shallow waters, less salty due to the proximity to the mouth of the river Ulla and with low ambient noise; the AGU raft is in more exposed, rougher and deeper waters, with higher ambient noise from

wind and waves. At both rafts, mussel behaviour jointly with acoustic field measurements were continuously monitored.

2.2. Mussel Behaviour Monitoring

The shell opening of each mussel was measured with a valvometry system [30] consisting of a Hall effect sensor (model 49E) and a small magnet (4.8 mm diameter x 0.8 mm high) attached with cyanoacrylate to each of the two valves, as shown in Figure 2. These two elements are placed at the furthest point from the umbo, where the opening is maximum. Both elements have negligible (non-invasive) weight for the individual: 0.1 g for the magnet and 0.5 g for the Hall sensor.



Figure 2. Valvometry sensor attached to the shell of a mussel in a lab setup: magnet (right valve) and Hall effect sensor (left valve).

The Hall sensor measures the magnetic flux density produced by the magnet, which is inversely proportional to the distance between the magnet and the sensor, so the more open the shell is, the smaller the signal generated by the sensor. This signal is acquired with a monitoring system based on an Arduino device. Hall sensors were attached to 26 mussels per raft (14 connected directly to the Arduino's ADCs, and 12 more through three external 4-channel analogue-to-digital converters (ADCs)), allowing to monitor gapping behaviour of 52 mussels in total at a sampling rate of 10 Hz. Each monitoring system also collects information on water temperature, the rope weight with a load cell sensor and the raft movement with an inertial measurement unit (IMU). A real-time clock (RTC) module timestamps each data record, which is stored on a microSD card. The information is sent every 20 minutes to our servers onshore via mobile telephony, and the local copy on the memory card is downloaded on monthly sampling visits. The monitoring systems are powered by batteries recharged by solar panels.

Figure 3 shows a schematic of the complete monitoring system. A HAT (stackable add-on functionality module) including the 3 external ADCs, load cell, fluorometer, IMU sensor, water temperature sensor, clock module and SD module is connected to the Arduino.




Figure 3. Schematic of the raft monitoring system.

Figures 4 and 5 shows photographs of the electronics, the Peli case containing it and the solar panels installed on the raft, the operation of attaching the sensorised mussels to a rope of the raft and a hydrophone mooring operation where the monitoring system installed in one of the rafts can be seen.

Curation of the valvometry data is performed by visually analysing the signals and comparing them with those previously obtained in laboratory experiments [15,28], in which the good condition of the cables, sensors and magnets could be visually verified. This procedure discards signals whose value ranges are out of the usual magnitude, or show anomalous behaviour, caused by a bad state of magnet or sensor fastening, the cutting of the wires or the detachment of the mussel clusters from the rope.






Figure 4. Top: electronics; Bottom: the Peli case with the electronics and the solar panels installed on the raft.




Figure 5. Left: Operation of attaching the sensorised mussels to a rope of the raft; Right: Hydrophone deployment operation, the monitoring system integrated in the Peli case with the electronics can be seen in the upper right corner, behind the solar panels.

The cured data are processed to convert the magnetic field values, provided by the sensors, into percentages of valve opening. This is done for each mussel individually by assigning the maximum magnetic field value (interpreted to be the 0% valve opening value), and the minimum (assumed to

be 100% opening). Finally, a 10-sample median filter is applied to reduce the sampling frequency to 1 Hz, and a 30-sample moving average smoothing is applied to reduce high-frequency valve closing unrelated to acoustic noise-induced mussel behaviour.

2.3. Acoustic Field Measurement

Systems for measuring particle motion, which is the acoustic variable that mussels perceive (accelerometers, geophones or hydrophone pairs) need to be installed in such a way that they move as little as possible, so that their own movement does not mask the movement of the water particles they are intended to measure. For this reason, they are usually installed on the seabed and are fitted with a calibrated suspension system that minimises the movement of the sensor in the water [25]. This deployment is incompatible with the measurement of the acoustic field affecting the mussels monitored in this study, because they hang from rafts at a considerable distance from the bottom. The chosen alternative is therefore to suspend a hydrophone, whose measurements are much more immune to raft movement and estimate particle movement from its measurements.

Given the characteristics of the experimental site (floating platforms not connected to land), and in order not to overload the power supply and the data storage and transmission capacity of the monitoring systems installed in the rafts, it was decided to use autonomous hydrophones with batteries and local data storage (Marsensing digitaHyd SR-1). They are deployed by hanging from a line from each raft, at a depth of 6 m, corresponding to half the mussel ropes' length. The line is suspended from a spring to minimise the hydrostatic noise produced by the vertical oscillations of the raft caused by waves and is ballasted at its lower end. During the month-long experiment, 4 trips were made to the two rafts to install and clean the hydrophones, and to replace batteries and data storage cards.

The hydrophones were programmed to record the sound in 3-minute files in wav format, with a gain of 1x, a resolution of 16 bits/sample and a sampling frequency of 52734 Hz. With this configuration the effective frequency range, between 63 Hz and 20 kHz, covers the significant sound emission from ship traffic, the main source of anthropogenic noise in the area.

The calculation of the sound pressure level from the sound recordings is performed in 1s segments, which are transformed to the frequency domain using the Fourier transform. The sensitivity curve of each hydrophone is then applied to obtain the spectral sound pressure density $P(f)$ in dB re 1μPa with a time resolution of 1 second and frequency resolution of 1 Hz.

The particle motion is described by the displacement, velocity, or acceleration of the water molecules due to the propagation of the sound wave. These three variables can be calculated, except for the direction information, from the measured sound pressure, under certain conditions. Thus, when the acoustic field can be properly approximated to a plane wave or spherical propagation from a monopole, then it is possible to calculate the magnitude of the particle velocity from a sound pressure measurement, using the expression [25]:

$$u(t) = \frac{p(t)}{\rho c_w}, \quad (1)$$

where u is the magnitude of the particle velocity as a function of time, p is the sound pressure, ρ is the density of water, and c_w is the speed of sound propagation in water.

The monopole condition (point source radiating equally in all directions) is difficult to fulfil in our case, since the sound sources are ships of several metres in length that can hardly be modelled as point sources. The plane wave hypothesis implies that the wave propagates far enough away from its source for the wavefront to be considered flat, and that the propagation occurs without interference from other waves, including reflections. The applicability of this hypothesis depends on the frequencies of interest and the properties of the waveguide. These properties are the water depth, the distance to the sound source, the type of source and the sound velocities in the water and sediment. In practice, the plane wave approximation is reasonable when two conditions are met [25]:

1) the sound source is at a distance greater than one wavelength at the lowest frequency of interest, and

2) the lowest frequency of interest is higher than the cut-off frequency of the waveguide formed by the bottom and the sea surface.

To specify when condition 1) is met, we start from the relationship between wavelength and frequency in a sound wave:

$$\lambda = \frac{c_w}{f}, \quad (2)$$

where λ is the wavelength and f the frequency.

The lowest frequency of interest in our experiment is 63 Hz, which is the lower cut-off frequency of the hydrophones used. At this frequency the wavelength is approximately 24 m, which means that ships passing at a distance from the hydrophone greater than 24 m produce an acoustic field that meets condition 1). In practice, during our experiment we applied this condition by discarding sound and valvometry records during ship passes at shorter distances, calculated from Automatic Identification System (AIS) data. In the case of the raft service vessels which usually sail in their vicinity and do not normally carry AIS, we determined by in-situ recordings that at distances greater than 24m they generate levels below 140 dB, and we eliminated from our analysis the records with levels above this value.

To determine when condition 2) is met, we start from the expression of the waveguide cut-off frequency:

$$f_0 = \frac{c_w}{4D \sqrt{1 - \left(\frac{c_w}{c_b}\right)^2}}, \quad (3)$$

where f_0 is the cut-off frequency, D the depth of the water channel, and c_b the sound propagation velocity in the bottom sediment.

The bottom of the experimental area is a mixture of mud and sand, so a reasonable value for c_b is 1600 m/s [31]. Taking the depth of the VIL raft, 20 m (see Table 1), which being in shallower water is the most unfavourable, and an average value of 1500 m/s for c_w , we obtain a waveguide cut-off frequency of 54 Hz, below the lower cut-off frequency of our equipment. Therefore condition 2) is fulfilled for all frequencies of interest.

As a conclusion, it is feasible to calculate the particle motion from the sound pressure measurement made with a single hydrophone, under the plane wave hypothesis, for frequencies above 63 Hz, provided that the ship producing the measured noise is at a distance of more than 24 m from the hydrophone. On the other hand, it must be taken into account that surface and bottom reflections influence the degree to which it is reasonable to assume that the plane wave condition is fulfilled. For this reason, a hydrophone installation depth of 6 m has been chosen, neither too close to the surface nor too close to the bottom.

In practice, the estimation of the magnitude of the particle motion from the sound pressure has been made in the frequency domain using expression (4), equivalent to (1),

$$U(f) = \frac{P(f)}{\rho c_w}, \quad (4)$$

where $U(f)$ is the magnitude of the particle velocity spectral density and $P(f)$ is the sound pressure spectral density.

Since displacement, velocity and acceleration are related to each other by derivative in time, the properties of the Fourier transform allow to express

$$\Delta(f) = \frac{U(f)}{2\pi i f}, \quad (5)$$

$$A(f) = 2\pi i f U(f), \quad (6)$$

where $\Delta(f)$ and $A(f)$ represent the particle displacement and acceleration spectral densities, and $i = \sqrt{-1}$.

The particle displacement variables estimated using this methodology have, like the sound pressure spectra from which they are derived, a time resolution of 1s, a frequency resolution of 1 Hz and a frequency range between 63 Hz and 20 kHz. The spectral densities of the four acoustic variables (sound pressure, acceleration, velocity and displacement) can be integrated in a certain frequency band, or in the complete frequency range to obtain their global value, in the time domain, with a sampling frequency of 1 Hz.

To estimate the robustness of the cause-effect relationship between particle motion and valve aperture we employed cross-correlation. This function measures the resemblance between two variables for different time offsets between them, to account for possible lags between cause and effect. The cross-correlation of two jointly stationary random processes x_n and y_n is given by:

$$R_{xy}(m) = E\{x_{n+m}y_n^*\} = E\{x_ny_{n-m}^*\}, \quad (7)$$

where $-\infty < n < \infty$, the asterisk denotes complex conjugation (which can be ignored in our case when working with real variables), m is the sample offset, and E is the expected value operator. Since we have only a finite segment of a random process realisation, we will estimate the cross-correlation by the expression

$$\hat{R}_{xy}(m) = \begin{cases} \sum_{n=0}^{N-m-1} x_{n+m}y_n^*, & m \geq 0, \\ \hat{R}_{yx}^*(-m), & m < 0, \end{cases} \quad (8)$$

where N is the length of both variables. To obtain a normalised estimator, we subtract from each variable its mean value and normalise the result of the calculation so that the autocorrelation for displacement $m=0$ is unity. We thus obtain the cross-correlation coefficients $R_{xy, \text{coeff}}(m)$, which take values between +1 and -1,

$$\hat{R}_{xy, \text{coeff}}(m) = \frac{1}{\sqrt{\hat{R}_{xx}(0)\hat{R}_{yy}(0)}} \hat{R}_{xy}(m). \quad (9)$$

The largest absolute value of the sequence $R_{xy, \text{coeff}}(m)$ is an estimate of the linear dependence (positive or negative) between x and y for any time shift between the two variables. In our case, we bound this shift to ± 600 samples (10 minutes) to accommodate the reaction time of the mussels, and possible slippage between the clocks of the two signals. We therefore define the linear dependence between x and y , LD_{xy} , as

$$LD_{xy} = \max(|\hat{R}_{xy, \text{coeff}}(m)|), \text{ for } -600 < m < 600. \quad (10)$$

3. Results

3.1. Description of the Acoustic Environment of the Experimental Sites

After the experimental phase, more than 50 hours of sound pressure measurements, corresponding to non-contiguous intervals on different days and times within the experimental period, are available for each of the experimental sites.

Based on these measurements, the acoustic environment can be described by sound power Spectral Probability Density and exceedance curves, which are shown in Figure 6 for both locations.

Comparison of graphs in Figure 6 shows that the noise environment at AGU is significantly louder than at VIL: the background noise level, described by the 1% curve, is about 20 dB higher at AGU than at VIL, while the 99% curve, associated with the noisiest events, is about 10 dB higher at AGU. These results are related to the higher wave noise at AGU, located very close to the northern mouth of the estuary. On the other hand, the VIL curves show tonal components of frequencies below

250 Hz, related to the engine and propeller noise caused by boat traffic, which is more intense and closer in VIL than in AGU.

The description of both acoustic environments in terms of the particle motion variables is shown in Figures 7 and 8 using the units recommended in [21]. They show, in boxplot and histogram format, the velocity, acceleration and displacement levels in full band (63 Hz-20 kHz) and in octave bands (from 63 Hz to 4 kHz) integrations of their respective spectra. In the calculation of these statistics, ship passes within 24 m of the hydrophones have been excluded in order to maximise the accuracy of the estimation of particle motion from sound pressure.




Figure 6. Sound power Spectral Probability Density and exceedance curves for 1, 5, 50, 95, 99% of the time in the two rafts: Left: VIL; Right: AGU.

As can be seen in Figures 7 y 8 the frequency distribution of the velocity is parallel to that of the sound pressure (Figure 6), as indicated by equation (4). In contrast, the acceleration has an increasing spectral tilt with frequency and the displacement a decreasing one, as indicated by equations (5) and (6). The range of full-band acceleration values is 25-99 dB re $\mu\text{m}^2/\text{s}^4$ with a mean value of 68 dB re $\mu\text{m}^2/\text{s}^4$ at VIL, and 56-103 dB re $\mu\text{m}^2/\text{s}^4$ and mean of 72 dB re $\mu\text{m}^2/\text{s}^4$ at AGU.




Figure 7. Boxplots and histograms of the full-band and octave-band for particle motion variables: velocity, acceleration, and displacement in VIL. The boxplot box is defined between the 25th and 75th percentiles, with whiskers up to 1.5 times the interquartile range; in red the median and outliers.




Figure 8. Boxplots and histograms of the full-band and octave-band particle motion variables: velocity, acceleration, and displacement in AGU. The boxplot box is defined between the 25th and 75th percentiles, with whiskers up to 1.5 times the interquartile range; in red the median and outliers.

An example of the behaviour of the acoustic variables over time is shown below. The first graph in Figure 9 shows the time evolution of the full-band sound pressure in a segment of a recording containing the sound of the engine of a raft service vessel very close to the hydrophone installed at AGU. The area marked in red corresponds to a sound pressure level higher than 140 dB, which indicates that the distance to the hydrophone was probably less than the 24 m that guarantees the fulfilment of the plane wave hypothesis. The following graphs show the velocity, acceleration and particle displacement estimated from this pressure record.





Figure 9. Time series of pressure and particle motion: velocity, acceleration, and displacement, for a close pass of a raft service vessel. In red SPL above 140 dB.

3.2. Mussel Reactions to Noise

At this point, our aim is to verify whether mussels react to the noise caused by passing ships by modifying their shell opening-closing rhythms. For this purpose, we manually select, with the help of AIS data, analysis intervals in which the movement of particles shows noticeable increases due to the passage of ships. We chose the acceleration variable for this analysis because it is, a priori, the one that best detects the bivalve statocyst [22]. In particular, we show the results in the 1 kHz band, as this is the band that most clearly describes the level changes produced by ship passages, although the temporal evolution of the correlation results with the valvar aperture are similar in other bands.

Figure 10 shows four examples of reactions of mussels in AGU to noise increases produced by ship passages. Examination of the reactions depicted in examples 1 and 4, show that the noise peaks causing significant changes in shell opening rhythm have levels around 50-60 dB $\mu\text{m}^2/\text{s}^4$ of particle acceleration in the 1 kHz band, corresponding in our recordings to 75-85 dB $\mu\text{m}^2/\text{s}^4$ in the full band (63 Hz to 20 kHz).






Figure 10. Examples of reactions of mussels to increased noise levels from passing ships.

Table 2 column 2 shows the linear dependence, LD, as defined in Eq. 10, between the valve opening and the particle acceleration for the four examples. The LD value between valve opening and temperature (column 3) and between valve opening and raft motion due to waves, measured with an accelerometer (column 4), are also given to consider the possibility that the mussel reactions are due to variations in these conditions. The LD values shown retain the sign of the $R_{xy,coeff}$ coefficient obtained in (10). For all values shown in the table, $p<0.05$ is satisfied.

Table 2. Linear dependence between valvar aperture and particle acceleration, water temperature and the modulus of raft acceleration.

	LD _{open, accel}	LD _{open, temp}	LD _{open, raftaccel}
Example 1	-0.59	0.55	-0.22
Example 2	-0.27	0.06	0.56
Example 3	-0.50	-0.47	0.26
Example 4	-0.64	-0.52	-0.40

The analysis of the LD values in Table 2 reveals significant negative correlation values between valve opening and particle acceleration, reflecting that the higher the particle acceleration level, the lower the valve opening, i.e. in these examples, the increased noise level causes partial valvar closure. On the other hand, the LD values between valve opening and the other two environmental variables (water temperature and raft movement) show both positive and negative values, which seems to show that either there is no clear cause-effect relationship between these variables and the valve opening in the examples studied, or the different variables studied interact with other factors not analysed, generating opposite effects.

4. Discussion

The aim of this work is to analyse the reactions of mussels to underwater noise caused by ship traffic establishing a new methodology of in situ monitoring based on more accessible technologies. Therefore, the experimental setup has been deployed at sea, rather than in a laboratory, as all previous studies on bivalves have done [9–19]. Experimenting at sea allows to accurately assess the impact of the particle motion generated by ship traffic noise, the main pollutant in mussel farming areas, which is not possible at a laboratory tank, but also to demonstrate the effectiveness of using HFNI valvometry as a monitoring system to detect stress in marine invertebrates caused by anthropogenic underwater noise.

An important aspect of the methodology followed in this project is the description of the reactions to the acoustic field in terms of particle motion variables rather than with sound pressure. However, suspending particle motion measuring instruments from the rafts makes it very difficult to obtain good quality measurements, because of the noise that would be introduced by the

movement of the sensor itself under these conditions [25,32]. In this study, we performed sound pressure measurements with hydrophones because it was important to sample the acoustic field in the vicinity of the mussels, which are suspended from floating rafts in areas up to 45m deep. Then, the hydrophone sound pressure measurements were transformed into particle movement variables using well-established equations.

The correct application of sound pressure to particle motion transformation requires the fulfilment of certain conditions. It must be estimated under the plane wave hypothesis [25], whose verification in this study is guaranteed by eliminating from the analysis the recordings corresponding to ship passages very close to the hydrophones (less than 24 m). The statistical description of the acoustic environment at the two experimental sites has been carried out in terms of the power spectral density of sound pressure and three particle motion variables (displacement, velocity and acceleration). This description contributes to the general understanding of the behaviour of these variables, which are particularly important for invertebrates and most fish species, and still little studied [22,32,33].

The difficulties inherent to an experiment in the sea are manifold. On the one hand, it is impossible to control environmental variables, which causes uncertainty as to what is the real cause of a change in mussel behaviour. In the examples analysed, we have been able to rule out the effect of raft movement and short-term variation in water temperature, but these are only two of the multiple stressors present in the habitat studied. However, in this regard, we can provide evidence that the maintenance of circadian rhythms of *Mytilus galloprovincialis* in a raft culture situation with respect to 'almost permanent' valvar opening has been previously described even though the set of abiotic/biological factors in the cultivation area fluctuates greatly (e.g. solar irradiance, primary production, temperature, salinity, etc.) [27].

Another difficulty of conducting experiments at sea has to do with the fact that the marine environment is very damaging to electronic equipment. During our experiment, the valvometry systems suffered numerous failures: cable breakage, detachment of sensors, failures in the data acquisition, storage and transmission equipment. It has not been possible to solve many of them because of the difficulty of access to the rafts, and the impossibility of handling the ropes on which the mussels studied are cultivated due to the risk of them becoming detached. In fact, detachments have been significant in the two experimental study stations, associated with the effect of the presence of biofouling and hydrodynamic conditions, correlations that are still in the analysis phase.

For these reasons, the number of correctly monitored specimens is small, and we have only been able to obtain evidence of a cause-effect relationship between the noise of a passing ship and partial valvar closure in a few cases, some of which are shown in this contribution. From these examples, we have been able to estimate particle acceleration levels that cause partial valvar closure for specific individuals (75-85 dB re $\mu\text{m}^2/\text{s}^4$ in the band from 63 Hz to 20 kHz). Previous laboratory studies reported mussels reacting to ship noise levels above 114 dB in the band between 63 Hz to 4kHz, with a lag of about 13 seconds after a sudden increase in sound level [15]. Nonetheless, we are aware that our results are not sufficient to draw valid conclusions for a cultivated biomass and communities, considering the high inter-individual variability of bivalve molluscs in general. The sample size to be tested needs to be significantly expanded to increase the statistical power and contemplate this inter-individual variability for a better understanding of the anthropogenic stressor thresholds that cause changes in the behaviour of cultured populations.

In future studies, to minimise failures in the valvometry systems, it would be advisable to look for an experimental site in protected waters and reduce the size of the colony in which the monitored specimens live. On the other hand, a site accessible from land would facilitate the tasks of monitoring the specimens and repairing the valvometry systems. In addition, an experimental deployment in shallower waters would enable the installation of equipment to measure particle movement, allowing a comparative study with particle movement estimated from sound pressure.

Author Contributions: Conceptualization, S.T.G and D.S.D.; methodology, S.T.G and D.S.D.; software, D.S.D and M.G.; validation, S.T.G and J.MF.B; formal analysis, S.T.G and D.S.D.; investigation, D.S.D and M.G.; resources, S.T.G and J.MF.B; data curation, D.S.D and M.G.; writing—original draft preparation, S.T.G; writing—review and editing, S.T.G., D.S.D, J.MF.B, L.G.P, M.G ; visualization, S.T.G, D.S.D, MG; supervision, S.T.G and J.MF.B; project administration, S.T.G and J.MF.B; funding acquisition, S.T.G and J.MF.B. All authors have read and agreed to the published version of the manuscript.

Funding: Please add: This research has been funded by the company Mensoft Consultores through the Movipart23 contract (CO-0047-23) with the University of Vigo, and has been carried out within the framework of the project Wave effects in the Rías Baixas Upwelling System: Surface dynamics in specific biological case studies (STRAUSS) financed by MCIN/AEI/10.13039/501100011033 (Ministry of Science and Innovation, Government of Spain). The APC was funded by the Galician Regional Government under project ED431B 2024/36 GPC.

Institutional Review Board Statement: Not applicable, molluscs bivalves are invertebrates with no specific regulation for behaviour monitoring.

Data Availability Statement: Data available upon request to interested researchers.

Conflicts of Interest: The authors declare no conflicts of interest. The funder Mensoft Consultores contributed to the design of the study. The funders had no role in the collection, analyses, or interpretation of data in the writing of the manuscript; or in the decision to publish the results.

References

1. Rako-Gospić, N.; Picciulin, M. Chapter Four - Addressing underwater noise: Joint efforts and progress on its global governance. In *Advances in Marine Biology*; Sheppard, C., Ed. Academic Press: London, UK, 2023; Volume 94, pp. 201–232. <https://doi.org/10.1016/bs.amb.2023.02.001>
2. Borja, A.; Elliott, M.; Andersen, J. H.; Cardoso, A. C.; Carstensen, J.; Ferreira, J. G.; ... & Zampoukas, N. Good environmental status of marine ecosystems: what is it and how do we know when we have attained it?. *Marine Pollution Bulletin* **2023**, 76(1-2), pp. 16-27. <https://doi.org/10.1016/j.marpolbul.2013.08.042>
3. Weilgart, L. S. A Brief Review of Known Effects of Noise on Marine Mammals. *International Journal of Comparative Psychology* **2007**, 20(2). <https://doi.org/10.46867/ijcp.2007.20.02.09>
4. Erbe, C.; Dunlop, R.; Dolman, S. Effects of Noise on Marine Mammals. In *Effects of Anthropogenic Noise on Animals*; Slabbekoorn, H.; Dooling, R.; Popper, A.; Fay, R.; Eds. Springer Handbook of Auditory Research, vol 66. Springer, New York, NY, 2018. https://doi.org/10.1007/978-1-4939-8574-6_10
5. Thomsen, F.; Mendes, S.; Bertucci, F.; Breitzke, M.; Ciappi, E.; Cresci, A.; Debusschere, E.; Ducatel, C.; Folegot, F.; Juretzek, C.; Lam, F-P.; O'Brien, J.; dos Santos, M. E. (2021) Addressing underwater noise in Europe: Current state of knowledge and future priorities. In *Future Science Brief 7*; Kellett, P., van den Brand, R.; Alexander, B.; Muniz Piniella, A.; Rodriguez Perez, A.; van Elslander, J.; Heymans, J. J.; Eds. European Marine Board: Ostend, Belgium, 2021, ISSN: 2593-5232. ISBN: 9789464206104. DOI: 10.5281/zenodo.5534224
6. Chen, E. Y.-S. (2021). Often Overlooked: Understanding and Meeting the Current Challenges of Marine Invertebrate Conservation. *Frontiers in Marine Science* **2021**, 8. p. 1161. <https://doi.org/10.3389/fmars.2021.690704>
7. Pourmozaffar, S.; Abrari Lazarjani, S.; Rameshi, H.; Behzadi, S.; Bagheri, T.; Zahedi, M. R.; Sadeghi, A.; Gozari, M.; & Tamadoni Jahromi, S. The role of salinity in physiological responses of bivalves. *Reviews in Aquaculture* **2019**, 12(3), pp. 1548–1566. <https://doi.org/10.1111/raq.12397>
8. Liu, Z.; Yi, Q.; Song, L.; Li, M.; & Wang, L. (2018). The Neuroendocrine-Immune Regulation in Response to Environmental Stress in Marine Bivalves. *Frontiers in Physiology* **2018**, 9, p. 1456. <https://doi.org/10.3389/fphys.2018.01456>
9. Roberts, L.; Cheesman, S.; Breithaupt, T.; Elliott, M. Sensitivity of the mussel *Mytilus edulis* to substrate-borne vibration in relation to anthropogenically generated noise. *Marine Ecology Progress Series* **2015**, 538, pp. 185-195. <https://doi.org/10.3354/meps11468>

10. Baltzer, J.; Maurer, N.; Schaffeld, T.; Ruser, A.; Schnitzler, J. G.; Siebert, U. Effect ranges of underwater noise from anchor vibration operations in the Wadden Sea. *Journal of sea research* **2020**, *162*, p. 101912. <https://doi.org/10.1016/j.seares.2020.101912>
11. Wale, M. A.; Briers, R. A.; Hartl, M. G.; Bryson, D.; Diele, K. From DNA to ecological performance: effects of anthropogenic noise on a reef-building mussel. *Science of the Total Environment* **2019**, *689*, pp. 126-132. <https://doi.org/10.1016/j.scitotenv.2019.06.380>
12. Hubert, J.; Booms, E.; Witbaard, R.; Slabbekoorn, H. Responsiveness and habituation to repeated sound exposures and pulse trains in blue mussels. *Journal of Experimental Marine Biology and Ecology* **2022**, *547*, pp. 151668. <https://doi.org/10.1016/j.jembe.2021.151668>
13. Hubert, J.; Moens, R.; Witbaard, R.; Slabbekoorn, H. Acoustic disturbance in blue mussels: sound-induced valve closure varies with pulse train speed but does not affect phytoplankton clearance rate. *ICES Journal of Marine Science* **2022**, *79*, no 9, pp. 2540-2551. <https://doi.org/10.1093/icesjms/fsac193>
14. Vazzana, M.; Celi, M.; Maricchiolo, G.; Genovese, L.; Corrias, V.; Quinci, E. M.; ... Filiciotto, F. Are mussels able to distinguish underwater sounds? Assessment of the reactions of *Mytilus galloprovincialis* after exposure to lab-generated acoustic signals. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **2016**, *201*, pp. 61-70. <https://doi.org/10.1016/j.cbpa.2016.06.029>
15. Torres-Guijarro, S.; Estevez-Marquez, P.; Babarro, J. M.; Silva-Caride, E.; Santos-Dominguez, D.; Comeau, L. A.; Gilcoto, M. Laboratory measurement of the effect of exposure to ship noise on mussels. In Proceedings of INTER-NOISE and NOISE-CON Congress and Conference, Nantes, France, 25-29 August 2024. https://doi.org/10.3397/IN_2024_2156
16. Peng, C.; Zhao, X.; Liu, S.; Shi, W.; Han, Y.; Guo, C.; ... Liu, G. Effects of anthropogenic sound on digging behavior, metabolism, Ca²⁺/Mg²⁺ ATPase activity, and metabolism-related gene expression of the bivalve *Sinonovacula constricta*. *Scientific reports* **2016**, *6*, no 1, p. 24266. <https://doi.org/10.1038/srep24266>
17. Júnior, U. G. D. M.; Xavier, F. C.; Campbell, D.; Silveira, N.; Versiani, L.; Cumplido, R.; ... Netto, E. B. F. Characterization of the acoustic activity of *Perna perna* (bivalve mollusc) under laboratory conditions. In Proceedings of 5th International Conference on the Effects of Noise on Aquatic Life, Den Haag, The Netherlands, 7-12 July 2019. <https://doi.org/10.1121/2.0001254>
18. Ledoux, T.; Clements, J. C.; Comeau, L. A.; Cervello, G.; Tremblay, R.; Olivier, F.; ... Lamarre, S. G. Effects of anthropogenic sounds on the behavior and physiology of the Eastern oyster (*Crassostrea virginica*). *Frontiers in Marine Science* **2023**, *10*, p. 1104526. <https://doi.org/10.3389/fmars.2023.1104526>
19. Charifi, M.; Khalifa, R.; Giraldez, B.W.; Sow, M.; Hizam, Z.; Carrara, M.; Maneux, E.; Hamza, S.; Bassères, A.; Blanc, P.; Leitão, A.; Massabuau, J.-C. Deep behavioral impairment in the pearl oyster *Pinctada radiata* exposed to anthropogenic noise and light stress. *Frontiers in Marine Science*, **2023**, vol. 10, p. 1251011. <https://doi.org/10.3389/fmars.2023.1251011>
20. Andrade, H.; Massabuau, J.-C.; Cochran, S.; Ciret, P.; Tran, D.; Sow, M.; Camus, L. High Frequency Non-invasive (HFNI) Bio-Sensors As a Potential Tool for Marine Monitoring and Assessments. *Frontiers in Marine Science*, **2016**, vol. 3, p. 187. <https://doi.org/10.3389/fmars.2016.00187>
21. ISO 18405:2017 Underwater acoustics - Terminology, International Organization for Standardization (ISO, Geneva, Switzerland). <https://www.iso.org/standard/62406.html>
22. Solé, M.; Kaifu, K.; Mooney, T. A.; Nedelec, S. L.; Olivier, F.; Radford, A. N.; ... André, M. Marine invertebrates and noise. *Frontiers in Marine Science* **2023**, *10*, p. 1129057. <https://doi.org/10.3389/fmars.2023.1129057>
23. Zhadan, P. M. Directional sensitivity of the Japanese scallop *Mizuhopecten yessoensis* and Swift scallop *Chlamys swifti* to water-borne vibrations. *Russian Journal of Marine Biology*, **2005**, *31*, pp. 28-35. <https://link.springer.com/article/10.1007/s11179-005-0040-7>
24. Santos-Domínguez, D. Metodologías de mapeo de ruido submarino en aguas someras. PhD Thesis, Universidade de Vigo, Vigo, Spain, 2020. <https://www.investigacion.biblioteca.uvigo.es/xmlui/handle/11093/1555>
25. Nedelec, S. L.; Ainslie, M. A.; Andersson, M. H.; Cheong, S. H.; Halvorsen, M. B.; Linné, M.; ... & Ward, J. Best Practice Guide for Underwater Particle Motion Measurement for Biological Applications; University

- of Exeter for the IOGP Marine Sound and Life Joint Industry Programme, Exeter, UK, 2021. <http://dx.doi.org/10.25607/OBP-1726>
26. Schnitzler, J. G.; Aguilar de Soto, N.; André, M.; van Benda-Beckmann, S.; Caro, P. P.; Findlay, C. R.; ... Siebert, U. How SATURN is studying the impact of ship noise on the behaviour, health, energetics, and populations of aquatic organisms. In Proceedings of INTER-NOISE and NOISE-CON Congress and Conference, Nantes, France, 25-29 August 2024. https://doi.org/10.3397/IN_2024_3568
 27. Comeau, L. A.; Babarro, J. M.; Longa, A.; Padin, X. A. Valve-gaping behavior of raft-cultivated mussels in the Ría de Arousa, Spain. *Aquaculture Reports* **2018**, *9*, pp. 68-73. <https://doi.org/10.1016/j.aqrep.2017.12.005>
 28. Comeau, L. A.; Babarro, J. M.; Riobó, P.; Scarratt, M.; Starr, M.; Tremblay, R. PSP-producing dinoflagellate *Alexandrium minutum* induces valve microclosures in the mussel *Mytilus galloprovincialis*. *Aquaculture* **2019**, *500*, pp. 407-413. <https://doi.org/10.1016/j.aquaculture.2018.10.025>
 29. Babarro, J. M.; Filgueira, R.; Padín, X. A.; Longa Portabales, M. A. A novel index of the performance of *Mytilus galloprovincialis* to improve commercial exploitation in aquaculture. *Frontiers in Marine Science* **2020**, *7*, p. 719. <https://doi.org/10.3389/fmars.2020.00719>
 30. Gil Coto, M.; Redondo Caride, W.; Silva Caride, E.; Velo Lanchas, A.; Comeau, L. A.; Filgueira, R.; Fernández Babarro, J. M. Arduino controlled valvometry equipment for mussel raft monitoring. 2023. Proceedings of 10th International Workshop on Marine Technology (MARTECH), Castellón de la Plana, Spain, 19-10 June 2023. <http://hdl.handle.net/2117/391983>
 31. Jensen, F. B.; Kuperman, W. A.; Porter, M. B.; Schmidt, H.; Tolstoy, A. *Computational ocean acoustics*, 1st ed.; Springer, New York, NY, EEUU, 2000; p.38.
 32. Popper, A. N.; Hawkins, A. D. The importance of particle motion to fishes and invertebrates. *The Journal of the Acoustical Society of America* **2018**, *143*, no 1, pp. 470-488. <https://doi.org/10.1121/1.5021594>
 33. Nedelec, S. L.; Campbell, J.; Radford, A. N.; Simpson, S. D.; Merchant, N. D. Particle motion: The missing link in underwater acoustic ecology, *Methods Ecol. Evol.* **2016**, *7*, no 7, pp. 836-842. <https://doi.org/10.1111/2041-210X.12544>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.