

Article

Not peer-reviewed version

Physiological, Biochemical, and Genetic Reactions of Winter Wheat to Drought Under the Influence of Plant Growth Promoting Microorganisms and Calcium

[Mariam Zareyan](#)*, [Rima Mockevičiūtė](#), [Sigita Jurkonienė](#)*, [Virgilija Gavelienė](#), Algimantas Paškevičius, [Vaidevutis Šveikauskas](#)

Posted Date: 28 March 2025

doi: 10.20944/preprints202503.2104.v1

Keywords: drought stress; *Triticum aestivum*; PGPM; *Bacillus subtilis*; calcium salts



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Article

Physiological, Biochemical, and Genetic Reactions of Winter Wheat to Drought Under the Influence of Plant Growth Promoting Microorganisms and Calcium

Mariam Zareyan *, Rima Mockevičiūtė, Sigita Jurkonienė *, Virgilija Gavelienė, Algimantas Paškevičius and Vaidevutis Šveikauskas

Laboratory of Plant Physiology and Laboratory of Biodeterioration Research, Nature Research Centre, Akademijos Str. 2, 08412 Vilnius, Lithuania

* Correspondence: mariam.zareyan@gamtc.lt (M.Z.); sigita.jurkoniene@gamtc.lt (S.J.)

Abstract: Improving wheat drought stress tolerance is a critical and challenging task, and more research is necessary since many parts of the world depend on this crop for food and feed. Our current work is focused on the influence of probiotic microorganisms in combination with calcium salts on the physiological and biochemical metabolic pathways that wheat uses when exposed to drought stress and on the analysis of gene expression levels that contribute to wheat drought tolerance. The research was conducted in the laboratory under controlled conditions, simulating a prolonged drought. Seedlings were treated with different microorganisms (*Bacillus subtilis*, *Lactobacillus paracasei*, and some yeast) in 10^5 CFU/ml concentrations for seed priming and later in the same concentration for seedling spraying. 70 g/m² CaCO₃ or 100 g/m² CaCl₂ was added to the soil before the sowing seeds. Almost all tested treatments improved plant growth and positively affected prolonged drought resistance in winter wheat. *Bacillus subtilis*, in combination with calcium salts, had the greatest effect on maintaining the relative leaf water content (RWC). Proline, MDA, and H₂O₂ tests proved a significant positive impact of the treatments on the plant's response at the biochemical level and growth parameters close to those of irrigated plants, along with lower levels of drought-induced gene expression. This study showed that tested microorganisms in combination with calcium salts can activate plants' defense reactions in response to drought. The practical significance of this study is that these ecological measures can be useful under field conditions.

Keywords: drought stress; *Triticum aestivum*; PGPM; *Bacillus subtilis*; calcium salts

1. Introduction

Drought is the most critical abiotic limiting factor for plant cultivation in many parts of the world [1–5]. Drought affects plants' morphological, physiological, biochemical, and molecular processes, resulting in growth inhibition [6–8]. A bioinformatic analysis of European crop losses over recent decades revealed that drought waves are responsible for greater yield loss in cereals (9% and 7.3%) compared to non-cereal crops (3.8% and 3.1%) [9]. Many studies have explored the genetic and molecular mechanisms behind drought resistance in cereals such as wheat, barley, maize, sorghum, and rice [10,11].

Wheat is the leading food for over a third of the world's population [12]. Wheat production in recent years (2018–2025) is lower than it used to be in the largest producing countries such as Canada, Iran, France, Germany, and Greece due to drought [13–15]. According to the European Drought Observatory (EDO), a drought warning was in effect for nearly half of the EU territory in 2022, with a red alert for 15%. By mid-January 2025, the Combined Drought Indicator (CDI) showed warning drought conditions in southern Italy, the eastern Baltic Sea region, eastern Poland, Belarus, central-eastern Ukraine, Greece, part of the Balkans, Cyprus, Malta, and other Mediterranean islands, as well

as Ireland, northern UK, and more than a half of Türkiye [16]. Therefore, developing new pathways to help plants overcome drought stress is essential.

Nutrient management and organic amendments can reduce the harmful effects of drought stress on crops [17]. As an essential plant nutrient, calcium plays a vital role in plant growth and development. Drought stress is known to alter plant mineral status and metabolic processes. Calcium (here and later Ca) is involved in plant signaling responses to drought [18] and thereby positively affects plant adaptation to drought stress [19]. Ca^{2+} can help plants adapt to drought quickly by adjusting their stomatal opening/closing, optimizing their gas exchange, and improving their photosynthetic efficiency [20]. Probiotic microorganisms increase plants' uptake of essential elements from the soil. For example, microbial biostimulants enhance plant growth under stress conditions by facilitating the uptake of insoluble elements and essential minerals [21]. Probiotic bacteria are also thought to induce drought tolerance in plants by altering root morphology, activating the antioxidant system, and promoting the expression of genes associated with abiotic stress tolerance [22]. Plant growth-promoting microorganisms (PGPMs) use different mechanisms that directly or indirectly help in crop plants' growth and development, increasing crop productivity [23]. *Bacillus subtilis* is one of the most attractive probiotics for developing natural plant protection products because it is generally considered safe for food use [24]. *B. subtilis* strains produce cytokinin that interferes with drought-induced suppression of shoot growth, thereby increasing plant growth during periods of drought [25].

Several drought tolerance indicators have been identified as reliable drought resilience measures, including relative water content (RWC), water use efficiency, stomatal conductance, photosynthesis, and drought-responsive genes. The RWC is considered a measure of plants' water status and the dehydration index. It allows you to assess plants' physiological states and shows their ability to retain water in unfavorable conditions [26,27]. Malondialdehyde (MDA) is commonly used to assess the status of redox and osmotic adaptation, which is important for plant adaptation to environmental stresses. MDA is a good marker of oxidative stress because ROS breaks down polyunsaturated lipids and forms MDA. The overall increase in membrane lipid peroxidation is proportional to the intensity of drought stress and may result from spontaneous reactions of ROS with organic molecules contained in membranes [28]. Among ROS species, hydrogen peroxide (H_2O_2) is an especially important one involved in plant responses to different environmental stressors. During stress, peroxide concentration increases because it works as a signaling molecule. However, increasing H_2O_2 concentration over time can harm plants (oxidative stress) [29]. In stressful conditions, plants accumulate metabolites, especially some amino acids such as proline. Proline plays the role of the osmolyte for plants; during drought stress, it can take the role of a signaling molecule, a metal chelator, and an antioxidant defense molecule. It is often used as a stress marker [30,31]. Plants respond to stressful environments by altering gene expression and protein production [6]. It is known that some genes that code late embryonic protein (LEA) are affected by drought [32,33]. LEA proteins are generally predicted to maintain intrinsically disordered proteins in a fully hydrated state, which can then fold under water-deficient conditions to adopt α -helical structures. Ordered LEAs can act as molecular chaperones to bind enzymes, membranes, water or ions, and reactive oxygen species (ROS) and, therefore, play a critical role in protein/membrane stabilization and maintaining cellular environmental homeostasis under drought stress [33].

The experiments were modeled and conducted with the microorganisms – *Bacillus subtilis*, *Lactobacillus paracasei*, *Zygosaccharomyces bailii*, *Geotrichum silvicola*. This study aimed to investigate how the effects of different microorganisms and Ca could modify the drought tolerance of wheat at the molecular, biochemical, and morphological levels. Further work focused on determining which of the selected microorganisms help the plant overcome drought stress, especially in the presence of added Ca in the soil. The objectives of this study were to (1) explore whether chosen microorganisms and Ca salts improve shoot growth under drought stress, (2) to study the changes in biochemical and molecular responses of treated winter wheat to drought, (3) to investigate which microorganisms work better in combination with Ca.

2. Materials and Methods

2.1. Plant Material, Growth Conditions, and Treatments

Wheat seeds (*Triticum aestivum* L. cv. 'Skagen') were sown in plastic cylindric pots (10 × 13 cm), 20 seeds per pot, in a peat moss substrate (pH 5.5 – 6.5). Plants were germinated and grown under controlled conditions of a constant temperature of 22 ± 1 °C, a photoperiod of 16/8 h, and a fluorescent light photon flux of $60 \mu\text{mol m}^{-2} \text{s}^{-1}$ at soil level (Figure 1). Soil moisture was maintained at 60 - 70%.



Figure 1. Growth conditions.

The following treatments were used for the drought stress control studies:

- a) Ca was added to the soil in the form of CaCO_3 at a rate of 1.3 g per pot, based on 70 g m^{-2} , and in the form of CaCl_2 prediluted in water at a rate of 1 g in 100 ml per pot.
- b) The following microorganisms - *Bacillus subtilis*, *Lactobacillus paracasei*, *Zygosaccharomyces bailii*, *Geotrichum silvicola* - were used for seed priming in 10^5 CFU/ml concentrations (microorganisms were grown in their specific liquid media (Nutrient, MRS, YPD) to the log phase and then diluted with distilled water to obtain the required concentration) and later in the same concentration for seedlings spraying in the 1–2 leaf stage (BBCH-scale 1–2) [34]

2.2. Experimental Design and Drought Conditions

Treatments were added according to the scheme (Table 1). Each treatment was carried out in 6 pots, divided into two groups: 3 for simulating drought and 3 for rational watering. Drought was initiated immediately after plants were sprayed with microorganisms at the 1-2 leaf stage as irrigation termination. As a result, the soil gradually dried, reaching 15 - 20 % humidity. Soil humidity of watered plants was kept at a 60 - 70% level. Soil moisture was measured using a soil moisture meter (Aicevoos, AS-PH3, China).

Table 1. Experimental design.

Treatment	CaCO ₃	CaCl ₂	Microorganisms
Control	-	-	-
CaCO ₃	+	-	-
CaCl ₂	-	+	-
<i>Bacillus subtilis</i>	-	-	+
<i>B. subtilis</i> + CaCO ₃	+	-	+
<i>B. subtilis</i> + CaCl ₂	-	+	+
<i>Geotrichum silvicola</i>	-	-	+
<i>G. silvicola</i> + CaCO ₃	+	-	+
<i>G. silvicola</i> + CaCl ₂	-	+	+
<i>Lactobacillus paracasei</i>	-	-	+
<i>L. paracasei</i> + CaCO ₃	+	-	+
<i>L. paracasei</i> + CaCl ₂	-	+	+
<i>Zygosaccharomyces bailii</i>	-	-	+
<i>Z. bailii</i> + CaCO ₃	+	-	+
<i>Z. bailii</i> + CaCl ₂	-	+	+

2.3. Sampling

Plant samples were taken for analysis when soil moisture in the drought treatments was 15 - 20%. The watered plants were sampled simultaneously (soil moisture 60 - 70%). Shoots of wheat seedlings were sampled for morphometrical measurements: 10 plants from each pot. For biochemical and molecular analysis, three independent replicates were carried out using the third leaves of wheat plants. For MDA, H₂O₂, proline assays, and RNA isolation, the samples were immediately frozen in liquid nitrogen and stored in a low-temperature freezer (Skadi Green line, International Labmate Ltd., St Albans, UK) at -80 °C until the analysis.

2.4. Morphometrical Measurements

Shoot length and biomass were measured using a ruler and balances (Kern EWJ).

2.5. RWC

RWC was determined according to Weng et al. [35]. Fresh wheat leaves were collected and weighed as fresh weight (FW). After that, the leaves were left in the water for 24 hours and weighed again to obtain a saturated weight (SW). The dry weight (DW) was obtained by drying the leaves in a drying chamber and weighted. RWC has been calculated according to the formula [35]:

$$RWC = [(FW - DW) / (SW - DW)] \times 100\%, \tag{1}$$

2.6. Assessment of Biochemical Parameters

2.6.1. H₂O₂

For analysis of H₂O₂ and MDA, leaf material (0.5 g) was homogenized using 5% trichloroacetic acid (TCA) (Sigma-Aldrich, St. Louis, MO, USA). The homogenates were centrifuged at 10000 × g for 15 min (centrifuge MPW-351 R). H₂O₂ content in leaves was determined according to Velikova et al. [36]. The supernatant was mixed with 10 mM, pH 7.0 potassium phosphate buffer (Alfa Aesar) and 1 M potassium iodide (Alfa Aesar) in a ratio of 1:1:2. The reaction solution was incubated for 30 min at 25° C in the dark. The absorbance of the supernatant was measured at 390 nm by a spectrophotometer (Analytik Jena Specord 210 Plus, Analytik Jena, Jena, Germany). The amount of H₂O₂ was calculated using a standard curve. The results are expressed in μmol g⁻¹ FW.

2.6.2. Lipid Peroxidation According to MDA

The method of Hodges et al. [37], with slight modifications, has been used to estimate MDA. The supernatant was added to 20% TCA containing 0.5% thiobarbituric acid (TBA) (Alfa Aesar, Haverhill, MA, USA). The homogenate was incubated in a heater (Blockthermostat BT 200) at 95 °C for 30 minutes and then cooled on the ice. The optical density was measured at 532 and 660 nm by a spectrophotometer (Analytik Jena Specord 210 Plus, Analytik Jena, Jena, Germany). The results were expressed in $\mu\text{mol g}^{-1}\text{FW}$ [37].

2.6.3. Proline

Carillo & Gibon's proline extraction and determination method was used to determine the free proline content [38]. Leaf material (0.5g) was homogenized using 40% ethanol and left overnight at +4 °C. The homogenates were centrifuged at $14000 \times g$ for 5 min (centrifuge MPW-351 R), and the supernatant was added to the reaction mix (1% ninhydrin, 60% acetic acid, 20% ethanol). The homogenate was incubated in a heater (Blockthermostat BT 200) at 95 °C for 20 min. The optical density was measured at 520 nm by a spectrophotometer (Analytik Jena Specord 210 Plus, Analytik Jena, Jena, Germany). The corresponding content of the proline was determined using the standard curve. The calculations were made using the equation given in the method [38]. The results were expressed in $\mu\text{mol g}^{-1}\text{FW}$.

2.7. Molecular Techniques

2.7.1. RNA Extraction and Reverse Transcription

Total RNA was extracted from 200 mg of plant leaf material using the PureLink RNA Mini Kit (Ambion, Waltham, MA, USA) following the manufacturer's recommendations and using Heraeus Fresco 21 Centrifuge (Thermo Scientific, Waltham, MA, USA). To avoid contamination with genomic DNA, extracted total RNA was treated with RapidOut DNA Removal Kit (Thermo Scientific). The concentration and purity of treated RNA were evaluated with the spectrophotometer NanoPhotometer P330 (IMPLEN, Westlake Village, CA, USA). DNase-treated RNA samples were reverse-transcribed using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Waltham, MA, USA) following the manufacturer's recommendations. The obtained cDNA was stored at -20 °C.

2.7.2. Real-Time Quantitative PCR

Real-time quantitative PCR was carried out using SYBR® Green Universal Master Mix kit (Applied Biosystems) following the manufacturer's recommendations and employing QuantStudio™ 5 real-time PCR system (Applied Biosystems). Two microliters of cDNA (the equivalent of 25 ng of total RNA) were used as a template for PCR. Cycling conditions comprised one cycle at 95 °C for 2 min and 40 cycles at 95 °C for 15 s followed by 60 °C for 1 min. After the PCR run, a melting curve was generated and analyzed each time with the Quant Studio Design & Analysis Software v.1.5.2 (Applied Biosystems). Gene expression was calculated using the $2^{-\Delta\Delta C_t}$ method [39].

2.7.3. Primers

The sequences of primers used in the work were based on the sequences of LEA protein genes and were taken from the publication of Ali-Benali et al. [40]. The wheat mitochondrial 26S ribosomal RNA gene was used as a housekeeping gene [40]. The primer concentrations in all cases were 200 nM except for the *Td27e*, where the concentration was 900 nM. The sequences of all primers used in the work are listed in Table 2.

Table 2. The sequences of primers used in the work.

Gene	LEA Proteins Group	Primer Pairs	Primer Sequences (5'-3')
26S		F/R	CCGGTTGTTATGCCAATAGCA/GCGGCGCAGCAGTTCT
Td11	2	F/R	AGGTGATCGATGACAACGGTG/ACCCTCGGTGTCCTTGTTGG
Td29b	4	F/R	CGCACCCAGCTAGTAAGTTTCG/CCCAGCCCAGTAATAACCCAT
Td27e	2	F/R	CAGCACTGAGCCGACGG/ACGTGGAAGTAGAAGGCATTGAC

2.8. Statistical Analysis

The results are presented as mean ± standard deviation (SD) of three independent experiments with at least three replicates. The data were analyzed using a two-way analysis of variance with replication (two-way ANOVA with replication). The significant differences between treatment means were determined using analysis of variance and mean separation at a 5% significance level ($p < 0.05$).

3. Results

3.1. Impact of PGPMs and Ca Salts on Morphometric Parameters of Wheat Seedlings Exposed to Prolonged Drought Stress

The shoot length and biomass of wheat were significantly higher when the plants were treated with probiotic microorganisms and grown in soil supplemented with Ca salts, especially in the ones with the combination of *B. subtilis* and CaCO_3 . This was seen in both drought-stressed and normally watered plants (Table 3).

Table 3. Effect of PGPMs and Ca salts on morphometric parameters of winter wheat seedlings (per plant). Different letters in columns designate statistically significant differences at $p < 0.05$.

	Average shoot length, cm		Average shoot biomass, g	
	Watered	Drought	Watered	Drought
Control	24.2 d	15.95 f	0.288 e	0.039 f
CaCO_3	26.95 d	17.05 e	0.3159 c	0.036 f
CaCl_2	24.55 c	17.75 e	0.267 f	0.0335 f
<i>Bacillus subtilis</i>	27.95 a	20.65 b	0.333 b	0.1278 b
<i>B. subtilis</i> + CaCO_3	27.55 a	21.65 a	0.345 a	0.1164 c
<i>B. subtilis</i> + CaCl_2	27.85 a	19.73 c	0.3197 c	0.1218 b
<i>Geotrichum silvicola</i>	26.6 b	20.5 b	0.29 e	0.107 d
<i>G. silvicola</i> + CaCO_3	26.9 b	19.56 c	0.297 d	0.109 d
<i>G. silvicola</i> + CaCl_2	27.1 ab	18.45 d	0.316 c	0.112 c
<i>Lactobacillus paracasei</i>	25.3 c	18.55 d	0.269 f	0.068 e
<i>L. paracasei</i> + CaCO_3	26.55 b	18.85 d	0.306 d	0.057 e
<i>L. paracasei</i> + CaCl_2	25.8 d	20.45 b	0.253 f	0.134 a
<i>Zygosaccharomyces bailii</i>	25.1 c	18.65 d	0.2846 e	0.072 e
<i>Z. bailii</i> + CaCO_3	26.45 b	16.4 f	0.309 d	0.035 f
<i>Z. bailii</i> + CaCl_2	27.4 a	20.5 b	0.289 e	0.1454 a

3.2. Effect of Used PGPMs and Ca Salts on RWC of Wheat Seedlings Exposed to Prolonged Drought

Wheat plants treated with PGPMs and Ca salts retained higher water content in their leaves after exposure to prolonged drought (Figure 2). The ones with a combination of *B. subtilis* and Ca salts had the best retention of leaf water content at ~ 80% (moderate stress). In comparison, the RWC of Control wheat leaves was 47% (high-stress level).




Figure 2. Impact of PGPMs and Ca salts on RWC of winter wheat leaves after drought stress. Error bars represent the standard deviation of the mean. Means with different letters in the same color columns are significantly different ($p < 0.05$).

3.3. Effect of PGPMs and Ca Salts on Biochemical Responses of Wheat Plants Exposed to Prolonged Drought

3.3.1. Hydrogen Peroxide (H₂O₂)

The levels of H₂O₂ increased in drought-affected plants (Figure 3). However, H₂O₂ content was found to be significantly lower in all variants treated with PGPMs and Ca salts in combinations. The ones treated with *B. subtilis* alone or with Ca salts had the lowest H₂O₂ content, 0.86 - 0.96 $\mu\text{mol g}^{-1}$ FW, compared to 3.85 $\mu\text{mol g}^{-1}$ FW for the Control. The difference is also significant when comparing plants only treated with *Bacillus subtilis*, *Lactobacillus paracasei*, or with Ca salts.




Figure 3. The effect of PGPMs and Ca salts application and prolonged drought stress on winter wheat H₂O₂ content. Error bars represent the standard deviation of the mean. Different lowercase letters in the same color columns indicate statistically significant differences ($p < 0.05$).

3.3.2. Malondialdehyde (MDA)

MDA levels that indicate the lipid peroxidation levels showed that although drought stress significantly increased MDA content in wheat leaves, the amount of MDA was considerably lower in treated plants, especially in plants with a combination of *B. subtilis* and Ca salts (18 - 22 $\mu\text{mol g}^{-1}$ FW) compared to the Control plants (72.4 $\mu\text{mol g}^{-1}$ FW). The difference is also significant when comparing plants only treated with *B. subtilis* (31.013 $\mu\text{mol g}^{-1}$ FW) or with Ca salts (47 - 49.4 $\mu\text{mol g}^{-1}$ FW). Combinations of *Lactobacillus paracasei* and CaCl₂ (28.47 $\mu\text{mol g}^{-1}$ FW) and *Zygosaccharomyces bailii* and CaCl₂ (21.96 $\mu\text{mol g}^{-1}$ FW) were also very effective in reducing MDA levels (Figure 4).




Figure 4. The effect of PGPMs and Ca salts application and prolonged drought stress on winter wheat MDA content. Error bars represent the standard deviation of the mean. Different lowercase letters in the same color columns indicate statistically significant differences ($p < 0.05$).

3.3.3. Free Proline

The amount of free endogenous proline in Control plants exposed to drought stress was 8.88 $\mu\text{mol g}^{-1}$ FW. In contrast, in plants treated with PGPMs and Ca salts, it ranged from 1.97 to 6.1 $\mu\text{mol g}^{-1}$ FW (except *Lactobacillus paracasei* + CaCO_3 , which was closer to control plants), the lowest in the samples treated with the combinations of *Bacillus subtilis* and CaCl_2 and *Zygoaccharomyces bailii* and CaCl_2 (Figure 5).






Figure 5. The effect of PGPMs and Ca salts application and prolonged drought stress on winter wheat proline accumulation. Error bars represent the standard deviation of the mean. Different lowercase letters in the same color columns indicate statistically significant differences ($p < 0.05$).

3.4. Effect of PGPMs and Ca Salts Application on Late Embryogenesis Abundant (lea) Genes Expression Levels of Wheat Plant Exposed to Prolonged Drought

Quantitative results of RT-qPCR analysis showed that among the three studied genes, the *Td27e* gene was more highly expressed in drought stress. The level of *Td27e* expression in the Control sample was significantly higher than in the all treated ones (Figure 6a). *Td29b* gene expression levels were markedly lower in all treated variants compared with Control (Figure 6b). The lowest levels of expression of the *Td11* gene were recorded in the samples of plants treated with the combinations of *Bacillus subtilis* and Ca salts and *Geotrichum silvicola* and Ca salts (Figure 6c).






Figure 6. The effects of PGPMs and Ca salts application and prolonged drought stress on winter wheat lea genes (a - *Td27e*, b - *Td29b*, c - *Td11*) expression level. Different lowercase letters in different gene fold changes indicate statistically significant differences ($p < 0.05$).

4. Discussion

According to the Population Division of the United Nations, the world human population will reach about 9.5 billion by 2050. Agriculture is the primary source of food production and must solve this problem by producing enough food for the global population [41].

This work is a logical continuation of the previous work [42]. Since a positive reaction of plants under drought conditions to the combination of commercial plant probiotics and CaCO₃ was shown, it was important to understand which PGPMs were at work here and what role Ca played. Therefore, experiments were modeled and conducted with the ingredient microorganisms (or related to them): *Bacillus subtilis*, *Lactobacillus paracasei*, *Zygosaccharomyces bailii*, *Geotrichum silvicola*. In addition to CaCO₃, CaCl₂ was added to the experimental model as a more easily absorbed form of Ca to compare which works better in combination with microorganisms. A series of experiments were performed to explore how the application of probiotic microorganisms and Ca affects the response of wheat to drought stress at physiological, biochemical, and molecular levels. A test experiment was designed and performed to exclude the possibility that minerals, sugars, or other components of microbial media have a more significant positive effect on plants than microorganisms. The results showed that microorganisms are essential for obtaining the previously observed positive impact. Further work focused on determining which of the selected microorganisms help the plant overcome drought stress, especially in the presence of added Ca in the soil.

Much research has been done to understand how plants respond to drought and how to optimize that response [28,29,31,43]. Thanks to previous research, we were able to select some of the most significant markers that show the level of stress in plants. For example, the RWC results showed that our treatments stimulated plants' ability to retain water and subsequently maintain their physiological state close to that of watered control plants. Proline, MDA, and H₂O₂ tests, in turn,

proved a significant positive impact of the treatments on the plant's response at the biochemical level. Some of the PGPMs used with Ca salts, maintained concentrations of these stress indicators at almost half those of untreated controls. The MDA concentration difference is significant when comparing plants only treated with *B. subtilis* or in combination with Ca salts, which indicates that exactly the combination of them is the most beneficial for the plant. The same is visible with H₂O₂ test results: when comparing plants only treated with *Bacillus subtilis*, *Lactobacillus paracasei*, or in combination with Ca salts and for proline test with the combinations of *Bacillus subtilis* and CaCl₂ and *Zygosaccharomyces bailii* and CaCl₂. The three genes that we used for our studies have a major role in wheat response to drought stress in the early stage of growth. *Td27e* and *Td11* code for group 2 LEA proteins, also named dehydrins. Dehydrins are produced by plants in response to cold and drought stress and are well-known as stress proteins [40,44]. *Td27e* gene encode YSK2-type dehydrin, *Td11* gene a SK3-type dehydrin. *Td29b* gene codes for a group 4 LEA protein and plays an important role in the early stages of seed development by protecting cells against damage caused by desiccation [40]. In this study, these genes were used as stress indicators to see the level of changes at the molecular level in plants and the effects of treatments at that level. As a result, their expression levels proved to be significantly less when plants were treated with chosen PGPMs and Ca salts.

Several studies have been conducted on using plant growth-promoting microorganisms under abiotic stress, especially from the *Bacillus* genus, and they have shown great potential for organic farming [5,45,46]. Previous research has already shown that *B. subtilis* plays an important role in wheat's response to adverse environmental factors [47]. Based on our experiments, we can confirm that *B. subtilis* helped winter wheat plants maintain growth even under prolonged drought conditions and maintained their biochemical and molecular homeostasis close to that of plants watered rationally. Moreover, we see that in combination with Ca salts, the positive effect of *B. subtilis* increases significantly.

In contrast to the *Bacillus* genus, several studies have been conducted in animal models that have linked the genus *Lactobacillus* and the efficiency of Ca absorption in their presence [48–50]. An interesting research connected the use of Ca supplements with *L. paracasei* colonization in the gut [51]. These studies and our results allow us to hypothesize that using *Lactobacillus* species in combination with Ca may also positively affect plants, but further research is necessary to clarify the essence of this combination.

Some other studies have linked yeast (*Zygosaccharomyces* genus) use, to improve yield growth under abiotic stress conditions [52,53]. Our results showed great potential for using *Zygosaccharomyces bailii* in combination with Ca salts to alleviate the impact of drought on winter wheat plants. *Geotrichum silvicola*, which isn't a well-investigated yeast yet, also showed this potential.

The results of our research allow us to propose several hypotheses of the connection of the use of PGPMs in combination with Ca for further investigation of possible pathways related to their effect on plants during drought. The current winter wheat study data showed that using these treatments can activate the defense responses of plants that can compensate for the negative impact of drought. This again proves the results of previous research [54,55] that plant growth-promoting microorganisms have big potential to be used in agriculture.

5. Conclusions

All treatments have a generally positive effect in drought conditions. Exogenous application of our PGPMs and Ca salts improved the prolonged drought tolerance of winter wheat. *B. subtilis*, in combination with Ca salts, had the most significant impact on maintaining the relative water content of the leaves and keeping plant growth parameters close to those of irrigated plants. This treatment also showed the best results for maintaining plants' biochemical and molecular homeostasis. The results of this study in winter wheat showed that using probiotic microorganisms in combination with Ca salts can activate defense reactions that can compensate for the harmful effects of drought stress. PGPMs and Ca salts used in this study are suitable for organic farming since they won't harm the environment.

Author Contributions: Conceptualization, M.Z. and V.Š.; methodology, S.J., A.P., R.M. and V.Š.; formal analysis, V.Š. and M.Z.; investigation, M.Z.; data curation, M.Z. and V.Š.; writing—original draft preparation M.Z.; writing—review and editing, S.J., V.G., R.M.; visualization, V.Š. and M.Z. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Nature Research Centre R & D II program.

Data Availability Statement: The data supporting the reported results can be found in the Nature Research Centre's archive of scientific reports.

Acknowledgments: We thank the staff of the Laboratory of Plant Physiology of the Nature Research Centre for their support and assistance and the Laboratory of Biodeterioration Research for providing the necessary microorganisms.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

The following abbreviations are used in this manuscript:

Ca	Calcium
MDA	Malondialdehyde
LEA	Late embryogenesis abundant
PGPM	Plant growth promoting microorganisms
RWC	Relative water content
ROS	Reactive oxygen species

References

1. Fathi, A.; Tari, D.B. Effect of drought stress and its mechanism in plants. *Int. J. Life Sci.* **2016**, *10*, 1–6.
2. Fahad, S.; Bajwa, A.A.; Nazir, U.; Anjum, S.A.; Farooq, A.; Zohaib, A.; Sadia, S.; Nasim, W.; Adkins, S.; Saud, S.; Ihsan, M.Z.; Alharby, H.; Wu, C.; Wang, D.; Huang, J. Crop production under drought and heat stress: plant responses and management options. *Front Plant Sci.* **2017**, *8*, 1147.
3. Zia, R.; Nawaz, M.S.; Siddique, M.J.; Hakim, S.; Imran, A. Plant survival under drought stress: implications, adaptive responses, and integrated rhizosphere management strategy for stress mitigation. *Microbiol. Res.* **2021**, *242*, 126626.
4. Todorova, D.; Sergiev, I.; Katerova, Z.; Shopova, E.; Dimitrova, L.; Brankova, L. Assessment of the biochemical responses of wheat seedlings to soil drought after application of selective herbicide. *Plants* **2021**, *10*, 733.
5. Chattaraj, S.; Samantaray, A.; Ganguly, A.; Thatoi, H. Employing plant growth-promoting rhizobacteria for abiotic stress mitigation in plants: with a focus on drought stress. *Discov. Appl. Sci.* **2025**, *7*, 68.
6. Nezhadahmadi, A.; Prodhan, Z.H.; Faruq, G. Drought tolerance in wheat. *Sci. World J.* **2013**, *12*, 1–4.
7. Seleiman, M.F.; Al-Suhaibani, N.; Ali, N.; Akmal, M.; Alotaibi, M.; Refay, Y.; Dindaroglu, T.; Abdul-Wajid, H.H.; Battaglia, M.L. Drought stress impacts on plants and different approaches to alleviate its adverse effects. *Plants (Basel)* **2021**, *10*, 259.
8. Oguz, M.C.; Aycan, M.; Oguz, E.; Poyraz, I.; Yildiz, M. Drought stress tolerance in plants: interplay of molecular, biochemical and physiological responses in important development stages. *Physiologia* **2022**, *2*, 180–197.
9. Brás, T.A.; Seixas, J.; Carvalhais, N.; Jägermeyr, J. Severity of drought and heatwave crop losses tripled over the last five decades in Europe. *Environ. Res. Lett.* **2021**, *16*, 065012.
10. Geng, L.; Li, M.; Zhang, G.; Ye, L. Barley: a potential cereal for producing healthy and functional foods. *Food Qual. Saf.* **2022**, *6*, fyac012.
11. Bafir Rusu, D.C.; Murariu, D.; Gheorghita, R.; Graur, M. Some nutritional value aspects of barley and oat and their impact in human nutrition and healthy life. *Plants* **2024**, *13*, 2764.
12. Sultan, M.A.R.F.; Hui, L.; Yang, L.Y.; Xian, Z.H. Assessment of drought tolerance of some *Triticum* L. species through physiological indices. *Czech J. Genet. Plant Breed.* **2012**, *48*, 178–184.

13. US Wheat Associates 2020. Available online: <https://www.uswheat.org/wheatletter/2020-global-wheat-harvest-update/> (accessed on 12 January 2025).
14. USDA1. World Agricultural Supply and Demand Estimates 2021. Available online: World Agricultural Supply and Demand Estimates (WASDE) | USDA Foreign Agricultural Service (accessed on 20 January 2025).
15. Eurostat. Available online: <https://ec.europa.eu/eurostat/web/agriculture/publications> (accessed on 15 February 2025).
16. European Drought Observatory. Available online: https://joint-research-centre.ec.europa.eu/european-and-global-drought-observatories/current-drought-situation-europe_en (accessed on 15 February 2025).
17. Farooq, S.; Ul-Allah, S.; Hussain, M. Drought stress in crop plants and its management. In *Disaster Risk Reduction in Agriculture*, Ahmed, M., Ahmad, S., Eds.; Publisher Springer, Singapore, 2023, pp. 21–42.
18. Cousson, A. Involvement of phospholipase C-independent calcium-mediated abscisic acid signaling during *Arabidopsis* response to drought. *Biol. Plant* **2009**, *53*, 53–62.
19. Hosseini, S.A.; Réthoré, E.; Pluchon, S.; Ali, N.; Billiot, B.; Yvin, J.C. Calcium application enhances drought stress tolerance in sugar beet and promotes plant biomass and beetroot sucrose concentration. *Int. J. Mol. Sci.* **2019**, *20*, 3777.
20. Feng, D.; Wang, X.; Gao, J.; Zhang, C.; Liu, H.; Liu, P.; Sun, X. Exogenous calcium: its mechanisms and research advances involved in plant stress tolerance. *Front. Plant Sci.* **2023**, *14*, 1143963.
21. Glick, B.R. Plant growth-promoting bacteria: mechanisms and applications. *Scientifica (Cairo)*, **2012**, *2012*, 963401.
22. Saberi Riseh, R.; Fathi, F.; Moradzadeh-Eskandari, M. The effect of some probiotic bacteria in induction of drought tolerance in cucumber plants. *Microbiol. Metabolites Biotechnol.* **2018**, *2*, 113–127.
23. Kumar, M.; Shamshad Ahmad, P.; Singh, R.P. Plant growth promoting microbes: diverse roles for sustainable and ecofriendly agriculture. *Energy Nexus*, **2022**, *7*, 100133.
24. Lubyanova, A.R.; Allagulova, C.R.; Lastochkina, O.V. The effects of seed pretreatment with endophytic bacteria *Bacillus subtilis* on the water balance of spring and winter wheat seedlings under short-time water deficit. *Plants* **2023**, *12*, 2684.
25. Etesami, H.; Jeong, B.R.; Glick, B.R. Potential use of *Bacillus* spp. as an effective biostimulant against abiotic stresses in crops—A review. *Curr. Res. Biotechnol.* **2023**, *5*, 100128.
26. Soltys-Kalina, D.; Plich, J.; Strzelczyk-Żyta, D.; Śliwka, J.; Marczewski, W. The effect of drought stress on the leaf relative water content and tuber yield of a half-sib family of 'Katahdin'-derived potato cultivars. *Breed Sci.* **2016**, *66*, 328–331.
27. González-Espíndola, L.Á.; Pedroza-Sandoval, A.; Trejo-Calzada, R.; Jacobo-Salcedo, M.d.R.; García de los Santos, G.; Quezada-Rivera, J.J. Relative water content, chlorophyll index, and photosynthetic pigments on *Lotus corniculatus* L. in Response to Water Deficit. *Plants* **2024**, *13*, 961.
28. Khaleghi, A.; Naderi, R.; Brunetti, C.; Maserti, B.E.; Salami, S.A.; Babalar, M. Morphological, physiochemical and antioxidant responses of *Maclura pomifera* to drought stress. *Sci. Rep.* **2019**, *9*, 19250.
29. Belkadi, A.; De Haro, A.; Soengas, P.; Obregon, S.; Cartea, M.E.; Chaibi, W.; Djebali, W. Salicylic acid increases tolerance to oxidative stress induced by hydrogen peroxide accumulation in leaves of cadmium-exposed flax (*Linum usitatissimum* L.). *J. Plant Interact.* **2014**, *9*, 647–654.
30. Hayat, S.; Hayat, Q.; Alyemeni, M.N.; Wani, A.S.; Pichtel, J.; Ahmad, A. Role of proline under changing environments: a review. *Plant Signal Behav.* **2012**, *7*, 1456–1466.
31. Liang, X.; Zhang, L.; Natarajan, S.K.; Becker, D.F. Proline mechanisms of stress survival. 2013, *19* (9), 998–1011. *Antioxid Redox Signal.* **2013**, *19*, 998–1011.
32. Pnueli, L.; Hallak-Herr, E.; Rozenberg M.; Cohen, M.; Goloubinoff, P.; Kaplan, A.; Mittler, R. Molecular and biochemical mechanisms associated with dormancy and drought tolerance in the desert legume *Retama raetam*. *Plant J.* **2002**, *31*, 319–330.
33. Li, Z.; Chi, H.; Liu, C.; Zhang, T.; Han, L.; Li, L.; Pei, X.; Long, Y. Genome-wide identification and functional characterization of *LEA* genes during seed development process in linseed flax (*Linum usitatissimum* L.). *BMC Plant Biol.* **2021**, *21*, 193.

34. Weber, E.; Bleiholder, H.; Lancashire, P.D.; Langelüddecke, R.; Stauss, R.; Van der Boom, T.; Witzten-Berger, A. Growth stages of mono- and dicotyledonous plants. In *BBCH Monograph.*, 1st ed.; Meier, U., Ed.; Julius Kühn-Institut: Quedlinburg, Germany, 2018; pp. 18–21.
35. Weng, M.; Cui, L.; Liu, F.; Zhang, M.; Shan, L.; Yang, S.; Deng, X.-P. Effects of drought stress on antioxidant enzymes in seedlings of different wheat genotypes. *Pak. J. Bot.* **2015**, *47*, 49–56.
36. Velikova, V.; Yordanov, I.; Edreva, A. Oxidative stress and some antioxidant systems in acid rain-treated bean plants: protective role of exogenous polyamines. *Plant Sci.* **2000**, *151*, 59–66.
37. Hodges, D.; DeLong, J.; Forney, C.; Prange, R.K. Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. *Planta* **1999**, *207*, 604–611.
38. Carillo, P.; Gibon, Y. Extraction and Determination of Proline. Available online: https://www.researchgate.net/publication/211353600_PROTOCOL_Extraction_and_determination_of_prolineproline (accessed on 10 July 2016).
39. Livak, K.J.; Schmittgen, T.D. Analysis of relative gene expression data using Real-Time Quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* **2001**, *25*, 402–408.
40. Ali-Benali, M.A.; Alary, R.; Joudrier, P.; Gautier, M.-F. Comparative expression of five LEA genes during wheat seed development and in response to abiotic stresses by real-time quantitative RT-PCR. *Biochim. Biophys. Acta, Gene Struct. Expression* **2005**, *1730*, 56–65.
41. Menendez, E.; Garcia-Fraile, P. Plant probiotic bacteria: solutions to feed the world. *AIMS Microbiol.* **2017**, *3*, 502–524.
42. Mockevičiūtė, R.; Jurkonienė, S.; Šveikauskas, V.; Zareyan, M.; Jankovska-Bortkevič, E.; Jankauskienė, J.; Kozeko, L.; Gavelienė, V. Probiotics, proline and calcium induced protective responses of *Triticum aestivum* under drought stress. *Plants* **2023**, *12*, 1301.
43. Dikilitas, M.; Şimşek, E.; Roychoudhury, A. Role of proline and glycine betaine in overcoming abiotic stresses. In *Protective Chemical Agents in the Amelioration of Plant Abiotic Stress: Biochemical and Molecular Perspectives*, 1st ed.; Roychoudhury, A., Tripathi, D.K., Eds.; John Wiley & Sons Ltd, 2020; pp. 1–23.
44. Yang, Y.; He, M.; Zhu, Z.; Li, S.; Xu, Y.; Zhang, C.; Singer, S.; Wang, Y. Identification of the dehydrin gene family from grapevine species and analysis of their responsiveness of various forms of abiotic and biotic stress. *BMC Plant Biol.* **2012**, *12*, 140.
45. Lalarukh, I.; Al-Dhumri, S.A.; Al-Ani, L.K.T.; Hussain, R.; Al Mutairi, K.A.; Mansoor, N.; Amjad, S.F.; Abbas, M.H.H.; Abdelhafez, A.A.; Poczar, P.; Meena, K.R.; Galal, T.M. A Combined use of Rhizobacteria and Moringa leaf extract mitigates the adverse effects of drought stress in wheat (*Triticum aestivum* L.). *Front Microbiol.* **2022**, *13*, 813415.
46. Yadav, V.K.; Yadav, R.C.; Choudhary, P.; Sharma, S.K.; Bhagat, N. Mitigation of drought stress in wheat (*Triticum aestivum* L.) by inoculation of drought tolerant *Bacillus paramycoides* DT-85 and *Bacillus paranthracis* DT-97. *J. Appl. Biol. Biotechnol.* **2022**, *10*, 59–69.
47. Xie, J.; Bai, L.; Zhou, W. Research progress of *Bacillus subtilis* in wheat growth promotion and disease control. *Int. J. Biol. Life Sci.* **2024**, *7*, 68–72.
48. Collins, F.L.; Rios-Arce, N.D.; Schepper, J.D.; Parameswaran, N.; McCabe, L.R. The potential of probiotics as a therapy for osteoporosis. *Microbiol Spectr.* **2017**, *5*, 10.1128/microbiolspec.bad-0015-2016.
49. Raveschot, C.; Coutte, F.; Frémont, M.; Vaeremans, M.; Dugersuren, J.; Demberel, S.; Drider, D.; Dhulster, P.; Flahaut, C.; Cudennec, B. Probiotic *Lactobacillus* strains from Mongolia improve calcium transport and uptake by intestinal cells in vitro. *Food Res. Int.* **2020**, *133*, 109201.
50. Harahap, I.A.; Suliburska, J. Probiotics and isoflavones as a promising therapeutic for calcium status and bone health: a narrative review. *Foods* **2021**, *10*, 2685.
51. Trautvetter, U.; Ditscheida, B.; Kiehn, M.; Jahreis, G. A combination of calcium phosphate and probiotics beneficially influences intestinal lactobacilli and cholesterol metabolism in humans. *Clin Nutr.* **2012**, *31*, 230–237.
52. Bilek, F.N.; Rezki, M.A.; Grondin, C.; Yahia, N.; Bekki, A. Plant growth promoting characteristics and stress tolerance of yeasts isolated from algerian agricultural soils. *South Asian J. Exp. Biol.* **2020**, *10*, 413–426.

53. Pérez-Moncada, U.A.; Santander, C.; Ruiz, A.; Vidal, C.; Santos, C.; Cornejo, P. Design of microbial consortia based on *Arbuscular Mycorrhizal* fungi, yeasts, and bacteria to improve the biochemical, nutritional, and physiological status of strawberry plants growing under water deficits. *Plants* **2024**, *13*, 1556.
54. Menendez, E.; Garcia-Fraile, P. Plant probiotic bacteria: solutions to feed the world. *AIMS Microbiol.* **2017**, *3*, 502–524.
55. Souza Vandenberghe, L.P.; Garcia, L.M.B.; Rodrigues, C.; Camara, M.C.; de Melo Pereira, G.V.; de Oliveira, J.; Socol, C.R. Potential applications of plant probiotic microorganisms in agriculture and forestry. *AIMS Microbiol.* **2017**, *3*, 629–648.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.