

*Article*

# Effect of Animal Welfare Quality on the Reproductive Performance of Extensive Pasture-based Beef Cows in New Zealand

Y. Baby Kaurivi <sup>1,\*</sup>, Richard Laven <sup>1</sup>, Tim Parkinson <sup>1</sup>, Rebecca Hickson <sup>2</sup>, and Kevin Stafford <sup>2</sup>

<sup>1</sup> School of Veterinary Medicine, Massey University, Private Bag 11 222, Palmerston North 4442, New Zealand; r.laven@massey.ac.nz (R.L.); t.j.parkinson@massey.ac.nz (T.P.)

<sup>2</sup> School of Agriculture and Environmental Management, Massey University, Private Bag 11 222, Palmerston North 4442, New Zealand; R.Hickson@massey.ac.nz (R.H.); K.J.Stafford@massey.ac.nz (K.S.)

\* Correspondence: Y.Kaurivi@massey.ac.nz; Tel.: +64-6350-5328

**Abstract:** One key area where animal welfare may relate to productivity is through reproductive performance. Welfare was assessed on 25 extensively managed pastoral New Zealand beef farms and the relationship between welfare and reproductive performance was explored. Relationships between welfare measures and key reproductive performance indicators (pregnancy rate, weaning rate, mating period and bull: cow ratio) were investigated using an exploratory Principal Components Analysis and linear regression model. Seven welfare measures (thinness, poor rumen fill, dirtiness, blindness, mortality, health checks of pregnant cows and yarding frequency/year) showed potential influence on reproductive performance, and lameness was retained individually as a potential measure. Mean pregnancy rates in both 2018 (PD18) and 2017 (PD17) were ~91% and mean weaning rate was 84%. Of the welfare measures, only lameness had a direct association with pregnancy rate, as well as confounding effect on the association between mating period and pregnancy rate. The bull: cow ration (mean 1:31) and reproductive conditions (dystocia, abortion, vaginal prolapse) did not influence pregnancy and weaning rates. In the study population there was no clear association between welfare and reproductive performance, except for the confounding effects of lameness.

**Keywords:** beef cow welfare, reproductive performance, New Zealand

---

## 1. Introduction

Beef productivity has generally been regarded as the outcome of genetic selection for production traits, such as growth rate and maternal milk yield, and management of nutrition and reproduction. Reproductive efficiency is a critical component of overall performance of cow-and-calf beef cattle enterprises, for which the key performance indicators (KPI) of reproductive efficiency are pregnancy rate, calving and weaning rates, and calving to conception interval [1-3]. Many of the factors that affect these KPIs have been extensively studied, including, for example, macro- and micro-nutrition, changes in cow bodyweight, the effects of topography and climate, the presence of disease, and the bull to cow ratio [1, 4, 5]. Hence, measures of good animal welfare such as low incidences of disease, excellent nutrition, environmental comfort and good stockpersonship are all expected to give a direct or indirect correlation with reproductive performance of cattle, including beef cows in extensive production systems.

In New Zealand, there is an emphasis upon improving reproductive performance in beef cattle. For example, the target is to have 95% of cows in the herd calving and 90% weaning a calf [6]. The proportion of herds achieving this KPI is relatively low [7], so a significant number of studies have examined the factors which allow beef cows to achieve this level of reproductive performance (e.g. [7, 8]). Although there is no standard recipe for achieving good reproductive performance, most New Zealand beef farms have adopted reproduction management plans, with monitoring of key

monitoring KPIs, especially pregnancy rates, calving rates and dates, and managing parturition with assistance when necessary. In addition, reproduction management allows for nutritional management of peri-parturient cows and adding value to planning for weaning and selling [9].

The relationship between the welfare of the cows and their reproductive performance has received relatively less attention than nutritional factors, but, more recently, evidence has started to accumulate that both production and reproduction are influenced by welfare [10-12]. There is strong evidence from other domestic species such as rabbit does [13] and sows [14, 15] that welfare is a significant factor for determining reproductive performance. Reproduction can have direct adverse effects upon animal welfare through, for example, catastrophes such dystocia, abortion and metritis. However, poor performance in other reproductive KPIs, such as low conception rates and long post-partum anoestrus, particularly where many cattle are affected, may also provide an indication that there is poor herd welfare [16]. Whilst it has not been clearly established whether, or to what extent, suboptimal animal welfare is related to suboptimal reproductive KPIs, a recent study by Irico [12] found a direct correlation between welfare measures and reproductive performance of confined beef cattle in two different housing systems.

Hence, the current paper aims to evaluate whether there is a relationship between the different animal welfare standards and the reproductive efficiency of beef cows in New Zealand managed in extensive pasture-based systems. It was postulated that farms with poor welfare indicator scores would be correlated with poor reproductive KPIs. The aim was achieved through a series of factor analysing and selecting the likely measures to predict welfare and reproductive performance. Results from this study are foreseen to fill a gap in the knowledge of how animal welfare quality holistically affects reproductive performance of beef cattle that are extensively reared.

## 2. Materials and Methods

### 2.1 Welfare assessment and reproductive data collection

Welfare of beef herds was assessed using the Welfare Assessment Protocol (WAP) developed by Kaurivi [17, 18]. The WAP was developed as an amalgamation of the best practicable measures in the Welfare Quality protocol [19] and the rangeland-based University of California Davis (UC Davis) Cow-Calf Health & Handling assessment protocol [20], with additional measures suitable for New Zealand beef systems.

The study was carried out on 25 extensive pasture-based beef herds in Waikato district of the North Island of New Zealand. This was a convenience sample of the beef cattle clients of a local rural practice, who used the practice for routine pregnancy diagnosis in autumn. All farms used rotational grazing as their main means of feeding cattle, with supplementary hay or silage given during winter when necessary. No housing or off-pasture feeding was used on any farm. Average herd size (cows mated) was 198 (range 18 - 541).

Each farm was visited twice. The first visit took place in the autumn (March/April 2018) at the time at which pregnancy diagnosis was undertaken. During the first visit, more than half of the animals in each herd were directly observed against the parameters of the WAP whilst they were yarded for pregnancy diagnoses. The second visit was a few months later, in winter, at a time when the animals were not yarded. This visit involved a farm resource evaluation and a questionnaire-guided interview to assess health and management of each herd in the last 12 months. The general cattle management and key health aspects on the farm (disbudding, castration, vaccination, diseases history, cattle deaths, access and type of water supply, feed/pasture condition, wintering practices) were recorded.

Pregnancy rates for 2018 (PD18) were determined at the same time as the animal welfare assessment was undertaken in the yard, while similar data for 2017 (PD17) were retrieved from the veterinary clinic records. The questionnaire-guided interview captured data of cows' reproductive conditions (retained fetal membranes, prolapse vagina/uterus, dystocia and abortion) and their potential causes and outcomes in the previous 12 months. Other reproductive data including bull: cow ratios, start dates of mating and durations of mating periods were summarised along with numbers of calves born, tagged and weaned from 2017 pregnancy testing. Calf mortality was

calculated from calved weaned from in-calf cows. From these data, reproductive KPIs were calculated using an adaption of Hewitt [3]: Table 1).

In order to interpret the collected data of animal welfare measures in a meaningful way, the measures were integrated into the four animal welfare principles (good feeding, appropriate environment, good health and appropriate stockpersonship: Table 2). For each farm, the welfare impact of each measure was categorised separately in 3 categories; 0: poor; 1: marginal and 2: poor/unacceptable welfare. All discrete data were measured according to percentage of cases and given a 3-point ordinal welfare score. Descriptions of how each measure was assessed and categorised was reported by Kaurivi [18].

**Table 1.** Calculation of reproductive key performance indicators

Reproductive KPI	Calculation	Raw data
Pregnancy rate	(A-E)/A %	(A) Number of cows mated
Barren rate	E/A %	(B) Number of bulls
Calving rate	F/A %	(C) Date bulls turned in to herd
Bull: cow ratio	B: A	(D) Date bulls removed from herd
Mating period	D-C days	(E) Number of non-pregnant cows
Weaning rate (weaning/mated cows)	H/A %	(F) Number of calves born
Weaned/known pregnant cows	H/Pregnant	(G) Number of calves tagged
Calf mortality to tagging	(F-G)/F %	(H) Number of calves weaned
Calf mortality tagging to weaning	(G-H)/F %	

**Table 2.** Ascribing welfare measures into welfare principles

Welfare principle	Welfare measures
<b>Good feeding</b>	Body condition, rumen fill, access to water
<b>Appropriate environment</b>	Dirtiness, short tail, hazards, shade, hazards
<b>Good health</b>	Swelling, abrasions, hair loss, blindness, ocular discharge, nasal discharges, diarrhoea, lameness, dystocia, mortality, castration, disbudding, ear tag/notching
<b>Appropriate stockpersonship</b>	Fearful/agitated behaviour, mis-catching, hitting, fall, run, stumble, health checks, yarding frequency, yard design

## 2.2 Data analysis

All data were analysed using SPSS IBM version 27. Descriptive statistics for continuous measures were mean, median, standard deviation, range and percentiles. Qualitative methods were used to analyse the frequency of ordinal measures. The welfare data were correlated with the KPI of each herd. In order to determine how welfare predicts reproductive performance, principal components from 32 welfare measures were used as predictors for reproduction performance. Relationships between reproductive variables (PD17 and PD18 rates, weaning rate, mating period and bull: cow ratio, dystocia, abortion, vaginal prolapse) and the welfare measures were investigated using an exploratory Principal Components Analysis (PCA). Seven variables were selected in the PCA; two were related to *good feeding* (thinness and poor rumen fill), one was related to *appropriate environment* (dirtiness), and two were related to *good health* (blindness, mortality) and two to *appropriate stockpersonship* (yarding frequency/year, health checks of pregnant cows). Principal Components were identified based on Eigenvalue  $>1$ . After extraction of the data, an Oblimin rotation was used to show a correlation [21]. An individual score for each Component was assigned to each measure. Based on explanatory components, the seven measures for PCA were chosen based on: 1) diversity across farms and 2) likely impact on reproductive performance. Lameness and dystocia were added individually as part of the measures predicting reproduction performance based on the likelihood of their affects.

Data were further analysed with linear regression model. The Component scores (BART factor scores) for the 7 selected welfare measures along with lameness, bull: cow ratio and mating period

were then used as predictor variables with one of the reproductive outcomes (PD18, PD17 and Weaning 17) as dependent variables in separate regression models. Backwards stepwise selection of variables was used (target P-value <0.05).

### 3. Results

#### 3.1. Reproductive data

Basic reproductive performance data for the 25 herds are summarised in Table 3. Qualitative data for reproductive conditions (pooled data for abortions, dystocia and vaginal prolapse) are summarised in Table 4. The mean pregnancy rate for 2018 was 91.4%, with only 3 farms achieving pregnancy rates of <85%. The PD18 was like PD17 with 91.9% mean and the same median at 92.7%. The number of calves born was recorded at only 4/25 farms, tagging record was for 18/25 farms (at 2-4 months) while 3/25 farms tagged calves only at weaning (mode 6 months). Calving rate (93%) could only be calculated at the 4/25 farms that recorded number of calves born. Mortality of calves from births to tagging (95.6%) was also only calculated at these 4 farms. Losses from tagging to weaning was calculated for 22 of the farms (excluded 3/25 farm that only tagged at weaning). Calf mortality varied from 0 to 9%, with an overall mean of 2.4%, and 10 (48%) farms achieving <2.5% losses. The mean weaning rate was 83.9%, with only 6 farms achieving ≥90%. The proportion of calves weaned from known pregnant cows was 92.3%. Bull: cow ratio was between 1:20 and 1:40 for all but two farms (one 1:12, the other 1:62). The mating period was between 42 and 72 days in 16 (64%) farms, 4 had short mating periods of 30-41 days, whilst two had mating periods of ≥100 days. There were relatively few reproductive disorders, with an overall average for all disorders in 2017 of 2.3% of the entire cohort of 4637 calving cows (Table 4). Vaginal prolapse was very rare (0.2% of calvings). Overall, 2.6% of calvings were assessed by the farm manager as dystocias, but these data are skewed by one farm that reported an incidence of 17%. All other farms had an incidence of <5.5%. Twelve farms reported no abortions, whilst 2 reported 9%. No cases of retained fetal membranes were reported at any of the farms.

**Table 3.** Summary of reproductive performance data of beef cows in 25 Waikato farms

	% Pregnant 2018	% Pregnant 2017	% Weaned 2017	Bull: Cow ratio	Mating period (d)
<b>Mean</b>	91.4	91.9	83.9	1:31.3	61.8
<b>Median</b>	92.7	92.6	83.3	1:29.7	61.0
<b>Range</b>	69.2-97.9	81.4-100	67.8-96.7	1:12-1:62	30-125
<b>Std dev</b>	5.8	5.1	7.7	9.3	21.4

**Table 4.** Summary of reproductive disorders experienced by beef cows (in-calf 2017 mating) in 25 Waikato farms

	Herd size	Vaginal prolapse (n/farm)	Dystocia (n/farm)	Abortion (n/farm)	Total n/farm
<b>Mean</b>	185.5	0.2	2.6	1.5	4.3
<b>Median</b>	153.0	0.0	1.9	0.6	2.5
<b>Proportion of entire calving cohort</b>		0.1%	1.4%	0.8%	2.3%

#### 3.2. Welfare data

Descriptive statistics for welfare measures are shown in Tables 5 and 6. For *good feeding*, the average of thin cows at 10.7% was skewed by one farm with 61%, whilst 50% of farms had  $\leq 5.7\%$  of thin cows. There was an average of 30.6% of cows with poor rumen fill. For *appropriate environment* 21.3% of cows were assessed as being dirty and 39.6% had faecal soiling. Dirtiness was correlated with the presence of faecal soiling ( $r=0.75$ ; [17]). For most *health-related measures*, the prevalence was low on most farms. The exceptions were lameness and mortality rate. There was an average of 2.7% lame cows (maximum: 12%). Mortality, as presented by accidental deaths, deaths due to diseases and culling for health issues at the herds were present at an average of 3.9% of cattle. Blindness was rare at the farms with an average 0.4% (0-4%). For *stockpersonship*, an average of 2.7% of cows displayed fearful/agitated behaviours (attempt to escape in the race, climbing/pushing others). Most farmers (11/25) did daily health checks on pregnant cows, whilst 9/25 inspected cows once or twice a week and others 5/25 do not inspect within a week. For yarding frequency, no farms yarded cows two times or less per year, and most farms (20/25) were yarded between 3-4 times per year. For painful management procedures, castration was performed on 20 of the 25 farms (mode and median 2 months of age; range 1 to 4 months). Only two farms disbudded calves (at 3 and 4 months respectively). Ear tagging was performed at all farms with median and mode of 2 months.

**Table 5** Descriptive statistics (from 25 Waikato beef farms) for measures recorded as percentage of observed animals. Description of assessments used to create these figures were reported by Kaurivi [18].

Animal Welfare Principles	Animal Welfare Measures	Descriptive Statistics					
		Min	Max	Mean	Percentiles		
					25	50	75
<b>Good Feeding</b>	*Thin cows	0	61	10.7	2.6	5.7	10.0
	*Poor rumen fill	0	68	30.6	15.5	29.9	45.7
<b>Good Environment</b>	Short tail	0	21	4.2	0.6	3.0	6.0
	*Dirtiness	4	50	21.3	10.7	20.6	29.4
	Diarrhoea (faecal staining)	15	87	39.6	24.0	35.7	48.5
<b>Good Health</b>	Swelling	0	5	0.7	0.0	0.0	1.1
	Hair loss	0	1	0.1	0.0	0.0	0.0
	Abrasion	0	2	0.1	0.0	0.0	0.0
	*Lameness	0	12	2.7	0.5	1.5	3.6
	*Blindness	0	4	0.4	0.0	0.0	0.0
	Ocular discharge	0	8	1.5	0.0	0.0	3.2
	Nasal discharge	0	13	1.2	0.0	0.0	1.3
	*Accidental deaths	0	2	0.6	0.0	0.4	1.2
	*Deaths from disease	0	7	2.0	0.9	1.6	2.9
	*Culling for health	0	6	1.2	0.0	0.8	2.0
<b>Stockpersonship</b>	Fearful/Agitate	0	7	2.7	1.3	2.3	4.1
	Fall/lie	0	8	0.9	0.0	0.0	0.8
	Stumble	0	21	1.6	0.0	0.0	1.7
	Run exit	0	51	13.0	2.6	7.8	15.1

\*Measures selected as having high potential effects on reproductive KPI

**Table 6.** Observed frequencies for categorical stockpersonship measures at the 25 Waikato beef farms.

Description of assessments used to create these figures were reported by Kaurivi [18].

Stockpersonship measures	Categories and number of farms in each category		
Mis-catch	No mis-catch 18	< 1% of cows mis 4	>1% of cows 3
Hitting	No hitting 18	Few cows hit 4	>10% hit 3
Noise of handlers	No noise 4	Minor audible noise 18	Noisy handlers 3
Noise of Equipment	No noise 9	Minor audible noise 6	Very noisy equipment 10
Dogs noise around the yard	No dogs around 7	Quiet dogs 8	Noisy dogs 10
*Health checks (pregnant)	Daily inspection 11	Once or twice a week 9	Less than weekly 5
*Yarding frequency	>4 times/year 5	3-4 times/year 20	Two times or less 0
Yard (design) flow of cows	Effective 13	Minor problems 7	Significant problems 5

\*Measures selected as having high potential effects on reproductive KPI

### 3.3. Prediction of welfare effects on reproduction variables

See Appendix A for descriptive statistics of measures selected for inclusion in the prediction of welfare effects on reproductive performance for each of the 25 farms in Waikato.

The seven welfare measures subjected to PCA, showed many coefficients of 0.3 and above. The Keiser-Meyer-Olkin (KMO) of 0.058 was acceptable threshold and Bartlett's tests reached a statistical significance of <0.001. The PCA revealed presence of three components with an eigenvalue of >1, explaining 73.9% of variance with 15%, 19.9% and 39% of the variances respectively. An inspection of the scree-plot revealed a break after Component 2, which was retained for further analysis. The highest extraction was for thin cows (0.929) and lowest for yarding frequency (0.265). The two components explained cumulative variance of 58.9% with 39% in Component 1 and 19.9% in Component 2. Component 1 loaded for thin, poor rumen fill and blindness and Component 2 was for mortality, yarding frequency/year and dirtiness (Figure 1). The rotated solution revealed the presence of a simple structure with components showing loadings and all variables loading on only one component, except health checks that loaded on both. The interpretation of the components shows a relationship between variables in each specific component. There was a slight positive correlation between the two factors  $r = 0.151$ . The results of this analysis showed a strong association between health-related measures (blindness, lameness), but mortality showed a weaker association. Health checks and yarding showed an association, although yarding frequency associated more with the health and feeding measures. Dirtiness showed a negative association with health and feeding measures. There was a strong correlation between thin cows and cows with poor rumen fill ( $r = 0.714$ ) as well as a strong correlation between thin cows and blindness ( $r = 0.892$ ). PD17 and PD18 were negatively correlated ( $r = -0.217$ ).


Linear regression modelling showed:

*PD18 dependent variable:* Mating period and lameness were the variables remaining at the end of the selection process. At a constant mating period, the coefficient showed that an increase of 1% in lameness was associated with an increase of 0.89% in PD18% (95% CI). At a constant lameness rate,

the increase of mating period by one day decreased PD18 by 0.24% (95% CI). A 1% increase in lameness was associated with 4 days longer mating period. Further analysis with Cooks Distance removed one influential farm that greatly affected the slope of the regression line. For PD18, lameness and mating period then became confounders.

*PD17 dependent variable:* Mating period was the only variable remaining at the end of the selection process. The coefficient indicated that removing lameness changed mating period from 0.13 to 0.08, a change of ~40% (95% CI), indicating that lameness was a confounder. So, a one day increase in mating period (before 2018 PDs) was associated with an increase of 1% in PD17. Having accounted for lameness prevalence, the effect of mating period on PD17 was 1.4% (95% CI). Having accounted for mating period, lameness reduced PD17 (but the data was compatible with a large negative impact -1.3 days per percent rise and a small positive benefit 0.24 for a 1% rise). There was no influential farm in PD17, and Cooks Distance was not analysed.

*Weaning 17 dependent variable:* Measures in Component 1 (thin and poor rumen fill cows, and blindness) were the variables remaining at the end of the selection process. For an increase of one unit ( $\equiv 1$  SD) in component score, weaning increased up by 3.5%. The weaning rate was higher with an increase in these factors but was negatively correlated with health checks.



**Figure 1. PCA: Components showing animal welfare measures rotation**

#### 4. Discussion

The current study postulated that animal welfare influences the reproductive performance of beef farms managed extensively on pasture. Of the many reproductive KPIs that can potentially be used to assess the performance of beef cattle, it was difficult to standardise these across the 25 farms in the present study because of incomplete record keeping. The most consistently available KPIs were pregnancy rates, weaning rates, bull: cow ratio and duration of the mating period. Moreover, the reproductive KPIs that are relevant and assessable within extensive, pastoral beef cow-calf systems differ markedly from those that are feasible in housed/feedlot systems [3]. Thus, in the latter systems, data such as calving intervals, calf to cow ratio per year, number of inseminations per conception, mean pregnancy to cow ration, age at culling and cattle culling % are readily available [12]; but none of these could be ascertained or were pertinent to bull-bred cows in extensive pastoral systems. Thus, direct comparison of reproductive performance and welfare effects with that study is not particularly meaningful, even though its overall conclusion that good welfare standards are related to improved reproductive performance in cattle was not clearly reinforced by the present results. It has been argued that welfare data should be compiled into an 'overall' score[22], but, on the other hand, others [23, 24] have rejected such compilation of scores as being complex and covering up important welfare

problems. Rather, the influence of welfare on reproductive performance was assessed through analysis of individual welfare measures and not as an overall farm score.

Average pregnancy rates (PD17: 91.9% and PD18: 91.3%) were consistent with other studies in New Zealand (91%: [7], 90.8-91.0%: [5], but were below the target of 95% set by the beef industry [6]. In terms of relationships with welfare measures, the present study found no relationship between thinness and poor rumen fill and pregnancy rate. However, Morris [25] found that cows with higher BCS at mating have better pregnancy rates than thinner cows, whilst experimentally, Weik [5] have shown that ideal body weight (e.g. 7 on 1-10 scale) at mating is associated with an improvement of pregnancy rates. Furthermore, access to adequate feed and water is widely recognised as influencing both the health and welfare [26] and productivity and fertility of cows. [25, 27]. The current findings probably mean that the nutritional status of cows was not sufficiently compromised to have had a negative impact on fertility: as also suggested by Probert [28]; Pleasants and Barton [29] and Hickson [30]. Indeed, in the extensive pasture-based systems of beef cattle in New Zealand, nutritional problems are only sufficiently severe to affect reproductive performance of cattle during drought periods [6, 28]. On the other hand, cattle in New Zealand are prone to copper deficiency (which contributes to lowering fertility and weaning: [6, 31], which may have confounded the relationship between BCS and fertility. Proving the micro-nutrient status of the cows might help to resolve this conundrum (see [32, 33]).

Lameness is a critical welfare compromise indicator, as it is both a painful condition and affects productivity and fertility [34, 35]. Increasing literature is mounting that suggest that lameness negatively impact on reproductive performance of cows. For example, various studies have shown that lame beef cows have an increased service period, increased calving to conception interval, increased number of services per conception, impaired follicular growth, ovulation and oestrous behaviour and, hence, less chance of getting pregnant [35-38]. In the present study, lameness was associated with a longer (4 day) mating period. It was also a confounder for mating period when PD17 was the predictive variable: i.e. indicating that lameness was associated with reduced PD17. Paradoxically, however, farms with high levels of lameness and long mating periods had high PD18 and vice versa. Taken together, these results suggest that, in herds with low levels of lameness, long mating periods indicate low pressure for pregnancy; whereas in herds with high levels of lameness, long mating periods reflect a response to try and get as many cows pregnant as possible.

Other welfare measures were unrelated or only weakly related to pregnancy rate. Dirtiness was unrelated, as, in the present study, it probably was only an indicator of the lushness of the pasture [18, 39]. This contrasts with the situation with housed cattle, where dirtiness is related to both risks of disease and of failure to conceive ([40]. Likewise, measures of stockpersonship, such as yarding frequency, cows' behaviour in the yard were unrelated in the present study to any reproductive parameters. Interestingly, the frequency of yarding was related to the incidence of fearful behaviour (i.e. cattle attempting to escape the race or climbing/pushing on others), as also previously recorded [41, 42]. Others have shown that fearful/agitated behaviours in the yards were associated with reduced fertility [11, 43], possibly due to stress-induced increases in plasma cortisol concentrations [44], so the lack of an association in the present study was not expected. However, it appears that, in the present study, the commonness of fearful/agitated behaviour was just a temporary indication of the lack of familiarity of extensively managed cattle with yarding and handling, and not necessarily a permanent situation.

The average weaning rate (proportion of weaned calves from mated cows) was 83.9%, which is below the industry target of 90% [6, 45]. However, [45] indicated a very similar figure (80-84%) in a survey of cow-calf herds in New Zealand. However, the proportion of calves weaned from recorded pregnant cows (i.e. rather than mated cows) was 92%. If the number of pregnant cows is used as a proxy for expected calves to be born, it indicates an effect of lower conception (~6% than calf survival to weaning). Thus, remedial strategies geared towards improving conception rate might be more pertinent to increase reproductive performance in the national cow-calf herds, than focusing on ensuring calf survival. There was, however, a paradoxical relationship between welfare measures and weaning rate in the present study, inasmuch as weaning rate positively related to an increase in

thinness, poor rumen fill and blindness in cows, but negatively to health checks. Most other reports e.g. [46–48], show the opposite: i.e. that good nutrition and health are associated with improved pregnancy and weaning rates. Whether this discrepancy can be attributed to peri-partum or pre-weaning mortality is a moot point, but the present results may merely indicate that the mortality rate (average 3.9%) was not high enough to influence reproductive performance.

The average duration of the mating period (62 days) was closely aligned with the industry standard of 63 days [45]. The target is to at least have 60% of cows calving in the first 21 days of calving in order to avoid a spread-out calving and to indicate a high reproductive performance of a herd [6]. Mating period has also been shown to be related to pregnancy rates and the incidence of post-partum anoestrus [7]. Ideally, the interval from calving to conception would have also been measured, as it has been indicated that this could be an indicator of animal welfare [49], but this is not systematically recorded in extensive beef systems (as also found by McFadden [7]). On the other hand, there is evidence that calving to conception interval may not be a particularly useful welfare indicator, inasmuch as even in dairy cows (in which this is routinely recorded), it is disregarded as a routine performance indicator for animal welfare [50].

The median bull: cow ratio was 1:30. Previous studies of reproduction management plans in New Zealand show that bulls in natural breeding systems can mate with 30–40 cows [6]. Whilst there was no relationship between bull: cow ratio in the present study, it was clear that farmers tended to reduce the number of cows per bull to reduce the risk of pregnancy failure in the herd (as also found by McFadden [7]). However, over the range of ratios of 1:25 to 1:50, the literature shows only a tenuous relationship between bull: cow ratio and pregnancy rate (e.g. [51]). Thus, the 1:30 ratio observed in the present study is probably a reflection of current recommendations regarding maximising the dissemination of ‘superior’ genetics ([6]) through optimising bull: cow ratio. Farmers also reported to purchasing bulls certified from reputable breeders to limit the use of low performing bulls, although bulls were not routinely tested before mating.

The average prevalence of dystocia in the present study was 2.6%; half the industry standard of 5% [45]. Dystocia has a very significant impact upon the welfare of affected animals (and upon calves born/stillborn as a result of dystocia), and, at high incidences, can markedly impair the productivity of the farm [30, 52]. Hence, our previous study [18] recommended a threshold of > 2% for dystocia as indicative of poor welfare that needs immediate intervention. However, the present study did not show a relationship between the prevalence of dystocia and reproduction outcomes. Interestingly, the prevalence of dystocia was unrelated to BCS, as also shown for pastoral beef cows by Hickson [53]. The prevalence of abortion was also low (1.5%), well below the 3% industry target [45] and was unrelated to any other parameters of reproductive performance. High incidences of abortion are, in extensive beef systems, usually related to diseases such as BVD infection and leptospirosis [54]. No individual herd in the present study had a high prevalence of abortion, and only one herd in this study reported abortion losses due to BVD, others were in individual cows due to unknown causes.

This was a relatively small study in terms of the number of herds enrolled, although very extensive welfare data were gathered from each herd. However, the results suggest no clear impact of welfare at a whole-of-study level, and only an individual farm association between animal welfare measures and reproductive performance. It was evident that variations may occur in farm reproductive performance across farms, as showed in the negatively correlated pregnancy rates in two consecutive years. The animal welfare data were also limited to one-year and further studies that consider multiple farms and enterprise goals are warranted to more fully evaluate the relationship between welfare standards and reproductive performance. Although not many KPI were feasible in the study, the study reinforced that routine pregnancy testing is critical to identify infertile cows and heifers in order to make appropriate management decision, such as culling of poor fertile cows. Hence, pregnancy rates are considered good indicators of cows’ reproductive performance in the country [6] and, coupled to weaning rate, these factors give an overall indication of the reproduction performance in beef herds.

## 5. Conclusions

The study embarked on investigating the relationship between animal welfare and the reproductive performance of extensive pasture-based beef cows in New Zealand. The use and analysis of a newly developed animal welfare assessment protocol [17, 18] for New Zealand cow and calf systems showed that seven welfare measures (thinness, poor rumen fill, dirtiness blindness, mortality, yarding frequency/year and health checks frequency of pregnant cows) may potentially influence reproductive KPI. Lameness was also retained to the potential measures. The key findings were that feasible KPI across the cow-calf farms were pregnancy and weaning rates, which coincided with results from previous studies. From the welfare measures, it was only lameness that showed a direct effect upon pregnancy rate, as well as showing a confounding increase in pregnancy rates with an increase in the mating period. The bull: cow ration and reproductive conditions (dystocia, abortion, vaginal prolapse) did not influence pregnancy and weaning rates. Overall, the study showed no clear evidence of an effect of animal welfare standards on the reproductive performance in the study population. Further research on multiple beef farms is warranted to clarify these findings. The integrated approach on how animal welfare directly or indirectly affects the reproductive performance of beef cattle in the extensive production system could be beneficial to address mitigation or intervention strategies for the improvement of reproductive efficiency in beef cattle.

**Author Contributions:** Y.B.K.—main researcher who did the project proposal and data collection, analysis and main write up. R.L.—main supervisor who did conceptualization, methodology, funding acquisition, formal analysis, review and editing. R.H.—conceptualization and editing. K.S.—conceptualization, methodology and editing and validation. T.P.—review and editing. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding but funded from Massey University IVABS.

**Acknowledgments:** The authors gratefully acknowledge farmers for participating in this study. We thank Dean Burman for technical support; VetEnt Waikato veterinary clinic, especially to Veterinarians Ben Hodgson and Will Cuttance for technical support on their scheduled farms visit for pregnancy scanning, and for Liz Dunn and Mark Oakes for technical support and organizing the farms. Finally, we acknowledge Prof Christopher Winckler for the original thought of combining the Welfare Quality protocol with the UC Davis Cow-Calf Health and Handling protocol.

**Conflicts of Interest:** The authors declare no conflict of interest.

## Appendix A

Descriptive statistics of measures selected for inclusion in the prediction of welfare effects on reproductive performance at the 25 Waikato co-calf farms

FARM	Cow		Cow		Wean	Vaginal			Bull: Cow	Mating period	Thin cows	Poor			Mortality rate	Yarding/ year	Health Checks	
	mated		mated			prolapse	Dystocia	Abortion				rumen fill	Dirty-ness	Blind-ness	Lame-ness			
	18	PD18	17	PD17		17												
F1	253	89.3	248	84.7	76.8	0	0	0	62	50	5.7	17	17.8	0	1.1	0.4	1	0
F2	410	92	539	90.7	81.6	0.2	0.4	0	39	62	5.7	13.9	28.7	0	0	0.8	1	0
F3	33	88	43	81.4	74.4	2.9	2.9	0	22	46	0	0	5.3	0	0	6.3	1	1
F4	97	82.5	100	100	76	0	0	0	33	69	17.9	26.9	13.4	0	0	5.7	1	2
F5	123	92.7	85	94.1	85.9	0	3.8	0	28	100	6.4	43.6	20.1	0	11.5	2	1	2
F6	212	96.5	369	88.9	82.7	0	1.8	0.6	31	30	0	4.9	11.9	0	0.6	6.8	1	2
F7	18	83	12	100	83.3	0	16.7	0	12	87	0	0	11.1	0	5.6	3.6	0	0
F8	318	91.8	338	91.4	85.1	0	0.3	0	38	62	1.8	21.3	20.6	0	1.1	3	0	0
F9	305	95.5	378	89.2	75.4	0	1.3	0.9	38	64	5.9	31.8	10.3	0	1.3	2.6	1	0
F10	241	90	170	91.8	88.8	0	1.9	0.6	21	45	4.4	25.6	41.7	0	2.2	5	1	2
F11	103	90.3	130	95.4	93.8	0.8	0.8	0.8	22	61	57.4	68.3	5	3	2	10.4	1	0
F12	154	92.9	134	91	85.1	0.8	2.5	0	34	60	5.3	23.3	39.1	0	4.5	1.1	0	1
F13	145	93.8	214	85	67.8	0	2.6	0	36	45	8	37.2	15.3	0	1.8	4.4	1	2
F14	162	94.4	253	96	92.9	0	4.1	0	25	62	6	18	32.3	0.8	0.8	1.3	1	1
F15	96	97.9	120	95	83.3	0	3.5	0	40	66	0	0	27.9	0	3.3	2.4	1	0
F16	30	69.2	29	96.6	96.6	0	0	3.6	29	125	60.7	67.9	7.1	3.6	3.6	2.7	1	0
F17	81	95.1	60	98.3	96.7	0	1.7	0	30	62	1.9	7.7	30.1	0	0	4.4	1	0
F18	293	95.2	294	92.9	90.1	0	5.5	1.8	29	82	5.2	47.7	6	0	10.3	2.2	1	0
F19	98	91.8	94	85.1	70.2	0	5	8.8	24	90	3.2	32.3	22.6	0	8.1	8.1	1	1
F20	464	89.9	432	92.6	83.3	0	2.5	4.8	36	59	3.2	34.4	35.7	0	0.3	5.8	1	1
F21	174	96	142	98.6	94.4	0	0.7	0.7	28	37	8.9	26.6	4	0	0.8	2.5	1	1

F22	232	94	230	94.3	87	0.9	0	2.3	33	50	27	54	25.3	0	1.5	4.7	1	1
F23	541	92.6	485	84.9	82.3	0	1.5	1.7	40	41	3.3	36.2	26.1	0	2.3	2.1	0	1
F24	306	94.1	265	85.3	77.4	0	2.2	8.8	29	31	11.1	60.8	25	0	3.5	6.1	0	1
F25	67	95.5	70	94.3	88.6	0	4.5	1.5	23	60	17.5	66.7	50.3	1.8	0	2.5	1	0
Sum	4956	2284	5234	2298	2100	5.6	66.2	36.9	782	1546	266.5	766.1	532.7	9.2	66.2	96.9		
Aver	198.2	91.4	209.4	91.9	84.0	0.2	2.6	1.5	31.3	61.8	10.7	30.6	21.3	0.4	2.7	3.9	0: >4	0: daily
Min	18	69.2	12	81.4	67.8	0	0	0	12	30	0	0	4	0	0	0.4	times	1: 2x weekly
Max	541	97.9	539	100	96.7	2.9	16.7	8.8	62	125	60.7	68.3	50.3	3.6	11.5	10.4	1: 3-4	2: < weekly
Medn	162	92.7	170	92.6	83.3	0	1.9	0.6	30	61	5.7	26.9	20.6	1.5	3	2.3		
Stdev	135.2	5.8	145.1	5.1	7.7	0.6	3.3	2.5	9.3	21.4	15.6	20.9	12.4	0.9	3.1	2.4		

## References

1. Burns, B. M.; Fordyce, G.; Holroyd, R. G. A review of factors that impact on the capacity of beef cattle females to conceive, maintain a pregnancy and wean a calf-Implications for reproductive efficiency in northern Australia. *Anim Reprod Sci.* **2010**, *122*, 1-22.
2. Caldow, G.; Hill, G.; Logan, R; Pritchard, I., A Guide to Improving Suckler Herd Fertility. Manuscript ed.; (QMS), Q. M. S., Ed. <http://www.qmsscotland.co.uk/breeding>; Ingliston, Newbridge, Scotland, **2016**; pp 1-40.
3. Hewitt, S.; Green, M.; Hudson, C. Evaluation of key performance indicators to monitor performance in beef herds. *Livestock.* **2018**, *23*, 72-78.
4. Bova, T. L.; Chiavaccini, L.; Cline, G. F.; Hart, C. G.; Matheny, K.; Muth, A. M., . . . Memili, E. Environmental stressors influencing hormones and systems physiology in cattle. *Reprod Biol Endocrinol.* **2014**, *12*, 1.
5. Weik, F.; Archer, J.; Morris, S.; Garrick, D.; Hickson, R. Relationship between body condition score and pregnancy rates following artificial insemination and subsequent natural mating in beef cows on commercial farms in New Zealand. *New Zealand Journal of Animal Science and Production* **2020**, *80*.
6. Geenty, K.; Morris, S. T., *Guide to New Zealand Cattle Farming*. Beef and Lamb New Zealand: Wellington, **2017**; p 1-129.
7. McFadden, A. M.; Heuer, C.; Jackson, R.; West, D. M.; Parkinson, T. J. Reproductive performance of beef cow herds in New Zealand. *N Z Vet J.* **2005**, *53*, 39-44.
8. Hickson, R. E.; Laven, R. L.; Lopez-Villalobos, N.; Kenyon, P. R.; Morris, S. T. Postpartum anoestrous interval in first-lactation beef and dairy-beef crossbred cows. *Animal Production Science.* **2012**, *52*, 478-482.
9. Morris, S. T.; Kenyon, P. R. Intensive sheep and beef production from pasture - A New Zealand perspective of concerns, opportunities and challenges. *Meat Science.* **2014**, *98*, 330-335.
10. Kramer, M.; Erbe, M.; Bapst, B.; Bieber, A.; Simianer, H. Estimation of genetic parameters for novel functional traits in Brown Swiss cattle. *J Dairy Sci.* **2013**, *96*, 5954-5964.
11. Cziszter, L. T.; Gavojdian, D.; Neamt, R.; Neciu, F.; Kusza, S.; Ilie, D. E. Effects of temperament on production and reproductive performances in Simmental dual-purpose cows. *Journal of Veterinary Behavior: Clinical Applications and Research.* **2016**, *1550-55*.
12. Irizo, L.; Tomassone, L.; Martano, G.; Gottardo, F.; Tarantola, M. Animal welfare and reproductive performance in two Piemontese housing systems. *Italian Journal of Animal Science.* **2018**, *17*, 499-504.
13. Castellini, C. Reproductive activity and welfare of rabbit does. *Italian Journal of Animal Science.* **2007**, *6*(SUPPL. 1), 743-747.
14. Einarsson, S.; Sjunnesson, Y.; Hultén, F.; Eliasson-Selling, L.; Dalin, A. M.; Lundeheim, N.; Magnusson, U. A 25 years experience of group-housed sows-reproduction in animal welfare-friendly systems. *Acta Vet Scand.* **2014**, *56*. (1).
15. Knox, R.; Salak-Johnson, J.; Hopgood, M.; Greiner, L.; Connor, J. Effect of day of mixing gestating sows on measures of reproductive performance and animal welfare. *J Anim Sci.* **2014**, *92*, 1698-1707.
16. OIE, Animal Welfare and Beef Cattle Production Systems. In *World Organisation for Animal Health(OIE) Terrestrial Animal Health Code*, 26 ed.; OIE: [http://www.oie.int/fileadmin/Home/eng/Health\\_standards/tahc/current/chapitre\\_aw\\_beef\\_catthe.pdf](http://www.oie.int/fileadmin/Home/eng/Health_standards/tahc/current/chapitre_aw_beef_catthe.pdf), **2017**; Vol. 1 Chapter 7.9.
17. Kaurivi, Y. B.; Laven, R.; Hickson, R.; Parkinson, T.; Stafford, K. Developing an Animal Welfare Assessment Protocol for Cows in Extensive Beef Cow-Calf Systems in New Zealand. Part 1: Assessing the Feasibility of Identified Animal Welfare Assessment Measures. *Animals.* **2020a**, *10*, 1597.

18. Kaurivi, Y. B.; Hickson, R.; Laven, R.; Parkinson, T.; Stafford, K. Developing an Animal Welfare Assessment Protocol for Cows in Extensive Beef Cow-Calf Systems in New Zealand. Part 2: Categorisation and Scoring of Welfare Assessment Measures. *Animals*. **2020**, *10*, 1592.
19. Welfare Quality, Welfare Quality Assessment Protocol for Cattle. In *Welfare Quality Assessment Protocol for Cattle (without Veal Calves)*, Welfare Quality®: Lelystad, Netherlands., **2009**; pp 1-142.
20. UC Davis University of California, Davis Cow-Calf Health and Handling Assessment. Available online <https://www.ucdcowcalfassessment.com/> (accessed on 12 August 2017).
21. Tabachnick, B. G.; Fidell, L. S.; Ullman, J. B., *Using multivariate statistics*. Pearson Boston, MA: **2007**; Vol. 5.
22. Botreau, R.; Bracke, M. B. M.; Perny, P.; Butterworth, A.; Capdeville, J.; Van Reenen, C. G.; Veissier, I. Aggregation of measures to produce an overall assessment of animal welfare. Part 2: analysis of constraints. *Animals*. **2007**, *1*, 1188-1197.
23. Mellor, D. Operational Details of the Five Domains Model and Its Key Applications to the Assessment and Management of Animal Welfare. *Animals*. **2017**, *7*, 60.
24. Sandøe, P.; Corr, S. A.; Lund, T. B.; Forkman, B. Aggregating animal welfare indicators: can it be done in a transparent and ethically robust way? *Anim Welf*. **2019**, *28*, 67-76.
25. Morris, S. T.; Morel, P. C. H.; Kenyon, P. R. The effect of individual liveweight and condition of beef cows on their reproductive performance and birth and weaning weights of calves. *N Z Vet J*. **2006**, *54*, 96-100.
26. Fraser, D.; Duncan, I. J. H.; Edwards, S. A.; Grandin, T.; Gregory, N. G.; Guyonnet, V., . . . Whay, H. R. General Principles for the welfare of animals in production systems: The underlying science and its application. *Vet J*. **2013**, *198*, 19-27.
27. Tait, I. M.; Morris, S. T.; Kenyon, P. R.; Garrick, D. J.; Pleasants, A. B.; Hickson, R. E., Effect of cow body condition score on inter-calving interval, pregnancy diagnosis, weaning rate and calf weaning weight in beef cattle. In *Proceedings of the New Zealand Society of Animal Production*, Rotorua, **2017**, *77*, pp 23-28.
28. Probert, A. D. A Study of aspects of beef cattle reproduction in the Manawatu district. Thesis, **1989**.
29. Pleasants, A. B.; Barton, R. A. In *Precalving nutrition of heavy two year old Angus heifers weighing 415 kg at calving*, Proceedings of the New Zealand Society of Animal Production, New Zealand, 1992; New Zealand, pp 303-5.
30. Hickson, R. E.; Morris, S. T.; Kenyon, P. R.; Lopez-Villalobos, N. Dystocia in beef heifers: A review of genetic and nutritional influences. *N Z Vet J*. **2006**, *54*, 256-264.
31. Morris, C. A. Managing mineral deficiencies in grazing livestock. *N Z J Agric Res*. **2011**, *54*, 353-354.
32. Enjalbert, F.; Lebreton, P.; Salat, O. Effects of copper, zinc and selenium status on performance and health in commercial dairy and beef herds: Retrospective study. *J Anim Physiol Anim Nutr (Berl)*. **2006**, *90*, 459-66.
33. Neville, G. D.; Knowles, S. O. Trace Element Supplementation of Livestock in New Zealand: Meeting the Challenges of Free-Range Grazing Systems. *Vet. Med. Int.* **2012**, 2012639472.
34. Lawrence, K. E.; Chesterton, R. N.; Laven, R. A. Further investigation of lameness in cows at pasture: An analysis of the lesions found in, and some possible risk factors associated with, lame New Zealand dairy cattle requiring veterinary treatment. *J Dairy Sci*. **2011**, *94*, 2794-2805.
35. Tyagi, K.; Lathwal, S.; Sharma, J.; Devi, I.; Gupta, R.; Patbandha, T.; Tewari, H. Lameness in crossbred cows: Its effect on productive and reproductive performance. *IJDS*. **2017**, *70*, 443-446.
36. Morris, M. J.; Kaneko, K.; Walker, S. L.; Jones, D. N.; Routly, J. E.; Smith, R. F.; Dobson, H. Influence of lameness on follicular growth, ovulation, reproductive hormone concentrations and estrus behavior in dairy cows. *Theriogenology*. **2011**, *76*, 658-668.

37. Somers, J. R.; Huxley, J.; Lorenz, I.; Doherty, M. L.; O'Grady, L. The effect of lameness before and during the breeding season on fertility in 10 pasture-based Irish dairy herds. *Ir Vet J.* **2015**, *68*, 14.

38. Ratanapob, N.; Thiangtum, W.; Rukkwamsuk, T.; Srisomrun, S.; Panneum, S.; Arunvipas, P. The relationship between lameness and reproductive performance in dairy cows raised in small holder farms, Thailand. *Songklanakarin J Sci Technol.* **2020**, *42*, 766-770.

39. Laven, R. A.; Fabian, J. Applying animal-based welfare assessments on New Zealand dairy farms: feasibility and a comparison with United Kingdom data. *N Z Vet J.* **2016**, *64*, 212-217.

40. Kara, N. K.; Galic, A.; Koyuncu, M. Effects of stall type and bedding materials on lameness and hygiene score and effect of lameness on some reproductive problems in dairy cattle. *J Appl Anim Res.* **2011**, *39*, 334-338.

41. Francisco, C. L.; Cooke, R. F.; Marques, R. S.; Mills, R. R.; Bohnert, D. W. Effects of temperament and acclimation to handling on feedlot performance of *bos taurus* feeder cattle originated from a rangeland-based cow-calf system. *J Anim Sci.* **2012**, *90*, 5067-5077.

42. Simon, G. E.; Hoar, B. R.; Tucker, C. B. Assessing cow-calf welfare. Part 1: Benchmarking beef cow health and behavior, handling, and management, facilities, and producer perspectives. *J Anim Sci.* **2016**, *94*, 3476-3487.

43. Rueda, P. M.; Sant'Anna, A. C.; Valente, T. S.; Paranhos da Costa, M. J. R. Impact of the temperament of Nellore cows on the quality of handling and pregnancy rates in fixed-time artificial insemination. *Livestock Science.* **2015**, *177*189-195.

44. Cooke, R. F.; Schubach, K. M.; Marques, R. S.; Peres, R. F. G.; Silva, L. G. T.; Carvalho, R. S., . . . Vasconcelos, J. L. M. Effects of temperament on physiological, productive, and reproductive responses in *Bos indicus* beef cows. *J Anim Sci.* **2017**, *95*, 1-8.

45. Morris, S. T.; Smeaton, D. C., Reproduction in the beef cow herd In *Profitable farming of beef cows- Beef + Lamb New Zealand*, Wells, D., Ed. Fusion Print Group Limited, NZ Beef Council: Hamilton, **2009**; pp 42-69.

46. Stalker, L. A.; Adams, D. C.; Klopfenstein, T. J.; Feuz, D. M.; Funston, R. N. Effects of pre- and postpartum nutrition on reproduction in spring calving cows and calf feedlot performance. *J Anim Sci.* **2006**, *84*, 2582-2589.

47. Diskin, M. G.; Kenny, D. A. Optimising reproductive performance of beef cows and replacement heifers. *Animal.* **2014**, *8 Suppl* 127-39.

48. D'Occhio, M. J.; Baruselli, P. S.; Campanile, G. Influence of nutrition, body condition, and metabolic status on reproduction in female beef cattle: A review. *Theriogenology.* **2019**, *125*277-284.

49. De Vries, M.; Bokkers, E. A. M.; Dijkstra, T.; van Schaik, G.; de Boer, I. J. M. Invited review: Associations between variables of routine herd data and dairy cattle welfare indicators. *J Dairy Sci.* **2011**, *94*, 3213-3228.

50. Brownlie, T. S.; Morton, J. M.; Heuer, C.; Hunnam, J.; McDougall, S. Reproductive performance of seasonal-calving, pasture-based dairy herds in four regions of New Zealand. *N Z Vet J.* **2014**, *62*, 77-86.

51. Neville, W. E., Jr.; Richardson, K. L.; Utley, P. R. Breeding Performance of Bulls Assigned to 40 or 50 Cows per Bull During the Breeding Period1. *J Anim Sci.* **1988**, *66*, 613-617.

52. Mason, W. A.; Cuttance, E. L.; Laven, R. A.; Phyn, C. V. C. Short communication: Replacement heifer mortality from weaning until second mating in seasonal-calving, pasture-based dairy herds in New Zealand. *J Dairy Sci.* **2020**, *103*, 902-908.

53. Hickson, R. E.; Kenyon, P. R.; Lopez-Villalobos, N.; Morris, S. T. Effects of liveweight gain during pregnancy of 15-month-old Angus heifers on dystocia and birth weight, body dimensions, estimated milk intake and weaning weight of the calves. *N Z J Agric Res.* **2008**, *51*, 171-180.

54. Sanhueza, J. M.; Heuer, C.; West, D. Contribution of *Leptospira*, *Neospora caninum* and bovine viral diarrhea virus to fetal loss of beef cattle in New Zealand. *Prev Vet Med.* **2013**, *112*, 90-98.