

Article

Not peer-reviewed version

Valproic Acid Releases Ca^{2+} from InsP_3 -Sensitive Ca^{2+} Stores

Ana Ruiz-Nuño and [María Francisca Cano-Abad](#)*

Posted Date: 12 December 2025

doi: 10.20944/preprints202512.1137.v1

Keywords: valproic acid; InsP_3 receptor activation; endoplasmic reticulum Ca^{2+} release; intracellular calcium signaling; cytosolic Ca^{2+} dynamics; ER calcium homeostasis; aequorin; fura-2



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Valproic Acid Releases Ca^{2+} from InsP_3 -Sensitive Ca^{2+} Stores

Ana Ruiz-Nuño ¹ and María F. Cano-Abad ^{1,2,*}

¹ Departamento de Farmacología, Facultad de Medicina. Universidad Autónoma de Madrid. Spain

² Instituto de Investigación Biomédica. Hospital Universitario de la Princesa. Spain

* Correspondence: maria.cano@uam.es; Tel.: +34-91-497-2766

Highlights

What this study is about

1. The study explores how the antiepileptic drug valproic acid (VPA) affects intracellular calcium (Ca^{2+}) homeostasis within the endoplasmic reticulum (ER).
2. Using aequorin-based Ca^{2+} imaging, the research examines how VPA modulates Ca^{2+} fluxes and their potential impact on neurotransmission regulation.

What the study found

1. VPA induces Ca^{2+} release from the ER in a concentration-dependent manner ($\text{IC}_{50} \approx 17 \mu\text{M}$), mimicking the inositol 1,4,5-trisphosphate (InsP_3)-triggered release.
2. Blocking InsP_3 receptors with heparin or 2-APB abolished VPA-evoked Ca^{2+} release, identifying InsP_3 -sensitive ER stores as its target.
3. In neuronal-like cells, VPA enhanced neurotransmitter (catecholamine) release, linking its intracellular Ca^{2+} effects to neurosecretory modulation.
4. This is the first evidence of an intracellular site of action for VPA, through activation of InsP_3 -sensitive Ca^{2+} stores.

Abstract

Calcium signaling dysfunction is a central contributor to neuronal hyperexcitability and seizure propagation in epilepsy, yet the intracellular mechanisms underlying the actions of valproic acid (VPA) remain incompletely understood. In this study, we investigated whether VPA modulates calcium homeostasis at the level of the endoplasmic reticulum (ER) and how this action influences cytosolic calcium dynamics associated with epileptiform activity. ER calcium levels were directly measured using ER-targeted aequorin in HeLa and PC12 cells, while cytosolic Ca^{2+} signals were monitored by fura-2 fluorescence imaging in bovine chromaffin cells exposed to veratridine, a model of sustained sodium channel activation and calcium oscillations. VPA induced a concentration-dependent release of Ca^{2+} from the ER, with an IC_{50} of approximately $17 \mu\text{M}$. This effect was preserved in permeabilized cells and exhibited activation kinetics comparable to those elicited by inositol 1,4,5-trisphosphate (InsP_3). Pharmacological inhibition of InsP_3 receptors (InsP_3Rs), but not ryanodine receptors or SERCA, abolished VPA-induced ER Ca^{2+} release, supporting a selective InsP_3R -mediated mechanism. Functionally, VPA suppressed the repetitive cytosolic Ca^{2+} oscillations induced by veratridine, while simultaneously producing a sustained elevation of cytosolic Ca^{2+} originating from ER stores and facilitating depolarization-evoked catecholamine secretion. Together, these results support the conclusion that VPA acts as a novel functional agonist of InsP_3Rs and identify ER Ca^{2+} mobilization as a previously unrecognized intracellular mechanism contributing to its modulatory effects on calcium signaling and excitability in epilepsy.

Keywords: valproic acid; InsP3 receptor activation; endoplasmic reticulum Ca^{2+} release; intracellular calcium signaling; cytosolic Ca^{2+} dynamics; ER calcium homeostasis; aequorin; fura-2

1. Introduction

Epilepsy is a chronic neurological disorder characterized by recurrent seizures arising from abnormal, synchronized neuronal discharges. The pathophysiology of epilepsy involves a complex interplay of excitatory and inhibitory mechanisms in the brain, where calcium ions (Ca^{2+}) play a fundamental regulatory role. Intracellular calcium fluctuations are essential for neuronal excitability, neurotransmitter release, gene expression, and synaptic plasticity processes such as long-term potentiation [1,2]. However, prolonged or excessive increases in intracellular calcium concentration ($[\text{Ca}^{2+}]_i$) can trigger excitotoxicity, mitochondrial dysfunction, and neuronal death—processes implicated in the onset and progression of epilepsy [3–5].

In neurons, Ca^{2+} homeostasis is tightly regulated through fluxes between the cytosol and internal stores, primarily the endoplasmic reticulum (ER). Two families of calcium-release channels coordinate these movements: the inositol 1,4,5-trisphosphate receptor (InsP_3R) and the ryanodine receptor (RyR) [6,7]. InsP_3R -mediated Ca^{2+} release plays a central role in the generation of intracellular Ca^{2+} waves, which influence synaptic activity, dendritic integration, and gene transcription [8]. Perturbations in InsP_3R and RyR function have been linked to epileptogenesis and neuronal hyperexcitability [9]. Despite this, the potential modulation of intracellular Ca^{2+} stores by antiepileptic drugs remains insufficiently explored.

Valproic acid (VPA) is one of the most widely prescribed broad-spectrum antiepileptic drugs, effective against generalized and focal seizures, as well as in bipolar disorder and migraine prophylaxis. Its primary mechanisms of action have traditionally been associated with the potentiation of γ -aminobutyric acid (GABA) neurotransmission [10,11], the modulation of glutamate uptake [12], and the inhibition of voltage-dependent sodium and T-type calcium channels [13–17]. Additional studies have also suggested neuroprotective roles for VPA in reducing ER stress and lipid accumulation [18,21]. Yet, despite over six decades of clinical use, the precise molecular targets that mediate its anticonvulsant and neuroregulatory properties remain incompletely defined.

The ER is increasingly recognized as a key modulator of neuronal Ca^{2+} signaling and excitability [22]. Among the intracellular signaling systems regulating Ca^{2+} dynamics, the inositol 1,4,5-trisphosphate receptor (IP_3R), located on the membrane of the endoplasmic reticulum (ER), plays a pivotal role in mobilizing Ca^{2+} from internal stores to the cytosol and nucleus [22–24]. The ER constitutes the major intracellular Ca^{2+} reservoir, maintaining distinct subcompartments of high and low Ca^{2+} concentrations that are tightly regulated to ensure proper signaling and neuronal viability [25–27].

The activation of IP_3Rs by phospholipase C-derived IP_3 results in finely tuned Ca^{2+} release events, which contribute to the generation of local and global Ca^{2+} waves that control neurotransmitter exocytosis and synaptic efficiency [28–30]. Moreover, the existence of different IP_3R isoforms with tissue-specific distribution and functional diversity further refines this regulation, allowing neurons to adapt their Ca^{2+} signaling patterns to physiological and pathological stimuli [31,32].

Dysregulation of IP_3 -mediated Ca^{2+} release has been implicated in several neuropathological conditions, including excitotoxicity, ER stress, and epileptogenesis. Alterations in ER Ca^{2+} homeostasis can modify synaptic strength and neuronal excitability, thereby contributing to seizure propagation. Understanding how pharmacological agents, particularly antiepileptic drugs, interact with IP_3R -dependent Ca^{2+} release pathways is therefore crucial for elucidating their cellular mechanisms of action and for developing novel therapeutic strategies targeting Ca^{2+} signaling in the nervous system.

Research using genetically encoded aequorin targeted to the ER has demonstrated that VPA induces Ca^{2+} release from InsP_3 -sensitive stores in a concentration-dependent manner, with an IC_{50} of approximately 17 μM , and kinetics strikingly similar to the second messenger InsP_3 [23–26]. Pharmacological blockade of InsP_3R with heparin or 2-aminoethyl diphenylborinate (2-APB) abolishes this effect, supporting the hypothesis that VPA acts as a novel InsP_3R agonist. Moreover, studies in PC12 neuronal-like cells and bovine chromaffin cells revealed that VPA enhances catecholamine release and modulates ER Ca^{2+} signaling without affecting RyR-mediated pathways [25–30]. These intracellular actions suggest that VPA can directly influence synaptic release machinery through modulation of ER Ca^{2+} dynamics.

Nevertheless, the relationship between VPA's intracellular effects and its clinical efficacy remains controversial. While therapeutic concentrations of VPA suppress epileptiform activity in some models [14], paradoxical pro-epileptic effects have been reported in others [31–33], potentially reflecting patient-dependent variations in InsP_3R sensitivity or downstream Ca^{2+} signaling. This duality may contribute to phenomena such as pharmacoresistance or idiosyncratic adverse responses in certain epileptic individuals.

Taken together, these findings highlight the need to further elucidate the intracellular mechanisms underlying VPA's actions. The present study was designed to investigate how VPA modulates Ca^{2+} homeostasis at the level of the ER, focusing on its interaction with InsP_3R and subsequent effects on neurotransmitter release. Understanding these mechanisms could provide a unifying framework linking VPA's antiepileptic efficacy, its side-effect profile, and the cellular basis of variable patient responses. As far as we know, VPA has not been found yet to be implicated directly in the regulation of Ca^{2+} homeostasis at the ER level. We have therefore approached such a study here. Our findings suggest that VPA caused the release of Ca^{2+} from intracellular stores through a mechanism reminding of InsP_3 .

2. Materials and Methods

HeLa Cell Culture and Transfection

HeLa cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% fetal calf serum (FCS). For transfection, cells were seeded onto 13 mm glass coverslips and grown to 60–70% confluence. Transfection was performed using 4 μg of plasmid DNA encoding the genetically engineered photoprotein aequorin. The mutated aequorin with low Ca^{2+} affinity targeted to the endoplasmic reticulum (erAEQ) was employed, as described previously [34]. Transfection was achieved using the calcium phosphate method [35]. Experiments aimed at measuring changes in endoplasmic reticulum calcium concentration ($[\text{Ca}^{2+}]_{\text{ER}}$) were carried out 36 h post-transfection.

PC12 Cell Culture and Transfection

PC12 cells were maintained in DMEM supplemented with 7.5% fetal calf serum, 7.5% horse serum, 2 mM glutamine, 25 U/mL penicillin, and 25 $\mu\text{g}/\text{mL}$ streptomycin. Cells were seeded on 13 mm poly-L-lysine-coated glass coverslips in 24-well plates and allowed to reach 60–70% confluence after 24 h at 37 °C in a humidified 5% CO_2 atmosphere. Transfection with the erAEQ plasmid was achieved using Metafectene (Biontex Laboratories) [36]. Measurements of $[\text{Ca}^{2+}]_{\text{ER}}$ were performed 36–48 h after transfection.

Bovine Adrenal Chromaffin Cell Culture

Bovine chromaffin cells (BCCs) were isolated according to standard procedures with minor [37]. Cells were suspended in DMEM containing 5% FCS, 50 IU/mL penicillin, and 50 $\mu\text{g}/\text{mL}$ streptomycin. For secretion experiments, 5×10^6 cells were plated in 5 cm Petri dishes and maintained at 37 °C in a 5% $\text{CO}_2/95\%$ air atmosphere. Cells were used between 1 and 5 days after plating.

Measurement of $[\text{Ca}^{2+}]_{\text{ER}}$ Changes with Aequorin

Two experimental conditions were used:

(a) Intact cells: The monolayer was superfused with Krebs–Hepes buffer for HeLa (KHBH) containing (in mM): 125 NaCl, 5 KCl, 1 Na₃PO₄, 1 MgSO₄, 5.5 glucose, and 20 HEPES, pH 7.4, at room temperature (24 ± 2 °C), supplemented with 1 mM CaCl₂.

(b) Permeabilized cells: An intracellular-like buffer (IB) was used containing (in mM): 140 KCl, 10 NaCl, 1 K₃PO₄, 10 HEPES, 1 MgCl₂, 1 ATP, 5 succinate, and 20 μM ADP, pH 7.0, supplemented with 0.5 μM CaCl₂.

Reconstitution of ER-targeted aequorin was achieved by incubating cells for 1–2 h in KHBH or KHBPC12 supplemented with 5 μM coelenterazine n, 5 μM ionomycin, and 600 μM EGTA. After loading, cells were washed with buffer containing 2% bovine serum albumin (BSA) and 1 mM EGTA. During experiments, 1 mM CaCl₂, histamine, VPA, CPA, DTN, caffeine, and 2-APB were added as indicated in figure legends. Permeabilization was performed using 100 μM digitonin for 30 s. IB buffer containing 0 Ca²⁺/100 μM EGTA was applied until stabilization, followed by IB containing 0.5 μM Ca²⁺. Luminescence was measured using a purpose-built luminometer. Calibration to [Ca²⁺] was achieved by adding excess Ca²⁺ (10 mM) in KHBH or KHBPC12 supplemented with 100 μM digitonin to expose the aequorin to maximal Ca²⁺.

On-Line Measurement of Catecholamine Release

Cells were gently detached using a rubber policeman and centrifuged at 800 rpm for 10 min. The pellet was resuspended in Krebs–Hepes buffer containing (in mM): 144 NaCl, 5.9 KCl, 1.2 MgCl₂, 11 glucose, 10 HEPES, and 1 Ca²⁺ (pH 7.4). The suspension was placed in a jacketed microchamber superfused at 2 mL/min at room temperature. Catecholamine secretion was continuously monitored “on-line” by an electrochemical detector (Metrohm AG CH-9100, Herisau, Switzerland) operating in amperometric mode[38]. Secretion was evoked by 5 s pulses of high K⁺ (75 mM) every 2 min.

Single-Cell [Ca²⁺]_i Measurements

Single-cell [Ca²⁺]_i was determined at room temperature in fura-2-loaded cells as described previously [39]. Excitation wavelengths were alternated between 340 and 380 nm, and emitted light at 520 nm was collected and analyzed using CellR® software (Olympus). Data were expressed as the fluorescence ratio F₃₄₀/F₃₈₀.

Chemicals

Coelenterazine n was obtained from Labnet Biotechnica (Madrid, Spain). CPA, histamine, InsP₃, VPA, veratridine (VTD), dantrolene (DTN), 2-APB, caffeine, and heparin were purchased from Sigma-Aldrich (Madrid, Spain). Fura-2 was from Molecular Probes. The cDNA encoding ER-targeted aequorin was a generous gift from Prof. Javier Alvarez.

Statistics

Values are expressed as mean ± SE. Statistical significance was determined using one-way ANOVA. Differences were considered significant at p < 0.05.

Results

Valproic Acid Induces Ca²⁺ Release from the Endoplasmic Reticulum

After aequorin reconstitution with coelenterazine in ER-depleted conditions, the experiments were initiated by perfusing HeLa cells with a medium containing 1 mM Ca²⁺ to allow ER refilling (Figure 1a). Complete ER refilling required 30–60 s, reaching a steady-state [Ca²⁺]_{ER} of 570.53 ± 17.52 μM. When histamine (100 μM) was applied for 2 min, the expected Ca²⁺ release from the ER occurred, decreasing [Ca²⁺]_{ER} from approximately 600 μM to 400 μM (Figure 1a). Application of VPA (3 μM) produced a similar effect, reducing [Ca²⁺]_{ER} from 600 to 470 μM (Figure 1b), corresponding to a 37.36

± 0.01% decrease. This effect was reversible. The concentration–response curve (Figure 1d) revealed a threshold at 3 μM and a maximal effect between 30–100 μM (approximately 55% ER release). The calculated IC₅₀ value was approximately 17 μM.




Figure 1. Panel a, shows the effects of histamine (100 μM) on $[Ca^{2+}]_{ER}$. This original trace was obtained in 23 experiments from 5 different batches of cells. Panels b and c illustrate VPA action at 3 and 10 μM, respectively, on $[Ca^{2+}]_{ER}$ in intact HeLa cells. Once the ER was refilled with 1 mM of Ca^{2+} (as shown by dots), drugs were applied as shown in the horizontal bars at the top of the figure. VPA experiments are representative of 25, 20, and 28 experiments of each type from 6, 5, and 7 different cell batches, respectively. Panel d shows a concentration-response curve for VPA. The average percentage of Ca^{2+} released from the ER induced by increased concentrations of VPA 3, 5, 10, 15, 30 and 100 μM. Data are means±s.e SE. Mean from 25, 12, 20, 12, 28, and 25 different experiments from 6, 3, 5, 3, 7, and 6 cell batches, respectively.

ER Ca^{2+} Release Elicited by VPA in Permeabilized HeLa Cells

To determine whether VPA acted via plasma membrane receptors or directly on intracellular targets, experiments were performed on digitonin-permeabilized HeLa cells expressing erAEQ. Following permeabilization with 100 μM digitonin, cells were superfused with intracellular buffer (IB) to obtain a stable baseline, after which ER refilling was achieved using 0.5 μM Ca^{2+} . Application of $InsP_3$ (5 μM) triggered a decrease in $[Ca^{2+}]_{ER}$ from 600 μM to 300 μM (Figure 2a). VPA (10 μM) also induced a reversible ER Ca^{2+} release (data not shown). Averaged results (Figure 2b) showed that $InsP_3$ and VPA released $49.26 \pm 0.01\%$ and ~25% of ER Ca^{2+} , respectively. The activation time constants (τ_{act}) for VPA- and $InsP_3$ -elicited Ca^{2+} release were comparable (Figure 2c), indicating similar kinetics.




Figure 2. [Ca²⁺]_{ER} in permeabilized HeLa cells. Panel a, shows original traces of InsP₃ 5 μM. After permeabilization, ER was refilled with 0.5 μM of Ca²⁺ as shown by dots. Drugs were applied as indicated in the horizontal bars at the top of the figure. Panel b represents average data expressed as % of Ca²⁺ release from ER due to VPA (3 and 10 μM) and InsP₃ (5 μM) application. Panel c shows average results expressed as % Ca²⁺ release. Thus, 3 and 10 μM of VPA demonstrated similar kinetics to InsP₃. Pooled data are means±s.e. mean of 12 and 15 experiments from 3 and 4 different cell batches (VPA 3 and 10 μM, respectively). Means±s.e. mean of InsP₃ came from 20 experiments from 5 different cell batches. # indicates significant differences between InsP₃ and VPA 3 μM. * indicates significant differences between InsP₃ and VPA 10 μM. n/s indicate not significant differences. An ANOVA test was performed.

Comparison of the Kinetics of ER Ca²⁺ Release Induced by CPA and VPA

To further understand the mechanism of VPA-induced ER Ca²⁺ release, its effect was compared with that of cyclopiazonic acid (CPA), a well-known inhibitor of the sarco/endoplasmic reticulum Ca²⁺-ATPase (SERCA). CPA (30 μM) decreased [Ca²⁺]_{ER} from 530 μM to 330 μM, corresponding to a 32.94 ± 0.10% release (Figure 3c). Lower concentrations of CPA (3 and 10 μM) elicited 11.74 ± 0.13% and 20.11 ± 0.15% Ca²⁺ release, respectively. In parallel experiments, VPA (3, 10, and 30 μM) produced 21.43 ± 0.10%, 37.36 ± 0.10%, and 51.10 ± 0.67% ER Ca²⁺ depletion, respectively (Figure 3b, c). Notably, VPA caused significantly faster ER Ca²⁺ release compared to CPA. The activation time constants (τ_{act}) for CPA were 94.41 ± 28.63 s (3 μM), 64.97 ± 2.89 s (10 μM), and 49.15 ± 1.08 s (30 μM), whereas for VPA they were 23.18 ± 2.12 s (3 μM), 25.66 ± 1.83 s (10 μM), and 32.06 ± 1.06 s (30 μM) (Figure 3d).




Figure 3. VPA follows a different mechanism from CPA. Original traces of CPA (30 μ M, panel a), a SERCA blocker, show different kinetics to VPA (30 μ M, panel b) in intact HeLa cells. Once the ER was refilled with 1 mM Ca^{2+} (as shown by dots), drugs were applied as shown in the horizontal bars at the top of the figure. Panel c represents average data expressed as % of Ca^{2+} release from ER, CPA, and VPA application. Panel d shows average results expressed as τ_{act} at increasing concentrations of VPA and CPA applications. For CPA 3, 10, and 30 μ M pooled data are means $\pm$ s.e. mean of 10, 10, and 30 experiments from 3, 3, and 7 different cell batches; for VPA 3, 10, and 30 μ M pooled data are means $\pm$ s.e. mean of 25, 20, and 28 experiments from 6, 5, and 7 different cell batches. # indicates significant differences between VPA and CPA in terms of Ca^{2+} release. * indicates significant differences between VPA and CPA when the τ_{act} is analyzed. An ANOVA test was performed.

[Ca^{2+}]_{ER} Is Unaffected by 2-APB and Heparin

To directly assess the involvement of the $InsP_3$ receptor ($InsP_3R$) in VPA-induced ER Ca^{2+} release, the $InsP_3R$ inhibitors 2-aminoethyl diphenylborinate (2-APB) and heparin were used. Pre-incubation with 10 μ M 2-APB for 2 min abolished the VPA (10 μ M)-induced ER Ca^{2+} release (Figure 4a). Similarly, in permeabilized HeLa cells, superfusion with 200 μ g/mL heparin for 2 min before VPA application prevented any detectable change in $[Ca^{2+}]_{ER}$ (Figure 4b). The inhibitory effects of both agents are summarized in Figure 4c, confirming that VPA acts through $InsP_3$ -sensitive Ca^{2+} stores.




Figure 4. Panel a shows HeLa cells perfused with 2-APB at 10 μ M before the addition of VPA 10 μ M. Panel b shows HeLa cells treated with VPA (10 μ M) and heparin (200 μ g/ml). Drugs were applied as indicated in the horizontal bars at the top of the figure. Panel d are pooled data from 12 and 15 experiments from 3 and 4 different cell batches. * Indicates significant differences between 2-APB and heparin with respect to VPA. An ANOVA test was performed.

VPA Releases Ca^{2+} via $InsP_3R$ in PC12 Cells

To examine the specificity of VPA action on $InsP_3R$, experiments were performed in PC12 cells, which express both $InsP_3R$ and RyR, as well as voltage-dependent calcium channels (VDCCs) of the L- and N-type. Activation of VDCCs by depolarization with high K^+ induces Ca^{2+} release through RyR. In ER-depleted PC12 cells expressing erAEQ, reintroduction of 1 mM Ca^{2+} restored $[Ca^{2+}]_{ER}$ (Figure 5a). Subsequent perfusion with high K^+ or VPA (10 μ M) elicited ER Ca^{2+} release via RyR or $InsP_3R$, respectively. Co-application of caffeine (CAF) and VPA further enhanced ER Ca^{2+} release (Figure 5b). Dantrolene (DTN) (100 μ M), an RyR inhibitor, abolished Ca^{2+} release triggered by K^+ depolarization (Figure 5c) but not that induced by VPA (Figure 5d), confirming that VPA acts selectively on $InsP_3R$ -mediated pathways.

Fig. 5




Figure 5. VPA releases Ca^{2+} from the ER in PC12 cells. Original traces of high K^+ (75 mM) and VPA (10 μM , panel a), the coapplication of CAF (1 mM) and VPA (10 μM) showed different Ca^{2+} release from the ER to panel b) in intact PC12 cells. Once the ER was refilled with 1 mM Ca^{2+} (as shown by dots), drugs were applied as shown in the horizontal bars at the top of the figure. The perfusion of DTN (100 μM) abolished Ca^{2+} release from the ER induced by high K^+ (75 mM) (panel c) but not VPA Ca^{2+} release (panel d). Pooled data are means $\pm$ s.e. mean of 10, from different cell batches (panel a); pooled data are means $\pm$ s.e. mean of 3 experiments from 2 different cell batches (panel b); pooled data are means $\pm$ s.e. mean of 9 experiments from 3 different cell batches (panel c, d).

VPA Mitigates Intracellular Ca^{2+} Oscillations Induced by Veratridine

Given VPA's clinical use in epilepsy, we investigated whether its ER Ca^{2+} -releasing effect modulates veratridine (VTD)-induced cytosolic Ca^{2+} oscillations, a cellular model of epileptiform activity. Application of VTD (50 μM) evoked rhythmic Ca^{2+} oscillations in fura-2-loaded bovine chromaffin cells (BCCs) (Figure 6a, b). Superfusion with VPA (30 μM) abolished these oscillations (Figure 6c) while simultaneously increasing basal $[\text{Ca}^{2+}]_{\text{cyt}}$. Thus, VPA suppresses VTD-induced Ca^{2+} oscillations, likely by releasing Ca^{2+} from ER stores.

Fig. 6




Figure 6. The effect of VPA on VTD epilepsy model in fura-2 loaded BCC.

Panel a shows a row of images with the region of interest (roi) of $[Ca^{2+}]_c$ peak caused upon high K^+ (75 mM) depolarization, followed by Ca^{2+} oscillations produced by VTD (as shown in horizontal bar) in the control solution. The pseudocolor scale indicates the increase in $[Ca^{2+}]_c$. Original traces in the absence (b) and in the presence of VPA (c) using the protocol described in panel a. Horizontal bars at the bottom of the figure represent the application of the compounds. Experiments are representative of 3 to 6 of each type.

VPA Facilitates Catecholamine Release in Bovine Chromaffin Cells

To evaluate the functional consequences of VPA-induced ER Ca²⁺ release on neurotransmitter exocytosis, catecholamine secretion was measured in fast-superfused BCCs. Depolarization with 35 mM K⁺ for 5 s every 2 min evoked reproducible catecholamine release peaks averaging 80–90 nA (Figure 7a). Following 15 min of Krebs–Hepes superfusion, the mean peak value was 122.19 ± 4.31 nA. In the presence of VPA (3 μM), the average peak amplitude increased significantly to 198.20 ± 4.81 nA (Figure 7b). Data from eight independent experiments confirmed that VPA potentiated K⁺-evoked catecholamine release (Figure 7c), suggesting that ER Ca²⁺ mobilization by VPA enhances vesicular exocytosis.

Fig. 7




Figure 7. VPA potentiates the release of catecholamines in BCC.

Panel a shows original traces of catecholamine secretory peaks induced by K⁺ (35 mM, during 5 s every 2 min) pulse in control cells. Panel b displays the enhanced response in the presence of VPA,

as shown in the horizontal bar at the bottom of the figure. Panel c illustrates the averaged results expressed as % of catecholamine secretion. Pooled data are means $\pm$ s.e. mean of 8 experiments from 3 different cell cultures. * Indicates significant differences of VPA with respect to control. An ANOVA test was performed.

4. Discussion

The major finding of the present study is that the antiepileptic drug valproic acid (VPA) acts as an agonist for the inositol 1,4,5-trisphosphate receptor (InsP₃R). This conclusion is supported by the observation that, in HeLa cells, VPA induced a concentration-dependent Ca²⁺ release from the endoplasmic reticulum (ER) (Figure 1d). Importantly, the effect was not mediated by plasma membrane receptors, as it persisted in digitonin-permeabilized cells (Figure 2). In the absence of second messengers such as InsP₃, VPA evoked Ca²⁺ release with kinetics remarkably similar to those triggered by InsP₃ itself. The activation time constant (τ_{act}) was almost identical for both compounds (Figure 2c), indicating that VPA mimics the physiological agonist of the InsP₃R. In contrast, the kinetics of ER Ca²⁺ release elicited by VPA and by cyclopiazonic acid (CPA)—a specific inhibitor of the sarco/endoplasmic reticulum Ca²⁺-ATPase (SERCA)—were clearly different. These data suggest that VPA does not act through SERCA inhibition, but rather directly targets the InsP₃R. This interpretation is further reinforced by the observation that the specific blockade of InsP₃R using either 2-APB [28] or heparin [29] completely abolished VPA-induced ER Ca²⁺ release.

VPA specifically targets InsP₃R but not RyR. The InsP₃R appears to be the principal intracellular target of VPA. Although HeLa cells express ryanodine receptors (RyR), they do not contribute significantly to Ca²⁺ signaling [7]. To confirm the specificity of VPA for InsP₃R, experiments were extended to PC12 cells, which express both InsP₃R and RyR [21,23]. In these cells, K⁺ stimulation and VPA each triggered ER Ca²⁺ release in a two-step sequence. However, the caffeine-induced Ca²⁺ release via RyR was not affected by VPA, indicating that VPA selectively targets InsP₃R-dependent Ca²⁺ stores.

VPA suppresses veratridine-induced Ca²⁺ oscillations. A particularly intriguing finding was that VPA abolished the cytosolic Ca²⁺ oscillations induced by veratridine (VTD) in chromaffin cells (Figure 6c). Because VTD is used as a cellular model of epileptiform activity [40–43], the ability of VPA to suppress these oscillations may have direct pathophysiological relevance. Interestingly, VPA itself increased cytosolic [Ca²⁺] (Figure 6c, right), likely reflecting InsP₃R-mediated ER Ca²⁺ release. Such modulation could influence vesicular trafficking and neurotransmitter release, contributing to the drug's antiepileptic effects. Previous studies have demonstrated that VPA can inhibit epileptiform discharges induced by VTD [41], yet paradoxically, VPA has also been reported to exert pro-epileptic effects under certain conditions [40]. Clinical reports describe cases of valproate-induced status epilepticus [10]. These divergent effects may depend on individual sensitivity of InsP₃R subtypes to VPA, offering a potential explanation for both pharmacoresistance and proconvulsive reactions in some patients.

To our knowledge, this is the first comprehensive study demonstrating that VPA directly triggers Ca²⁺ release from the ER through InsP₃R activation and enhances exocytosis. Although other authors have shown that VPA modulates insulin secretion [17], the mechanism underlying this effect was not linked to ER Ca²⁺ dynamics. Similarly, Yamamoto et al. (1997) reported that chronic VPA exposure upregulated sodium channels and increased catecholamine secretion in adrenal chromaffin cells, but without considering ER Ca²⁺ homeostasis.

Furthermore, several studies have explored the role of VPA in bipolar disorder by analyzing its effects on Ca²⁺ signaling pathways [2,15,19]. Akimoto and colleagues found that VPA inhibited serotonin-induced Ca²⁺ responses in human platelets in a concentration-dependent manner, possibly involving protein kinase C (PKC) modulation. However, none of these investigations examined whether VPA acts directly on ER Ca²⁺ stores, as revealed in our present study.

We propose that VPA-induced Ca²⁺ release from the ER via InsP₃R may represent an effective mechanism to regulate secretion and synaptic transmission. If similar phenomena occur in neurons,

VPA-triggered ER Ca^{2+} mobilization could contribute to synaptic plasticity and neurotransmitter modulation. Under pathological conditions such as epilepsy, this mechanism might help attenuate the propagation of epileptic discharges by enhancing GABA release, thereby counteracting neuronal hyperexcitability [16–43].

In conclusion, our results reveal a novel intracellular mechanism of action for the classical antiepileptic drug VPA. The drug behaves as an InsP_3R agonist, causing Ca^{2+} release from ER stores. Considering the central role of InsP_3 -dependent Ca^{2+} signaling in neuronal excitability and epileptogenesis, this mechanism offers promising insights into the pharmacological action of VPA. Future studies should focus on identifying the molecular determinants of VPA– InsP_3R interaction and their implications for antiepileptic resistance and neuronal calcium homeostasis in chronic epilepsy.

Author Contributions: For research articles with several authors, a short paragraph specifying their individual contributions must be provided. The following statements should be used “Conceptualization, MFCA and ARN; methodology, MFCA and ARN; software, MFCA and ARN.; validation, MFCA and ARN; formal analysis, ARN; investigation, MFCA; resources, MFCA and ARN; data curation, ARN; writing—original draft preparation, MFCA and ARN; writing—review and editing, MFCA and ARN; visualization, MFCA; supervision, MFCA; project administration, MFCA and ARN; funding acquisition, MFCA and ARN. All authors have read and agreed to the published version of the manuscript.”

Funding: This work was supported the by following grants: (1) Ministerio de Sanidad y Consumo FIS (nº: 041665); (2) from Fundación La Caixa (nº: BN05-32-0); (3) Ministerio de Educación: Acciones integradas and (4) Ramon y Cajal Programme, to MFCA; ARN was supported by FIS (nº: PI052124; CP04/00132).

Acknowledgments: The generous gift of er-AEQ by Prof. Javier Alvarez is gratefully acknowledged.

Conflicts of Interest: Declare conflicts of interest or state “The authors declare no conflicts of interest.”

References

1. Malenka, R. C. and Nicoll, R. A. Long-term potentiation a decade of progress?. *Science* **1999**, *285*, 1870–1874
2. Akimoto, T., Kusumi, I., Suzuki, K., Masui, T. and Koyama T. Effects of valproate on serotonin-induced intracellular calcium mobilization in human platelets. *J Psychiatry Neurosci.* **2007**, *32*, 17-22. PMID: 17245470
3. Pal, S., Limbrick Jr., D. D., Rafiq, A. and DeLorenzo, R. J. Induction of spontaneous recurrent epileptiform discharges causes long term changes in intracellular calcium homeostatic mechanisms. *Cell Calcium* **2000**, *28*, 181– 193
4. Pal, S., Sun, D., Limbrick, D., Rafiq, A. and DeLorenzo, R. J. Epileptogenesis induces long-term alterations in intracellular calcium release and sequestration mechanisms in the hippocampal neuronal culture model of epilepsy. *Cell Calcium.* **2000**, *30*, 285– 296
5. DeLorenzo, R. J., Sun, D. A. and Deshpande, L.S. Cellular mechanisms underlying acquired epilepsy: The calcium hypothesis of the induction and maintenance of epilepsy. *Pharmacol. and Therap.* **2005**, *105*, 229– 266
6. Ehrlich, BE., Kaftan, E., Bezprozvannaya, S. and Bezprozvanny, I. The pharmacology of intracellular Ca^{2+} release channels. *Trends Pharmacol.* **1994**, *15*, 145–149
7. Bennett, D. L., Cheek, T. R., Berridge, M. J., De Smedti H., Parysi, J. B., Missiaeni L. and Bootman, M. D. Expression and function of ryanodine receptors in nonexcitable cells. *J Biol. Chem.* **1996**, *271*, 6356-6362
8. Berridge MJ. Neuronal calcium signaling. *Neuron* **1998**, *21*, 13–26
9. Royero PX, Higa GSV, Kostecki DS, Dos Santos BA, Almeida C, Andrade KA, Kinjo ER, Kihara AH. Ryanodine receptors drive neuronal loss and regulate synaptic proteins during epileptogenesis. *Exp Neurol.* **2020** 327:113213. doi: 10.1016/j.expneurol.2020.113213.
10. Capocchi, G., Balducci, A., Cecconi, M., Pelli, MA., Picchiarelli, A., Silvestrelli, G. and Zampolini, M. Valproate-induced epileptic tonic status. *Seizure.* **1998**, *7*, 237-241

11. Ekwuru, MO. and Cunningham, JR. Phaclofen increases GABA release from valproate treated rats. *Br. J. Pharmacol.* **1990**, *99*, suppl., 251P.
12. Gram, L., Larsson, OM., Johnsen, AH. and Schousboe, A. Effects of valproate, vigabatrin and aminooxyacetic acid on release of endogenous and exogenous GABA from cultured neurons. *Epilepsy Res.* **1988**, *2*, 87-95
13. Kelly, KM., Gross, RA. and Macdonald, RL. Valproic acid selectively reduces the low-threshold (T) calcium current in rat nodose neurons. *Neurosci. Lett.* **1990**, *116*, 233-8
14. Kim, A.J., Shi, Y., Austin, RC. and Werstuck, GH. Valproate protects cells from ER stress-induced lipid accumulation and apoptosis by inhibiting glycogen synthase kinase-3. *J. Cell Sci.* **2005**, *118*, 89-99
15. Kurita, M., Nishino, S., Ohtomo, K., Rai, M., Shirakawa, H., Mashiko, H., Niwa, S. and Nakahata, N. Sodium valproate at therapeutic concentrations changes Ca^{2+} response accompanied with its weak inhibition of protein kinase C in human astrocytoma cells. *Prog Neuropsychopharmacol. Biol. Psychiatry.* **2007**, *31*, 600-4
16. Löscher, W. Basic pharmacology of valproate: A review after 35 years of clinical use for the treatment of epilepsy. *CNS drugs.* **2002**, *16*, 669-694
17. Luef, GJ., Lechleitner, M., Bauer, G., Trinka, E., and Hengster, P. Valproic acid modulates islet cell insulin secretion: a possible mechanism of weight gain in epilepsy patients. *Epilepsy Res.* **2003**, *55*, 53-57
18. Vreugdenhil, M., van Veelen, CW., van Rijen, PC., Lopes da Silva, FH. and Wadman, WJ. Effect of valproic acid on sodium currents in cortical neurons from patients with pharmaco-resistant temporal lobe epilepsy. *Epilepsy Res.* **1998**, *32*, 309-320
19. Suzuki, K., Kusumi I., Akimoto, T., Sasaki, Y. and Koyama, T. Effects of lithium and valproate on agonist-induced platelet intracellular calcium mobilization: relevance to myosin light chain kinase. *Prog Neuropsychopharmacol Biol Psychiatry.* **2004**, *28*, 67-72
20. Pizzo, P., Fasolato, C. and Pozzan, T. Dynamic properties of an inositol 1,4,5-trisphosphate- and thapsigargin-insensitive calcium pool in mammalian cell lines. *J Cell Biol.* **1997**, *136*, 355-366
21. Clementi, E., Scheer, H., Zacchetti, D., Fasolato, C., Pozzan, T. and Meldolesi, J. Receptor-activated Ca^{2+} influx. Two independently regulated mechanisms of influx stimulation coexist in neurosecretory PC12 cells. *J Biol Chem.* **1992**, *267*, 2164-2172
22. Ehrlich, BE., Kaftan, E., Bezprozvannaya, S. and Bezprozvanny, I. The pharmacology of intracellular Ca^{2+} release channels. *Trends Pharmacol.* **1994**, *15*, 145-149
23. Fasolato, C., Zottini, M., Clementi, E., Zacchetti, D., Meldolesi, J. and Pozzan, T. Intracellular Ca^{2+} pools in PC12 cells. *J. Biol. Chem.* **1991**, *266*, 20159-20167
24. García-Sancho, J., Montero, M. and Alvarez, J. Ca^{2+} -induced Ca^{2+} release in chromaffin cells seen from inside the ER with targeted aequorin. *J Cell Biol.* **1999**, *144*, 241-54.
25. Montero, M., Alvarez, J., Scheenen, WJ.J., Rizzuto, R., Meldolesi, J. and Pozzan, T. Ca^{2+} Homeostasis in the endoplasmic reticulum: coexistence of high and low $[\text{Ca}^{2+}]$ subcompartments in intact HeLa cells. *J. Cell. Biol.* **1997**, *139*, 601-611
26. Moreno, A. Lobatón CD., SantoDomingo J., Vay L., Hernandez-SanMiguel E., Rizzuto, R., Montero, M. and Alvarez, J. Calcium dynamics in catecholamine containing secretory vesicles. *Cell Calcium.* **2005**, *37*, 555-564
27. Mori, F., Okada, M., Tomiyama, M., Kaneko, S. and Wakabayashi, K. Effects of ryanodine receptor activation on neurotransmitter release and neuronal cell death following kainic acid-induced status epilepticus. *Epilepsy Res.* **2005**, *65*, 59-70
28. Zima, AV., Bare, DJ., Mignery, GA. and Blatter, LA. IP_3 -dependent nuclear Ca^{2+} signalling in the mammalian heart. *J Physiol.* **2007**, *584*, 601-611
29. Ramos-Franco, J., Fill M. and Mignery G. A. Isoform-specific function of single inositol 1,4,5-trisphosphate receptor channels. *Biophysic. J.* **1998**, *75*, 834-839.
30. Rizzuto, R. Intracellular Ca^{2+} pools in neuronal signalling. *Curr Opin Neurobiol.* **2001**, *11*, 306-311
31. Yamamoto, R., Yanagita, T., Kobayashi, H., Yokoo, H. and Wada, A. Up-regulation of sodium channel subunit mRNAs and their cell surface expression by antiepileptic valproic acid: activation of calcium channel and catecholamine secretion in adrenal chromaffin cells. *J. Neurochem.* **1997**, *68*, 1655-1662

32. Zona, C. and Avoli, M. Effects induced by the antiepileptic drug valproic acid upon the ionic currents recorded in rat neocortical neurons in cell culture. *Exp. Brain. Res.* **1990**, *81*, 313-317
33. Ootom, S.A. and Alkadhi, K.A. Valproic acid intensifies epileptiform activity in the hippocampal pyramidal neurons. *Neurosci Res.* **1999**, *35*, 299-307
34. Alvarez, J. and Montero, M. Measuring Ca^{2+} in the endoplasmic reticulum with aequorin. *Cell Calcium.* **1999**, *32*, 251-260
35. Chen, CA. and Okayama, H. Calcium phosphate-mediated gene transfer: a highly efficient transfection system for stably transforming cells with plasmid DNA. *Biotechniques.* **1988**, *6*, 632-638
36. Díaz-Prieto N, Herrera-Peco I, de Diego AM, Ruiz-Nuño A, Gallego-Sandín S, López MG, García AG, Cano-Abad MF. Bcl2 mitigates Ca^{2+} entry and mitochondrial Ca^{2+} overload through downregulation of L-type Ca^{2+} channels in PC12 cells. *Cell Calcium.* **2008**, *44*:339-52. doi: 10.1016/j.ceca.2008.01.007.
37. Moro, MA., Lopez, MG., Gandia, L., Michelena, P. and Garcia, AG. Separation and culture of living adrenal- and noradrenalina-containing cells from bovine adrenal medullae. *Anal. Biochem.* **1990**, *185*, 243-8
38. Borges, R., Sala, F. and Garcia, AG. Continuous monitoring of catecholamine release from perfused cat adrenals. *J. Neurosci. Methods.* **1986**, *16*, 289-300
39. Cano-Abad, MF., Lopez, MG., Hernandez-Guijo, JM., Zapater, P., Gandia, L., Sanchez-Garcia, P. and Garcia, AG. Effects of the neuroprotectant lubeluzole on the cytotoxic actions of veratridine, barium, ouabain and 6-hydroxydopamine in chromaffin cells. *Br. J. Pharmacol.* **1998**, *124*, 1187-1196.
40. Ootom, S., Tian, LM. and Alkadhi, KA. Veratridine-treated brain slices: a cellular model for epileptiform activity. *Brain Res.* **1998**, *789*, 150-156.
41. Ootom, S.A. and Alkadhi, K.A. Epileptiform activity of veratridine model in rat brain slices: effects of antiepileptic drugs. *Epilepsy Res.* **2000**, *38*, 161-170
42. Conceição I. M., Lebrun I., Cano-Abad M. F., Gandía L., Hernández-Guijo J. M., López M. G., Villarroja M., Jurkiewicz A. and García A. G. Synergism between toxin gamma from the brazilian scorpion *Tityus serrulatus* and veratridine in chromaffin cells. *Am. J. Physiol.* **1998**, *274*: 1745-1754
43. Cano-Abad, MF., Villarroja, M., Garcia, AG., Gabilan, NH. and Lopez, MG. Calcium entry through L-type calcium channels causes mitochondrial disruption and chromaffin cell death. *J. Biol. Chem.* **2001**, *276*, 39695-39704

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.