

Article

Not peer-reviewed version

Impact of Epithelial Inflammation of the Membrane Remnant on the Outcome of Tympanoplasty

María Fernanda Galindo-Tapia , Alejandro Esteban Deras-Quiñones , Itzel Maria Montoya-Fuentes , [Eduardo Osiris Madrigal-Santillán](#) , [Angél Morales-González](#) , [Naria A. Flores-Fuentes](#) , Liliana Anguiano-Robledo , [Raúl Rojas-Martínez](#) , [Beatriz Montaña-Velázquez](#) ^{*} , [José A. Morales-González](#) ^{*}

Posted Date: 14 April 2025

doi: 10.20944/preprints202504.1056.v1

Keywords: Chronic otitis media; tympanic perforation; tympanoplasty; epithelial inflammation



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Article

Impact of Epithelial Inflammation of the Membrane Remnant on the Outcome of Tympanoplasty

María Fernanda Galindo-Tapia ¹, Alejandro Esteban Deras-Quñones ²,
Itzel Maria Montoya-Fuentes ², Eduardo Osiris Madrigal-Santillán ³, Ángel Morales-González ⁴,
Naria A. Flores-Fuentes ⁴, Liliana Anguiano-Robledo ⁵, Raúl Rojas-Martínez ³,
Beatriz Montaña-Velázquez ^{1,*} and José A. Morales-González ^{3,*}

¹ Instituto Mexicano del Seguro Social, Centro Médico Nacional La Raza. Hospital General "Dr. Gaudencio González Garza", servicio de Otorrinolaringología. Ciudad de México

² Instituto Mexicano del Seguro Social, Centro Médico Nacional La Raza. Hospital de especialidades "Dr. Antonio Fraga Mouret", servicio de Patología. Ciudad de México, México

³ Laboratorio de Medicina de conservación, Escuela Superior de Medicina, Instituto Politécnico Nacional, Plan de San Luis y Díaz Mirón, Col. Casco de Santo Tomás, Alcaldía Miguel Hidalgo, Ciudad de México, C.P. 11340, México

⁴ Escuela Superior de Cómputo, Instituto Politécnico Nacional, Unidad Profesional "A. López Mateos", Ciudad de México 07738, Mexico

⁵ Laboratorio de Farmacología Molecular, Escuela Superior de Medicina, Instituto Politécnico Nacional, Plan de San Luis y Díaz Mirón, Col. Casco de Santo Tomás, Del. Miguel Hidalgo, Mexico City 11340, Mexico

* Correspondence: beamont_2000@yahoo.com.mx (B.M-V.), jmorales101@yahoo.com.mx (J.A.M-G.)

Abstract: Background: Chronic otitis media (COM) with tympanic perforation sometimes requires tympanoplasty. Many factors can interfere with surgical success; however, the histological status of the remaining epithelium of the perforation has not been studied as a risk factor for surgical failure. **Methods:** Observational, longitudinal, analytical study in patients with COM, candidates for tympanoplasty who met the inclusion criteria between August and December 2024. Tympanoplasty was performed and the tympanic ring epithelium was sent for histological analysis. After 30 days, closure or non-closure of the perforation was determined and the results were collected. Descriptive and analytical statistics were performed according to the data distribution using the SPSS 26.0 statistical package. **Results:** Twenty subjects were included; 80% with tubal dysfunction, 60% central perforation, and 65% medium-sized. 13 were successful and 7 failed. Histopathological analysis revealed dystrophic calcification, chronic lymphocytic infiltrate, histiocytic infiltrate, fibrosis, loose keratin sheets, metaplasia, and spongiosis. The logistic regression model showed an OR of 7.3 for marginal perforation and 3.4 for the OPSS score. Of the patients with surgical failure, 57.4% had epithelial inflammation. **Conclusions:** Epithelial inflammation affected surgical success in more than 50%.

Keywords: Chronic otitis media; tympanic perforation; tympanoplasty; epithelial inflammation

1. Introduction

Chronic otitis media (COM) is a very common condition in our environment. It is defined as an inflammatory process of the middle ear mucosa that lasts more than 3 months (1), with purulent discharge and perforation of the tympanic membrane (2)(3). Tympanic perforation is defined as a rupture of the tympanic membrane, creating a connection between the external auditory canal and the middle ear (4), making the ear more susceptible to infection and hearing loss. The prevalence of COM ranges from 1% to 6% of the world's population, up to 11% in developing countries such as Mexico and 57% in countries with extreme poverty (5–7). There are more than 150 million people with tympanic perforation worldwide (5), with a global incidence of 4.8 cases per 1000 people per year. Tympanic perforation varies between 5% and 30% (4). In Mexico, it is estimated that 0.5% of patients over 15 years of age have chronic otitis media and 4% have tympanic perforation (1). In the Otolaryngology and Head and Neck Surgery Department of the General Hospital of the National Medical Center "La Raza", 82% of consultations are for COM.

The main cause of perforation is middle ear infection, followed by trauma (87.5% vs. 12.5%) (8). Recurrent infections, bacterial colonization, aural ventilation dysfunction, tubal dysfunction, or

immunodeficiency can affect closure (9), as can myringotomy, ventilation tubes, and hyperbaric oxygen therapy (10). In rarer cases, it can be due to otic herpes zoster, tuberculosis, and aggressive otitis externa (11).

The tympanic membrane separates the middle ear from the external ear. It is oval in shape, with a vertical diameter of 9-10 mm and a horizontal diameter of 8-9 mm. Macroscopically, it is divided into two parts: the pars flaccidum superior to the malleus, and the pars tensa inferiorly. It is composed of three layers (12): an outer layer composed of simple stratified squamous epithelium, which is continuous with the skin of the ear canal; a middle fibrous layer called the lamina propria, formed by radial and circular fibers; and finally, the inner layer composed of pseudostratified low cuboidal epithelium, which is continuous with the mucosa of the middle ear (13).

When the tympanic membrane is perforated, it tends to close spontaneously. Unlike the skin, it closes from the outside in, due to the proliferation of stratified squamous epithelium in the lateral portion, which migrates from the remnant to the central portion of the perforation (12,14). Once the outer layer closes the defect, the fibrous layer and mucosa regenerate beneath it (15). This regeneration can be affected by several factors, including chronic inflammation, epithelialization of the perforation margins, insufficient blood supply, or growth factor deficiency (16). If a tympanic membrane perforation does not heal on its own within 12 weeks, it is unlikely to heal and should be treated with tympanoplasty (7,9), which is the reconstruction of the tympanic membrane to prevent disease progression and, to a lesser extent, to restore hearing.

Most studies on surgical success report success rates between 80% (17) and 90% in ears with normal ventilation (9); however, in cases with chronic inflammation of the middle ear, the outcome tends to be less successful. Moyakan et al. mention that 20% of unsuccessful tympanoplasties are associated with incomplete integration, graft retraction, or graft displacement (17).

Throughout history, studies have been conducted on the histological changes in the ear mucosa resulting from chronic otitis media. In 1970, the presence of mucosal hyperplasia and mucus hypersecretion was demonstrated (18). In 2013, mucosal hyperplasia, edema, polypoid changes, tympanosclerosis, and mucosal granulomas were identified (7). However, there are few histological studies of the epithelium of the perforation remnant. In 2022, a study was conducted on the epithelial remnant of the perforation and the surrounding middle ear mucosa. The study found normal epithelium in 10.26% of cases, hypertrophied epithelium with chronic inflammatory infiltrate in 48.72%, tympanosclerosis with chronic inflammatory infiltrate in 20.51%, edematous hypertrophic mucosa in 17.95%, and keratosis with vascular stroma in 2.56%. With a surgical success rate of 84.62% at 3 months (6). However, data on inflammation were not cross-referenced with surgical success. This is the reason for the interest in conducting this study, to determine whether there was a relationship between the presence of epithelial inflammation and the success of surgical closure.

2. Materials and Methods

This was an observational, longitudinal, analytical study with non-probability convenience sampling. It was conducted in the otolaryngology department of the General Hospital "Dr. Gaudencio González Garza" at the National Medical Center "La Raza" of the Mexican Social Security Institute.

Patients were selected following inclusion criteria: a diagnosis of chronic otitis media with tympanic membrane perforation, no prior surgical treatment, age over 18 years, preoperative CT and audiometric studies, and no requirement for mastoidectomy or any other procedure. Once selected, the informed consent form was read, reviewed, and signed by each individual. Between August 2024 and December 2024, the tympanoplasties were performed, all endoscopic transcanal, using tragus cartilage grafts, by the same surgeon.

During the procedure, tissue was collected from the remnant of the perforation using the following technique: using a 45-degree incision, the epithelium surrounding the perforation was incised approximately 1 mm in the shape of a postage stamp, the incisions were then joined and the fibrous ring was detached, which was extracted with alligator forceps. Then, using a Rosen knife, the surrounding epithelium was removed 2 mm from the outer layer by gently scraping the membrane, the tissue was removed with alligator forceps.

Once the tissue was collected, it was placed in a sealed container immersed in 10% formaldehyde for fixation for at least 24 hours while it was sent to the pathology department. Once there, each sample was passed through an alcohol train for dehydration, embedded, and cut into paraffin blocks approximately 3 micrometers thick. The slides were mounted and stained with hematoxylin and

eosin, then covered with coverslips for microscopic analysis using 4x, 10x, and 40x objectives to evaluate the presence of inflammatory cells per field. Sample analysis was performed by two pathologists in a blinded manner, with a pre-standardized approach for reporting histopathological changes.

Patients were seen again at 10 and 30 days. At the first appointment, the sutures were removed from the graft harvesting site, and the success or failure of the tympanoplasty was determined at the second appointment. Histopathology results were also collected to determine the presence or absence of epithelial inflammation in the perforation remnant.

Once the data were collected, statistical analysis was performed. Mean and standard deviation were used for normally distributed quantitative variables; median and interquartile ranges were used for freely distributed variables. Finally, qualitative variables were presented as frequencies and percentages. Bivariate analysis was performed using the chi-square test to compare the presence or absence of epithelial inflammation versus tympanoplasty success or failure. The different characteristics between the two study groups were compared using the Student t test for normally distributed quantitative variables; the Mann-Whitney U test was used for freely distributed variables; and the chi-square test for dichotomous qualitative variables. Variables that showed statistical significance were included in a linear regression model to determine risk factors for epithelial inflammation and those that directly impacted surgical success. This model was performed using the SPSS 26.0 statistical package, and p values < 0.05 were considered significant.

This study was evaluated and approved by the local Research and Ethics Committee. The study adhered to the General Health Law and its Regulations on Health Research, as well as the Standards of the Mexican Social Security Institute. It complies with the provisions of the National Institute for Transparency, Access to Information, and Protection of Personal Data, as well as NOM 012-SSA3-2012. The study also adhered to the current ethical considerations contained in the Nuremberg Code, the Council for International Organizations of Medical Sciences (CIOMS), the Belmont Report, the Declaration of Helsinki, as well as the international guidelines for medical research involving human subjects adopted by the WHO and the Council for International Organizations for Research Involving Human Subjects. There are no conflicts of interest for the conduct of this study or its publication.

3. Results

Twenty patients were included, 15 women (75%) and 5 men (25%). The mean age was 56 years with a SD of 10.9 years, with a minimum age of 37 and a maximum of 78. The OOPS index, which indicates the state of the middle ear mucosa by means of a score, presented a median of 1, with a IQR of 0 and 3. The total presurgical hearing average presented a mean of 45.5 dB (SD 15) with a minimum of 23 dB and a maximum of 86 dB. Regarding tubal dysfunction, 16 patients presented it (80%), 50% of the affected ears were right and 50% left and 12 (60%) presented central perforation. About of size, 3 (15%) had a small perforation (<30%), 13 (65%) had a medium perforation (30-60%), and 4 (20%) had a subtotal perforation (>60%). Half of the subjects (n=10) had well pneumatized mastoids and the rest (n=10) had sclerotic mastoids (Table 1).

Table 1. Baseline characteristics of patients with tympanic perforation.

Variable	n=20
Sex, women, N (%)	15 (75)
Age, years, mean, (SD)	56.7 (10.9)
ATA, dB, mean (SD)	45.5 (15)
OOPS index, median (IQR)	1 (0, 3)
Tubal dysfunction, present, N (%)	16 (80)
Ear, right, N (%)	10 (50)
Perforation site, central, N (%)	12 (60)
Perforation size	
< 30% N (%)	3 (15)
30-60% N (%)	13 (65)
>60% N(%)	4 (20)

Mastoids, well pneumatized, N (%)	10 (50)
IQR: interquartile ranges (25,75). SD: standard deviation. ATA: auditory tonal average. dB:decibel	

Regarding postoperative care, 18 subjects (90%) reported adequate postoperative dry ear care, while 2 of them experienced water ingress. Thirteen subjects (65%) were successful, and 7 (35%) had surgical failure within the first 30 days. Histopathological analysis of the samples reported 6 cases (30%) with dystrophic calcification, 3 (15%) with leukocytic infiltrate, 3 (15%) with histiocytic infiltrate, 1 patient presented metaplasia of squamous epithelium with cuboidal epithelium and 1 additional patient presented spongiosis; the Figure 1 show the rest of the results with fibrosis, loose keratin sheets, normal epithelium. Cases with leukocyte infiltrate, spongiosis and presence of histiocytes were classified as having inflammation, the rest were considered without epithelial inflammation (Figure 1).




Figure 1. Histopathological results.

Table 2 summarizes the baseline characteristics of the patients according to the presence or absence of epithelial inflammation of the remnant of the tympanic perforation, where no statistically significant differences were observed between the groups.

Table 2. Baseline characteristics of patients with epithelial inflammation vs patients without epithelial inflammation.

Variable	Epithelial inflammation		p
	Absent	Present	
	n=12	n=8	
Sex, women, N (%)	10 (83.33)	5 (62.5)	.296`
Age, years, mean, (SD)	52 (13.3)	52.6 (14.6)	.427**
ATA, dB, mean (SD)	50.8 (15.8)	42.4 (12.9)	.313**
OOPS index, median (IQR)	0 (0-0)	2 (0-2)	.191**
Tubal dysfunction, present, N (%)	10 (83.33)	6 (75)	.535`
Ear, right, N (%)	7 (58.33)	3 (37.5)	.325`
Perforation site, central, N (%)	7 (58.33)	5 (62.5)	.612`
Perforation size			

< 30% N (%)	2 (16.66)	1 (12.5)	.651°
30-60% N (%)	8 (66.66)	5 (62.5)	
>60% N (%)	2 (16.66)	2 (25)	
Mastoids, well pneumatized, N (%)	5 (41.66)	5 (62.5)	.325`
IQR: interquartile ranges (25,75). SD: standard deviation. ATA: auditory tonal average. dB:decibel. HP: Histopathological results., **U mann whitney. `Fisher test. °Chi²			

Cross-tabulating the presence of epithelial inflammation of the perforation remnant versus surgical outcome showed that 50% of patients with epithelial inflammation had surgical failure, while only 25% of those without inflammation of the remnant had failure. Of the total patients with surgical failure, 57.4% had epithelial inflammation.

Regarding the various histological changes reported, the most successful was dystrophic calcification, with 31% of the total successful cases and categorized as absence of epithelial inflammation. While the most unsuccessful was histiocytosis, with 29% of the failed cases and categorized as epithelial inflammation. (Figure 2).




Figure 2. Histopathological results vs surgical outcome.

Fisher's exact test to calculate the relationship between the inflammatory status of the epithelial remnant and surgical outcome yielded a p value of 0.356, with no statistical significance. However, when calculating the OR, the presence of inflammation in the remnant yielded a result of 1.5 (95% CI 0.697–3.22).

The multivariate logistic regression model to determine predictors of surgical failure showed the presence of epithelial inflammation to be the primary variable, with an OR of 3.46 (95% CI 0.37–31.94), followed by median perforation with an OR of 2.6 (95% CI 0.158–43.75), and sclerotic mastoid with an OR of 2.4 (95% CI 0.158–39.09)(Table 3).

Table 3. Multivariate model of factors associated with successful tympanoplasty outcome.

	Beta	Standard error	Odds ratio	CI 95% lower	CI 95% upper
Central perforation	-1,979	1,564	,138	,006	2,962
30-60% perforation	,966	1,435	2,627	,158	43,759
Sclerotic mastoid	,912	1,405	2,489	,158	39,092
Presence of inflammation	1,244	1,133	3,469	,377	31,940

A logistic regression model was performed to determine the predictive factors for the presence of inflammation of the epithelial remnant of the tympanic perforation, finding that the variables that presented a higher probability of risk for presenting inflammation were marginal perforation with an OR of 7.3 (95% CI .145-371.58) and the high OOPS index with an OR of 3.4 (95% CI .65-18.712) (Table 4).

Table 4. Multivariate model for factors associated with the presence of epithelial remnant inflammation.

	Beta	Standard error	Odds ratio	CI 95% lower	CI 95% upper
OOPS Index	1,249	,857	3,489	,650	18,712
Tubal dysfunction present	-3,353	2,638	,035	,000	6,163
Sclerotic mastoid	-4,044	2,545	,018	,000	2,569
Left ear affected	-2,130	1,558	,119	,006	2,518
Marginal perforation	1,994	2,002	7,342	,145	371,586
Perforation 30-60%	-7,506	4,873	,001	,000	7,731
Perforation >60%	-3,854	2,867	,021	,000	5,850
Inadequate ear care	-4,025	3,017	,018	,000	6,606

4. Discussion

The primary objective of this study was to determine whether there was a relationship between epithelial inflammation of the perforation remnant and surgical success. We also evaluated the success rate and factors associated with surgical failure.

Regarding surgical success, our overall success rate was 65%, similar to that reported by Fukuchi (19); however, it is lower than that reported in most studies. Asfaha et al. conducted a study with 92 patients, obtaining an overall success rate of 89.1%. However, in tympanoplasties performed with cartilage grafts, they reported a failure rate of 31.3% (20). This failure rate is higher than ours, which was 25%, considering that all tympanoplasties were performed with cartilage grafts. Chen Chin Kuo evaluated the success of endoscopic tympanoplasty with cartilage, a technique similar to ours, reporting a success rate of 86.3% (21), higher than ours. Age and sex were excluded from the analysis because they were not associated or statistically significant with the surgical outcome or with the inflammatory status of the tympanic membrane. Similarly, Momin et al., in their study of risk factors for the success of tympanoplasty, reported that age was not associated with closure of the perforation (OR 1.04, P=0.31) (22). Our study presented 7 patients with surgical failure, 85% of whom presented tubal dysfunction. Moneir et al. studied the correlation between tubal dysfunction and the results of type 1 tympanoplasty, finding that 60% of patients with tubal dysfunction presented graft retraction and residual perforations, while 20% persisted with the same perforation (23). Among other risk

factors, the presence of postoperative infection was found; the 2 patients who presented this condition suffered surgical failure, while Asfaha reported that of the patients who presented postoperative otorrhea, 28.6% had surgical failure (20). In their logistic regression model, they demonstrated that the most important predictor of failure was the presence of postoperative infection with an odds ratio of 9.6 (95% CI 1.1–88.9) (20), meaning that patients with postoperative otorrhea had an 8.6-fold higher risk of surgical failure compared to those without otorrhea. Fukuchi mentions that the site and size of the perforation did not present statistical significance as a risk factor for the success of tympanoplasty (19), as did Abdelhameed, who studied whether the size of the perforation and graft had an impact on the surgical result by dividing the sample according to the size of the perforation into 3 groups, and then correlating them obtaining non-significant p values (24). In contrast to these results, when performing the multivariate logistic regression model, we found that medium-sized tympanic perforation (30% to 60%) did represent a risk factor for surgical success with an OR of 2.62 (95% CI .158 – 43.7). Regarding epithelial inflammation of the perforation remnant, it was found that of the 7 patients who presented surgical failure at 30 days, 57.4% presented epithelial inflammation, which is close to the hypothesis proposed in the present study, in which it was postulated that the presence of inflammation of the epithelium of the tympanic perforation would influence the success of tympanoplasty in at least 60% of patients with chronic otitis media. On the other hand, Mili et al. reported chronic inflammatory infiltrate in 69.23%, with a surgical success rate of 84.6% (6). Apart from this study, no similar studies were found that evaluated the impact of epithelial inflammation on the surgical outcome of tympanoplasty. However, Leffers, in his article on the immunomodulatory response of epithelial cells in otitis media, evaluated the immunohistochemical changes that occur in the epithelium after inoculation with *Haemophilus influenzae*, finding an increase in IL-1 β , IL-6, IL-8 and TNF α (25). This could open a new line of research in which immunohistochemical studies of the epithelial remnant of the perforation can be performed.

The multivariate logistic regression model showed that the factor that most influenced the success of tympanoplasty was the presence of epithelial inflammation of the perforation remnant, with an odds ratio of 3.4 (95% CI 0.37–31.9), indicating that the presence of epithelial inflammation increases the likelihood of surgical failure by 2.4 times.

In turn, the regression model for predictors of epithelial inflammation indicated that the factors with the greatest risk were the OOPS score, which indicates the state of the mucosa, and the presence of marginal perforation.

Regarding the state of the mucosa, we standardized it with the OOPS score, in which a lower score indicated better condition of the middle ear mucosa. our study showed low scores, that is, ears with little mucosal damage derived from the previous selection of patients, however, when analyzing its association with the presence of inflammation of the remnant of the perforation, it showed an odds ratio of 3.4 (95% CI 0.65-18.7), which means that for each unit that the OOPS index increases, the probability of having inflammation of the epithelial remnant increases 2.4 times. For his part, Han Yu compared the impact of different states of the mucosa with the result of tympanoplasty, he did so by classifying patients into 4 groups: group 1 healthy mucosa, group 2 mucosa with little edema, group 3 with moderate edema and group 4 with severe edema, finding that most cases of residual perforation occurred in groups 3 and 4 with a percentage of 91.1% and 84.6% (26) respectively.

5. Conclusions

Epithelial inflammation of the perforation remnant affects surgical success in more than 50% of patients. However, the wide confidence intervals suggest that the sample size is insufficient, so further research could be carried out and the sample expanded to obtain more statistically reliable results.

Among the factors that favor the presence of epithelial inflammation of the perforation remnant are the presence of a marginal perforation, a perforation size greater than 30%, and a high OOPS index. This leads us to conclude that the presence of epithelial inflammation of the perforation remnant negatively impacts surgical outcome. However, it is not the only factor involved in tympanoplasty failure. It would be worth extending the study and taking into account factors such as disease progression time and the presence of comorbidities that were not considered.

References

- Campos N LA, Barron S MB, Fajardo D G, México DF. Otitis media aguda y crónica, una enfermedad frecuente y evitable. *Rev Fac Med UNAM*. 2014;57(1):5–14.
- Schilder AGM, Chonmaitree T, Cripps AW, Rosenfeld RM, Casselbrant ML, Haggard MP, et al. Otitis media. *Nature* [Internet]. 2016 [citado el 26 de febrero de 2023];2:1–18. Disponible en: www.nature.com/nrdp
- Fukuchi I, Cerchiari DP, Garcia E, Rezende CEB, Rapoport PB. Tympanoplasty: Surgical results and a comparison of the factors that may interfere in their success. *Braz J Otorhinolaryngol*. marzo de 2006;72(2):267–71.
- Marchisio P, Esposito S, Picca M, Baggi E, Terranova L, Orenti A, et al. Prospective evaluation of the aetiology of acute otitis media with spontaneous tympanic membrane perforation. *Clin Microbiol Infect* [Internet]. 2017 [citado el 17 de octubre de 2023];23:486.e1e-486.e6. Disponible en: <http://dx.doi.org/10.1016/j.cmi.2017.01.010>
- World Health Organization. Geneva World Health Organ . 2004 [citado el 20 de octubre de 2023]. Chronic suppurative otitis media: burden of illness and management options. Disponible en: https://iris.who.int/handle/10665/42941?search-result=true&query=tympanic+perforation&scope=&rpp=10&sort_by=score&order=desc
- Mili MK, Hoque SR, Sarmah M, Phookan J. Results of tympanoplasty in reference to the condition of middle ear mucosa and edge of perforation in safe chronic otitis media. *International Journal of Otorhinolaryngology and Head and Neck Surgery*. el 25 de enero de 2022;8(2):124–8.
- Sharma K, Manjari M, Salaria N. Middle Ear Cleft in Chronic Otitis Media: A Clinicohistopathological Study. *Indian Journal of Otolaryngology and Head & Neck Surgery* [Internet]. el 7 de diciembre de 2013 [citado el 11 de febrero de 2023];65(S3):493–7. Disponible en: <http://link.springer.com/10.1007/s12070-011-0372-x>
- Coronado Terrazas Javier. Eficacia del cierre de perforación con injerto de pericondrio en pacientes sometidos a timpanoplastía. (Trabajo de grado de especialización) Universidad Nacional Autónoma de México, Coordinación General de Estudios de Posgrado, UNAM. 2019;
- Luers JC, Hüttenbrink KB. Surgical anatomy and pathology of the middle ear. *J Anat*. el 1 de febrero de 2016;228(2):338–53.
- Ma Y, Zhao H, Zhou X. Topical treatment with growth factors for tympanic membrane perforations: progress towards clinical application. *Acta Otolaryngol* [Internet]. 2002 [citado el 17 de octubre de 2023];122(6):586–99. Disponible en: <https://pubmed.ncbi.nlm.nih.gov/12403120/>
- Gladstone HB, Jackler RK, Varav K. Tympanic membrane wound healing. An overview. *Otolaryngol Clin North Am* [Internet]. 1995 [citado el 20 de octubre de 2023];28(5):913–32. Disponible en: <https://pubmed.ncbi.nlm.nih.gov/8559580/>
- Sainsbury E, Amaral R do, Blayney AW, Walsh RMC, O'Brien FJ, O'Leary C. Tissue engineering and regenerative medicine strategies for the repair of tympanic membrane perforations. *Biomaterials and Biosystems*. el 1 de junio de 2022;6:100046e1-10046e10.
- Baudouin R, Denoyelle F, Simon F. Shedding light on the tympanic membrane: A brief history of the description and understanding of its anatomy. *Journal of Laryngology and Otology* [Internet]. el 25 de febrero de 2022 [citado el 17 de noviembre de 2023];136(2):97–102. Disponible en: <https://hal.science/hal-03625841>
- Johnson A, Hawke M. The function of migratory epidermis in the healing of tympanic membrane perforations in guinea-pig. A photographic study. *Acta Otolaryngol* [Internet]. 1987 [citado el 20 de octubre de 2023];103(1–2):81–6. Disponible en: <https://pubmed.ncbi.nlm.nih.gov/3564931/>
- Reijnen CJ, Kuijpers W. The healing pattern of the drum membrane. *Acta Otolaryngol Suppl* [Internet]. 1971 [citado el 20 de octubre de 2023];287:1–74. Disponible en: <https://pubmed.ncbi.nlm.nih.gov/4338658/>
- Wang AY, Shen Y, Wang JT, Friedland PL, Atlas MD, Dilley RJ. Animal models of chronic tympanic membrane perforation: A 'time-out' to review evidence and standardize design. *Int J Pediatr Otorhinolaryngol*. el 1 de diciembre de 2014;78(12):2048–55.
- Mokoyan Z, Svistushkin V, Zolotova A, Svistushkin M. Chronic tympanic membrane perforation: Histopathological evidence of the experimental model. *Int J Pediatr Otorhinolaryngol* [Internet]. el 1 de

- diciembre de 2021 [citado el 17 de octubre de 2023];151:110964. Disponible en: <https://www.sciencedirect.com/science/article/abs/pii/S0165587621003578?via%3Dihub>
18. Sadé J, Eliezer N. Secretory otitis media and the nature of the mucociliary system. *Acta Otolaryngol.* 1970;70(5–6):351–7.
 19. Fukuchi I, Cerchiari DP, Garcia E, Rezende CEB, Rapoport PB. Tympanoplasty: Surgical results and a comparison of the factors that may interfere in their success. *Braz J Otorhinolaryngol.* marzo de 2006;72(2):267–71.
 20. Asfaha FG, Tesfa BB, Gebremariam LW, Geremew A, Temam S, Abreha GF. Tympanoplasty in Northern Ethiopia: success rates, failure factors, and audiometric improvements. *BMC Res Notes.* el 3 de febrero de 2025;18(1):52.
 21. Chen CK, Hsu HC, Wang M. Endoscopic tympanoplasty with post-conchal perichondrium in repairing large-sized eardrum perforations. *European Archives of Oto-Rhino-Laryngology.* el 1 de diciembre de 2022;279(12):5667–74.
 22. Momin ZK, Yancey KL, Mitton TJ, Kutz JW. Patient Factors Predicting Success in Lateral Graft Tympanoplasty. *Otology & Neurotology Open.* septiembre de 2022;2(3):e015.
 23. Moneir W, El-Kholy NA, Ali AI, Abdeltawwab MM, El-Sharkawy AAR. Correlation of Eustachian tube function with the results of type 1 tympanoplasty: a prospective study. *European Archives of Oto-Rhino-Laryngology.* el 1 de abril de 2023;280(4):1593–601.
 24. Abdelhameed W, Rezk I, Awad A. Impacto do tamanho do enxerto de cartilagem no sucesso da timpanoplastia. *Braz J Otorhinolaryngol.* el 1 de septiembre de 2017;83(5):507–11.
 25. Leffers D, Penxova Z, Kempin T, Därr M, Fleckner J, Hollfelder D, et al. Immunomodulatory Response of the Middle Ear Epithelial Cells in Otitis Media. *Otology and Neurotology.* el 1 de marzo de 2024;45(3):E248–55.
 26. Han Y, Yang R, Mao X, Li R, Song Y, Shi H, et al. Comparison of the Impacts of Different Middle Ear Mucosal Conditions on Type I Tympanoplasty Outcomes. *J Otolaryngol Head Neck Surg.* el 1 de enero de 2024;53.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.