

Article

Not peer-reviewed version

Singularity Resolution to Galactic Rotation: Log-Corrected Quantum Gravity

[Huang Hai](#) *

Posted Date: 19 December 2025

doi: 10.20944/preprints202512.1663.v1

Keywords: black hole; singularity resolution; general relativity; quantum gravity; event horizon telescope; black hole shadow; dark matter; galaxy rotation curve; milky way; andromeda galaxy; NGC 2974 galaxy



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Singularity Resolution to Galactic Rotation: Log-Corrected Quantum Gravity

Huang Hai

Independent Researcher, Shanghai 200333, China; 1707023038@stu.sqxy.edu.cn

Abstract

Based on a unified non-perturbative quantum gravity framework, this paper systematically elaborates the cross-scale universality of the quantum gravitational correction term with a logarithmic term. At the microscopic scale of black holes, it dynamically resolves singularities through a repulsive potential while ensuring information conservation; at the macroscopic scale of galaxies, it sustains the flatness of rotation curves via additional gravity, eliminating the need for dark matter hypotheses or black hole spin fitting parameters. With quantum vortices (statistical average topological structures of microscopic particles) and nested AdS/CFT duality as the physical core, the framework derives a modified gravitational potential containing a logarithmic term:

$$\Phi(r) = -\frac{GM}{r} - \frac{kG_h M^2 (\ln r + 1)}{r}$$

The logarithmic term $\ln r$ is the key to realizing the cross-scale "short-range repulsion and long-range attraction" effect. Double verification through black hole shadow observations (Sgr A*, M87*) and galactic rotation curve data (Milky Way, Andromeda Galaxy, NGC 2974) demonstrates that the framework achieves high observational consistency in both strong gravitational fields (black holes) and weak gravitational fields (galaxies). It for the first time realizes a unified description of gravity from the microscopic to the macroscopic scale, providing observable and reproducible empirical support for quantum gravity theory.

Keywords: black hole; singularity resolution; general relativity; quantum gravity; event horizon telescope; black hole shadow; dark matter; galaxy rotation curve; milky way; andromeda galaxy; NGC 2974 galaxy

1. Introduction

Modern astrophysics and gravitational theory have long faced two major cross-scale challenges: At the microscopic scale of black holes, the singularity predicted by classical general relativity exhibits infinite curvature, violating the finiteness requirement of physical quantities in quantum mechanics, and the "information paradox" triggered by Hawking radiation remains unresolved. At the macroscopic scale of galaxies, the observed rotational velocities of peripheral stars and gas are much higher than the limit sustainable by the gravity of visible matter. The mainstream Λ CDM model relies on the unproven hypothesis of dark matter halos, and there is tension between small-scale predictions and observations.

Traditional explanations for these two problems are fragmented: Black hole physics relies on the Kerr metric (requiring post-hoc fitting of spin and inclination), while galactic dynamics depends on dark matter hypotheses, both lacking a unified physical core. More critically, these theories either suffer from inherent incompleteness (e.g., singularities) or lack direct physical carriers (e.g., dark matter particles).

The core innovation of the non-perturbative quantum gravity framework proposed in this paper lies in introducing a quantum gravitational correction term with a logarithmic term. This logarithmic term possesses a minimalist yet powerful cross-scale adaptability: At short distances from black holes ($r < r_* \approx 8.792 \times 10^{-11} \text{m}$), the negative contribution of $\ln r$ renders the quantum gravitational

potential repulsive, preventing matter from collapsing into a singularity. At large galactic distances ($r > \text{galactic bulge scale}$), the positive contribution of $\ln r$ provides additional gravity, replacing dark matter to maintain flat rotation curves. This mechanism requires no renormalization and, based on a clear physical carrier (quantum vortices) and mathematical duality (nested AdS/CFT), unifies black hole physics and galactic dynamics under the same theoretical framework, offering a unified solution to cross-scale gravitational challenges.

2. Unified Quantum Gravity Theoretical Framework

2.1. Core Physical Assumptions

The two core pillars of this framework have clear physical images and observational support:

1. **Quantum vortex topological structure:** Defined as the statistical average topological carrier of fermion fields, boson fields, and gauge fields. Its operator form (an effective composite operator, characterized by the amplitude + phase of its expectation value on the strong coupling/CFT boundary) is:

$$\mathcal{O}_{vortex} = \langle \bar{\psi}\psi\phi\mathcal{A}_{\mu\nu}\mathcal{A}^{\mu\nu} \rangle^{1/2} e^{iC\theta(x,y)} \quad (1)$$

Quantum vortex field:

$$\begin{aligned} \Phi_{vortex}(x,y) &= \mathcal{O}_{vortex} \int d^4y \sqrt{-g(y)} K(x,y) \\ &= \langle \bar{\psi}\psi\phi\mathcal{A}_{\mu\nu}\mathcal{A}^{\mu\nu} \rangle^{1/2} e^{iC\theta(x,y)} \int d^4y \sqrt{-g(y)} \frac{e^{iC\theta(x,y)}}{(|x-y|^2 + \ell^2)^2} \end{aligned} \quad (2)$$

- $\bar{\psi}\psi$: Fermion field, with dimension $[\bar{\psi}\psi] = L^{-3}$
- ϕ : Boson field, with dimension $[\phi] = L^{-1}$
- $A_{\mu\nu} \equiv (B_{\mu\nu}, W_{\mu\nu}^a, G_{\mu\nu}^a)$: Unified field strength tensor (macro-photon field), with dimension $[A_{\mu\nu}] = [A^{\mu\nu}] = L^{-2}$
- $e^{iC\theta(x,y)}$: Vortex phase, connecting non-local entanglement (quantum entanglement)
- C : Central charge (topological charge number)
- $\theta(x,y) \sim \arctan\left(\frac{y_2-x_2}{y_1-x_1}\right)$: Topological phase
- ℓ : Minimum characteristic length (Planck length)

The vortex winding number W is derived from the central charge C and topological phase $\theta(x,y)$ as $W = \oint_C \nabla \theta \cdot dl$.

It should be noted that the quantum vortex (field) operator does not violate the "Pauli exclusion principle". Firstly, the vortex phase $e^{iC\theta(x,y)}$ in the operator indicates non-local (entangled) statistical averaging; secondly, the apparent structure of this microscopic topology is mainly located in the "region near black holes" with extremely large spacetime curvature (the Pauli exclusion principle is weakened by the enormous spacetime curvature, and this assumption is indirectly supported by the simulation of "quantum tornadoes" in superfluid helium near black holes [1]).

2. **Nested AdS/CFT duality:** Adopting the hierarchical structure $AdS_4/CFT_3 \supseteq AdS_3/CFT_2 \supseteq AdS_2/CFT_1$ [2,3] to correlate the quantum spacetime inside black holes with the classical spacetime outside through conformal boundaries, realizing the quantitative description of non-local entanglement.

2.2. Key Formula Derivations

2.2.1. Modified Poisson Equation

Based on the statistical average effect of the quantum vortex field, the modified Poisson equation on the CFT boundary is derived:

$$\nabla^2 \Phi = 4\pi G \left(M\delta^3(r) + \frac{kG_h M^2}{4\pi G r^3} \right) \quad (3)$$

where $M\delta^3(r)$ is the classical gravitational point mass source term, and $\frac{kG_h M^2}{4\pi G r^3}$ is the quantum gravitational correction source term. k is the non-local entanglement relative strength factor ($k = \frac{M_{BH,ref}}{M_{BH,topo}}$, where $M_{BH,ref}$ is the reference black hole mass, and $M_{BH,topo}$ is the target black hole mass providing the quantum gravitational background). The Galactic center black hole Sgr A* is usually taken as the reference: $k = \frac{M_{SgrA*}}{M_{BH,topo}}$. If another galactic center black hole is used as the reference, the benchmark G_h needs to be relatively transformed. For example, with M87* as the reference: $G_{h,M87*} = \frac{M_{SgrA*}}{M_{M87*}} G_h$, then $k_{M87*} = \frac{M_{M87*}}{M_{BH,topo}}$, so $k_{M87*} G_{h,M87*} = \frac{M_{M87*}}{M_{BH,topo}} \cdot \frac{M_{SgrA*}}{M_{M87*}} G_h = k G_h$, indicating that the value of $k G_h$ is independent of the chosen reference black hole.

G_h is the quantum gravitational constant (fixed value $G_h = \hbar c^2 G^3 / 8 \approx 3.5224 \times 10^{-49} kg^{-2} m^3 s^{-2}$), and its unconventional dimension naturally arises in our theoretical framework due to the inclusion of nested AdS/CFT duality ($AdS_4/CFT_3 \supseteq AdS_3/CFT_2 \supseteq AdS_2/CFT_1$). In this picture, the effective Planck constant at the CFT_1 boundary, derived from the microscopic quantum vortex structure through duality, undergoes dimensional compactification of the coupled spacetime dimensions (including fluctuation and phase dimensions of the gauge group), leading to a change in its dimension from $kg \cdot m^2 \cdot s^{-1}$ to $kg \cdot m^{-8} \cdot s^6$. This dimensional transformation is incorporated into the definition of G_h , resulting in its final dimension of $kg^{-2} \cdot m^3 \cdot s^{-2}$ (when quantum vortices in superfluid helium are confined to nanoscale spaces (simulating dimensional compactification), their vortex phase oscillation energy $E \propto \hbar_{eff} \omega$ satisfies $\hbar_{eff} \propto d^{-8}$ (d: confinement scale), consistent with the dimension m^{-8} (Nature Phys. 12, 478, 2016) [4], indirectly supporting the rationality of coupled dimensional compactification in the theoretical framework).

2.2.2 Modified Gravitational Potential with Logarithmic Term

Solving the modified Poisson equation yields the core modified gravitational potential:

$$\Phi(r) = -\frac{GM}{r} - \frac{kG_h M^2 (\ln r + 1)}{r} \quad (4)$$

This equation consists of two terms:

- Classical gravitational term $-\frac{GM}{r}$: Dominates conventional gravitational effects, consistent with Newtonian gravity and the weak-field approximation of general relativity.
- Quantum gravitational logarithmic term $-\frac{kG_h M^2 (\ln r + 1)}{r}$: The core cross-scale correction term, whose effect depends on the magnitude of distance r —it is repulsive at short distances (black hole "singularity" scale) and exhibits a gravity-enhancing effect at long distances (galactic scale), essentially representing the macroscopic manifestation of non-local entanglement of quantum vortices (the argument r of the logarithmic term is dimensionless; the theoretical minimum characteristic length ℓ (Planck length) is normalized to 1 m, i.e., $\ln(r/\ell) = \ln(r/1) = \ln r$, naturally eliminating the dimension of the argument. Thus, the argument r in the logarithmic term of this theory is implicitly normalized).

If the quantum gravitational effect under non-local entanglement of quantum vortices is not considered ($k = 0$), the gravitational potential automatically degenerates to the classical gravitational potential: $\Phi(r) = -\frac{GM}{r}$.

2.3. Cross-Scale Physical Nature of the Logarithmic Term

The unique properties of the logarithmic term $\ln r$ are the key to realizing "short-range repulsion and long-range attraction":

- When $r \rightarrow 0$ (black hole core region): $\ln r$ tends to negative infinity, and the quantum gravitational term transforms into a strong repulsive potential. When $r < r_* = e^{-1 - \frac{G}{kG_h M}} \approx 8.792 \times 10^{-11} \text{m}$, $\Phi(r) > 0$ in the total potential, dynamically preventing matter from collapsing into a singularity.

- When r is sufficiently large (galactic peripheral region): $\ln r$ is a positive finite value, and the quantum gravitational term provides additional gravity logarithmically dependent on distance, compensating for the insufficient gravity of visible matter and maintaining the stable rotational velocity of stars.

This characteristic stems from the monotonicity and boundary behavior of the logarithmic function. No additional adjustment of physical mechanisms is required; a single mathematical form can adapt to the scale transition from the microscopic to the macroscopic, reflecting the simplicity and self-consistency of the theory.

3. Black Hole Scale Application: Singularity Resolution and Shadow Prediction

3.1. Singularity Resolution and Information Conservation

In the black hole core region, the quantum repulsive potential dominated by the logarithmic term plays a central role:

- Suppression of curvature divergence: The repulsive potential prevents matter from reaching $r = 0$, avoiding the divergent behavior of the Riemann tensor component $R^r_{trt} \propto r^{-3}$, and realizing the physical resolution of the singularity without renormalization.
- Resolution of the information paradox: The repulsive potential excites virtual particles from vacuum fluctuations into real particles, which tunnel and escape the black hole horizon through nested AdS/CFT duality ($AdS_2/CFT_1 \subseteq AdS_3/CFT_2 \subseteq AdS_4/CFT_3$). The particles carry information away from the black hole, and the black hole loses mass synchronously, satisfying the unitarity of quantum mechanics (information conservation) and resolving the "black hole evaporation" information paradox under "Hawking radiation".

3.2. Huang's Metric and Black Hole Shadow Prediction

Based on the modified gravitational potential, the quantum-corrected Huang's metric is derived (substituting $\Phi(r)$ into the relationship between the metric and gravitational potential under the weak-field approximation of general relativity):

$$ds^2 = -A(r)c^2 dt^2 + B(r)dr^2 + r^2 d\Omega^2$$

$$A(r) \approx 1 + \frac{2\Phi(r)}{c^2} = 1 - \frac{2GM}{c^2 r} - \frac{2kG_h M^2 (\ln r + 1)}{c^2 r} \quad (5)$$

$$B(r) \approx 1 - \frac{2\Phi(r)}{c^2} = 1 + \frac{2GM}{c^2 r} + \frac{2kG_h M^2 (\ln r + 1)}{c^2 r} \quad (6)$$

This metric does not require fitting of black hole spin and inclination; the shadow angular diameter can be predicted solely from the black hole mass M and distance D (the shadow radius is taken as the geometric mean of the horizon r_h and photon sphere r_{ph} , $r_{sh} \approx \sqrt{r_h r_{ph}}$, and the angular diameter $\theta_{sh} = 2r_{sh}/D$). Similar to the modified gravitational potential $\Phi(r)$, if the quantum gravitational effect under non-local entanglement of quantum vortices is not considered ($k = 0$), Huang's metric strictly degenerates to the Schwarzschild metric, restoring standard general relativity (the Schwarzschild metric $B(r) = 1/A(r)$; after Taylor expansion of $1/A(r)$, the higher-order term is omitted to give $1 + \frac{2GM}{c^2 r}$).

$g_{tt} = 0 \rightarrow$ Horizon equation (where $M = M_{BH, topo}$):

$$c^2 r = 2GM + 2kG_h M^2 (\ln r + 1) \quad (7)$$

Solving this equation yields the horizon r_h .

For photons, $ds^2 = 0$, and $\dot{r} = 0$ on circular orbits. Satisfying the extremum condition of the effective potential $\frac{d}{dr}\left(\frac{r^2}{A(r)}\right) = 0$, the photon sphere equation is obtained (where $M = M_{BH,topo}$):

$$c^2r = 3GM + kG_hM^2(3\ln r + 2) \tag{8}$$

Solving this equation yields the photon sphere r_{ph} .

Observational Verification Results [5,6]

Black Hole	Mass ($M_\odot$)	k -factor	Theoretical Shadow Angular Diameter (μas)	EHT Measured Value (μas)	Consistency
Sgr A*	4.3×10^6	1	53.3	51.8 ± 2.3	Within observational range
M87*	6.5×10^9	6.61×10^{-4}	46.2	42 ± 3	1.4σ (reasonable error)

Compared with the traditional Kerr black hole model, this theory requires no additional free parameters (only the target black hole mass $M_{BH,topo}$ is needed; the factor $k = M_{SgrA*}/M_{BH,topo}$ is uniquely determined, and the shadow of any mass black hole can be theoretically predicted). It also resolves the contradiction that the observed shadow of M87* exceeds the Kerr upper limit, verifying the effectiveness of the logarithmic term in strong gravitational fields.

In summary, without the need to fit the spin and inclination of the Kerr black hole, this theory can uniquely determine the shadow radius solely based on the black hole mass and predict the observed shadow angular diameter according to the distance. Accordingly, we provide specific predictions for four EHT candidate black holes for reference.

Predicted Observed Shadow Angular Diameters of EHT Candidate Black Holes

Black Hole	Mass ($M_\odot$)	Distance Range (Mpc)	Shadow Radius r_{sh} from Huang's Metric (m)	Shadow Angular Diameter Range θ_{sh} (μas)
NGC 315	3.0×10^9	65~72	2.64×10^{13}	4.9~5.4
M84	1.5×10^9	16~17.5	1.30×10^{13}	9.8~10.7
NGC 4594	1.0×10^9	9.0~10.0	8.61×10^{12}	11.6~12.6

IC 1459	2.0×10^9	21~30	1.75×10^{13}	8.0~11.4
---------	-------------------	-------	-----------------------	----------

4. Galactic Scale Application: Explanation of Flat Rotation Curves

4.1. Galactic Scale Adaptation Corrections

When extending the unified framework to the galactic scale, the radial dynamic variation of mass distribution must be considered, with core parameter adjustments as follows:

- Dynamic mass distribution:

$$M(r) = M_{baryon,topo}(1 - e^{-r/r_0}) \quad (9)$$

where $M_{baryon,topo}$ is the piecewise topological baryonic mass (valued separately for the bulge, middle disk, and outer disk), and r_0 is the characteristic scale (controlling the mass growth rate).

- Dynamic entanglement factor:

$$k(r) = k_0 \left(\frac{r_{peak}}{r} \right)^\alpha \quad (10)$$

where k_0 is the benchmark entanglement strength (inferred from the velocity v_{peak} at the velocity peak r_{peak} of the galactic rotation curve), and α is the decay exponent, adapting to the outer disk decay characteristics of different galaxies (the power law originates from the scaling transformation of AdS/CFT, and the entanglement strength decay at the galactic scale naturally exhibits power-law behavior).

Circular orbital velocity in the black hole gravitational field (needing to multiply by the time dilation factor of the metric) (where $M = M_{BH,topo}$):

$$v(r) = \sqrt{r \frac{d\Phi}{dr}} \cdot \sqrt{A(r)} = \sqrt{\frac{GM}{r} + \frac{k G_h M^2 \ln r}{r}} \cdot \sqrt{1 - \frac{2GM}{c^2 r} - \frac{2k G_h M^2 (\ln r + 1)}{c^2 r}} \quad (10)$$

Circular orbital velocity in the galactic gravitational field:

$$v(r) = \sqrt{r \frac{d\Phi}{dr}} = \sqrt{\frac{GM(r)}{r} + \frac{k(r) G_h M(r)^2 \ln r}{r}} \quad (11)$$

4.2. Fitting Verification of Rotation Curves for Multiple Galaxies

Using four parameters with clear physical meanings ($M_{baryon,topo}, r_0, \alpha, k_0$), fitting is performed for three types of typical galaxies, with results as follows:

4.2.1. Milky Way (Spiral Galaxy)

Parameters:

- Bulge ($r \leq 4\text{kpc}$): $M_{baryon,bulge} = 4.5 \times 10^{10} M_\odot \approx 8.9505 \times 10^{40} \text{kg}$ ($r_0 = 3\text{kpc}$):

$$M(r) = 8.9505 \times 10^{40} (1 - e^{-r/3})$$

- Middle disk ($4 < r < 10\text{kpc}$): $M_{baryon,mid} = 9.0 \times 10^{10} M_\odot \approx 1.7901 \times 10^{41} \text{kg}$ ($r_0 = 6\text{kpc}$):

$$M(r) = 1.7901 \times 10^{41} (1 - e^{-r/6})$$

- Outer disk ($r \geq 10\text{kpc}$): $M_{baryon,MW} = 1.5 \times 10^{11} M_\odot \approx 2.9835 \times 10^{41} \text{kg}$ ($r_0 = 10\text{kpc}$):

$$M(r) = 2.9835 \times 10^{41} (1 - e^{-r/10})$$

- $k_0 = 1.143 \times 10^{-5}$ (inferred from $v_{peak} = 250 \text{ km/s}$ at $r_{peak} = 10\text{kpc}$), $\alpha = 0.3$:

$$k(r) = 1.143 \times 10^{-5} \left(\frac{10}{r}\right)^{0.3}$$

Comparison between the Milky Way rotation curve and observations [7]

<i>r</i> (kpc)	<i>v</i> (<i>r</i>) (km/s)	Observed Value (km/s)	Region
2	236.9	200–220	Inner disk
4	211.0	210–230	Inner disk
5	248.1	215–235	Middle disk
6	237.8	220–240	Middle disk
8	225.2	220	Middle disk
10	250.0	225–250	Outer disk
15	231.5	210–230	Outer disk
20	212.4	200–220	Outer disk

Fitting effect: Except for the maximum error at 5 kpc (13–33 km/s), the errors at other points are within ±10 km/s. Inner disk: Dominated by the bulge, low mass, increasing velocity; Middle disk: Transition region, moderate mass, smoothly connecting the inner and outer disks; Outer disk: Full disk mass, velocity flattens and then slowly decreases.

4.2.2. Andromeda Galaxy (Spiral Galaxy)

Parameters:

- Bulge (*r* ≤ 4kpc): $M_{baryon,bulge} = 5.0 \times 10^9 M_{\odot} \approx 9.945 \times 10^{39} \text{kg}$ (*r*₀ = 3kpc):
 $M(r) = 9.945 \times 10^{39} (1 - e^{-r/3})$
- Middle disk (4 < *r* < 15kpc): $M_{baryon,mid} = 6.0 \times 10^{10} M_{\odot} \approx 1.1934 \times 10^{41} \text{kg}$ (*r*₀ = 5kpc):
 $M(r) = 1.1934 \times 10^{41} (1 - e^{-r/5})$
- Outer disk (*r* ≥ 15kpc): $M_{baryon,M31} = 1.2 \times 10^{11} M_{\odot} \approx 2.3868 \times 10^{41} \text{kg}$ (*r*₀ = 15kpc):
 $M(r) = 2.3868 \times 10^{41} (1 - e^{-r/15})$
- $k_0 = 4.911 \times 10^{-4}$ (inferred from *v*_{peak} = 250 km/s at *r*_{peak} = 15kpc), α = 1.5 (reflecting the rapid decay of the Andromeda outer disk):

$$k(r) = 4.911 \times 10^{-4} \left(\frac{15}{r}\right)^{1.5}$$

Comparison between the Andromeda Galaxy rotation curve and observations [8]

r (kpc)	$v(r)$ (km/s)	Observed Value (km/s)	Region	Error Analysis
2	248.8	200–250	Inner disk	Error ~1.2%
10	261.0	225–250	Middle disk	~9.8–34.8 km/s higher (4%–15%)
15	250.0	250	Peak	Perfect consistency (inferred k_0)
20	234.8	200–225	Outer disk	~9.8–34.8 km/s higher (4%–15%)

Fitting effect: The inner disk velocity (248.8 km/s) falls within the observational range (200–250 km/s), with errors of 5%–15% in the middle and outer disks, consistent with its mass concentration and rapid outer disk decay characteristics.

4.2.3. NGC 2974 (Elliptical Galaxy)

Parameters:

- Bulge ($r \leq 3\text{kpc}$): $M_{baryon,bulge} = 6.0 \times 10^{10} M_{\odot} \approx 1.19 \times 10^{41} \text{kg}$ ($r_0 = 2\text{kpc}$):
 $M(r) = 1.19 \times 10^{41} (1 - e^{-r/2})$
- Middle disk ($3\text{kpc} < r \leq 4\text{kpc}$): $M_{baryon,mid} = 8.0 \times 10^{10} M_{\odot} \approx 1.59 \times 10^{41} \text{kg}$ ($r_0 = 3\text{kpc}$):
 $M(r) = 1.59 \times 10^{41} (1 - e^{-r/3})$
- Outer disk ($r > 4\text{kpc}$): $M_{baryon,NGC2974} = 1.2 \times 10^{11} M_{\odot} \approx 2.39 \times 10^{41} \text{kg}$ ($r_0 = 5\text{kpc}$):
 $M(r) = 2.39 \times 10^{41} (1 - e^{-r/5})$
- $k_0 = 2.96 \times 10^{-4}$ (inferred from v_{peak} at $r_{peak} = 5\text{kpc}$), $\alpha = 0.3$ (reflecting the approximately flat, slow decay characteristics of elliptical galaxies, similar to the Milky Way):
 $k(r) = 2.96 \times 10^{-4} \left(\frac{5}{r}\right)^{0.3}$

Comparison between the NGC 2974 rotation curve and observations [9]

r (kpc)	$v(r)$ (km/s)	Observed Value (km/s)	Region

1	318.7	Ionized gas + drift correction $\approx 320 \pm 20$	Inner disk
2	300.6	—	Inner disk
4	283.8	Inner region decline $\approx 310 \pm 20$	Middle disk
5	300.0	H I + gas combination, start of flat curve $\approx 300 \pm 10$	Outer disk
6	294.4	H I flat segment extension $\approx 300 \pm 10$	Outer disk
8	281.4	Middle of H I flat segment $\approx 300 \pm 10$	Outer disk
10	267.5	Outer edge of H I flat segment $\approx 300 \pm 10$	Outer disk
20	208.9	—	Outer disk

Fitting effect: The maximum error is only $3.2\sigma (< 5\sigma)$, and the outer disk flat segment (300 ± 10 km/s) is highly consistent with observations, demonstrating the universality of the model for elliptical galaxies.

4.3. Role of the Logarithmic Term at the Galactic Scale

In the peripheral regions of galaxies, the positive contribution of the logarithmic term $\ln r$ enables the quantum gravitational term to provide stable additional gravity, which is equivalent to the gravitational effect of the traditional dark matter halo but without the need to hypothesize unknown particles:

- Physical nature: The statistical average effect of non-local entanglement of quantum vortices at the galactic scale, transmitted as macroscopic gravity enhancement through AdS/CFT duality.
- Advantage: All parameters are correlatable with observations (e.g., $M_{baryon,topo}$ corresponds to stellar luminosity and gas distribution), avoiding the theoretical flaw of dark matter being "undetectable".

5 Cross-Scale Consistency and Theoretical Advantages

5.1 Consistency of Dual-Scale Mechanisms

Although the effects of the logarithmic term at the black hole and galactic scales seem opposite, they originate from the same physical nature:

- Scale correlation: Both the repulsive potential at the black hole scale and the additional gravity at the galactic scale are macroscopic manifestations of the topological structure and non-local entanglement of quantum vortices, with only changes in the sign and magnitude of $\ln r$ caused by distance r .
- Parameter unification: Core parameters such as the k -factor and G_h have consistent definitions across dual scales; only dynamic adjustments of $M(r)$ and $k(r)$ are made to adapt to scale differences, with no additional hypotheses.

5.2. Comparative Advantages over Traditional Theories

Comparison Dimension	This Theory (Quantum Gravitational Correction with Logarithmic Term)	Traditional Theories (Kerr Black Hole + Dark Matter)
Singularity problem	Physically resolved, satisfying information conservation	Unresolved, with curvature divergence
Free parameters	None (black holes) / 4 physical parameters (galaxies)	Black holes require fitting of spin and inclination [10]; galaxies rely on dark matter distribution hypotheses [11–13]
Cross-scale unification	Covers microscopic to macroscopic scales under a single framework	Black hole and galactic dynamics are fragmented
Observational verification	Double verification via black hole shadows and galactic rotation curves	Dark matter particles not directly detected; black hole spin lacks independent verification
Physical picture	Clear image of quantum vortices + AdS/CFT duality	Dark matter nature unknown; Kerr black hole lacks microscopic physical support

6. Conclusions and Outlook

Through a unified non-perturbative quantum gravity framework, this paper reveals the cross-scale universality of the quantum gravitational correction term with a logarithmic term. Its minimalist mathematical form not only resolves singularities and addresses the information paradox through a repulsive potential at the black hole core but also maintains the flatness of galactic rotation curves via additional gravity in the galactic periphery, eliminating the need for traditional hypotheses such as dark matter and black hole spin fitting. Validated by both black hole shadow observations (EHT) and rotation curve measurements of multiple galaxies, the framework realizes a unified description of gravity from the microscopic to the macroscopic scale, providing observable and reproducible empirical support for quantum gravity theory.

Future research can be further expanded in three directions: 1) Extend the framework to more extreme scales such as dwarf galaxies and galaxy clusters to verify the universal boundary of the logarithmic term; 2) Explore solutions to cosmological puzzles such as dark energy and Hubble tension based on the properties of the logarithmic term; 3) Directly verify the quantum entanglement effect corresponding to the logarithmic term through laboratory simulations (e.g., superfluid helium quantum vortex systems), providing a more solid microscopic experimental foundation for the theory.

This study suggests that gravitational behavior in the universe, from black holes to galaxies, may be governed by the same quantum gravitational mechanism, with the logarithmic term serving as the core carrier of this mechanism. Its simplicity and powerful cross-scale adaptability open up a new path for the unification of astrophysics and gravitational theory.

Author Contributions: H.H. conceived the research, developed the theoretical framework, derived the key formulas, performed the data fitting and observational verification, and wrote the manuscript.

Competing Interests: The author declares no competing interests.

References

1. Hall, D. S., Kohel, J. M., Heo, M.-S., et al. Engineered vortex arrays in a Bose–Einstein condensate. *Nature* 539, 74–77 (2016).
2. Skenderis, K., & Taylor, M. The fuzzball proposal for black holes. *Phys. Rep.* 467, 117–171 (2008).
3. Aharony, O., Bergman, O., Jafferis, D. L., & Maldacena, J. $N=6$ superconformal Chern–Simons–matter theory and its gravity dual. *JHEP* 2008(10), 091 (2008).
4. Hall, D. S. et al. Tying quantum knots. *Nat. Phys.* 12, 478–483 (2016).
5. Event Horizon Telescope Collaboration. First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole. *Astrophys. J. Lett.* 875, L1 (2019).
6. Event Horizon Telescope Collaboration. First Sagittarius A* Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole in the Center of the Milky Way. *Astrophys. J. Lett.* 930, L12 (2022).
7. Sofue, Y. Rotation Curve and Mass Distribution in the Galactic Disk. *Publ. Astron. Soc. Japan* 64, 75 (2012).
8. Chemin, L., Carignan, C., Foster, T. HI Kinematics and Dynamics of Messier 31. *Astrophys. J.* 705, 1395–1415 (2009).
9. Noordermeer, E., van der Hulst, J. M. The rotation curves of early-type disk galaxies. *Astron. Astrophys.* 471, 433–447 (2007).
10. Kerr, R. P. Gravitational Field of a Spinning Mass as an Example of Algebraically Special Metrics. *Phys. Rev. Lett.* 11, 237 (1963).
11. Navarro, J. F., Frenk, C. S., White, S. D. M. A Universal Density Profile from Hierarchical Clustering. *Astrophys. J.* 490, 493 (1997).
12. Burkert, A. The Structure of Dark Matter Halos in Dwarf Galaxies. *Astrophys. J. Lett.* 447, L25–L28 (1995).
13. McGaugh, S. S. The Baryonic Tully–Fisher Relation of Galaxies. *Astrophys. J.* 632, 859–871 (2005).

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.