

Article

Not peer-reviewed version

---

# About Asymptotic Properties of Bilinear Difference Equations Under Stochastic Perturbations

---

[Leonid Shaikhet](#) \*

Posted Date: 22 August 2025

doi: 10.20944/preprints202508.1521.v1

Keywords: zero and nonzero equilibria; stochastic perturbations; stability in probability; asymptotic mean square stability; numerical simulation; unsolved problem



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

# About Asymptotic Properties of Bilinear Difference Equations Under Stochastic Perturbations

Leonid Shaikheth

Department of Mathematics, Ariel University, Ariel 40700, Israel; leonid.shaikheth@usa.net

## Abstract

During the last two decades, rational difference equations, in particular, rational bilinear difference equations, have become very popular in research. In this paper the asymptotic properties of one rational bilinear difference equation are first studied under stochastic perturbations. It is assumed that stochastic perturbations are directly proportional to the deviation of the current value of the equation solution from one of its equilibria (the zero or nonzero). Some conditions are obtained by which the equilibrium under consideration is stable in probability or unstable. The obtained results are illustrated by numerical simulation of solutions of the considered stochastic difference equation. It is noted, that the research method used here can be applied to stability investigation of many other types of nonlinear difference equations with the order of nonlinearity higher than one. Some directions for further development of research are proposed. To readers attention some discussion about an unsolved problem of stabilization by noise for stochastic difference equations is also proposed. This problem, well known already for more than 50 years for stochastic differential equations, has not yet been solved until now for stochastic difference equations.

**Keywords:** zero and nonzero equilibria; stochastic perturbations; stability in probability; asymptotic mean square stability; numerical simulation; unsolved problem

## 1. Introduction

During the last two decades, rational difference equations, in particular, rational bilinear difference equations have become very popular in research (see, for instance, [1–17] and references therein). Various interesting results related to the solvability and stability of equations of this type are obtained. In particular, stability of the bilinear difference equation

$$\begin{aligned} x_{n+1} &= ax_n + \frac{bx_n x_{n-1}}{cx_n + dx_{n-1}}, \quad n = 0, 1, \dots, \\ x_n &= \phi_n, \quad n = -1, 0, \end{aligned}$$

with positive parameters  $a, b, c, d$  and initial values  $\phi_{-1}, \phi_0$  is studied in [7].

It is clear that, without loss of generality, one of the parameters  $b, c, d$  in this equation can be equated to 1. Let  $d = 1$ . So, we will consider the bilinear difference equation

$$\begin{aligned} x_{n+1} &= ax_n + \frac{bx_n x_{n-1}}{cx_n + x_{n-1}}, \quad n = 0, 1, \dots, \\ x_n &= \phi_n, \quad n = -1, 0, \end{aligned} \tag{1}$$

with positive  $a, b, c$  and  $\phi_{-1}, \phi_0$ .

The following two statements, obtained in [7], for the Equation (1) take the form:

- Statement 1** If  $(1 - a)(c + 1) \neq b$  then the unique equilibrium of the Equation (1) is  $x^* = 0$ .  
**Statement 2** If  $(1 - a)(c + 1) > b$  then the zero solution of the Equation (1) is locally asymptotically stable.

Below, the bilinear difference Equation (1) is studied under stochastic perturbations that are directly proportional to the deviation of the current value of the solution of the stochastic bilinear difference equation under consideration from the zero or nonzero equilibrium of the Equation (1). To the best of author's knowledge stochastic bilinear difference equations have not yet been considered.

Based on the obtained results, a hypothesis is proposed and discussed about the possibility of "stabilization by noise" for stochastic difference equations. The well-known effect of "stabilization by noise" for stochastic differential equations still has no analogue for stochastic difference equations. Here, this hypothesis is tested by numerical simulation and confirmed, but the formal proof of this hypothesis is still an unsolved problem.

### 1.1. Equilibria of the Equation (1)

Putting  $x_n = x^*$  for  $n \geq -1$ , we obtain that the equilibrium  $x^*$  of the Equation (1) is defined by the equation

$$x^* \left( 1 - a - \frac{b}{c+1} \right) = 0. \quad (2)$$

It is clear that if

$$a + \frac{b}{c+1} \neq 1 \quad (3)$$

then  $x^* = 0$  only can be the equilibrium of the Equation (1) (that coincides with **Statement 1**).

From the other hand, by the assumption

$$a + \frac{b}{c+1} = 1 \quad (4)$$

each  $x^* \in \mathbf{R}$  is a solution of the Equation (2) and  $x_n = x^*$ ,  $n \geq -1$ , is a solution of the Equation (1).

**Remark 1.** Note that by the condition (4) from the equality  $x_m = x_{m-1} = x^*$  for some  $m \geq 0$  from (1) it follows that  $x_n = x^*$  also for all  $n > m$ .

Below stability of the zero and nonzero equilibria of the Equation (1) is investigated under stochastic perturbations.

## 2. Some Transformations of the Initial Equation

Let  $n \in Z \cup Z_0$  be a discrete time,  $Z = \{0, 1, \dots\}$ ,  $Z_0 = \{-1, 0\}$ . Let  $\{\Omega, \mathfrak{F}, \mathbf{P}\}$  be a basic probability space,  $\mathfrak{F}_n \in \mathfrak{F}$ ,  $n \in Z$ , be a nondecreasing family of  $\sigma$ -algebras,  $\mathbf{E}$  be the expectation,  $(\xi_n)_{n \in Z}$  be a sequence of  $\mathfrak{F}_n$ -adapted mutually independent identically distributed random variables such that  $\mathbf{E}\xi_n = 0$ ,  $\mathbf{E}\xi_n^2 = 1$ ,  $n \in Z$  [18].

Note that via Remark 1 by the condition (4) and  $x_0 = x_{-1}$  the Equation (1) has the equilibrium  $x^* = x_0$ . Let us assume that the Equation (1) is exposed to stochastic perturbations that are directly proportional to the deviation of the solution  $x_n$  from the equilibrium  $x^*$ , i.e., the Equation (1) takes the form of the stochastic difference equation [18]

$$x_{n+1} = ax_n + \frac{bx_n x_{n-1}}{cx_n + x_{n-1}} + \sigma(x_n - x^*)\xi_{n+1}, \quad n = 0, 1, \dots, \quad (5)$$

where  $\sigma$  is a constant. By that the solution  $x_n = x^*$  of the Equation (1) is also the solution of the Equation (5).

**Remark 2.** Note that stochastic perturbations of the type (5) were first used for a system of stochastic delay differential equations in [19] and later in many other different research both for differential and for difference equations (see, for instance, [18,20] and references therein).

Putting in (5)  $x_n = y_n + x^*$ , we obtain

$$y_{n+1} = a(y_n + x^*) - x^* + \frac{b(y_n + x^*)(y_{n-1} + x^*)}{c(y_n + x^*) + y_{n-1} + x^*} + \sigma y_n \xi_{n+1}$$

or

$$y_{n+1} = ay_n + (a-1)x^* + \frac{b[(x^*)^2 + (y_n + y_{n-1})x^* + y_n y_{n-1}]}{(c+1)x^* + cy_n + y_{n-1}} + \sigma y_n \xi_{n+1}. \quad (6)$$

**Lemma 1.** Let the condition (4) holds and  $x^* \neq 0$ . Then the linear part of the Equation (6) has the form

$$\begin{aligned} z_{n+1} &= Az_n + Bz_{n-1} + \sigma z_n \xi_{n+1}, \quad n = 0, 1, \dots, \\ z_n &= \phi_n, \quad n = -1, 0, \end{aligned} \quad (7)$$

where

$$A = a + \frac{b}{(c+1)^2}, \quad B = \frac{bc}{(c+1)^2}, \quad A + B = 1. \quad (8)$$

**Proof.** Using the equality

$$\frac{1}{p+y} = \frac{1}{p} - \frac{y}{p^2} + o(y), \quad \text{where } p \neq 0 \text{ and } \lim_{y \rightarrow 0} \frac{o(y)}{y} = 0,$$

we have

$$\begin{aligned} & \frac{b[(x^*)^2 + (y_n + y_{n-1})x^* + y_n y_{n-1}]}{(c+1)x^* + cy_n + y_{n-1}} \\ &= b[(x^*)^2 + (y_n + y_{n-1})x^* + y_n y_{n-1}] \left( \frac{1}{(c+1)x^*} - \frac{cy_n + y_{n-1}}{(c+1)^2(x^*)^2} + o(y) \right) \\ &= \frac{bx^*}{c+1} - \frac{b}{(c+1)^2}(cy_n + y_{n-1}) + \frac{b}{c+1}(y_n + y_{n-1}) + o(y). \end{aligned} \quad (9)$$

Substituting (9) into (6), neglecting the nonlinear terms and using (2), we obtain

$$\begin{aligned} z_{n+1} &= az_n + (a-1)x^* + \frac{bx^*}{c+1} - \frac{b}{(c+1)^2}(cz_n + z_{n-1}) + \frac{b}{c+1}(z_n + z_{n-1}) + \sigma z_n \xi_{n+1} \\ &= \left( a - 1 + \frac{b}{c+1} \right) x^* + \left( a + \frac{b}{c+1} - \frac{bc}{(c+1)^2} \right) z_n + \left( \frac{b}{c+1} - \frac{b}{(c+1)^2} \right) z_{n-1} \\ &\quad + \sigma z_n \xi_{n+1} \\ &= \left( a + \frac{b}{(c+1)^2} \right) z_n + \frac{bc}{(c+1)^2} z_{n-1} + \sigma z_n \xi_{n+1}, \end{aligned}$$

that via (8) gives (7). The proof is completed.  $\square$

### 3. Stability

Consider the scalar stochastic difference equation [18]

$$\begin{aligned} x_{n+1} &= ax_n + bx_{n-1} + \sigma x_n \xi_{n+1}, \quad i \in \mathbb{Z}, \\ x_n &= \phi_n, \quad n \in \mathbb{Z}_0, \end{aligned} \quad (10)$$

where  $a, b, \sigma$  are constants and  $\xi_{n+1}$  is described above.

**Definition 1.** The zero solution of the Equation (10) is called stable in probability if for any  $\varepsilon > 0$  and  $\varepsilon_1 \in (0, 1)$  there exists a  $\delta > 0$  such that the solution  $x_n$  of the Equation (10) satisfies the inequality  $\mathbf{P}\{\sup_{n \in \mathbb{Z}} |x_n| > \varepsilon / \mathfrak{F}_0\} < \varepsilon_1$  for any initial function  $\phi$  such that  $\mathbf{P}\{\max_{n \in \mathbb{Z}_0} |x_n| < \delta\} = 1$ .

**Definition 2.** The zero solution of the Equation (10) is called:

- mean square stable if for each  $\varepsilon > 0$  there exists a  $\delta > 0$  such that  $\mathbf{E}x_n^2 < \varepsilon$ ,  $n \in \mathbb{Z}$ , for any initial function  $\phi(n)$  such that  $\|\phi\|^2 = \max_{n \in \mathbb{Z}_0} \mathbf{E}|\phi(n)|^2 < \delta$ ;
- asymptotically mean square stable if it is mean square stable and for each initial function  $\phi(n)$  such that  $\|\phi\|^2 < \infty$  the solution  $x_n$  of the Equation (10) satisfies the condition  $\lim_{n \rightarrow \infty} \mathbf{E}x_n^2 = 0$ .

**Remark 3.** It is clear that stability of the equilibrium  $x^*$  of the Equation (5) is equivalent to stability of the zero solution of the Equation (6). It is known [18] that the investigation of stability in probability of the zero solution of a nonlinear stochastic difference equation with an order of nonlinearity higher than one can be reduced to the investigation of asymptotic mean square stability of the zero solution of the linear part of this equation. So, to get conditions for stability in probability of the equilibrium  $x^*$  of the nonlinear stochastic difference Equation (5) it is enough to get conditions for asymptotic mean square stability of the zero solution of the linear stochastic difference Equation (7) that is the linear part of the nonlinear difference Equation (6).

**Remark 4.** Note that the method of studying the stability in probability of nonlinear stochastic difference equations, presented in Remark 3, is also applicable to stochastic nonlinear differential equations with the order of nonlinearity higher than one (see [20–23]).

**Lemma 2.** If

$$|a| + |b| < \sqrt{1 - \sigma^2} \quad (11)$$

then the zero solution of the Equation (10) is asymptotically mean square stable.

**Lemma 3.** The inequalities

$$|b| < 1, \quad |a| < 1 - b, \quad \sigma^2 < \frac{1+b}{1-b} \left[ (1-b)^2 - a^2 \right], \quad (12)$$

are the necessary and sufficient conditions for asymptotic mean square stability of the zero solution of the Equation (10).

**Remark 5.** Note that the stability conditions (11) and (12) are obtained via the general method of Lyapunov functionals construction [18].

**Theorem 1.** If

$$a + \frac{b}{c+1} < \sqrt{1 - \sigma^2} \quad (13)$$

then the equilibrium  $x^* = 0$  of the Equation (5) is stable in probability.

**Proof.** From the condition (3) it follows that by the condition (13) the Equation (5) has the equilibrium  $x^* = 0$  only. Note also that

$$a + \frac{b}{(c+1)^2} + \frac{bc}{(c+1)^2} = a + \frac{b}{c+1}.$$

So, for the Equation (7) the condition (11) takes the form (13). Via Lemma 2 it means that by the condition (13) the equilibrium  $x^* = 0$  of the Equation (7) is asymptotically mean square stable and via Remark 3, the equilibrium  $x^* = 0$  of the Equation (5) is stable in probability. The proof is completed.  $\square$

**Remark 6.** Note that the conditions (12) for the Equation (7) take the form

$$a + \frac{b}{c+1} < 1, \quad \sigma^2 < \frac{(c+1)^2 + bc}{(c+1)^2 - bc} \left[ \left( 1 - \frac{bc}{(c+1)^2} \right)^2 - \left( a + \frac{b}{(c+1)^2} \right)^2 \right]. \quad (14)$$

Thus, the inequalities (14) are necessary and sufficient conditions for asymptotic mean square stability of the zero solution of the Equation (7) and, therefore, via Remark 3, sufficient conditions for stability in probability of the solution  $x^*$  of the Equation (5).


**Remark 7.** Note that by  $\sigma = 0$  each from the conditions (13) and (14) coincides with the inequality in Statement 2.

**Remark 8.** Note that in all examples below for numerical simulation of solutions of the Equation (5) the random value  $\xi_{n+1}$  is used in the form  $\xi_{n+1} = \sqrt{12}(\eta - 0.5)$ , where  $\eta$  is a random value uniformly distributed on the interval  $[0,1]$  with  $E\eta = 0.5$  and  $D\eta = 1/12$ . So,  $E\xi_{n+1} = 0$ ,  $D\xi_{n+1} = E\xi_{n+1}^2 = 1$ .

**Example 1.** Put  $a = 0.5$ ,  $b = 0.7$ ,  $c = 1$ ,  $\sigma = 0.5$ . The condition (13) holds:  $a + \frac{b}{c+1} = 0.85 < \sqrt{1 - \sigma^2} = 0.866$ . Besides, the conditions (13) and (14) give respectively  $\sigma^2 < 0.28$  and  $\sigma^2 < 0.32$ . Via Theorem 1 the zero solution of the Equation (5) with  $x^* = 0$  is stable in probability. In Figure1 500 trajectories (blue) of the solution of the Equation (5) are shown with the initial conditions  $x_{-1} = 0.8$ ,  $x_0 = 1.8$ , all trajectories converge to zero. The red line corresponds to the deterministic case, i.e.,  $\sigma = 0$ .


**Example 2.** Put now  $\sigma = 0.95$  with the same values of all other parameters as in Example 1. The conditions (13) and (14) do not hold. In Figure2 500 trajectories (blue) of the solution of the Equation (5) with  $x^* = 0$  are shown with the initial conditions  $x_{-1} = 0.01$ ,  $x_0 = 0.02$ . The zero solution is unstable and the trajectories fill the whole space. The red line corresponds to the deterministic case, i.e.,  $\sigma = 0$ .

**Example 3.** Put  $a = 0.65$ ,  $b = 0.7$ ,  $c = 1$ ,  $\sigma = 0.15$ ,  $x^* = 2.7$ . Wherein  $a + \frac{b}{c+1} = 1$ , therefore, the condition (4) holds, the conditions (13) and (14) do not hold. In Figure3 the red straight line corresponds to the constant solution  $x_n = x^*$ ,  $n \geq -1$ , 500 trajectories (blue) of the solution of the Equation (5) are shown with the initial conditions  $x_{-1} = x^* - 0.1 = 2.6$ ,  $x_0 = x^* - 0.2 = 2.5$ . The solution  $x_n = x^*$  is unstable, so, the trajectories fill the whole space.




**Figure 1.** 500 trajectories (blue) of the solution  $x_n$  of the Equation (5) with  $x^* = 0$ ,  $a = 0.5$ ,  $b = 0.7$ ,  $c = 1$ ,  $\sigma = 0.5$ ,  $x_{-1} = 0.8$ ,  $x_0 = 1.8$ . The red line corresponds to the deterministic case, i.e.  $\sigma = 0$ .





**Figure 2.** 500 trajectories (blue) of the solution  $x_n$  of the Equation (5) with  $x^* = 0, a = 0.5, b=0.7, c = 1, \sigma = 0.95, x_{-1} = 0.01, x_0 = 0.02$ . The red line corresponds to the deterministic case, i.e.  $\sigma = 0$



**Figure 3.** 500 trajectories (blue) of the solution  $x_n$  of the Equation (5) with  $x^* = 2.7, a = 0.65, b=0.7, c = 1, \sigma = 0.15, x_{-1} = 2.6, x_0 = 2.5$ . The red straight corresponds to the constant solution  $x_n = x^*, n \geq -1, \sigma = 0$

#### 4. About One Unsolved Problem of Stabilization by Noise

More than 50 years ago Khasminskii shows [24] that unstable by the conditions  $a > 0$  and  $\sigma = 0$  the zero solution of the differential equation

$$dx(t) = ax(t)dt + \sigma x(t)dw(t) \quad (15)$$

becomes stable by the presence of a big enough level of noise. More exactly, by the condition

$$0 < 2a < \sigma^2 \quad (16)$$

so-called "stabilization by noise" occurs and the zero solution of the Equation (15) becomes stable in probability.

Really, let  $L$  be the generator [20,24,25] of the stochastic differential Equation (15). For the Lyapunov function  $v(x)$  it has the form

$$Lv(x) = v'(x)ax + \frac{1}{2}v''(x)\sigma^2x^2.$$

So, via (16) for the Lyapunov function

$$v(x) = |x|^\nu, \quad \nu = 1 - \frac{2a}{\sigma^2} \in (0,1),$$

we have


$$\begin{aligned} Lv(x) &= \nu|x|^{\nu-1}ax + \frac{1}{2}\nu(\nu-1)|x|^{\nu-2}\sigma^2x^2 \\ &\leq a\nu|x|^\nu \left(1 + (\nu-1)\frac{\sigma^2}{2a}\right) = 0. \end{aligned}$$

From the obtained condition  $Lv(x) \leq 0$  it follows that the zero solution of the Equation (15) is stable in probability [20,24].

As it is noted in [26] any similar result for the stochastic difference Equation (10) is absent until now even in the case  $b = 0$ . But the following example shows that the effect of stabilization by noise can take a place for difference equations too.


**Example 4.** Consider Example 3 again with  $\sigma = 0.45$  and the same values of all other parameters. In Figure 4 one can see that all 500 trajectories converge to the solution  $x_n = x^* = 2.7$  of the Equation (5). Putting  $\sigma = 0.75$ , i.e., increasing once more the level of noise, we obtain (see Figure 5) that all 500 trajectories converge to the solution  $x_n = x^* = 2.7$  of the Equation (5) faster than in Figure 4. So, the solution  $x_n = x^* = 2.7$  of the Equation (5), that is unstable by the small level of noise (Figure 3,  $\sigma = 0.15$ ), becomes stable by increasing the level of noise.

Thus, the Hypothesis about stabilization by noise for stochastic difference equations may well take a place. However, the proof of this Hypothesis is currently an unsolved problem.



**Figure 4.** 500 trajectories (blue) of the solution  $x_n$  of the Equation (5) with  $\sigma = 0.45$  and the same values of all other parameters as in Figure 3.





**Figure 5.** 500 trajectories (blue) of the solution  $x_n$  of the Equation (5) with  $\sigma = 0.75$  and the same values of all other parameters as in Figure 3.

## 5. Conclusions

In conclusion, one would like to note that the study of the asymptotic properties of a bilinear difference equation under stochastic perturbations is carried out here for the first time, and that the research method used here can be extended to the following natural directions of research development:

- 1) Bilinear difference equations of other forms, for instance, obtained from this one

$$x_{n+1} = ax_n \pm \frac{bx_n x_{n-1}}{cx_n \pm dx_{n-1}}, \quad n = 0, 1, \dots,$$

$$x_n = \phi_n, \quad n = -1, 0,$$

using different combinations of plus and minus. In each case, one can expect the appearance of new features of the solution behavior.

- 2) Different mathematical models in various applications, described by nonlinear stochastic difference equations of a more complex form.

- 3) Formulate and prove for difference equations a statement similar to Khasminsky's statement about "stabilization by noise", which has been well known in the theory of stochastic differential equations for more than 50 years, but still has no analogue for stochastic difference equations.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The author declares no conflict of interest.

## References

1. Abo-Zeid, R.; Kamal, H. On the solutions of a third order rational difference equation. *Thai Journal of Mathematics*, **2020**, 18(4), 1865-1874.
2. Aloqeili, M. Dynamics of a rational difference equation. *Applied Mathematics and Computation*, **2006**, 176(2), 768-774.
3. Camouzis, E.; Ladas, G.; Voulou, H.D. On the dynamics of  $x_{n+1} = \frac{a + \gamma x_{n-1} + \delta x_{n-2}}{A + x_{n-2}}$ . *Journal of Difference Equations and Applications*, **2003**, 9(8), 731-738.
4. Chatterjee, E.; Grove, E.A.; Kostrov, Y.; Ladas, G. On the trichotomy character of  $x_{n+1} = \frac{a + \gamma x_{n-1}}{A + Bx_n + x_{n-2}}$ . *Journal of Difference Equations and Applications*, **2003**, 9(12), 1113-1128.

5. Cinar, C. On the positive solutions of the difference equation  $x_{n+1} = \frac{ax_{n-1}}{1 + bx_n x_{n-1}}$ . *Applied Mathematics and Computation*, **2004**, 156, 587-590.
6. Elabbasy, E.M.; El-Metwally, H.; Elsayed, E.M. On the difference equations  $x_{n+1} = ax_n - \frac{bx_n}{cx_n - dx_{n-1}}$ . *Advances in Difference Equations*, **2006**, (2006), Article ID 82579, 1-10.
7. Elsayed, E.M. Qualitative behavior of a rational recursive sequence. *Indagationes Mathematicae, N.S.*, **2008**, 19(2), 189-201.
8. Grove, E.A.; Kulenovic, M.R.S.; Ladas, G. Progress report on rational difference equations. *Journal of Difference Equations and Applications*, **2004**, 10(13-15), 1313-1327.
9. Halim, Y.; Touafek, N.; Yazlik, Y. Dynamic behavior of a second-order nonlinear rational difference equation. *Turkish Journal of Mathematics*, **2015**, 39(6), 1004-1018. <https://doi.org/10.3906/mat-1503-80>
10. Ibrahim, T.F.; Touafek, N. On a third order rational difference equation with variable coefficients. *Dynamics of Continuous, Discrete and Impulsive Systems, Series B: Applications & Algorithms*, **2013**, 20(2), 251-264.
11. Papaschinopoulos, G.; Stefanidou, G. Asymptotic behavior of the solutions of a class of rational difference equations. *International Journal of Difference Equations*, **2010**, 5(2), 233-249.
12. Stevic, S. On the system  $x_{n+1} = \frac{y_n x_{n-k}}{y_{n-k+1}(a_n + b_n y_n x_{n-k})}$ ,  $y_{n+1} = \frac{x_n y_{n-k}}{x_{n-k+1}(c_n + d_n x_n y_{n-k})}$ . *Applied Mathematics and Computation*, **2013**, 219, 4526-4534.
13. Stevic, S. Representations of solutions to linear and bilinear difference equations and systems of bilinear difference equations. *Advances in Difference Equations*. **2018**, Vol. 2018, Article No.474, 21 pages.
14. Stevic, S.; Diblík, J.; Iricanin, B.; Smarda, Z. On a solvable system of rational difference equations. *Journal of Difference Equations and Applications*, **2014**, 20(5-6), 811-825.
15. Stevic, S., Iricanin, B., Kosmala, W., Smarda, Z. Note on a solution form to the cyclic bilinear system of difference equations. *Applied Mathematics Letters*, **2021**, 111, 106690.
16. Taskara, N.; Tollu, D.T.; Yazlik, Y. Solutions of rational difference system of order three in terms of Padovan numbers. *Journal of Advanced Research in Applied Mathematics*, **2015**, 7(3), 18-29.
17. Yazlik, Y. On the solutions and behavior of rational difference equations. *Journal of Computational Analysis and Applications*, **2014**, 17(3), 584-594.
18. Shaikhet, L. *Lyapunov functionals and stability of stochastic difference equations*. Springer Science & Business Media: London, UK, 2011.
19. Beretta, E.; Kolmanovskii, V.; Shaikhet, L. Stability of epidemic model with time delays influenced by stochastic perturbations. *Mathematics and Computers in Simulation (Special Issue "Delay Systems")*, **1998**, 45(3-4), 269-277. <https://www.sciencedirect.com/science/article/pii/S0378475497001067>
20. Shaikhet, L. *Lyapunov functionals and stability of stochastic functional differential equations*. Springer Science & Business Media: Berlin/Heidelberg, Germany, 2013.
21. Shaikhet, L. About one method of stability investigation for nonlinear stochastic delay differential equations. *International Journal of Robust and Nonlinear Control*, **2021**, 31(8), 2946-2959, <https://onlinelibrary.wiley.com/doi/10.1002/rnc.5440>
22. Shaikhet, L. Some generalization of the method of stability investigation for nonlinear stochastic delay differential equations. *MDPI, Symmetry*, **2022**, 14(8), 1734, 1-13, <https://www.mdpi.com/2073-8994/14/8/1734/pdf>
23. Shaikhet L. About stability of nonlinear stochastic differential equations with state-dependent delay. *MDPI, Symmetry*, **2022**, 14(11), 2307, 1-12, <https://www.mdpi.com/2073-8994/14/11/2307/pdf>
24. Khasminskii, R.Z. *Stochastic stability of differential equations*; Springer: Berlin, Germany, 2012 (in Russian, Moscow, Nauka, 1969).
25. Gikhman, I.I., Skorokhod, A.V. *Stochastic Differential Equations*; Springer: Berlin, Germany, 1972.
26. Shaikhet, L. Some unsolved problems in stability and optimal control theory of stochastic systems. Special Issue "Models of Delay Differential Equations - II", *MDPI, Mathematics*, **2022**, 10(3), 474, 1-10, <https://doi.org/10.3390/math10030474>.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.