

Article

Not peer-reviewed version

---

# Sex-Specific Effects of Long-Term Antipsychotic Drug Medication on Adipocyte Tissue and the Crosstalk to Liver and Brain in Rats

---

[Karin Fehsel](#)<sup>\*</sup> and [Marie-Luise Bouvier](#)

Posted Date: 19 January 2024

doi: 10.20944/preprints202401.1508.v1

Keywords: perirenal adipose tissue, clozapine, haloperidol, lipid, glucose, perilipin-A hormone sensitive lipase, glucose transporter-4, insulin receptor- $\beta$



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

# Sex-Specific Effects of Long-Term Antipsychotic Drug Medication on Adipocyte Tissue and the Crosstalk to Liver and Brain in Rats

Karin Fehsel and Marie-Luise Bouvier

Department of Psychiatry and Psychotherapy, Medical Faculty, Heinrich-Heine-University, Bergische Landstraße 2, 40629 Düsseldorf, Germany

\* Correspondence: fehsel@hhu.de

**Abstract:** Antipsychotic drug (APD) medication can lead to metabolic dysfunctions and weight gain, which together increase morbidity and mortality. Metabolically active visceral adipose tissue (VAT) in particular plays a crucial role in the etiopathology of these metabolic dysregulations. Here we studied the effect of 12 weeks drug medication by daily oral feeding of clozapine and haloperidol on the perirenal fat tissue as part of VAT of male and female Sprague Dawley rats in the context of complex former investigations on brain, liver, and blood. Adipocyte area values were determined as well as triglycerides, non-esterified fatty acids (NEFAs), glucose, glycogen, lactate, malondialdehyde equivalents, ferric iron and protein levels of Perilipin-A, hormone-sensitive-lipase (HSL), hepcidin, glucose transporter-4 (Glut-4) and insulin receptor- $\beta$  (IR- $\beta$ ). We found increased adipocyte mass in males with slightly higher adipocyte area values in both males and females under clozapine medication. Triglycerides, NEFAs, glucose and oxidative stress in the medicated groups were unchanged or slightly decreased. In contrast to controls and haloperidol medicated rats, perirenal adipocyte mass and serum leptin levels were not correlated under clozapine. Protein expressions of perilipin-A, Glut-4 and HSL were decreased under clozapine medication. IR- $\beta$  expression changed sex-specifically in the clozapine medicated groups associated with higher hepcidin levels in the perirenal adipose tissue of clozapine-medicated females. Taken together, clozapine and haloperidol have a smaller effect than expected on perirenal adipose tissue. The perirenal adipose tissue shows only weak changes in lipid and glucose metabolism. The main changes can be seen in the proteins examined and probably in their effect on liver metabolism.

**Keywords:** perirenal adipose tissue; clozapine; haloperidol; lipid; glucose; perilipin-A hormone sensitive lipase; glucose transporter-4; insulin receptor- $\beta$

## 1. Introduction

Weight gain under second-generation antipsychotic (SGA) medication associated with metabolic dysfunctions leads to increased morbidity and mortality with potential non-compliance in patients. The risk of weight gain and obesity appears to be the highest with olanzapine and clozapine [1]. An increase in weight has also been reported in individuals treated with first-generation antipsychotics (FGAs) [2]. Especially adipose tissue as a target of SGAs seems to be involved in the etiopathology of obesity and metabolic abnormalities like type 2 diabetes (T2D) or insulin resistance including glucose and lipid dysregulation [3]. In murine adipocytes, SGAs increase both adipogenesis of preadipocytes (hyperplasia) [4] and/or lipid accumulation (hypertrophy) in mature adipocytes [5].

Energy balance includes energy intake, energy expenditure and energy storage. When energy intake is equal to energy expenditure, body weight remains stable [6]. The key role of white adipose tissue (WAT) consists in the storage of energy and in the protection from excessive glucose in the form of triacylglycerols (TAG) similarly named triglycerides (TG) in lipid droplets organelles. Perilipin is an intracellular protein that is hormonally regulated and is localized exclusively on the surface of lipid droplets. When energy is required, perilipin is phosphorylated by protein kinase-A. Once activated, perilipin changes its conformation and allows lipolysis by the three lipases adipocyte-

triglyceride-lipase (ATGL), the rate-limiting enzyme for TAG hydrolysis, hormone-sensitive-lipase (HSL) and monoacylglycerol-lipase (MGL) which degrade TAGs sequentially resulting in the release of free fatty acids and glycerol [7]. The fatty acids are then bound to albumin and transported to the target tissue. Glycerol can be isomerised in the liver to glyceraldehyde-3-phosphate and used for gluconeogenesis or degraded to pyruvate by glycolysis. Consequently, production of glycerol or lactate is an important primary function of adipocytes to lower the negative effects of excess glucose [8].

Increased VAT is associated with metabolic dysfunctions like T2D, insulin resistance, inflammatory processes and cardiometabolic disease [9]. In contrast to subcutaneous adipose tissue (SCAT), visceral preadipocytes originate from the mesothelium [10]. VAT is located around internal organs as uterus and ovaries in females (gonadal), epididymis and testes in males (epididymal), kidneys (perirenal), heart (epicardial) and intestines (retroperitoneal, mesenteric, and omental). Perirenal VAT has a close association to obesity and cardiovascular disease [11]. SCAT is located under the skin and serves primarily as a body temperature insulator and energy store. VAT adipocytes are more metabolically active, more sensitive to lipolysis and more insulin-resistant than SCAT adipocytes [12]. SGA therapy increases both subcutaneous and visceral adipose tissue in humans and rodents [3]. In recent years, VAT has been recognized as a major endocrine organ that secretes hormones like leptin, adiponectin, tumor necrosis factor- $\alpha$  (TNF $\alpha$ ), interleukin-6 (IL-6) and estradiol controlling whole body metabolism, insulin sensitivity, and energy homeostasis [13]. Leptin passes through the brain barrier and acts on neurons in the hypothalamic nuclei regulating food intake and energy metabolism by inhibiting the neuropeptides NPY (Neuropeptid Y) and agouti-related peptide (AGRP) [14]. It also induces lipolysis in adipocyte tissue, mediated by sympathetic neurons which innervate adipocytes in VAT [15]. Adiponectin is involved in the regulation of glucose levels as well as fatty acid oxidation. In humans and rodents, higher levels of the anti-inflammatory and anti-atherogenic adiponectin counteracts insulin resistance by reducing TG levels in muscle and liver cells [16]. However, it is also involved in the pathogenesis of obesity and non-alcoholic fatty liver disease (NAFLD), the hepatic manifestation of metabolic syndrome, characterized by increased hepatic lipid accumulation [17,18]. NAFLD covers a wide spectrum of disorders ranging from steatosis with a benign non-progressive clinical course, to finally advanced fibrosis and cirrhosis [19]. Ghrelin is mainly secreted by the gastric fundus mucosa and pancreatic cells. Circulating levels of ghrelin are increased in response to fasting or prolonged food restriction [20]. In addition to its stimulation of appetite and food intake by activating NPY and AGRP in the hypothalamus, ghrelin stimulates adipogenesis and suppresses lipolysis in rat visceral adipocytes [21]. The insulin-dependent glucose transporter-4 (Glut-4) is primarily expressed in adipocytes, skeletal and cardiac muscle cells. The binding of insulin to its receptor triggers kinase cascades that lead to translocation, docking and fusion of Glut-4 containing vesicles into the plasma membrane, allowing more glucose to enter the cell [22]. Through this glucose uptake by Glut-4 blood glucose levels can be lowered rapidly and efficiently, thereby reducing the harmful effect of excessive glucose levels.

Previously, we have examined the effects of clozapine and haloperidol on weight, food and water intake, and serum parameters [23], hepatic [24], hypothalamic [25] blood and iron parameters [26] in a rat model of metabolic syndrome. Finally, in this study, we examine the effects of the two antipsychotic drugs on basal metabolism of perirenal adipocytes.


## 2. RESULTS

In short, male and female rats received daily medication of clozapine or haloperidol by oral feeding in ground pellets over 12 weeks. Serum levels of haloperidol, clozapine, and its metabolite N-desmethylclozapine, body weight, motor activity, food and water intake (raw data and calculated on 1 kg body weight), and metabolic parameters were reported previously [23–26]. Male and female rats with haloperidol gained less weight than the control groups, associated with reduced food and water intake. Male rats under clozapine medication showed significantly increased body weight, and

liver mass, although their food and water intake were not increased. Higher fasting glucose in serum and decreased glycogen in liver were found in female rats medicated with clozapine.

### 2.1. Effect of haloperidol and clozapine on percentage body weight gain after 12-week medication of clozapine and haloperidol on male and female Sprague Dawley rats.

Percentage weight gain differed for MEDICATION [ $F(2,54)=27.94$   $p=0.0005$ ] with significant effect between male controls and clozapine medicated rats ( $p=0.022$ ) (Figure 1).




**Figure 1.** Significant increase of percental weight gain between male controls and clozapine-medicated Sprague Dawley rats with  $p=0.022$  and  $n=10$  animals per group (m=male, f=female, Contr=Control, Halo=haloperidol medicated, Cloz=clozapine medicated).

### 2.2. Effect of haloperidol and clozapine on adipocyte mass, neutral fat content, adipocyte area value, and ferric iron content of perirenal VAT

We found differences in adipocyte mass (Table 1) for MEDICATION [ $F(2,54)=8.90$   $p=0.00046$ ], SEX [ $F(1,54)=111.10$   $p\leq 0.000001$ ] and MEDICATION $\times$ SEX [ $F(2,54)=5.41$   $p=0.007$ ]. Male clozapine treated animals showed higher adipocyte mass than male controls ( $p=0.047$ ) and male haloperidol treated animals ( $p=0.001$ ). Perirenal VAT of female haloperidol medicated rats differed from clozapine treated animals ( $p=0.020$ ). adipocyte mass of males differed significantly from females in all groups ( $p=0.00006$  for controls;  $p=0.003$  for haloperidol medicated rats;  $p<0.000001$  for clozapine medicated rats).

Neutral fat content in adipose tissue (Table 1) did not differ between the groups, but differed for SEX [ $F(1,54)=111.10$   $p=0.001$ ] with differences between male and female controls ( $p=0.021$ ) and male and female haloperidol medicated animals ( $p=0.035$ ).

The mean adipocyte area value of 200 calculated perirenal adipocytes from each of two slices (Table 1 and Figure 2) differed for SEX [ $F(1,53)=13.66$   $p=0.00047$ ] with differences between male and female controls ( $p=0.041$ ) and clozapine medicated rats ( $p=0.018$ ).



**Figure 2.** Exemplary presentation of 20 µm frozen fat sections of male and female Sprague Dawley rats after 12-week medication with haloperidol or clozapine, stained with Mayer's hematoxylin. Two hundred adipocytes from each of two slices were counted (marked by X), their area was calculated with Axiovision (Zeiss, Germany) and the values were averaged. Values are given in Table 1 as "adipocytes area value". (m=male, f=female, Control=controls, Halo=haloperidol medicated, Cloz=clozapine medicated).

We did not find ferric iron deposits in adipose tissue, although the Perl's Prussian blue stain is an extremely sensitive method.

**Table 1.** Main metabolic parameters of perirenal VAT and selected metabolic parameters of liver and serum in male and female control, haloperidol and clozapine medicated Sprague Dawley rats after 12 week trial period.

	Contr m	Halo m	Cloz m	Contr f	Halo f	Cloz f
<b>adipocytes</b>						
percentage weight gain	27.55±2.38	22.73±2.68	36.13±1.89*	13.29±1.34	12.35±1.07	14.07±0.81
adipocytes mass [g]	27.55±2.38	22.73±2.60	36.13±1.89*	13.29±1.34	12.35±1.07	14.07±0.81
neutral fat content [g/g fat tissue]	0.51±0.04	0.58±0.05	0.43±0.02	0.33±0.06	0.41±0.04	0.36±0.04
adipocytes area value [µm <sup>2</sup> ]	1475632 ±100845	1342769 ±86813	1635892 ±88177	1115800 ±62396	1109041 ±63796	1230149 ±74687
triglycerides [mg/dl]	236.7±19.5	190.0±11.2	222.9±19.8	268.6±19.9	258.8±26.6	237.6±26.7
NEFAS [mmol/l]	0.54±0.04	0.41±0.01*	0.47±0.04	0.55±0.02	0.52±0.03	0.56±0.04
glucose [ng/µl]	15.71±2.2	12.13±2.4	11.89±2.0	13.99±1.9	12.69±1.6	12.52±2.3
glycogen [µg/ml]	7.62±1.36	4.80±0.70	6.51±0.78	6.43±1.36	8.68±1.40	11.31±1.82
lactate [mmol/l]	0.63±0.04	0.85±0.13	0.64±0.04	0.73±0.05	0.86±0.15	0.89±0.09
malondialdehyde [nm/ml]	32.72±5.2	17.97±2.3	20.76±2.9	18.42±3.5	21.52±3.7	20.83±4.1
<b>liver</b>						
NEFAS [mmol/l]	6.37±0.33	6.46±0.42	7.14±0.54	7.26±0.28	7.48±0.24	6.85±0.75
cholesterol/HDL-cholesterol ratio	7.73±0.36	8.05±0.37	8.50±0.38	7.33±0.31	8.18±0.52	7.50±0.58
lactate [mmol/l]	0.44±0.03	0.47±0.03	0.65±0.04*	0.82±0.05	0.81±0.04	0.77±0.05
pyruvate [mmol/l]	0.41±0.03	0.48±0.04	0.57±0.05*	0.52±0.03	0.56±0.06	0.48±0.05
<b>serum</b>						
NEFAS [mmol/l]	0.61±0.05	0.63±0.05	0.64±0.06	0.70±0.08	0.63±0.03	0.63±0.04
cholesterol/HDL-cholesterol ratio	3.39±0.07	3.27±0.07	3.86±0.13*	3.18±0.08	3.07±0.09	3.45±1.0
LDL-cholesterol/HDL ratio-cholesterol	1.81±0.06	1.73±0.11	2.28±0.11*	1.90±0.09	1.77±0.10	2.16±0.10
fasting glucose [ng/µl]	47.5±5.44	64.1±4.58	57.2±3.81	34.4±1.59	40.3±5.45	45.4±2.99**
lactate [mmol/l]	2.34±0.22	2.48±0.16	2.01±0.13	2.92±0.16	2.81±0.14	2.96±0.17
estrogen [pg/ml]				132.6±5.0	140.1±16.8	130.8±7.8

Data presented as group mean±SEM \*denotes significance of p<0.05, \*\* denotes significance of p<0.005

with n=10 animals per group.

NEFAS=non-esterified fatty acids; HDL=high density lipoprotein; LDL=low density lipoprotein; m=male; f=female; Contr=controls; Halo=haloperidol medicated; Cloz=clozapine medicated

Serum leptin levels are positively correlated with adipocyte mass in male ( $\rho=0.830$   $p=0.003$ ) and female ( $\rho=0.794$   $p=0.006$ ) controls, male ( $\rho=0.733$   $p=0.016$ ) and female ( $\rho=0.883$   $p=0.002$ ) haloperidol medicated animals, but not under clozapine medication (Supplement Figure 2A-D). Serum adiponectin levels were negatively correlated with percental adipocyte mass in the male ( $\rho=-0.661$

p=0.038) and positively in the female ( $\rho=0.667$  p=0.050) haloperidol medicated group (Supplement Figure 2 E.-F.)

### 2.3. Effect of haloperidol and clozapine on triglycerides, and NEFAs in rat perirenal adipose tissue as well as serum cholesterol/HDL ratio and LDL/HDL ratio, and hepatic Chol/HDL ratio

The amount of triglycerides (Table 1) in perirenal adipocytes did not differ between the groups but differed for SEX [F(1,51)=5.01 p=0.03] between male and female haloperidol medicated rats (p=0.034). Hepatic and serum NEFAs (Table 1) did not differ between the groups, but in adipocytes for MEDICATION [F(2,51)=3.38 p=0.042] and SEX [F(1,51)=8.37 p=0.006] with decreased levels in male haloperidol medicated rats compared to male controls (p=0.041). Only male haloperidol medicated animals differed significantly from the females (p=0.002).

In the serum, LDL/HDL ratios (Table 1) differed for MEDICATION [F(2,52)=12.71 p=0.000032], and the Chol/HDL ratios (Table 1) for MEDICATION [F(1,52)=15.07 p=0.0000071], and SEX [F(1,52)=13.39 p=0.001]. Male clozapine treated animals showed higher LDL/HDL and cholesterol/HDL ratios than male controls (LDL/HDL p=0.011; cholesterol/HDL p=0.007), and haloperidol treated animals (LDL/HDL p=0.002; cholesterol/HDL p=0.001). Female clozapine medicated rats differed from female haloperidol medicated rats (LDL/HDL p=0.032; cholesterol/HDL p=0.025). The Chol/HDL ratio of male clozapine treated rats differed significantly from females (p=0.020).

### 2.4. Effect of haloperidol and clozapine on glucose, glycogen, and lactate contents in rat perirenal VAT


In perirenal adipose tissue glucose, glycogen and lactate did not differ significantly between the groups (Table 1). However, glycogen shows differences for SEX [F(1,53)=5.77 p=0.02], and MEDICATION $\times$ SEX [F(2,53)=3.22 p=0.048]. Male and female controls did not differ, but females under haloperidol (p=0.023) and clozapine (p=0.033) medication had higher glycogen levels than males. We found no correlation between glucose and glycogen levels.

### 2.5. Effect of haloperidol and clozapine on oxidative stress of rat perirenal VAT and on plasma estrogen level in female rats

We found no difference between the groups and for sex of malondialdehyde equivalents in perirenal VAT (Table 1). Estrogen plasma levels did not differ between the female groups (Table 1).

### 2.6. Protein expression of Perilipin-A, HSL, hepcidin, Glut-4, IR- $\beta$ in perirenal adipocytes of haloperidol- and clozapine-medicated rats.

The protein level of perilipin-A in adipocytes (Figure 2A) varied significantly for MEDICATION [F(2,51)=5.91 p=0.005], in which clozapine medicated females differed significantly from the control group (p=0.006). Levels of HSL (Figure 2B) differed significantly for MEDICATION [F(2,53)=14.45 p=0.00001] and SEX [F(1,53)=20.43 p=0.000035]. Male clozapine medicated rats showed significantly lower levels than haloperidol medicated animals (p=0.015), whereas females under clozapine medication had lower levels than haloperidol medicated females (p=0.001) and female controls (p=0.012). All male groups had higher HSL levels than the respective female group (p=0.032 for controls, p=0.032 for the haloperidol medicated groups, p=0.005 for the clozapine medicated groups). A positive correlation between perilipin-A and HSL was found in the male ( $\rho=0.001$ ) and a negative correlation in the female ( $\rho=-0.636$  p=0.048) control group (Supplement Figure 2G-H). Hepcidin (Figure 2C) differed for MEDICATION [F(2,53)=3.94 p=0.025], SEX [F(1,53)=5.50 p=0.0023] and MEDICATION $\times$ SEX [F(2,53)=3.90 p=0.026]. Female clozapine medicated rats had higher hepcidin levels than female controls (p=0.002) and haloperidol medicated females (p=0.027). Only male controls had higher levels than female controls (p=0.001). The corresponding autoradiographs of the stripped and reprobed membranes are presented in Supplement Figure 1a in combination with the Ponceau S red stained membranes (Supplement Figure 1b).



**Figure 3.** Protein expression of A. perilipin A , B. hormone-sensitive lipase (HSL), C. hepcidin, D. glucose transporter 4 (Glut 4), E. insulin receptor  $\beta$  (IR  $\beta$ ) in male and female control, haloperidol, and clozapine medicated Sprague Dawley rats with \* $p \leq 0.05$ , \*\* $p \leq 0.001$  and \*\*\* $p \leq 0.0001$  as significant (n=10 animals per group). The autoradiographs and the Ponceau stained membranes are seen in Supplement Figure 1 A. and B.

Examining Glut-4 (Figure 2D) with two-way ANOVA showed differences for MEDICATION [F(2,53)=17.39  $p=0.000002$ ], SEX [F(1,53)=12.26  $p=0.001$ ] and MEDICATION $\times$ SEX [F(2,53)=6.94  $p=0.002$ ]. Male haloperidol ( $p=0.004$ ) and male clozapine ( $p=0.002$ ) medicated rats had significantly lower levels of Glut-4 than controls. Female clozapine ( $p=0.001$ ) treated rats had also lower levels, whereas female haloperidol ( $p=0.001$ ) medicated animals showed higher expression of Glut-4 than clozapine treated animals. Only male and female controls differed significantly from each other ( $p=0.0003$ ).

Two-way ANOVA of insulin receptor- $\beta$  (IR- $\beta$ ) expression (Figure 2E) showed differences for MEDICATION [F(2,51)=4.08  $p=0.023$ ] and MEDICATION $\times$ SEX [F(2,51)=16.23  $p=0.000004$ ]. IR- $\beta$  was significantly increased in male clozapine medicated rats, compared to controls ( $p=0.007$ ), whereas females under clozapine medication had lower levels than female controls ( $p=0.009$ ) and female haloperidol medicated rats ( $p=0.00039$ ). In clozapine medicated females, we found a nonparametric negative correlation between IR- $\beta$  and Glut-4 ( $\rho=-0.714$   $p=0.047$ ).

### 3. DISCUSSION

One of the main findings of our study is that perirenal visceral adipose tissue (VAT) - at least in rats- does not significantly contribute to the weight changes under SGA or FGA treatment. In a previous publication [23], we were able to show that male Sprague Dawley rats under clozapine gained significantly more weight over the entire duration of the trial, while males under haloperidol significantly lost weight. Percentage weight also increases significantly in male rats under clozapine medication (Figure 1) Although perirenal adipocytes mass is significantly increased in the male clozapine-treated group (Table 1), adipocytes mass related to 1 kg body weight is not significantly elevated [23]. Neutral fat, triglycerides and NEFAs in perirenal adipocytes are not increased (Table 1). Liver mass also increases significantly in clozapine-medicated males, as does the liver mass related to 1 kg body weight [24]. However, significant weight gain of male clozapine-treated rats or weight loss of haloperidol-medicated rats persists when fat mass and/or liver mass are subtracted from final body weight (Supplement Figure 3). Therefore, we conclude that there is at least one other parameter, such as subcutaneous adipose tissue (SCAT), that plays a role in weight gain with clozapine and weight loss with haloperidol. The sedative effect of haloperidol causes reduced food and water intake and thus weight loss in our animals [23]. In schizophrenic patients, treatment with SGAs results in an increase in both VAT and SCAT [27]. Due to its promotion of inflammation, the accumulation of VAT is considered harmful, whereas the beige adipocytes in SCAT seem to have neuroprotective and anti-inflammatory effects [28]. Further studies need to clarify whether SCAT, which was neglected in our study because it is difficult to determine in rats, has a higher impact on weight gain or loss under antipsychotic drug medication than VAT. An answer to this question would certainly influence the clinical assessment of obesity in patients.

In murine cells, SGAs seem to increase both adipocyte number, due to proliferation and differentiation of progenitor cells in the adipose tissue (hyperplasia), and adipocyte size (hypertrophy), due to lipid accumulation in mature adipocytes [4]. Hypertrophy of adipocytes is associated with dysfunction of adipocytes and metabolic effects like altered lipid and energy metabolism, increased pro-inflammatory cytokine secretion, and insulin resistance [9,29]. In humans, hypertrophy of VAT but not SCAT is associated with a more atherogenic lipid profile and several functional aspects of adipocyte activity relate to its size [9]. In our study, the increased adipocyte area value under clozapine points to hypertrophy (Table 1). Significant increases in leptin, triglycerides, insulin, intra-abdominal fat, adipocyte area and HSL activity are found in a metabolic syndrome model of male Wistar rats. This suggests that changed lipogenesis and lipolysis, hyperinsulinemia, and release of NEFAs cause a positive feedback loop that contributes to adipocyte hypertrophy [30]. But no direct influence of clozapine on fat storage and expression levels of an early marker of adipogenesis peroxisome-proliferator-activated-receptor-gamma (PPAR- $\gamma$ ) and a late marker of adipogenesis fatty-acid-binding-protein-4 (FABP-4) is found in human preadipocytes and mature adipocytes in vitro [31]. Unlike clozapine, which does not seem to have a significant effect on lipogenesis in mature adipocytes, haloperidol inhibits probably the lipogenic capacity of adipocytes in females, contributing to its lower propensity to induce weight gain [32].

Adipose tissue dysfunction as an intermediate between obesity and resulting inflammation and insulin resistance appears to be characterised by an increase in pro-inflammatory adipokines and a decrease in anti-inflammatory adipokines such as adiponectin [33]. Additionally, iron plays a role in the regulation of adipose tissue function, especially in the expression of adipokines [34]. In contrast to liver [24], we do not find ferric iron deposits in adipocytes. Hepatic ferric iron accumulation is associated with oxidative stress, which is increased in liver [24] but not in adipose tissue (Table 1). Low cellular iron level leads to obese-resistant white adipose tissue and limits lipid uptake in enterocytes, reducing hypertrophy, lipid peroxidation and inflammation of adipose tissue, thereby reducing the development of hepatic steatosis and insulin resistance [35]. Hpcidin is considered the key regulator of iron metabolism. Haloperidol or clozapine medicated female rats have higher serum ferritin levels and higher hepcidin levels in liver [26] and adipose tissue (Figure 2C). In mouse models of obesity, increased adiposity leads to iron deposits only in the epididymal adipose tissue


(eAT) of male mice. This tissue-specific iron accumulation causes local adipose tissue insulin resistance, macrophage accumulation, and finally inflammation [36].

Clozapine might play an important role in the regulation of the adipokines leptin and adiponectin, and total ghrelin [37]. In healthy individuals, leptin and total body adipose mass are positively correlated [38], but we find this correlation for leptin and perirenal VAT only in the control- and haloperidol-medicated groups (Supplement Figure 2 A-D). A negative correlation between obesity and circulating adiponectin is also well-established [9], and adiponectin concentrations increase concomitantly with weight loss, whereas decreased adiponectin levels are associated with insulin resistance and hyperinsulinemia [39]. In our study, we find an increase of serum adiponectin levels in all drug-medicated groups, highest in the two clozapine groups [23] and like expected, a negative correlation between serum adiponectin and perirenal VAT for male haloperidol-medicated rats (Supplement Figure 2 E). After binding to its receptors, Adipo-R1 and R2, adiponectin initiates the phosphorylation of adenosine-monophosphate-activated-protein-kinase (AMPK), a key regulator of hepatic energy and lipid homeostasis [40]. Activation of AMPK suppresses fatty acid and cholesterol biosynthesis by inhibiting the activity of acetyl-CoA-carboxylase and 3-hydroxy-3-methyl-glutaryl-CoA-reductase. Furthermore, activation of AMPK inactivates the SREBPs, the transcriptional regulators of lipogenesis, thereby inhibiting lipogenic genes. Clozapine medication suppresses AMPK-activity in male SD rats [41], thereby allowing upregulation of SREBP-1c in male and SREBP-2 in female rats [24]. SREBP-1c stimulates hepatic triglyceride accumulation in males [24], and SREBP-2 induces serum cholesterol production in females and partially in males [23]. Haloperidol has no effect on the SREBPs [24] and may inhibit cholesterol biosynthesis [42]. In healthy animals, glycogen-bound subunit AMPK- $\beta$  suppresses activation of AMPK [43]. Reducing the glycogen stores allows AMPK to inhibit lipogenesis and to direct cell metabolism towards fatty acid oxidation [44]. Downregulation of glycogen synthesis – probably by GSK3- $\beta$  activation – subsequently leads to hyperglycemia in female clozapine-treated rats [23]. In male rats medicated with clozapine, the glycogen stores are only slightly reduced. The both atherogenic serum indices cholesterol/HDL and LDL/HDL ratios (Table 1) are increased significantly in the male and weakly in the female clozapine-medicated groups. The same is seen in schizophrenic patients [45]. Therefore, we hypothesize that the increase of adiponectin levels counteracts cholesterol-regulated lipogenesis by stimulating the hepatic adiponectin receptors, which activate AMPK as seen in C2C12 myotube cells [46]. AMPK activation induces peroxisome proliferator-activated-receptor- $\alpha$  (PPAR- $\alpha$ ) signalling, which in turn induces anti-inflammatory signalling, reduces adipose tissue derived factors that could stimulate steatosis [18], and inhibits SREBP activity, thereby reducing the development of hepatic steatosis [47]. Activation of PPAR- $\alpha$  causes a reduction in circulating triglycerides and an increase in the uptake of free fatty acids. Surprisingly, chronic treatment with clozapine or haloperidol lowers adipocyte NEFAs in the male medicated groups as seen under olanzapine medication in mice and humans, but not in serum or liver (Table 1), whereby insulin resistance, adiposity, and diabetes are typically associated with elevated free fatty acids [48]. Ghrelin is elevated during fasting periods and decreases rapidly after food intake. Ghrelin not only promotes adiposity by activating hypothalamic orexigenic neurons, but also directly stimulates fat storage-related proteins in adipocytes like Acetyl-CoA-carboxylase, fatty-acid-synthase, lipoprotein-lipase and perilipin, thereby stimulating intra-cytoplasmic lipid accumulation [49]. In humans, perilipin is elevated in obese subjects and its level is enhanced in steatotic hepatocytes [50]. In our study the protein expression of perilipin-A and HSL in adipose tissue is decreased under clozapine medication in both sexes (Figure 2A.-B.), indicating no effect of the antipsychotic drug on lipolysis and lipogenesis. Therefore, we suspect that the serum ghrelin levels are elevated due to the 12-hour starvation period of the rats before blood and organ collection.

In adipocytes, we find slightly decreased glucose in all medicated groups, decreased glycogen in males and increased glycogen in females (Table 1) associated with decreased protein levels of the insulin-dependent Glut-4 in the clozapine-medicated groups. Under conditions of low insulin, most Glut-4 is sequestered in intracellular vesicles in muscle cells and adipocytes. Increasing insulin levels, induced by increasing blood glucose, cause the uptake of glucose into the adipocytes by Glut-4 which

is quickly incorporated into the plasma membrane of the cell when insulin binds to its membrane receptors. Thereby, insulin-stimulated glucose uptake enhances glycolysis, tricarboxylic-acid-cycle-flux and oxidative phosphorylation with mitoROS (reactive oxygen species) production in a glucose-dependent manner. Downregulation of insulin receptors as well as insulin resistance protect against glucose oversupply [51]. Haloperidol slightly increases serum insulin levels [23] associated with an increase of IR- $\beta$  in both sexes and a decrease of Glut-4 in adipose tissue of males. Clozapine treatment leads to a lower protein level of Glut-4 in males and females associated with an increase in IR- $\beta$  in males and a decrease in females (Figure 2 D-E). Under normal conditions, the insulin level should rise dramatically -at least in females under clozapine medication- due to the excessively elevated serum glucose level, with the result that more glucose would be taken up via Glut-4 into the adipocytes and would be stored there as triglycerides. However, despite strongly increased serum glucose levels, no more insulin is secreted. The fact that NEFAs are not elevated in any of the tissues examined also argues against a classic insulin resistance (Table 1). Murashita et al. [52] observed elevated blood glucose levels also in male clozapine-treated SD rats. Clozapine directly inhibits insulin secretion through membrane depolarisation and activation of protein kinase C in pancreatic cells [53]. Smith et al. [54] found that SGA medication in rats can interfere with glucose metabolism, not by acutely inducing insulin resistance but by increasing pancreatic glucagon secretion and thus stimulating hepatic glucose production. Hepatic glucose production derives from glycogen breakdown (glycogenolysis) and from de novo synthesis of glucose (gluconeogenesis). We found liver glycogenolysis in females under clozapine [24] and the significantly increased hepatic pyruvate and lactate levels in males favour suspected gluconeogenesis in males (Table 1). Glut-4 expression in insulin-sensitive tissues decreases in association with whole-body insulin resistance and high levels of inflammatory markers in a rat model of metabolic syndrome [55]. Clozapine is known to stimulate proinflammatory gene transcription in human adipocytes [56]. Studies implicate a complex network by which glucose sensing through Glut-4 in muscle and adipocytes may operate to integrate whole-body energy metabolism [57], and adipocyte insulin sensitivity appears critical for whole-body homeostasis.

We have investigated the impact of clozapine and haloperidol on metabolically active tissues such as liver and VAT and their neuronal control in the hypothalamus, the control center of body weight and homeostasis. It has been shown that the affected organs do not act independently of each other, but rather influence each other in a sex-specific manner (Figure 4). Under clozapine medication, increased hypothalamic lactate level in females points to enhanced glycolysis with a higher glucose demand to produce the required energy. The two insulin-independent glucose transporters Glut-1 and Glut-3 were upregulated and this refers to glucose deficiency in endothelial cells and ultimately in neurons and astrocytes. We found excessive fasting glucose in the serum of female rats medicated with clozapine. The glucose probably originates from the hepatic glycogen depot, which was either no longer replenished or had been emptied. Glycogenolysis in female liver contrasts with the findings in male rats, which compensate their hypothalamic glucose deficiency through gluconeogenesis and also carry out increased glycolysis with high hepatic pyruvate and lactate values. In contrast to VAT, the liver shows high oxidative stress for both sexes. In females, hemosiderin is deposited in the liver, while males show an iron deficiency, which is reflected in a low hemoglobin content and a reduced cytochrome level [26]. Males have extremely high hepatic triglyceride levels and increased lipid droplets. This is consistent with the upregulation of SREBP-1c in males and SREBP-2 in females, which have high serum cholesterol, HDL-cholesterol and LDL-cholesterol levels. Based on high triglyceride values, high NEFAs levels, increased liver mass and the accumulation of lipid droplets, we assume the onset of steatosis in male rats under clozapine.



**Figure 4.** Schematic diagram of the organ connections (crosstalk) derived from the results of the serum, liver, hypothalamus, and adipose tissue examinations under clozapine medication. The explanations can be found in the text below.

Unexpectedly, adipose tissue shows a much less disturbed lipid and glucose homeostasis without oxidative stress than the liver. Low cellular iron levels limit the absorption of lipids, thereby reducing hypertrophy, reduce lipid peroxidation and inflammation and decrease the development of steatosis and insulin resistance in the liver. Clozapin suppresses AMPK-activity and upregulates SREBP-1c in males, leading to hepatic triglyceride accumulation, and SREBP-2 in females increasing

the cholesterol level in serum. The increased release of adiponectin counteracts this effect and activates the hepatic AMPK and reduces fatty acid and cholesterol biosynthesis. In addition, AMPK can inhibit lipogenesis by depleting glycogen depots, as we were able to show in female rats. Clozapine does not cause classic insulin resistance in our animals. Despite the very high serum glucose level in female animals, insulin secretion did not raise.

#### 4. Limitations

The limitations of our studies, such as the social isolation of the test animals or the small group size, have already been discussed in the four previous publications [23–26]. Among the major limitations of our study on adipose tissue, is the fact that we only studied perirenal VAT, although this only accounts for 20% of the total adipose tissue. [9]. Clozapine has no acute effect on food intake in rats [Sorge [23,58]], unlike in patients, in whom, however, no precise studies exist on the amount of food ingested and the amount of energy consumed under clozapine or haloperidol. The cells of smaller mammals such as rats or mice consume more oxygen and nutrients per unit time. This, as well as the metabolic differences between humans and rats, leads to the assumption that the results from our animal model can only be transferred to humans to a limited extent and do not always mimic clinical findings [59], although rodent adipose tissue shows great similarity to human adipose tissue [60]. Nevertheless, the results of our study show many similarities with the clinical findings under antipsychotic medication and some results can only be obtained in an animal model. In our studies, male and female animals were tested. It has been shown that there are many sex-related differences. Increasing clinical research shows also that male and female patients differ in their metabolic expression of medication [61].

#### 5. CONCLUSION

The examination of the perirenal adipose tissue shows lower metabolic changes under haloperidol and clozapine than in serum and especially in liver. The differences between male and female animals were also smaller in adipocytes (Table 1). Lactate or pyruvate levels in liver and hypothalamus of male rats indicate a shift to glycolysis under clozapine, but this is also less pronounced in adipose tissue, probably because adipocyte metabolism depends on oxidative phosphorylation. Under clozapine medication, glucose seems to be stored in the male liver as triglyceride deposits rather than in adipose tissue. Microscopic examination of adipose cells indicates slight hypertrophy but without increased food intake [23]. However, the increase in adipose and liver tissue does not explain total weight gain, suggesting an additive involvement of subcutaneous adipose tissue. Increased protein expression of the IR- $\beta$  and decreased Glut-4 expression indicate insulin resistance under haloperidol and in clozapine-medicated males but not in females under clozapine treatment. Lipolysis or lipogenesis seem to be unaffected in adipocytes, while liver shows adipogenic transformation in male clozapine-treated rats leading to steatosis. Clozapine induces hepatic SREBP-1c expression in male rats [24], the upstream regulator of fatty-acid-synthase. Female rats are protected against fatty liver by low glycogen levels. Downregulation of glycogen synthesis and upregulation of glycogenolysis – probably by GSK3- $\beta$  activation – subsequently leads to higher serum glucose levels in female clozapine-treated rats. It is still an open question whether the glucose deficit of the neuronal cells in the hypothalamus [25] or the glycogen depletion in liver cells is responsible for the wide range of peripheral metabolic changes. The disturbance of glucose and lipid metabolism is tissue- and sex-dependent and obesity shows sexual dimorphism as seen in Palmer and Clegg [61].

#### 6. Methods

##### 6.1. Ethics statement

All experiments were carried out in accordance with the laws of the local authorities for animal experiments approved by the Landesamt für Natur, Umwelt- und Verbraucherschutz NRW, Recklinghausen, Germany; Reference number 9.93.2.10.34.07.227. The feeding experiment took place

in the rooms of the animal testing facility of the Heinrich Heine University, Düsseldorf. The study was conducted in compliance with ARRIVE guidelines 2.0.

### 6.2. Experimental design

30 male and 30 female healthy 10-weeks-old Sprague Dawley rats from Taconic (Denmark) housed in 2x5 groups of 6 same-gender animals (sibling group) from the same litter with free access to water and ground food pellets (maintenance diet (Altromin Spezialfutter GmbH, Germany) with 19% crude protein, 4% crude fat and additionally 15% fat to facilitate drug uptake). The animals were maintained on a 12:12 h light/dark cycle at 21°C and 60% humidity. For acclimatization, the animals were handled every day and weighted 2 times per week from postnatal day (PD) 71. On PD 78 the rats were separated from each other and individually housed during the test period. Two animals per sibling group were divided into the control, haloperidol-medicated and clozapine-treated groups, resulting in 10 animals per group (n=10). To avoid puberty effects antipsychotic drug treatment started at PD 85 (week 13) to PD 165 (week 25). Male (n=10 per group) or female (n=10 per group) rats were fed orally each day with clozapine (Leponex®, Novartis, Germany with 20mg/1kg body weight (BW)/day in ground pellets), corresponding to an effective average dose rate to  $18.5 \pm 0.26$  mg/kg BW for males and  $17.7 \pm 0.38$  mg/kg BW for females [23]. Haloperidol (Haloneurol®, Hexal, Germany) was fed with 1mg/1kg BW/day in ground pellets, corresponding to an effective daily average dose of  $0.8 \pm 0.03$  mg/kg BW for males and  $0.6 \pm 0.08$  mg/kg BW for females [23]. Results were compared to those in male and female control groups fed only with ground pellets. On PD 169 (week 25) twelve hours after food removal, the animals were anaesthetized by Pentobarbital (Narcoren, Merial, Germany), and perirenal fat pads as part of the intra-abdominal VAT were removed, weighted, and stored at -80°C in an ultra-low temperature cooler. All metabolic were determined for all test animals.

### 6.3. Determination of percentage weight, neutral fat content and adipocyte area value

To calculate the percentage weight, we took the difference between the final and initial weight and calculated its percentage of total weight.

0.5 g adipose tissue was homogenized on ice in 1 ml RIPA buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1% Na-deoxycholate, 0.1 % LDS, 1 mM EDTA, 3% Triton with protease inhibitor (complete mini, EDTA free, Roche Diagnostics, Germany) overnight at 4°C and centrifuged at 1000 g for 10 min at 4°C. The pellet, containing blood vessels and collagen, was discarded and the aqueous supernatant was collected for measurement of triglycerides, NEFAS, glucose, glycogen and lactate. The fat phase was dissolved by shaking with 1 ml diethyl ether overnight at 39.5°C. The liquid phase was transferred in weighted tubes, the volatile diethyl ether was removed, and the weight of neutral fat was determined.

The adipocyte area value was determined microscopically. 20 µm sections of frozen perirenal VAT were fixed with 4 % buffered neutral formalin at room temperature for 5 min, washed three times with distilled water and air-dried overnight. The slices were stained in Mayers hematoxylin (Roth, Germany) for 5 min, rinsed in cool running water for 5 min, stained in 0.5% aqueous eosin-G solution (Roth, Germany), rinsed with distilled water, dehydrated in Roti-Histol (Roth, Germany), and mounted in Roti-Histokitt (Roth, Germany). Two clusters on two different sections of each animal, each cluster enclosing 200 adipocytes, were marked by x, the two areas were calculated with Axiovision (Zeiss, Germany) and averaged.

### 6.4. Determination of ferric iron by Perl's Prussian blue staining of frozen fat tissue sections

We analysed ferric iron (Fe<sup>3+</sup>) by Perl's Prussian blue staining [62]. 30 µm sections of frozen adipose tissue were fixed with 4 % buffered neutral formalin at room temperature for 5 min, washed three times with distilled water and air-dried. After descending alcohol series, the tissue was prestained in 10% potassium-ferrocyanide for 5 min and stained with equal parts of a mixture of 2% potassium-ferrocyanide solution and 1% hydrochloric acid for 30 min at 37°C. After washing with

distilled water, the slides were counterstained with nuclear fast red aluminium sulphate solution (Roth, Germany) for 5 min, rinsed with distilled water, dehydrated in Roti-Histol (Roth, Germany) and mounted in Roti-Histokitt (Roth, Germany).

#### *6.5. Determination of triglycerides and non-esterified fatty acids (NEFAs) in adipocytes, cholesterol/HDL-cholesterol ratio in liver and serum and LDL-cholesterol/HDL-cholesterol ratio in serum*

Triglyceride levels were determined by automated counter in the clinical laboratory of the LVR Klinikum Düsseldorf. NEFAs were determined in serum, liver, and adipocyte lysate by NEFA-HR (2) assay kit (436-91995, Fujifilm, Germany) following manufacturer's recommendations. The two atherogenic indices cholesterol/HDL-cholesterol ratio and LDL-cholesterol/HDL-cholesterol ratio were calculated from the values for cholesterol, HDL-cholesterol and LDL-cholesterol in the serum and cholesterol/HDL-cholesterol ratio in liver.

#### *6.6. Determination of glucose, glycogen, and lactate in lysates of fat tissue*

Glucose levels in adipocyte lysate were determined in the whole cell lysate by glucose colorimetric assay kit (MAK263, Sigma-Aldrich, Germany) following manufacturer's recommendations. Lactate was determined in the whole cell lysate by a lactate LOX-PAP test (LT-LC0056, Labor+Technik, Eberhard Lehmann GmbH, Germany) following manufacturer's recommendations.

Glycogen content was determined by the anthron method. 0.25 g adipose tissue was mixed with 1.25 ml HClO<sub>4</sub> (6%, ice cooled) and heated for 10 min at 80°C. The solution was centrifuged for 5 min at 960 rpm and 4°C. 2.5 ml methanol was added to 1.0 ml supernatant, heated for 30 min at 37°C and centrifuged for 10 min at 8000 rpm and room temperature. The pellet was dissolved in 200 µl aqua dest. mixed with 400 µl anthron solution (0.2 % anthron (Sigma Aldrich, Germany) in concentrated H<sub>2</sub>SO<sub>4</sub>) and heated for 15 min at 95°C. After cooling to room temperature, the extinction of the green solution could be measured at 620 nm and the levels of glycogen were calculated in mg/g of fat tissue using a standard curve.

#### *6.7. Determination of oxidative stress in perirenal fat of all rats and of estrogen levels in serum*

Lipid peroxidation in the adipose tissue was determined through the production of TBARS as previously described by [63]. 50 µl SDS (8.1%), 375 µl acetic acid (20%, adjusted to pH 3.5 with 1 N NaOH) and 375 µl TBA (0.8% aqueous solution of thiobarbituric acid) were added to 50 µl cell lysate and filled-up to 1 ml with aqua dest. The solution was boiled for 60 min and cooled down to room temperature. 0,5 ml n-butanol + pyridine (15:1) were added, thoroughly mixed, and centrifuged for 10 min at 2000 rpm at room temperature. Spectral absorption was measured at 532 nm and malondialdehyde equivalents were calculated in µmol/ml by a standard curve.

Estrogen levels were measured in female rat serum with a rat estrogen ELISA kit (EKR3010, Nordic Biosite, Germany) following manufacturer's recommendations.

#### *6.8. Western blot analysis of adipocyte lysates of the drug-medicated rats and controls*

Protein expression was identified by Western blot analysis in the whole cell lysate. 0.5 g fat tissue was homogenized on ice in 1.0 ml tissue extraction buffer I (50 mM Tris, pH 7.4, 250 mM NaCl, 5mM EDTA, 2mM Na<sub>3</sub>VO<sub>4</sub>, 1mM NaF, 20 mM Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub>, Thermo Fisher scientific, Germany) with protease inhibitor cocktail (complete mini, Roche, Germany) shaken overnight at 4°C and centrifuged at 13000 g for 20 min at 4°C. Total protein content was measured using the DC Protein Assay from Bio Rad (Germany). 20 µg protein from each animal was separated on NuPAGE™ 4-12% Bis-Tris gels (Thermo Fisher scientific, Germany) and blotted onto Invitrolon membranes (Thermo Fisher scientific). Membranes were then blocked with Roti Block (Carl Roth, Germany) in TBS-T (0.5:10) at RT for 30 min. The blocked PVDF membranes were incubated overnight at 4°C either with rabbit polyclonal perilipin A antibody (62 kilodalton (kDa), 1:1000, PA1-1051, Thermo Fisher Scientific, Germany), rabbit polyclonal HSL antibody (80 kDa, 1:1000, PA1-16966, Thermo Fisher Scientific,

Germany), goat polyclonal hepcidin antibody (9 kDa; 1:1000, sc-240553, Santa Cruz Biotechnology, Germany), rabbit polyclonal Glut-4 antibody (54 kDa, 1:1000, ab216661, Abcam, Germany) and mouse monoclonal IR- $\beta$  antibody (95 kDa, 1:1000, sc-373975, Santa Cruz Biotechnology, Germany) in TBS-T and then probed at room temperature for 1 h with the HRP-conjugated secondary antibody (against rabbit 1:20000, ab97051, Abcam, mouse 1:20000, sc-2004, Santa Cruz, or goat 1:20000, sc-2064, Santa Cruz respectively). Resulting autoradiographs by autoradiography films (Santa Cruz Biotechnology, Germany) (Supplement Figure 1A) were measured by densitometry using Gene Snap and Gene Tools software (Syngene, Synoptics Ltd, United Kingdom). The membranes were routinely stained with Ponceau-S (Supplement Figure 1B) before immunostaining to confirm correct sample loading and the transfer of equivalent amounts of protein [64].

### 6.9. Statistical analysis

Our study is an exploratory study. Therefore, we chose 10 animals per group (male and female controls, male and female haloperidol-medicated rats, male and female clozapine-medicated animals) to obtain statistically valid conclusions.

We have re-analysed the following previously published results [23;24]: Final body weight of male clozapine-treated animals; data of serum leptin, adiponectin, ghrelin; serum and liver cholesterol, HDL-cholesterol and LDL-cholesterol.

Statistical analysis was performed using IBM SPSS version 25 for windows. All data were represented as mean  $\pm$  SE. The data were examined by two-way ANOVA with the between-subject factors MEDICATION and SEX and one-way ANOVA with MEDICATION as factor. In case of significant group effects appropriate posthoc tests were carried out between the groups. Sex-dependent effects were examined by t-tests for independent samples. Due to the small number of animals per group and the explorative character of the study,  $p$ -values  $\leq 0.05$  were considered as statistically significant and multiple testing was presented without  $\alpha$ -adjustment. A bivariate correlation procedure with Spearman-Rho coefficient was carried out to test the relationship between HSL and perilipin-A, between serum adiponectin, leptin and body fat mass related to 1 kg BW, ghrelin and perilipin-A, IR- $\beta$  and Glut-4 with  $p \leq 0.05$  as significant. Significant correlations are displayed as simple scatterplots performed by spss in the Supplement Figure 2.

**Supplementary Materials:** Supplement Figure 1a. and b. Supplement Figure 2. Supplement Figure 3a. and b.

**Author Contributions:** MLB engaged in study conception and in data collection, analyzed the data and wrote introduction, material and result part. KF and MLB wrote abstract, discussion and conclusion. All authors have read and approved the manuscript.

**Funding:** All costs were borne by the general research budget of the LVR-Klinikum/ Heinrich-Heine-University, Duesseldorf. Study design, data collection, data analysis, interpretation of data and manuscript preparation were not influenced by the funding body. Third-party funds were not raised. .

**Institutional Review Board Statement:** All animal experiments were carried out in accordance with the laws of the local authorities for animal experiments approved by the LANUV (Landesamt für Natur, Umwelt- und Verbraucherschutz NRW, Recklinghausen) with the following reference number 9.93.2.10.34.07.227. The LANUV is assigned to the Landesministerium für Umwelt und Naturschutz, Landwirtschaft und Verbraucherschutz, Duesseldorf. The LANUV, department 81, animal experimentation matters, grants approval of an animal experiment in connection with the animal care and use committee. The animal husbandry before and during the feeding experiment, as well as the killing, took place in the rooms of the animal experiment facility (TVA) of the Heinrich Heine University, Düsseldorf under the control of Dr. Treiber, director of the TVA.

**Data Availability Statement:** The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

**Acknowledgments:** We thank Dr. M. Jänner, LVR Klinikum, Duesseldorf, who kindly shared her expertise in statistical analysis of our results. Kirsten Nürenberg and her assistants kindly carried out the measurement of triglycerides in adipose tissue.

**Conflicts of Interest:** All authors declare that they have no competing interests.

## Abbreviations

**ACC:** Acetyl-CoA carboxylase; **AGRP:** agouti related peptide; **AMPK:** AMP-activated protein kinase; **APD:** antipsychotic drug; **ATGL:** adipocyte triglyceride lipase; **eAT:** epididymal adipose tissue; **FABP-4:** fatty acid binding protein-4; **FGA:** first-generation antipsychotic; **GLUT:** Glucose transporter; **HMGR:** 3-hydroxy-3-methyl glutaryl CoA reductase; **HSL:** Hormonsensitive lipase; **IR- $\beta$ :** Insulin receptor- $\beta$ ; **IL-6:** interleukin-6; **kDa:** kilodalton; **MGL:** monoacylglycerol lipase; **NAFLD:** non-alcoholic fatty liver disease; **NEFAs:** non-esterified fatty acids; **NPY:** neuropeptide Y; **PK-B:** protein kinase-B; **PD:** postnatal day; **PPAR- $\alpha$ :** peroxisome proliferator-activated receptor- $\alpha$ ; **PPAR- $\gamma$ :** peroxisome proliferator-activated receptor- $\gamma$ ; **ROS:** reactive oxygen species; **SCAT:** subcutaneous adipose tissue; **SGA:** second generation antipsychotic; **SREBP:** sterol regulatory element-binding protein; **TAG:** triacylglycerols; **TNF- $\alpha$ :** tumor necrosis factor- $\alpha$ ; **TG:** triglyceride; **T2D:** type 2 diabetes; **VAT:** visceral adipose tissue; **WAT:** white adipose tissue

## References

- Dayabandera, M.; Hanwella, R.; Ratnatunga, S.; Seneviratne, S.; Suraweera, C.; de Silva, V.A. Antipsychotic-associated weight gain: management strategies and impact on treatment adherence. *Neuropsychiatr Dis Treat.* 2017, 13,2231-2241. PMID: 28883731 PMCID: PMC5574691 DOI: 10.2147/NDT.S113099
- Alonso-Pedrero, L.; Bes-Rastrollo, M.; Marti, A. Effects of antidepressant and antipsychotic use on weight gain: a systematic review. *Obesity Rev.* 2019,20(12),1680-1690. PMID: 31524318 DOI: 10.1111/obr.12934
- Ferreira, V.; Grajales, D.; Valverde, A.M. Adipose tissue as a target for second generation (atypical) antipsychotics: a molecular review. *BBA Mol Cell Biol Lipids.*2020,1865(2),158534. PMID:31672575 doi: 10.1016/j.bbalip.2019.158534
- Spalding, K.L.; Arner, E.; Westermark, P.O.; Bernard, S.; Buchholz, B.A.; Bergmann, O.; Blomquist, L.; Hoffstedt, J.; Näslund, E.; Britton, T.; Concha, H.; Hassan, M.; Ryden, M.; Frisen, J.; Arner P. Dynamics of fat cell turnover in human. *Nature.* 2008,453(7196),783-787. PMID:18454136doi: 10.1038/nature06902
- Vestri, H.S.; Maianu, L.; Moellering, D.R.; Garvey, W.T. Atypical antipsychotic drugs directly impair insulin action in adipocytes: effects on glucose transport, lipogenesis, and antilipolysis. *Neuropsychopharmacol.* 2006,32(4),765-772. PMID:16823387 doi: 10.1038/sj.npp.1301142
- Wilson, J.L.; Enriori, P.J. A talk between fat tissue, gut, pancreas and brain to control body weight. *Mol Cell Endocrinol.* 2015,418(2),108-119. PMID:26316427 doi: 10.1016/j.m.c.2015.08.22
- Yang, A.; Mottillo, E.P. Adipocyte lipolysis: from molecular mechanisms of regulation to disease and therapeutics. *Biochem J.* 2020,477(5),985-1008. PMID:32168372 PMCID: PMC7187988 doi:10.1042/BCJ20190468
- Rotondo, F.; Ho-Palma, A.C.; Remesar, X.; Fernandez-Lopez, J.A.; del Mar Romero, M.; Alemany, M. Glycerol is synthesized and secreted by adipocytes to dispose of excess glucose, via glycerogenesis and increased acyl-glycerol turnover. *Scientific Reports.* 2017,7,8983. PMID:28827624 PMCID:PMC5567128 doi: 10.1038/s41598-017-09450-4
- Meyer, L.K.; Ciaraldi, T.P.; Henry, R.R.; Wittgrove, A.C.; Phillips, S.A. Adipose tissue depot and cell size dependency of adiponectin synthesis and secretion in human obesity. *Adipocyte.* 2013,2(4),217-226. PMID:24052897 PMCID:PMC3774697 doi:10.4161/adip.24953
- Chau, Y.Y.; Bandiera, R.; Serrels, A.; Martinez-Estrada, O.M.; Qing, W.; Lee, M.; Slight, J.; Thornburn, A.; Berry, R.; McHaffie, S.; Stimson, R.H.; Walker, B.R.; Chapuli, R.M.; Schedl, A.; Hastie, N. Visceral and subcutaneous fat have different origins and evidence supports a mesothelial source. *Nat Cell Biol.* 2014,16(4),367-375. PMID:24609269 PMCID: PMC4060514 doi:10.1038/ncb2922
- Huang, N.; Mao, E.W.; Hou, N.N.; Liu, Y.P.; Han, F.; Sun, X.D. Novel insight into perirenal adipose tissue: a neglected adipose depot linking cardiovascular and chronic kidney disease. *World J Diabetes.* 2020,11(4),115-125. PMID:32313610 PMCID:PMC7156295 doi:10.4239/wjd. v11.i4.115
- Ibrahim, M.M. Subcutaneous and visceral adipose tissue: structural and functional differences. *Obes Rev.* 2010,11(1),11-18. PMID:19656312 doi:10.1111/j.1467-789X.2009.00623.x
- Booth, A.; Magnuson, A.; Fouts, J.; Foster, M.T. Adipose tissue: an endocrine organ playing a role in metabolic regulation. *Horm Mol Biol Clin Invest.* 2016,26(1),25-42. PMID:26910750 doi:10.1515/hmbci-2015-0073
- Friedman, J.M. Leptin and the regulation of body weight. *Keio J Med.* 2011,60(1),1-9. PMID:21460597 doi: 10.2302/kjm.60.1
- Stern, J.H.; Rutkowski, J.M.; Scherer, P.E. Adiponectin, leptin, and fatty acids in the maintenance of metabolic homeostasis through adipose tissue crosstalk. *Cell Metab.* 2016,23(5),770-784. PMID:27166942 PMCID:PMC4864949 doi: 10.1016/j.cmet.2016.04.011

16. Achari, A.E.; Jain, S.K. Adiponectin, a therapeutic target for obesity, diabetes, and endothelial dysfunction. *Int J Mol Sci.* 2017,18,1321. PMID:28635626 PMCID:PMC5486142,doi:10.3390/ijms18061321
17. Ukkola, O.; Santaniemi, M. Adiponectin: a link between excess adiposity and associated comorbidities? *J Mol Med.* 2002,80,606-702. PMID:12436346 doi.org/10.1007/s00109-002-0378-7
18. Shabalala, S.C.; Dlodla, P.V.; Mabasa, L.; Kappo, A.P.; Basson, A.K.; Pfeiffer, C.; Johnson, R. The effect of adiponectin in the pathogenesis of non-alcoholic fatty liver disease (NAFLD) and the potential role of polyphenols in the modulation of adiponectin signaling. *Biomed Pharmacotherapy.* 2020,131,110785. PMID:33152943 doi: 10.1016/j.Biopha.2020.110785
19. Gatselis, N.K.; Ntaios, G.; Makaritsis, K.; Dalekos, G.N. Adiponectin: a key playmaker adipocytokine in non-alcoholic fatty liver disease. *Clin Exp Med.* 2014,14(2),121-131. PMID:23292294 doi: 10.1007/s10238-012-0227-0
20. Thompson, N.M.; Gill, D.A.S.; Loveridge, N.; Houston, P.A.; Robinson, I.C.A.F.; Wells, T. Ghrelin and desoctanoyl ghrelin promote adipogenesis directly in vivo by a mechanism independent of the type 1a growth hormone secretagogue receptor. *Endocrinol.* 2004,145(1),234-242. PMID: 14551228 doi:10.1210/en.2003-0899
21. Miao, H.; Pan, H.; Wang, L.; Yang, H.; Zhu, H.; Gong, F. Ghrelin promotes proliferation and inhibits differentiation of 3L3-L1 and human primary preadipocytes. *Front Physiol.* 2019,article10,1296. PMID:31681009 PMCID:PMC6798085 doi: 10.3389/fphys.2019.01296
22. Carvalho, E.; Schellhorn, S.E.; Zabolotny, J.M.; Martin, S.; Tozzo, E.; Peroni, O.D.; Houseknecht, K.L.; Mundt, A.; James, D.E.; Kahn, B.B. GLUT4 overexpression or deficiency in adipocytes of transgenic mice alters the composition of GLUT4 vesicles and the subcellular localization of GLUT4 and insulin-responsive aminopeptidase. *J Biol Chem.* 2004,279(20),21598-21605. PMID: 14985357 doi:10.1074/jbcM312269200
23. von Wilmsdorff, M.; Bouvier, M.L.; Henning, U.; Schmitt, A.; Schneider-Axmann, T.; Gaebel, W. The sex-dependent impact of chronic clozapine and haloperidol treatment on characteristics of the metabolic syndrome in a rat model. *Pharmacopsychiatry.* 2013,46(1),1-9. PMID: 22915487 doi: 10.1055/s-0032-1321907
24. von Wilmsdorff, M.; Bouvier, M.L.; Henning, U.; Schmitt, A.; Schneider-Axmann, T.; Gaebel, W. Sex-dependent metabolic alterations of rat liver after 12-week exposition to haloperidol or clozapine. *Horm Metab Res.* 2014,46(11),782-788. PMID: 25105542 doi:10.1055/s-0034-1384605
25. Bouvier, M.L.; Fehsel, K.; Schmitt, A.; Meisenzahl-Lechner, E.; Gaebel, W.; von Wilmsdorff, M. Sex-dependent alterations of dopamine receptor and glucose transporter density in rat hypothalamus under long-term clozapine and haloperidol medication. *Brain Behav.* 2020,e01694. PMID: 32525610. doi: 10.1002/brb3.1694
26. Bouvier, M.L.; Fehsel, K.; Schmitt, A.; Meisenzahl-Lechner, E.; Gaebel, W., von Wilmsdorff, M. Sex-dependent effects of long-term clozapine or haloperidol medication on red blood cells and liver iron metabolism in Sprague Dawley rats as a model of metabolic syndrome. *BMC Pharmacol and Toxicol.* 2022,23,8. PMID: 35033194 doi:10.1186/s40360-021-00544-4
27. Manu, P.; Dima, L.; Shulman, M.; Vancampfort, D.; De Hert, M.; Correll, C.U. Weight gain and obesity in schizophrenia: epidemiology, pathobiology, and management. *Acta Psychiatr Scand.* 2015,132,97-108. PMID:26016380 doi:10.1111/acps.12445
28. Guo, D.H.; Yamamoto, M.; Hernandez, C.M.; Khodadadi, H.; Baban, B.; Stranahan, A.M. Beige adipocytes mediate the neuroprotective and anti-inflammatory effects of subcutaneous fat in obese mice. *Nature Communications.* 2021,12,4623. PMID:34330904 PMCID:PMC8324783 doi: 10.1038/s41467-021-24540-8
29. Engfeldt, P.; Arner, P. Lipolysis in human adipocytes, effects of cell size, age and regional differences. *Horm Metabol Res Suppl.* 1988,19,26-29. PMID:3069692
30. Perez-Torres, I.; Gutierrez-Alvarez, Y.; Guarner-Lans, V.; Diaz-Diaz, E.; Pech, M.L.; Del Carmen Caballero-Chacon, S. Intra-abdominal fat adipocyte hypertrophy through progressive alteration of lipolysis and lipogenesis in metabolic syndrome rats. *Nutrients.* 2019,11,1529-1540. PMID:31284400 PMCID:PMC6683042 doi: 10.3390/nu11071529
31. Löffler, D.; Landgraf, K.; Körner, A.; Kratzsch, J.; Kirkby, K.C.; Himmerich, H. Modulation of triglyceride accumulation in adipocytes by psychopharmacological agents in vitro. *J Psychiatric Res.* 2016,72,37-42. PMID:26524413 doi: 10.1016/j.psychires.2015.10.008
32. Sertié, A.L.; May Suzuki, A.; Sertié, R.A.L.; Andreotti, S.; Lima, F.B.; Passos-Bueno, M.R.; Gattaz, W.F. Effects of antipsychotics with different weight gain liabilities on human in vitro models of adipose tissue differentiation and metabolism. *Prog Neuro-Psychopharm Biol Psych.* 2011,35,1884-1890, PMID:21840366 doi: 10.1016/j.pnpbp.2011.07.017
33. Westerink, J.; Olijhoek, K.; Koppen, A.; Faber, D.R.; Kalkhoven, E.; Monajemi, H.; van Asbeck, B.S.; van der Graaf, Y.; Visseren, F.L.J. The relation between body iron stores and adipose tissue function in patients with manifest vascular disease. *Eur J Clin Invest.* 2013,43(12),240-249. PMID:24245570 doi: 10.1111/eci.12165
34. Britton, L.; Jaskowski, L.A.; Bridle, K.; Secondes, E.; Wallace, D.; Santrampurwala, N.; Reiling, J.; Miller, G.; Mangiafico, S.; Andrikopoulos, S.; Subramaniam, V.N.; Crawford, D. Ferroporin expression in adipocytes

- does not contribute to iron homeostasis or metabolic responses to a high calorie diet. *Cell Mol Gastroenterol Hepatol.* 2018,5(3),319-331. PMID: 29552621 doi:10.1016/j.jcmgh.2018.01.005
35. Zhang, Z.; Funcke, J.B.; Zi, Z.; Zhao, S.; Straub, L.G.; Zhu, Y.; Zhu, Q.; Crewe, C.; An, Y.; Chen, S.; Li, N.; Wang, M.; Ghaben, A.L.; Lee, C.; Gautron, L.; Engelking, L.; Raj, P.; Deng, Y....Scherer, P.E. Adipocyte iron levels impinge on a fat-gut crosstalk to regulate intestinal lipid absorption and mediate protection from obesity. *Cell Metabol.* 2021,33(8),1624-1639.e9 PMID:34174197 PMID:PMC8338877 doi: 10.1016/j.cmet.2.06.001
  36. Ma, X.; Pham, V.T.; Mori, H.; MacDougald, O.A.; Shah, Y.M.; Bodary, P.F. Iron elevation and adipose tissue remodelling in the epididymal depot of a mouse model of polygenic obesity. *PLOS One.* 2017,12(6),e179889. PMID:28651003 PMID:PMC5484604 doi: 10.1371/journal.pone.0179889
  37. Lu, M.L.; Wang, T.N.; Lin, T.Y.; Shao, W.C.; Chang, S.H.; Chou, J.Y.; Ho, Y.F.; Liao, Y.T.; Chen, V.C. Differential effects of olanzapine and clozapine on plasma levels of adipocytokines and total ghrelin. *Prog Neuro-Psychopharmacol Biol Psychiatry.* 2015,58,47-50. PMID:25496829 doi: 10.1016/j.pnpbp.2014.12.001
  38. Shimizu, H., Shimomura, Y., Hayashi, R. et al. Serum leptin concentration is associated with total body fat mass, but not abdominal fat distribution. *Int J Obes* 1997; 21: 536–541. PMID:22128212 PMID:PMC3208009 doi:10.1038/sj.ijo.0800437
  39. Meier, U.; Gressner, A.M. Endocrine regulation of energy metabolism: review of pathobiochemical and clinical chemical aspects of leptin, ghrelin, adiponectin, and resistin. *Clin Chem.* 2004,50(9),1511-1525. PMID:15265818 doi:10.1373/clinchem.2004.032482
  40. Fang, H.; Judd, R.L. Adiponectin regulation and function. *Compr Physiol.* 2018,8(3),1031-1063. PMID: 29978896 doi:10.1002/cphy.c170046
  41. Liu, Z.; Cui, C.; Xu, P.; Dang, R.; Cai, H.; Liao, D.; Yang, M.; Feng, Q.; Yan, X.; Jiang, P. Curcumin activates AMPK pathway and regulates lipid metabolism in rats following prolonged clozapine exposure. *Front Neurosci.* 2017,11,article 558. PMID:29046626 PMID:PMC5632657 doi:10.3389/fnins.2017.00558
  42. Canfran-Duque, A.; Casado, M.E.; Pastor, O.; Sanchez-Wandelmer, J.; de la Pena, G.; Lerma, M.; Mariscal, P.; Bracher, F.; Lasuncion, M.A.; Busto, R. Atypical antipsychotics alter cholesterol and fatty acid metabolism in vitro. *J Lipid Res.* 2013,52,310-324. PMID:23175778 PMID:PMC3588861 doi:10.1194/jlr.M026948
  43. McBride, A.; Ghilagaber, S.; Nikolaev, A.; Hardie, D.G. The glycogen-binding domain on the AMPK beta subunit allows the kinase to act as a glycogen sensor. *Cell Metab.* 2009,9(1),23-34. PMID:19117544 doi:10.1016/j.cmet.2008.11.008
  44. Izumida, Y.; Yahagi, N.; Takeuchi, Y.; Nishi, M.; Shikama, A.; Takarada, A.; Masuda, Y.; Kubota, M.; Matsuzaka, T.; Nakagawa, Y.; Iizuka, Y.; Itaka, K.; Kataoka, K.; Shioda, S.; Nijima, A.; Yamada, T.; Katagiri, H.; Nagai, R.; Yamada, N.; Kadowaki, T.; Shimano, H. Glycogen shortage during fasting triggers liver-brain-adipose neurocircuitry to facilitate fat utilization. *Nature Commun.* 2013,4,2316. PMID:23939267 PMID:PMC3753545\_ doi:10.1038/ncomms3316
  45. Idonije, O.B.; Festus, O.O.; Akpamu, U.; Ohiaia, O.; Iribhogbe, O.I.; Iyalomhe, G.B.S. A comparative study of the effects of clozapine and risperidone monotherapy on lipid profile in Nigerian patients with schizophrenia. *Int J Pharmacol.* 2012,8(3),169-176. doi:10.3923/ijp.2012.169.176
  46. Kim, J.H.; Lee, J.O.; Lee, S.K.; Jung, J.; You, G.Y.; Park, S.H.; Park, M.; Kim, S.D.; Kim, H.S. Clozapine activates AMP-activated protein kinase (AMPK) in C2C12 myotube cells and stimulates glucose uptake. *Life Sci.* 2010,87(1-2),42-48. PMID:20515698 doi: 10.1016/j.lfs.2010.05.017
  47. Li, Y.; Xu, S.; Mihaylova, M.; Zheng, B.; Hou, X.; Jiang, B.; Park, O.; Luo, Z.; Lefai, E.; Shyy, J.; Gao, B.; Wierzbicki, M.; Verbeuren, T.; Shaw, R.; Cohen, R.; Zang, M. AMPK phosphorylates and inhibits SREBP activity to attenuate hepatic steatosis and atherosclerosis in diet-induced insulin resistant mice. *Cell Metab* 2011; 13(4): 376-388. PMID:21459323 PMID:PMC3086578 doi: 10.1016/j.cmet.2011.03.009
  48. Albaugh, V.L.; Vary, T.C.; Ilkayeva, O.; Wenner, B.R.; Maresca, K.P.; Joyal, J.L.; Breazeale, S.; Elich, T.D.; Lang, C.H.; Lynch, C.J. Atypical antipsychotics rapidly and inappropriately switch peripheral utilization to lipids, impairing metabolic flexibility in rodents. *Schizophr Bull.* 2012,38(1),153-166. PMID:0494946 PMID:PMC3245588\_ doi: 10.1093/schbul/sbq053
  49. Rodriguez, A. Novel molecular aspects of ghrelin and leptin in the control of adipobiology and the cardiovascular system. *Obes Facts.* 2014,7,82-95. PMID:24685565 PMID:PMC5644879 doi:10.1159/000360837
  50. Straub, B.K.; Stoeffel, P.; Heid, H.; Zimbelmann, R.; Schirmacher, P. Differential pattern of lipid droplet-associated proteins and de novo perilipin expression in hepatocyte steatogenesis. *Hepatology.* 2008,47,1936-1946. PMID:18393390 doi: 10.1002/hep.22268
  51. Krycer, J.R.; Elkington, S.D.; Diaz-Vegas, A.; Cooke, K.C.; Burchfield, J.G.; Fisher-Wellman, K.H.; Cooney, G.J.; Fazakerley, D.J.; James, D.E. Mitochondrial oxidants, but not respiration, are sensitive to glucose in adipocytes. *J Biol Chem.* 2020,295(1),99-110. PMID: 31744882 doi:10.1074/jbc.RA119.011695

52. Murashita, M.; Kusumi, I.; Hosoda, H.; Kangawa, K.; Koyama, T. Acute administration of clozapine concurrently increases blood glucose and circulating plasma ghrelin levels in rats. *Psychoneuroendocrinology*. 2007,32,777-784. PMID:17629416 doi:10.1016/j.psyneuen.2007.05.007
53. Sasaki, N.; Iwase, M.; Uchizono, Y.; Nakamura, U.; Imoto, H.; Abe, S.; Iida, M. The atypical antipsychotic clozapine impairs insulin secretion by inhibiting glucose metabolism and distal steps in rat pancreatic islets. *Diabetologia*. 2006,49(12),2930-2938. PMID:17072584 doi:10.1007/s00125-006-0446-6
54. Smith, G.C.; Chaussade, C.; Vickers, M.; Jensen, J.; Shepherd, P.R. Atypical antipsychotic drugs induce derangements in glucose homeostasis by acutely increasing glucagon secretion and hepatic glucose output in the rat. *Diabetologia*. 2008,51(12),2309-2317. PMID:18843478 doi:10.1007/s00125-1152-3
55. Leguisamo, N.M.; Lehnen, A.M.; Machado, U.F.; Okamoto, M.M.; Markoski, M.M.; Pinto, G.H.; Schaan, B.D. Glut4 content decreases along with insulin resistance and high levels of inflammatory markers in rats with metabolic syndrome. *Cardiovascular Diabetology*. 2012,11,100-110. PMID:22897936 PMID:PMC3439702 doi:10.1186/1475-2840-11-100
56. Sarvari, A.K.; Vereb, Z.; Uray, I.P.; Fesüs, L.; Balajthy, Z. Atypical antipsychotics induce both proinflammatory and adipogenic gene expression in human adipocytes in vitro. *Biochem Biophys Res Commun*. 2014,450,1383-1389. PMID:25019983 doi:10.1016/j.bbrc.2014.07.005
57. Chadt, A.; Al-Hasani, H. Glucose transporters in adipose tissue, liver, and skeletal muscle in metabolic health and disease. *Eur J Physiol*. 2020,472,1273-1298. PMID:32591906 PMID:PMC7462924 doi:10.1007/s00424-020-02417-x
58. Sorge S. Der Einfluss des atypischen Neuroleptikums Clozapin auf das Verhalten und das Immunsystem der Ratte. Thesis, München 2003.
59. Cooper, G.D.; Harrold, J.A.; Halford, J.C.G.; Goudie, A.J. Chronic clozapine treatment in female rats does not induce weight gain or metabolic abnormalities but enhances adiposity: implications for animal models of antipsychotic-induced weight gain. *Prog Neuropsychopharmacol Biol Psychiatry*. 2008,32(2),428-436. PMID:17933447 doi:10.1016/j.npbp.2007.09.012
60. Chusyd, D.E.; Wang, D.; Huffman, D.M.; Nagy, T.R. Relationships between rodent white adipose fat pads and human white adipose fat depots. *Front Nutr*. 2016,3,10 PMID:27148535 PMID:PMC4835715 doi:10.3389/fnut.2016.00010
61. Palmer, B.F.; Clegg, D.J. The sexual dimorphism of obesity. *Mol Cell Endocrinol*. 2015,402,113-119. PMID:25578600 doi:10.1016/j.mce.2014.11.029
62. Romeis, B. *Mikroskopische Technik* (17. neubearbeitete und erweiterte Auflage, herausgegeben von P. Böck). München – Wien – Baltimore 1989. Urban und Schwarzenberg. ISBN: 3-541-11227-1.
63. Ohkawa, H.; Ohishi, N.; Yagi, K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analyt Biochem*. 1979,95,351-358. PMID:36810 doi:10.1016/0003-2697(79)90738-3
64. Romero-Calvo, I.; Ocon, B.; Martinez-Moya, P.; Suarez, M.D.; Zarzuelo, A.; Martinet-Augustin, O.; de Medina, F.S. Reversible Ponceau staining as a loading control alternative to actin in Western blots. *Anal Biochem*. 2010, 401(2),318-320. PMID:20206115 doi: 10.1016/j.ab.2010.02.036.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.