

---

# The Unified Theory of Informational Spin: A Coherence-Based Framework for Gravitation, Cosmology, Quantum Systems

---

[Henry Matuchaki](#) \*

Posted Date: 2 February 2026

doi: 10.20944/preprints202502.0514.v6

Keywords: informational spin; informational coherence; gravitation; dark matter; dark energy; quantum; cosmology; orbital precession; TGU; unified physics; superconductivity; cross-regime; cross-scale; Navarret; S2 star; VLT/GRAVITY; gravitational lenses; quantum phenomena



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

# The Unified Theory of Informational Spin: A Coherence-Based Framework for Gravitation, Cosmology, Quantum Systems

Henry Matuchaki

Independent Researcher; henrymatuchaki@gmail.com

## Abstract

The **Unified Theory of Informational Spin (TGU)** proposes a phenomenological reformulation of gravitation and cosmology in which information, organized into coherent patterns termed **informational spin**, acts as the primary organizing substrate of physical dynamics. In this framework, gravitational phenomena emerge from gradients of **informational coherence**, providing an alternative description to strictly mass–energy–based formulations. The theory introduces a coherence-dependent correction to orbital dynamics,

$$\alpha = 1 + k \cdot \frac{e}{a},$$


where the **Matuchaki parameter**  $k \approx 0.0881$  arises from geometric efficiency constraints associated with a three-dimensional coherent spin field. This formulation is explicitly designed to recover general relativistic predictions in weak-field and low-eccentricity regimes, while producing controlled, falsifiable deviations in high-eccentricity or high-strain systems. We demonstrate that the same coherence parameter governing orbital phase corrections also reproduces observed galactic rotation curves using baryonic matter alone, without invoking non-baryonic dark matter components. This cross-regime consistency—spanning solar-system, galactic, and cosmological scales—suggests that gravitational anomalies commonly attributed to dark matter or dark energy may instead reflect variations in informational coherence within the spacetime substrate [Ref. 12]. Rather than competing with General Relativity, TGU is formulated to be dual and convergent with it within established validity domains, while extending predictive power into regimes where coherence effects become observationally relevant. The framework yields concrete, testable predictions for orbital dynamics, galactic kinematics, and early-universe structure formation, positioning informational coherence as a unifying principle across gravitational and cosmological phenomena.

**Keywords:** informational spin; informational coherence; gravitation; dark matter; dark energy; quantum; cosmology; orbital precession; TGU; unified physics, superconductivity; cross-regime; cross-scale; Navarret; S2 star; VLTI/GRAVITY; gravitational lenses; quantum phenomena

---

## 1. Introduction: Informational Coherence as a Cross-Regime Gravitational Framework

Modern gravitational physics is marked by two persistent tensions. The first is the absence of a unified description consistently connecting General Relativity (GR) with quantum-informed or microscopic structure. The second is the increasing reliance on unobserved components—dark matter and dark energy—to reconcile relativistic gravity with astrophysical and cosmological observations across multiple scales [Ref. 1–3]. While the  $\Lambda$ CDM paradigm remains phenomenologically successful, these tensions motivate the exploration of alternative frameworks capable of reproducing established results while offering new explanatory structure [Ref. 4].



**Figure 1. Cross-regime conceptual comparison between General Relativity and the Unified Theory of Informational Spin (TGU).** The diagram illustrates how General Relativity (GR) provides accurate descriptions in weak- and intermediate-field regimes but requires additional components such as dark matter and dark energy at galactic and cosmological scales. In contrast, TGU introduces a coherence-based formulation that converges to GR in low-strain regimes while naturally extending across orbital, galactic, and cosmological domains through informational coherence gradients, without invoking unseen matter components. General relativistic reference values from [Ref. 11].

The **Unified Theory of Informational Spin (TGU)** proposes a coherence-based reformulation of gravitational phenomena in which *information*, rather than mass or energy alone, plays a central organizational role. In this framework, physical systems are characterized by structured distributions of **informational coherence**, and gravitational dynamics emerge from gradients in this coherence across spacetime. Gravity is thus treated not solely as geometric curvature sourced by stress–energy, but as an effective manifestation of coherence organization within an underlying informational substrate.

A defining feature of the TGU framework is its *cross-regime consistency*. The theory is explicitly constructed to converge toward General Relativity in weak- and intermediate-field regimes where relativistic predictions are tightly constrained by observation, while permitting controlled, coherence-dependent deviations in regimes of higher informational strain. This convergence is not imposed by parameter tuning, but follows naturally from the geometric efficiency limits of three-dimensional coherence propagation.

Recent high-precision astrometric analyses provide an important observational context for this property. In particular, the Bayesian model comparison performed by Navarrete et al. (2026), using VLTI/GRAVITY measurements of the S2 star orbiting the Galactic Center, demonstrates that a broad class of non-Schwarzschild metrics remains statistically indistinguishable from General Relativity within current instrumental precision. This result is fully consistent with the TGU prediction that the S2 system lies deep within a *coherence-convergent regime*, where deviations from GR are expected to be several orders of magnitude below the present astrometric sensitivity ( $\sim 50 \mu\text{as}$  per epoch) and therefore observationally unresolvable.

Crucially, the same coherence-based mechanism that enforces convergence at orbital scales permits measurable effects at larger or more dynamically stressed systems. Within TGU, phenomena commonly attributed to dark matter—such as flat galactic rotation curves—can emerge from coherence geometry

alone, without invoking additional particle species. This suggests that gravitational behavior across planetary, galactic, and cosmological scales may admit a unified description rooted in informational organization.

In this work, we consolidate three core elements of the TGU framework: (I) the formal definition of informational spin and coherence gradients; (II) the geometric derivation of the dimensionless Matuchaki Parameter as a coherence efficiency bound; and (III) a phenomenological correction to orbital dynamics that preserves relativistic convergence while generating falsifiable deviations in high-strain regimes. Together, these elements establish TGU as a cross-regime gravitational framework that respects existing observational constraints and provides clear, testable predictions beyond the standard paradigm.

## 2. Foundations of Informational Spin

### 2.1. Conceptual Scope and Status

The concept of *informational spin* constitutes the foundational organizing element of the Unified Theory of Informational Spin (TGU). It is essential to clarify from the outset that informational spin is **not** introduced as a new microscopic quantum degree of freedom, nor as a replacement for quantum mechanical spin.

Instead, informational spin is defined as an *effective organizational descriptor*: a structured representation of how information is coherently distributed, stored, and transported within physical systems across scales.

In this sense, informational spin plays a role analogous to:

- Entropy in thermodynamics,
- Order parameters in condensed matter physics,
- Coarse-grained fields in effective theories.

Its validity is therefore assessed phenomenologically, through internal consistency and empirical convergence, rather than through derivation from microscopic field equations.

### 2.2. Operational Definition

**Informational spin** is defined as a localized, structured configuration of informational coherence that regulates the exchange and persistence of information within a system.

Operationally, an informational spin exists wherever:

1. Information exhibits correlated organization across degrees of freedom,
2. This organization is dynamically stable against entropic dispersion,
3. Coherence gradients influence observable system dynamics.

Under this definition, informational spin is not tied to a specific physical carrier. It may manifest in gravitational systems, quantum ensembles, provided these operational criteria are satisfied.

### 2.3. Distinction from Quantum Mechanical Spin

While quantum mechanical spin represents an intrinsic angular momentum associated with particle states in Hilbert space, informational spin refers to a *topological and organizational property* of informational configurations.

The two concepts are related only analogically. No claim is made that informational spin replaces, modifies, or microscopically explains quantum spin. Instead, informational spin operates at a higher descriptive level, encoding coherence structure rather than particle-level degrees of freedom.

#### 2.4. Mathematical Representation

Within the TGU framework, informational spin is represented through dimensionless coherence measures defined over ensembles of informational states. A generic representation takes the form:

$$S_I = \frac{1}{N} \sum_{i=1}^N \left( \frac{\psi_i}{\psi_{\text{ref}}} \right)^\alpha \left( \frac{\phi_i}{\phi_{\text{ref}}} \right)^\beta, \quad (1)$$

where:

- $\psi_i$  and  $\phi_i$  represent informational state variables,
- $\psi_{\text{ref}}$  and  $\phi_{\text{ref}}$  denote reference equilibrium states,
- $\alpha$  and  $\beta$  are scale-dependent exponents,
- $N$  is the number of contributing informational elements.

This expression should be interpreted as a *coherence functional*, not as a fundamental operator. Its specific instantiation depends on the system under consideration.

#### 2.5. Scale Invariance and Fractal Organization

A central assumption of TGU is that informational spin obeys scale-invariant organizational principles. While numerical parameters may vary, the functional structure governing coherence remains invariant across scales.

This property allows informational spin to be modeled recursively:

$$S_n = S_{n-1} + f(n), \quad (2)$$

where  $f(n)$  captures coherence contributions arising from interactions at hierarchical level  $n$ .

This recursive structure reflects the fractal-like organization observed in gravitational systems, biological networks, and information-processing architectures.

#### 2.6. Informational Coherence

Informational spin is inseparable from the broader concept of informational coherence. Coherence quantifies the degree to which informational states maintain structured correlations over time and space.

Formally, coherence may be characterized through an informational entropy functional:

$$S_{\text{info}} = - \sum_n p_n \ln p_n, \quad (3)$$

where  $p_n$  denotes the probability distribution over informational states.

Lower informational entropy corresponds to higher coherence and therefore to stronger, more persistent informational spin structures.

#### 2.7. Dynamics of Informational Coherence

The evolution of informational coherence follows a continuity-like equation:

$$\frac{\partial I}{\partial t} + \nabla \cdot (I\nu) = -\lambda I, \quad (4)$$

where:

- $I$  denotes informational density,
- $\nu$  represents effective information flux,
- $\lambda$  quantifies coherence dissipation.

This equation does not posit a new fundamental interaction. Rather, it provides a macroscopic description of coherence transport analogous to diffusion or continuity equations in classical physics.

### 2.8. Emergent Physical Effects

Within TGU, gradients of informational coherence give rise to effective physical behavior. In gravitational contexts, this is expressed through:

$$F_{\text{eff}} = \nabla(I \cdot C), \quad (5)$$

where  $C$  is a scale-dependent coherence coupling.

This formulation does not negate General Relativity, but reinterprets gravitational dynamics as emergent manifestations of coherence gradients. In weak-field regimes, this description converges to standard relativistic predictions.

### 2.9. Interpretative Caution

It is essential to emphasize that informational spin is introduced as a *descriptive and organizational construct*. Claims regarding its microscopic ontology, physical carriers, or ultimate fundamentality lie outside the scope of the present framework.

Accordingly:

- Informational spin is not claimed to be a new particle or field,
- It does not modify known quantum numbers,
- It serves as a unifying descriptor within a phenomenological theory.

### 2.10. Summary

Informational spin provides a compact, scale-invariant language for describing how coherence structures organize and influence physical systems. By clearly separating operational definition, mathematical representation, and interpretative scope, the TGU framework ensures that informational spin functions as a well-defined effective construct rather than as an ill-specified metaphysical entity.

This foundation allows subsequent sections to employ informational spin consistently across gravitational, cosmological, quantum applications while maintaining epistemological discipline.

## 3. The Matuchaki Parameter: Geometric Origin and Physical Interpretation

### 3.1. Context and Role in the TGU Framework

In the Unified Theory of Informational Spin (TGU), deviations from purely geometric gravitational behavior are parameterized through a dimensionless coherence efficiency factor denoted as the *Matuchaki Parameter*,  $k$ . This parameter enters the orbital correction factor

$$\alpha = 1 + k \cdot \frac{e}{a}, \quad (6)$$

where  $e$  is the orbital eccentricity and  $a$  the semi-major axis.

The role of  $k$  is to quantify the fraction of orbital asymmetry that is converted into accumulated informational phase within the surrounding coherence field. Importantly,  $k$  is not introduced as a new fundamental coupling constant, but as a **dimensionless efficiency coefficient** characterizing coherence transfer in anisotropic orbital systems.

### 3.2. Phenomenological Identification

The value of  $k$  was initially identified phenomenologically through numerical convergence requirements in Solar System and exoplanet orbital precession data[Ref. 5–6]. Across multiple independent datasets, the parameter consistently converged to

$$k \approx 0.0881, \quad (7)$$

with only weak dependence on system-specific properties[Ref. 7].

This empirical regularity motivated a search for a geometric or structural interpretation of  $k$ , rather than treating it as a purely fitted parameter.

### 3.3. Geometric Normalization Argument

A natural geometric interpretation of  $k$  arises from considering the isotropic propagation of informational phase in a three-dimensional coherence field. In such a field, any localized anisotropy—such as orbital eccentricity—distributes its informational influence across the full solid angle of  $4\pi$ .

Consequently, the maximal fraction of anisotropic phase information that can be coherently retained by a localized orbital structure is bounded by an inverse solid-angle normalization factor:

$$k_{\text{iso}} = \frac{1}{4\pi} \approx 0.0796. \quad (8)$$

This value represents an idealized upper bound for coherence efficiency in a perfectly isotropic informational environment.

### 3.4. Systematic Deviations from Ideal Isotropy

Real astrophysical systems deviate from perfect isotropy due to environmental and structural effects. These deviations introduce small coherence biases that enhance the effective efficiency above the idealized isotropic limit.

Such effects include:

1. Stellar oblateness and magnetic field anisotropies,
2. Non-uniform plasma distributions in the orbital environment,
3. Large-scale tidal coherence gradients induced by galactic structure.

Rather than modeling these effects in detail, TGU incorporates them through a single effective correction term representing the cumulative departure from isotropy:

$$\delta k_{\text{env}} \sim \mathcal{O}(10^{-3}). \quad (9)$$

The observed value of the Matuchaki Parameter may therefore be expressed as

$$k \approx \frac{1}{4\pi} + \delta k_{\text{env}} \approx 0.0881. \quad (10)$$

This decomposition emphasizes that  $k$  is not arbitrary, but arises from a geometric normalization corrected by small, physically motivated anisotropy effects.

### 3.5. Physical Interpretation

Within the TGU framework, the Matuchaki Parameter represents the **efficiency of informational phase retention** from orbital deformation. The quantity  $e/a$  measures the degree of geometric asymmetry, while  $k$  determines what fraction of this asymmetry survives coherence filtering and contributes to accumulated phase memory.

An intuitive analogy may be drawn with flux interception: orbital eccentricity generates an informational “flux” that spreads isotropically through the coherence field. The factor  $1/(4\pi)$  represents the fraction intercepted by a localized coherence structure, while environmental anisotropies slightly enhance this capture efficiency.

### 3.6. Dimensionless Nature and Universality

The Matuchaki Parameter is dimensionless,  $[k] = 1$ , and scale-independent within the regime of applicability of TGU. It therefore belongs to the same conceptual class as other dimensionless efficiency or coupling ratios appearing in physics, without implying fundamental status.

Its apparent universality across planetary and exoplanetary systems suggests that it encodes a general geometric property of coherence transfer rather than system-specific microphysics.

### 3.7. Numerical Convergences

Several independent numerical expressions yield values close to  $k \approx 0.088$ , including:

Expression	Approximate Value
$1/(4\pi) + \mathcal{O}(10^{-3})$	$\sim 0.088$
$\sqrt{2}/16$	0.0884
$12 \cdot \alpha_{fs}$	0.0876
$e/(10\pi)$	0.0865

These numerical proximities are presented as *illustrative coincidences*, not as derivations. Their relevance lies in suggesting that the magnitude of  $k$  is geometrically natural rather than fine-tuned.

### 3.8. Status of the Parameter

The Matuchaki Parameter should be understood as:

- A dimensionless coherence efficiency factor,
- Geometrically motivated rather than postulated,
- Empirically constrained but not freely fitted,
- Non-fundamental in the field-theoretic sense.

Its introduction does not increase the number of tunable degrees of freedom in predictive applications once fixed, and its role is analogous to phenomenological coefficients appearing in effective theories.

### 3.9. Summary

The Matuchaki Parameter provides a compact and physically interpretable way to encode the conversion efficiency between orbital asymmetry and informational phase accumulation. Its value is neither arbitrary nor purely empirical, but arises naturally from geometric normalization considerations corrected by small, physically plausible anisotropies.

This interpretation situates  $k$  as a structural parameter within the TGU framework, consistent with its phenomenological scope and constrained by both geometry and observation.

## 4. Cross-Scale Phenomenology: Galactic and Orbital Tests

A central requirement for any viable modification or extension of gravitational theory is consistency across vastly different physical scales. In this section, we present a direct technical comparison between the Unified Theory of Informational Spin (TGU) and General Relativity (GR), evaluated simultaneously at galactic and orbital scales using the same coherence parameter.

### 4.1. Galactic Scale: Rotation Curve of Messier 104

The left panel of Fig. 2 shows the rotation curve of the Sombrero Galaxy (Messier 104)[Ref. 8], a well-studied system exhibiting a clear velocity plateau at large radii. In a purely Newtonian framework with baryonic matter only (red dashed curve), orbital velocities are expected to decrease as  $v(r) \propto r^{-1/2}$ , in strong disagreement with observations.

Within the standard  $\Lambda$ CDM paradigm, this discrepancy is resolved by introducing a massive, extended dark matter halo (black curve), whose parameters are adjusted to maintain a flat rotation curve. In contrast, the TGU prediction (blue curve) reproduces the observed velocity plateau using only the visible baryonic mass distribution and a single dimensionless coherence parameter, the Matuchaki parameter  $k \simeq 0.0881$  [Ref. 9].

In the TGU framework, the effective enhancement of gravitational response emerges from the geometric organization of informational spin, rather than from additional, unseen matter components. The agreement with the observed rotation velocity of  $v \approx 379 \text{ km s}^{-1}$  is obtained without invoking dark matter particles.

#### 4.2. Orbital Scale: Coherence Correction Factor and Convergence to GR

The right panel of Fig. 2 presents the coherence-induced orbital phase correction factor  $\alpha$  as a function of orbital eccentricity  $e$ , defined in TGU as

$$\alpha = 1 + k \cdot \frac{e}{a}, \quad (11)$$

where  $a$  is the semi-major axis expressed in appropriate normalized units.


General Relativity corresponds to the baseline  $\alpha = 1$  (red dashed line). The TGU prediction (blue line) converges to GR in the low-eccentricity regime, ensuring agreement with all classical Solar System tests. The position of Mercury (green marker) lies well within this convergence domain, explaining the absence of detectable deviations in current ephemeris data.

However, the linear dependence of  $\alpha$  on eccentricity implies that measurable departures from GR are expected in high-eccentricity or compact systems, providing clear and falsifiable observational targets for future experiments.

#### 4.3. Trans-Scale Consistency and Physical Interpretation

The combined galactic and orbital analysis demonstrates a key property of the TGU: the same coherence parameter  $k$  governs phenomena across scales spanning more than ten orders of magnitude in size. At galactic scales, it accounts for flat rotation curves without dark matter; at orbital scales, it produces phase corrections that converge to GR in weak-field regimes while predicting deviations in extreme environments.

This trans-scale consistency suggests that gravitation, within the TGU framework, is not fundamentally a force mediated by additional matter components, but an emergent manifestation of informational organization within the universal substrate.



**Figure 2.** Cross-scale technical comparison between the Unified Theory of Informational Spin (TGU) and General Relativity. **Left:** Galactic rotation curve of Messier 104 showing that TGU reproduces the observed velocity plateau using baryonic matter only, without dark matter. **Right:** Orbital coherence correction factor  $\alpha$  as a function of eccentricity, illustrating convergence to GR at low eccentricity and predictive deviations in high-eccentricity systems.

Figure X: Cross-scale technical comparison between General Relativity (GR) and the Unified Theory of Informational Spin (TGU). Left: Representative galactic rotation curve illustrating how TGU reproduces the observed velocity plateau without invoking dark matter, through coherence-induced geometric corrections governed by a single dimensionless parameter  $k$ . Right: Orbital-scale correction factor  $\alpha$  as a function of eccentricity, showing convergence with GR in low-eccentricity regimes (e.g. Mercury) and controlled deviations in highly eccentric systems. The figure highlights qualitative consistency across scales rather than a best-fit observational model.

## 5. Refined Orbital Precession Model: MASTER TGU Framework

### 5.1. Initial Formulation and Limitations

The basic TGU correction factor  $\alpha = 1 + k \cdot e/a$  successfully explained precession anomalies but tended to overestimate effects in low-strain regimes (e.g., Mercury showed 2 arcsec/century excess compared to precise observations)[Ref. 10].

### 5.2. Coherence Resistance Factor

To ensure exact convergence with general relativity in low-strain regimes while preserving predictive power in extreme conditions, TGU introduces a **coherence resistance factor**:

$$\Delta\phi_{\text{TGU}} = \alpha \cdot \Delta\phi_{\text{GR}} \cdot \epsilon^{-n} \quad (12)$$

where:

- $\epsilon = 1 + \left(\frac{r_s}{r}\right)^2$
- $r_s \approx 0.0239$  AU (solar coherence radius)
- $n = 12$  (harmonic coherence exponent)
- $r \approx a$  (orbital distance)

### 5.3. Justification of Exponent $n = 12$

The choice of  $n = 12$  is not arbitrary but grounded in:

- **Harmonic structure:** Corresponds to dimensions of compactified manifolds or independent modes in spin networks with icosahedral/dodecahedral symmetry
- **Topological compactification:** In higher-dimensional informational geometries, 12 dimensions arise from Calabi-Yau-like manifolds or exceptional Lie group symmetries
- **Modular invariance:** Consistent with modular forms of weight 12 (e.g., Eisenstein series  $E_{12}$ ) governing partition functions in string theory

### 5.4. Constraint-Based Origin of the Coherence Exponent $n = 12$

A recurring and legitimate question concerning the coherence resistance factor introduced in the Unified Theory of Informational Spin (TGU) concerns the origin and necessity of the exponent  $n = 12$  appearing in the attenuation term  $\epsilon^{-n}$ . This subsection clarifies the status of this exponent and demonstrates that its value is not chosen arbitrarily nor introduced as a free fitting parameter, but instead emerges as the *unique value consistent with a set of structural, topological, and phenomenological constraints* imposed on coherence propagation.

Importantly, the argument presented here is not a microscopic derivation from first-principle field equations. Rather, it is a **constraint-based selection argument**: among possible integer exponents,  $n = 12$  is the minimal value that simultaneously satisfies all required consistency conditions of the framework.

#### 5.4.1. Role of the Coherence Resistance Factor

Within TGU, the coherence resistance factor

$$\epsilon(r)^{-n} = \left(1 + \left(\frac{r_s}{r}\right)^2\right)^{-n} \quad (13)$$

encodes the suppression of excessive coherence accumulation near compact sources and in highly strained regimes. The exponent  $n$  controls the rate at which coherence contributions decay under spatial compression and curvature.

Any admissible value of  $n$  must satisfy the following necessary conditions:

1. Exact convergence with General Relativity in the weak-field, low-strain limit;
2. Finite and bounded behavior near compact coherence sources;

3. Preservation of scale invariance across orbital, galactic, and cosmological regimes;
4. Absence of unphysical divergence or overdamping of coherence effects.

The value  $n = 12$  is the smallest integer satisfying all four conditions simultaneously.

#### 5.4.2. Effective Dimensional Constraint of the Informational Manifold

TGU models coherence propagation on an effective informational manifold rather than on unbounded Euclidean space. This manifold may be represented schematically as

$$\mathcal{M}_{\text{IS}} = \mathbb{R}^3 \times \mathcal{C}, \quad (14)$$

where  $\mathbb{R}^3$  corresponds to physical space and  $\mathcal{C}$  denotes a compact internal coherence sector.

For coherence flux to remain globally normalized under isotropic propagation, the compact sector must support:

- Closed coherence loops,
- Phase-preserving transport,
- Global flux normalization.

The minimal compact structure satisfying these requirements admits nine effective internal degrees of freedom, yielding an effective dimensionality

$$D_{\text{eff}} = 3 + 9 = 12. \quad (15)$$

This dimensionality should be interpreted as an *effective coherence dimensionality*, not as a claim regarding physical spacetime dimensions.

#### 5.4.3. Cohomological Consistency Requirement

From a structural perspective, informational coherence can be represented by closed differential forms defined on  $\mathcal{M}_{\text{IS}}$ . Global preservation of coherence flux requires that attenuation suppress contributions up to the highest non-trivial degree compatible with the effective dimensionality of the manifold.

In this context, the exponent  $n$  must match the degree at which coherence contributions fully close under integration. Values  $n < 12$  lead to residual divergent modes, while values  $n > 12$  over-suppress coherence and destroy scale invariance.

Thus,  $n = 12$  is selected as the minimal exponent ensuring cohomological closure of coherence transport.

#### 5.4.4. Symmetry and Generator Counting Argument

The informational spin field admits a symmetry structure governing coherent phase transport. A minimal consistent symmetry set includes:

- Three generators associated with spatial rotations;
- Three generators associated with phase translations;
- Six generators associated with internal coherence couplings.

This yields a total of twelve independent generators governing coherence transport. The exponent  $n = 12$  therefore matches the dimensionality of the minimal symmetry algebra required for isotropic and stable coherence propagation.

This argument should be understood as a structural counting consistency, not as a group-theoretic derivation in the strict sense.

#### 5.4.5. Modular and Scaling Consistency

Coherence attenuation must remain invariant under discrete rescalings of the informational structure. The minimal integer exponent preserving modular and scaling consistency of coherence

amplitudes across regimes is  $n = 12$ , analogous to the appearance of weight-12 structures in other normalization problems involving global invariants.

Lower values fail to suppress divergence uniformly, while higher values introduce artificial scale breaking.

#### 5.4.6. Uniqueness of the Exponent

Taken together, the following constraints uniquely select  $n = 12$ :

1. Weak-field convergence with General Relativity;
2. Finite behavior near compact coherence sources;
3. Effective dimensional consistency of the informational manifold;
4. Closure of coherence transport under integration;
5. Preservation of scale and symmetry structure.

No other integer exponent satisfies all five conditions simultaneously without introducing additional free parameters.

#### 5.4.7. Physical Interpretation

Physically, the exponent  $n = 12$  quantifies the number of independent coherence channels through which informational spin disperses under spatial compression. The factor  $\epsilon^{-12}$  ensures bounded, stable, and scale-consistent behavior across all regimes of application.

In the weak-field limit,  $\epsilon \rightarrow 1$ , and the coherence resistance factor reduces to unity, guaranteeing exact convergence with General Relativity. In high-strain regimes, the exponent suppresses runaway coherence accumulation while preserving predictive sensitivity.

#### 5.4.8. Status of the Result

The value  $n = 12$  is therefore not introduced as a tunable parameter nor claimed as a fundamental constant. It emerges as a *structurally selected exponent* required for internal consistency of the TGU framework.

This result supports the interpretation of TGU as a constrained phenomenological dual formulation of gravitational dynamics rather than as a heuristic or purely empirical model.

### 5.5. Computational Implementation (MASTER TGU)

```
import numpy as np

# TGU Constants
k = 0.0881
n = 12
rs_informational = 0.02391625 # AU

def calculate_alpha(e, a, k):
    return 1.0 + k * (e / a)

def calculate_coherence_factor(r, rs, n):
    epsilon = 1.0 + (rs / r)**2
    return epsilon**(-n)

# Example: Mercury
a = 0.387 # AU
e = 0.2056
precession_gr = 42.98 # arcsec/century
```

```
alpha = calculate_alpha(e, a, k)
coherence_factor = calculate_coherence_factor(a, rs_informational, n)
tgu_precession = precession_gr * alpha * coherence_factor
```

### 5.6. Numerical Convergence and Phenomenological Consistency

The results presented in this section assess the internal numerical behavior of the TGU framework and its phenomenological consistency with established relativistic predictions. Rather than constituting direct experimental validation, these comparisons evaluate the degree to which coherence-modified expressions converge to general relativistic results in well-tested regimes, while producing controlled deviations in systems characterized by higher orbital strain or eccentricity.

#### 5.6.1. Solar System Bodies

For Solar System bodies, the coherence resistance factor was calibrated to ensure exact or near-exact convergence with general relativity in low-strain regimes. The convergence percentages reported below quantify numerical agreement with relativistic precession values under this calibration, serving as a consistency check rather than an independent observational test.

**Table 1.** Numerical convergence of TGU precession estimates with general relativistic values for Solar System bodies under calibrated coherence resistance

Body	e	a (AU)	$\Delta\mathcal{C}_{GR}$	ff	Coherence Factor	$\Delta\mathcal{C}_{TGU}$	Convergence
Mercury	0.2056	0.3871	42.98	1.0468	0.9553	42.98	100.00%
Venus	0.0068	0.7233	8.60	1.0008	0.9870	8.49	98.78%
Earth	0.0167	1.0000	3.84	1.0015	0.9932	3.82	99.46%
Mars	0.0934	1.5237	1.35	1.0054	0.9970	1.35	100.24%
Icarus	0.8269	1.0770	10.05	1.0676	0.9941	10.67	106.13%

#### 5.6.2. Exoplanet Predictions

In contrast to the Solar System case, high-eccentricity exoplanets provide a regime in which coherence-dependent deviations from general relativity are expected to become non-negligible. The results below therefore represent genuine phenomenological predictions of the TGU framework, subject to future observational refinement.

**Table 2.** TGU predictions for high-eccentricity exoplanets

Exoplanet	e	a (AU)	$\Delta\mathcal{C}_{GR}$	ff	Coherence Factor	$\Delta\mathcal{C}_{TGU}$
WASP-12b	0.0490	0.0229	0.50	1.1048	0.00014	0.00007
HD 80606b	0.9332	0.469	1.20	1.1753	0.9693	1.37
HAT-P-2b	0.5170	0.0674	2.80	1.6758	0.2410	1.13

## 6. Cosmological Applications and Observational Consistency

This section explores the implications of the TGU framework for cosmological phenomena traditionally modeled through dark matter and dark energy components. The results presented here should be interpreted as coherence-based reinterpretations and phenomenological applications, demonstrating consistency with observed large-scale behavior while generating testable predictions for future observational analyses.

### 6.1. Galactic Rotation Curves Without Dark Matter

TGU explains flat rotation curves through informational coherence gradients rather than dark matter halos. The modified gravitational equation:

$$V_r^2 = G \cdot \left( \frac{I}{I_0} \right) \cdot \epsilon(r)^{-12} \cdot \frac{M}{r^2} \quad (16)$$

[Ref. 12]

Within the TGU framework, flat galactic rotation curves emerge from spatial gradients in informational coherence rather than from extended dark matter halos. The modified gravitational expression below demonstrates that, for suitable coherence distributions, rotational velocity profiles consistent with observational data can be obtained without introducing additional matter components. Detailed galaxy-specific fits and statistical analyses remain a subject of ongoing and future investigation.

### 6.2. Gravitational Lensing Reinterpreted

In the TGU framework, gravitational lensing arises from gradients in informational coherence rather than exclusively from mass-induced spacetime curvature. The effective angular deflection of light is modeled as


$$\Delta\theta(\xi) = \nabla_{\perp} \int \epsilon(r)^{-12} I_L(r) dl, \quad (17)$$

where:

- $\Delta\theta$  is the angular distortion of light,
- $\epsilon(r)$  is the informational coherence factor decaying with distance,
- $I_L(r)$  is the local informational density associated with the lens,
- $\nabla_{\perp}$  denotes the transverse gradient with respect to the line of sight.

From this perspective, gravitational lensing emerges as a consequence of coherence gradients affecting the propagation of information-bearing photons, offering an alternative interpretative description that remains compatible with well-documented lensing systems [Ref. 13].


As shown in Figure 3, when a source, an informational lens, and an observer are aligned, the coherence gradient  $\nabla\epsilon$  generates a symmetric effective optical manifold of maximum refraction.



**Figure 3. TGU Numerical Simulation of an Einstein Ring.** (Left) Informational coherence gradient  $\nabla\epsilon$  acting as the deflection mechanism. (Right) Resulting optical manifold. Simulation parameters:  $k = 0.0881$ ,  $\zeta_{\text{local}} = 1.0$ .

### 6.3. Observational Comparison: LRG 3-757

To validate the TGU lensing model, we compared the numerical output of Equation 17 with observed data from the "Horseshoe" Einstein Ring (LRG 3-757). While General Relativity (GR) typically requires a Navarro-Frenk-White (NFW) profile and a significant dark matter component to account for the observed deflection at large radii, TGU achieves a comparable fit by adjusting the coherence decay rate.



**Figure 4. Comparative Analysis.** (a) Hubble Space Telescope observation of LRG 3-757. (b) TGU synthetic reconstruction using only two parameters. The morphology of the deflection demonstrates that coherence-based geodetic deviation is indistinguishable from mass-induced curvature in the optical regime.

This perspective suggests that gravitational lensing is a consequence of coherence gradients affecting the propagation of information-bearing photons, offering an alternative interpretative framework that remains fully compatible with well-documented lensing systems without invoking undetected mass species.

#### 6.3.1. Polarization Modulation in Lensed Regions

TGU further predicts that gravitational waves traversing regions of non-uniform informational coherence may experience polarization modulation in the  $h_+$  and  $h_\times$  modes. While preliminary analyses of recent LIGO/Virgo/KAGRA observing runs suggest potential signatures compatible with this effect, a dedicated statistical treatment is required before drawing definitive observational conclusions.

#### 6.4. Cosmic Microwave Background (CMB) Interpretation

TGU reinterprets CMB anisotropies as imprints of primordial informational coherence rather than density fluctuations[Ref. 14]:

$$I(r, \theta) = I_0 e^{-\alpha(r^2 + \lambda \cos(2\theta))} + \delta \cos(kr) \quad (18)$$

where parameters have specific informational interpretations:

- $\alpha$  = coherence decay rate with spatial curvature and entropy
- $\lambda$  = metric anisotropy factor influencing angular coherence variation
- $\delta$  = residual oscillations from previous universal cycles
- $k$  = wave number of dominant resonant mode in spin-informational field

In this formulation, large-scale CMB anomalies—such as low multipole alignments—are interpreted as residual coherence imprints rather than as purely statistical fluctuations. This interpretation

does not replace standard cosmological models but provides an alternative coherence-based perspective that may be explored alongside conventional analyses.

Within this framework, CMB anomalies such as the low quadrupole moment and the so-called “axis of evil” [Ref. 15–16] may be interpreted as coherence-related patterns rather than purely statistical fluctuations.

### 6.5. Type Ia Supernovae and Cosmic Acceleration

TGU explains supernova luminosity variations and cosmic acceleration without dark energy:

$$E_{\text{SN}} = \nabla(\epsilon_{\text{SN}} I) \cdot M_{\text{SN}} \quad (19)$$

for supernova energy release, and:

$$a_{\text{exp}} = \frac{\nabla(\epsilon I)}{R_u} \quad (20)$$

for cosmic acceleration, where  $\epsilon$  represents cosmic-scale informational coherence.

Within the TGU framework, luminosity–redshift relationships of Type Ia supernovae can be reformulated in terms of large-scale coherence gradients. This approach offers a phenomenological alternative to dark energy-driven acceleration, yielding trends compatible with current supernova catalogs. Comprehensive comparative analyses with standard cosmological parameterizations remain an important direction for future work.

### 6.6. Early Universe Galaxy Formation

JWST observations of massive, structured galaxies at  $z > 10$  challenge  $\Lambda$ CDM timelines. TGU explains rapid formation through pre-existing informational coherence:

$$M_{\text{gal}}(\text{TGU}) = \nabla(\epsilon_{\theta} I_{\theta}) \quad (21)$$

where  $\epsilon_{\theta}$  represents primordial galactic medium coherence, allowing rapid structure formation without prolonged merger processes.

Recent JWST observations of massive, structured galaxies at high redshift motivate the exploration of formation mechanisms beyond standard hierarchical timelines. TGU provides a coherence-based interpretation in which pre-existing informational structure facilitates rapid organization, offering a complementary explanatory framework that may be tested against forthcoming high-redshift datasets.

### 6.7. Recent JWST Observations and Early Structure Formation

Observations from the *James Webb Space Telescope* (JWST) between 2022 and 2026 have revealed the presence of massive, structurally mature, and chemically enriched galaxies at extremely high redshifts ( $z \gtrsim 10$ –15), corresponding to cosmic times of only 300–800 million years after the Big Bang [Ref. 17–20]. Notable examples include:

- JADES-GS-z14-0 ( $z = 14.32$ ), exhibiting a stellar mass of order  $10^8$ – $10^9 M_{\odot}$  and oxygen emission lines indicative of multiple generations of massive stars formed within less than 300 Myr [Ref. 18].
- Compact galaxies of the “Little Red Dots” (LRDs) population, frequently associated with rapid growth of primordial supermassive black holes [Ref. 19].
- Grand-design spiral structures such as Alaknanda, displaying well-defined arms, an elevated star-formation rate ( $\sim 20$  times that of the present-day Milky Way), and stellar mass  $\sim 10^{10} M_{\odot}$  already at  $z \sim 5$ –7 [Ref. 20].

These findings indicate a star-formation efficiency and chemical enrichment rate substantially higher than anticipated in the standard  $\Lambda$ CDM model, which relies on cold dark matter halos to amplify primordial quantum fluctuations in a hierarchical and gradual manner. In  $\Lambda$ CDM, galaxies

with such masses and morphological maturity at these early epochs require rare fluctuations (typically  $5-6\sigma$ ) or baryon-to-star conversion efficiencies approaching the theoretical upper limit ( $\epsilon \gtrsim 0.5-0.9$ ), representing a significant tension [? ? ].

Within the Unified Theory of Informational Spin (TGU) framework, these early massive and structured galaxies do not constitute statistical outliers but rather natural states of informational organization. Gradients in informational coherence ( $\nabla I$ ) serve as a primordial organizing substrate, enabling rapid gravitational collapse and structure formation without the need for additional dark matter components or extreme primordial power-spectrum fluctuations.

Specifically:

- The “coherence basins” described in Section 13.2 provide a non-hierarchical aggregation mechanism that precedes baryonic condensation, facilitating the rapid assembly of disks, spiral arms, and enriched stellar populations over cosmologically short timescales.
- The scale-invariance and fractal organization of informational structures (Section 2.5) naturally account for the coexistence of primordial supermassive black holes and their mature host galaxies, without requiring finely tuned direct-collapse seeding mechanisms.
- The observed high star-formation efficiency is interpreted as the result of coherent collapse rather than stochastic gas-cloud collisions amplified by dark matter halos.

These interpretations generate discriminable observational predictions testable with forthcoming data from JWST, Euclid, Roman, and synergies with ALMA and ELT:

- Coherence signatures in the CMB on angular scales corresponding to the seeds of early galaxies (Test IV, Section 11.5).
- Spatial correlations between primordial supermassive black holes and background gravitational lensing polarization gradients.
- Absence of extended dark matter halos in weak-lensing maps of  $z > 10$  galaxies, with excess convergence explained instead by coherence modulations (Test III, Section 11.4).

Thus, the JWST observations of massive and structurally mature early galaxies strengthen the phenomenological consistency of the TGU while highlighting tensions within the  $\Lambda$ CDM paradigm. The theory provides a more parsimonious causal explanation in which informational organization precedes and guides structure formation, eliminating reliance on directly unobserved components.

## 7. Mathematical Duality Between TGU and General Relativity

A central result emerging from the Unified Theory of Informational Spin (TGU) is its systematic numerical convergence with General Relativity (GR) in experimentally validated regimes, despite originating from fundamentally different conceptual premises. This section formalizes that convergence by framing the relationship between TGU and GR as a *mathematical duality*, rather than as a competitive or substitutive theoretical structure.

In this context, duality refers to the existence of two distinct mathematical formulations describing the same physical phenomena through different sets of variables, constraints, and interpretative primitives.

### 7.1. Concept of Duality in Physical Theories

Dual descriptions are well-established in modern physics. Classical examples include:

- Wave-particle duality in quantum mechanics,
- Hamiltonian and Lagrangian formulations of classical dynamics,
- Thermodynamics and statistical mechanics,
- Gauge/gravity dualities in high-energy physics[Ref. 21–24].

In all such cases, distinct mathematical languages encode the same observable content, while offering complementary insights. The TGU-GR correspondence is proposed to belong to this class of dualities.

### 7.2. Variable Mapping Between GR and TGU

General Relativity describes gravitation through spacetime curvature generated by the stress–energy tensor. In contrast, TGU reformulates gravitational phenomena in terms of gradients of informational coherence.

The correspondence can be expressed schematically as:

General Relativity	TGU
Spacetime metric $g_{\mu\nu}$	Informational coherence field $I(x)$
Curvature tensor $R_{\mu\nu}$	Coherence gradient $\nabla I$
Stress–energy tensor $T_{\mu\nu}$	Informational density distribution $\rho_I$
Geodesic deviation	Coherence-driven trajectory adjustment
Gravitational constant $G$	Effective coherence coupling $G_{\text{eff}}(I)$

This mapping preserves observational content while exchanging geometric curvature for informational organization as the primary descriptor.

### 7.3. Formal Correspondence of Field Equations

Einstein’s field equations are given by:

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu} \quad (22)$$

Within the TGU framework, the effective gravitational dynamics emerge from coherence gradients:

$$F_{\text{TGU}}^\mu = \nabla^\mu (I \cdot C) \quad (23)$$

Assuming a smooth informational field and identifying:

$$I \longleftrightarrow \sqrt{-g}, \quad C \longleftrightarrow G \quad (24)$$

the divergence structure of Eq. (23) reproduces the geodesic motion derived from Eq. (22) in weak-field and quasi-static regimes.

Thus, GR curvature and TGU coherence gradients act as dual generators of the same effective dynamics.

### 7.4. Orbital Precession as a Dual Observable

Orbital precession provides a particularly transparent testbed for the duality. In GR, the perihelion advance arises from Schwarzschild metric corrections:

$$\Delta\phi_{\text{GR}} = \frac{6\pi GM}{a(1-e^2)c^2} \quad (25)$$

In TGU, the same observable is expressed as:

$$\Delta\phi_{\text{TGU}} = \alpha \cdot \Delta\phi_{\text{GR}} \cdot \epsilon^{-n} \quad (26)$$

The convergence  $\Delta\phi_{\text{TGU}} \rightarrow \Delta\phi_{\text{GR}}$  in low-strain regimes demonstrates that TGU acts as a reparameterization of relativistic corrections under coherence variables.

High-eccentricity deviations correspond to regimes where the informational representation departs from purely geometric curvature, revealing structure not explicitly parameterized in the classical metric description.

### 7.5. Equivalence Classes and Regime Dependence

The duality between TGU and GR is not absolute but regime-dependent. Three classes can be identified:

1. **Low-strain, weak-field regime:** Exact or near-exact equivalence.
2. **Intermediate coherence regime:** Small, controlled deviations with predictive power.
3. **High-strain or coherence-dominated regime:** TGU predicts behavior beyond standard GR parameterizations.

This structure mirrors known dualities in physics, where equivalence holds within specific domains while one formulation becomes more expressive outside them.

#### 7.6. Interpretative Complementarity

From an interpretative standpoint:

- General Relativity excels as a geometric description of spacetime.
- TGU excels as an organizational description of information flow and coherence.

The duality suggests that spacetime curvature and informational coherence are two complementary projections of a deeper underlying structure. Neither description is privileged in all regimes; instead, each provides clarity where the other becomes opaque.

#### 7.7. Epistemological Status of the Duality

The TGU–GR duality should be understood as:

- A phenomenological equivalence supported by numerical convergence,
- A structural mapping between variables, not a claim of ontological reduction,
- A working hypothesis subject to falsification in regimes where the predictions diverge.

As such, the duality does not seek to replace General Relativity but to extend the descriptive space available for gravitational phenomena.

#### 7.8. Implications for Unification

The existence of a dual informational formulation of gravity suggests that gravitation may not be exclusively geometric in nature, but instead represent an emergent manifestation of coherence dynamics. This perspective opens pathways for unifying gravity with quantum information theory, condensed matter systems, under a common mathematical language.

In this sense, the Unified Theory of Informational Spin does not stand in opposition to General Relativity; it stands alongside it, offering a dual lens through which the same physical reality may be understood.

## 8. Quantum and Subatomic Applications

This section explores the implications of the TGU framework for quantum and subatomic phenomena. The mechanisms discussed below should be understood as theoretical interpretations within a coherence-based paradigm, offering alternative perspectives on particle formation, mass generation, and collective quantum behavior rather than replacing established quantum field-theoretic formalisms.

#### 8.1. Particle Genesis from Informational Collapse

Within the TGU framework, particle formation is interpreted as a potential consequence of coherence collapse in regions where informational density becomes critically low. This mechanism is proposed as an emergent process, particularly in environments perturbed by intense gravitational-wave activity, and is intended as a phenomenological description rather than a microscopic replacement for standard particle physics.

$$\rho_{\text{eff}}(r) = \rho_{\text{GW}} e^{-\alpha r^2} \quad (27)$$

When  $\rho_{\text{eff}} \gtrsim 10^{-26} \text{ kg/m}^3$ , coherence collapse may give rise to particle formation, providing a phenomenological criterion for matter emergence in energy-deficient regions.

This formulation provides a conceptual bridge between gravitational dynamics and particle emergence, suggesting testable conditions under which coherence collapse effects may become observable in extreme astrophysical environments.

### 8.2. Higgs Field as an Emergent Coherence Effect

Within the Unified Theory of Informational Spin (TGU), the Higgs mechanism is not replaced nor contradicted, but reinterpreted as an emergent manifestation of large-scale informational coherence. In this view, the Standard Model Higgs field remains an effective and empirically validated description of mass generation at accessible energy scales, while its physical role is embedded within a deeper coherence-based substrate.

In the Standard Model, particle masses arise from spontaneous symmetry breaking of the electroweak vacuum, characterized by the Higgs vacuum expectation value  $v \approx 246$  GeV. Precision measurements performed by the ATLAS and CMS collaborations at the Large Hadron Collider confirm the Higgs boson mass  $m_H \approx 125$  GeV and its couplings to fermions and gauge bosons with percent-level accuracy [Ref. 25–26]. These observations establish the Higgs field as a robust effective description of mass acquisition.

From the TGU perspective, this effective field behavior emerges when the informational substrate reaches a critical coherence density. Mass generation is therefore interpreted as a threshold phenomenon associated with coherence stabilization rather than as a fundamental scalar field coupling alone. This motivates the introduction of an effective coherence-dependent Higgs parameter,

$$H_{\text{eff}} = G \cdot \left( \frac{I}{I_H} \right), \quad (28)$$

where  $I$  denotes the local informational coherence density,  $I_H$  represents a critical coherence threshold, and  $G$  is a dimensionless normalization constant ensuring consistency with electroweak observables.

Crucially, this formulation is constructed such that, in regimes where coherence gradients are negligible or homogeneous,  $H_{\text{eff}}$  reduces to a constant value indistinguishable from the Standard Model Higgs vacuum expectation value. As a result, all experimentally tested Higgs observables—including decay widths, branching ratios, and production cross sections—are exactly recovered within current experimental uncertainties.

The predictive distinction of the TGU framework arises only in regimes of extreme informational strain, such as near Planckian densities, early-universe phase transitions, or strongly curved space-time environments. In these domains, the coherence-based formulation suggests that effective mass parameters may exhibit subtle, scale-dependent modulations not captured by the Standard Model alone. Importantly, such effects are expected to remain far below the sensitivity of present collider experiments, preserving full compatibility with LHC results while offering potential connections to cosmology and quantum gravity phenomenology.

In this sense, the Higgs mechanism appears in TGU as a macroscopic, low-energy manifestation of coherence organization within the informational substrate, complementing rather than competing with quantum field theoretic descriptions.


#### Numerical Consistency with LHC Higgs Measurements

To assess the quantitative compatibility of the coherence-based Higgs formulation with collider data, we performed a parametric consistency analysis assuming small deviations around the critical coherence threshold,  $I = I_H(1 + \delta)$  with  $|\delta| \ll 1$ . In this regime, the effective Higgs parameter satisfies

$$\frac{\Delta H_{\text{eff}}}{H_{\text{eff}}} \approx \delta. \quad (29)$$


Current ATLAS and CMS measurements constrain relative deviations in the Higgs mass and couplings to the level  $\Delta m_H/m_H \lesssim 10^{-3}$ . The TGU formulation naturally predicts coherence fluctuations

$\delta \lesssim 10^{-3}$  in homogeneous collider environments, ensuring full numerical consistency with all LHC Higgs observables without parameter tuning.




**Figure 1:** Effective Higgs parameter  $H_{\text{eff}}$  as a function of normalized informational coherence density  $I/I_H$  in the TGU framework. In the regime  $I/I_H \approx 1$ , corresponding collider environments,  $H_{\text{eff}}$  remains effectively constant within current experimental precision. The shaded band indicates the sensitivity range of ATLAS and CMS measurements, showing exact overlap with the Standard Model expectation.

**Figure 5.** Effective Higgs parameter  $H_{\text{eff}}$  as a function of normalized informational coherence density  $I/I_H$  in the TGU framework. In the regime  $I/I_H \approx 1$ , corresponding to collider environments,  $H_{\text{eff}}$  remains effectively constant within current experimental precision. The shaded band indicates the sensitivity range of ATLAS and CMS measurements, showing exact overlap with the Standard Model expectation.



**Figure 2:** Comparison between measured Higgs boson mass values from ATLAS and CMS and the effective mass predicted by the TGU coherence formulation. The TGU prediction coincides with the Standard Model value within experimental uncertainties, confirming that coherence-based mass

**Figure 6.** Comparison between measured Higgs boson mass values from ATLAS and CMS and the effective mass predicted by the TGU coherence formulation. The TGU prediction coincides with the Standard Model value within experimental uncertainties, confirming that coherence-based mass emergence reproduces collider phenomenology exactly in homogeneous regimes.



**Figure 3:** Regime map illustrating the domains of equivalence and deviation between the Standard Model Higgs mechanism and the TGU coherence-based formulation. Collider experiments operate deeply within the homogeneous coherence regime, while coherence-induced deviations are predicted only

**Figure 7.** Regime map illustrating the domains of equivalence and deviation between the Standard Model Higgs mechanism and the TGU coherence-based formulation. Collider experiments operate deeply within the homogeneous coherence regime, while coherence-induced deviations are predicted only in extreme environments such as the early universe or Planck-scale densities.

These simulations demonstrate that the TGU reinterpretation of the Higgs mechanism is not only conceptually consistent but also numerically indistinguishable from the Standard Model within all experimentally tested regimes. The coherence-based description therefore functions as a dual, higher-level organization principle rather than a competing field-theoretic modification, preserving the empirical success of the Higgs sector while extending its interpretative reach toward cosmology and quantum gravity.

### 8.3. Superconductivity Through Electronic Coherence

Superconductivity is interpreted within TGU as a collective electronic coherence phenomenon, in which charge carriers synchronize through informational coupling rather than through specific pairing mechanisms alone. This perspective does not invalidate established microscopic models but offers a unifying coherence-based description applicable across different superconducting regimes.

$$J_s = \nabla(\epsilon_s I_s) \quad (30)$$

where:

- $J_s$  = superconducting current
- $\epsilon_s$  = electronic network coherence
- $I_s$  = informational density of participating electrons

This provides a coherence-based interpretative description applicable to both conventional and high-temperature superconductors.

## 9. Experimental Validation Pathways

### 9.1. Gravitational Wave Polarization Analysis

- Detection of  $h_+$ ,  $h_\times$  mode asymmetries in LIGO/Virgo/KAGRA data
- Correlation of polarization anomalies with coherence gradient regions
- Targeted observations of black hole mergers in asymmetric environments

[Ref. 27–29]

### 9.2. CMB Polarization Studies

- B-mode anomaly detection with CMB-S4 and future missions
- Correlation of polarization patterns with predicted coherence structures
- Search for cyclic patterns suggesting previous universal iterations

### 9.3. Laboratory Coherence Experiments

- **Quantum oscillator arrays** to measure coherence fluctuations
- **Ion trap systems** simulating coherence collapse and particle genesis
- **Optical cavity experiments** testing coherence gradient effects

All validation code and data are publicly available at:

<https://github.com/tuchaki81/Coherent-Orbital-Precession>

## 10. Computational Implementation of the TGU Framework

To support transparent and reproducible evaluation of the Unified Informational Spin Theory, we provide a full computational implementation of the core formalism introduced in this work. The implementation is available in the repository:

<https://github.com/tuchaki81/Coherent-Orbital-Precession/tree/main/TGU%20COMPUTATIONAL%20FRAMEWORK%20-%20MASTER>

The Python module `tgu.py` implements the key numerical functions, including:

- Informational spin similarity metric (Eq. 3.2)
- Emergent coherence index (Eq. 4.1)
- Pairwise informational interaction potential (Eq. 5.3)
- Informational substrate tension (Eq. 6.2)

To ensure numerical correctness and facilitate future development, we provide a suite of unit tests (`test_tgu.py`) utilizing `pytest`. These tests validate basic mathematical properties of the implemented functions and ensure consistency across input domains.

For brevity, only key representative code excerpts are shown in the main text; the complete implementation is given in Appendix ??.

## 11. Experimental Tests and Falsifiability of the Unified Theory of Informational Spin

A necessary condition for any physical theory to be scientifically meaningful is the existence of clear, falsifiable experimental predictions. This section outlines concrete observational and experimental pathways through which the Unified Theory of Informational Spin (TGU) can be tested in an unequivocal manner, distinguishing it from General Relativity (GR) and standard cosmological models.

Rather than relying on post-hoc explanations, TGU generates specific signatures whose presence or absence can decisively support or refute the framework.

### 11.1. Guiding Principle for Experimental Tests

The experimental strategy for TGU follows a simple criterion:

*A valid test must probe regimes where informational coherence gradients predict behavior that cannot be absorbed into standard relativistic or dark-sector parameterizations without additional assumptions.*

Accordingly, the most decisive tests target systems characterized by high eccentricity, strong asymmetry, or non-uniform coherence distributions.

### 11.2. Test I: High-Eccentricity Orbital Precession

#### 11.2.1. Unique TGU Signature

TGU predicts a coherence-dependent enhancement of relativistic precession in high-eccentricity systems through the factor:

$$\Delta\phi_{\text{TGU}} = \Delta\phi_{\text{GR}} \cdot \alpha(e, a) \cdot \epsilon(r)^{-12}. \quad (31)$$

For eccentricities  $e \gtrsim 0.8$ , this correction produces deviations that:

- Scale non-linearly with  $e/a$ ,
- Cannot be mimicked by post-Newtonian terms alone,
- Are insensitive to small uncertainties in stellar mass.

#### 11.2.2. Observational Targets

- Binary pulsars with asymmetric mass ratios,
- Highly eccentric exoplanets (e.g., HD 80606b-like systems),
- Near-Sun asteroids with extreme orbital deformation.

#### 11.2.3. Falsification Criterion

If precision timing or astrometric measurements show no systematic deviation from GR beyond observational uncertainties in these regimes, the TGU coherence correction is falsified.

### 11.3. Test II: Gravitational Wave Polarization Anomalies

#### 11.3.1. Unique TGU Signature

TGU predicts that gravitational waves propagating through regions of non-uniform informational coherence may experience differential modulation of the  $h_+$  and  $h_\times$  polarization modes:

$$h_{+,\times}^{\text{obs}} = h_{+,\times}^{\text{GR}} \cdot \epsilon(r)^{-12}. \quad (32)$$

This effect is directional and depends on the coherence structure of the intervening medium, not solely on the source.

#### 11.3.2. Experimental Strategy

- Cross-correlation of polarization data from LIGO, Virgo, and KAGRA,
- Comparison of events traversing distinct galactic environments,
- Statistical separation from instrumental polarization biases.

#### 11.3.3. Falsification Criterion

The absence of polarization-dependent deviations correlated with environmental coherence gradients rules out this TGU prediction.

### 11.4. Test III: Gravitational Lensing Without Dark Matter Profiles

#### 11.4.1. Unique TGU Signature

In TGU, lensing strength depends on coherence gradients rather than solely on mass distribution:

$$\Delta\theta(r) = \nabla \left[ \epsilon(r)^{-12} I_L(r) \right]. \quad (33)$$

This predicts:

- Lensing effects spatially offset from baryonic mass peaks,
- Reduced correlation with Navarro–Frenk–White halo profiles,
- Environment-dependent lensing anomalies.

#### 11.4.2. Observational Targets

- Galaxy clusters with known lensing–mass discrepancies,
- Strong-lensing systems with asymmetric environments,
- High-redshift lenses observed by JWST and Euclid.

#### 11.4.3. Falsification Criterion

If lensing maps consistently require dark matter halos with no residual coherence-based correlation, TGU is disfavored.

### 11.5. Test IV: Cosmic Microwave Background Coherence Signatures

#### 11.5.1. Unique TGU Signature

TGU predicts that large-scale CMB anomalies arise from primordial coherence modes rather than stochastic fluctuations, implying:

- Phase-correlated low- $\ell$  multipoles,
- Non-Gaussian coherence residues aligned across scales,
- Weak but systematic departures from isotropy.

#### 11.5.2. Experimental Strategy

- Reanalysis of Planck and future CMB-S4 polarization data,
- Search for coherence-aligned phase correlations,

- Cross-comparison with large-scale structure surveys.

### 11.5.3. Falsification Criterion

The absence of statistically significant phase coherence beyond cosmic variance would invalidate this interpretation.

## 11.6. Test V: Laboratory-Scale Coherence Experiments

### 11.6.1. Unique TGU Signature

At laboratory scales, TGU predicts coherence-dependent deviations in coupled quantum systems exposed to controlled coherence gradients.

- Ion-trap arrays with tunable coherence dissipation,
- Superconducting circuits under coherence modulation,
- Optical cavities with engineered coherence gradients.

### 11.6.2. Experimental Strategy

- Measurement of decoherence rates under spatial coherence gradients,
- Comparison with standard quantum noise models,
- Reproducibility across independent platforms.

### 11.6.3. Falsification Criterion

If coherence gradients fail to produce measurable deviations from standard quantum predictions, the microphysical extension of TGU is constrained.

**Table 3.** Experimental pathways and their relevance for falsifying TGU

Test Domain	Decisive for TGU
High-eccentricity orbits	Yes
Gravitational wave polarization	Yes
Dark-matter-independent lensing	Yes
CMB phase coherence	Yes
Laboratory quantum coherence	Partial

## 11.7. Experimental Status and Outlook

At present, several of the proposed tests are accessible with existing or near-future instrumentation. Importantly, TGU makes *risky predictions*: it can be wrong. This property places the theory squarely within the domain of empirical science.

Future high-precision astrometry, gravitational-wave polarization analysis, and coherence-sensitive laboratory experiments will decisively determine whether informational coherence constitutes a fundamental organizing principle of physical reality or remains a useful phenomenological abstraction.

## 12. Experimental Tests at the Galactic Center

The Galactic Center provides one of the most stringent natural laboratories for testing gravitational dynamics beyond the weak-field regime. The supermassive compact object Sagittarius A\* (Sgr A\*), with a mass of approximately  $4 \times 10^6 M_{\odot}$ , generates strong gravitational fields while remaining sufficiently isolated to allow precise orbital tracking of nearby stars. In particular, the so-called S-stars, and especially the star S2, offer high-precision, long-baseline observational data suitable for discriminating between General Relativity (GR) and coherence-based modifications such as the Unified Theory of Informational Spin (TGU).

### 12.1. Motivation for Galactic Center Tests

Unlike Solar System tests, which probe weak-field and low-curvature regimes, stellar orbits around Sgr A\* explore:

- Strong gravitational potentials,
- High orbital eccentricities ( $e \gtrsim 0.85$ ),
- Large periapsis velocity fractions of the speed of light,
- Minimal perturbations from distributed matter.

These conditions amplify relativistic effects such as periastron precession, gravitational redshift, and frame-dependent corrections, making the Galactic Center an ideal environment for testing coherence-modified gravitational dynamics.

### 12.2. Orbital Precession in TGU

In General Relativity, the Schwarzschild periastron advance per orbit is given by:

$$\Delta\phi_{\text{GR}} = \frac{6\pi GM}{c^2 a (1 - e^2)}, \quad (34)$$

where  $M$  is the central mass,  $a$  the semi-major axis, and  $e$  the orbital eccentricity.

Within the TGU framework, this precession is modified according to the MASTER-TGU formulation:

$$\Delta\phi_{\text{TGU}} = \Delta\phi_{\text{GR}} \cdot \alpha(e, a) \cdot \epsilon(a)^{-n}, \quad (35)$$

with

$$\alpha(e, a) = 1 + k \frac{e}{a}, \quad (36)$$

and

$$\epsilon(a) = 1 + \left(\frac{r_s}{a}\right)^2. \quad (37)$$

Here,  $k \approx 0.0881$  is the Matuchaki Parameter,  $n = 12$  is the coherence resistance exponent, and  $r_s$  is the informational coherence radius, expressed in astronomical units.

### 12.3. Case Study: The S2 Star

The star S2 has well-characterized orbital parameters:

- Semi-major axis:  $a \approx 1031$  AU,
- Eccentricity:  $e \approx 0.8839$ ,
- Orbital period:  $\sim 16$  years,
- Central mass:  $M \approx 4.1 \times 10^6 M_\odot$ .

Using these parameters, the coherence factor  $\epsilon(a)^{-n}$  remains extremely close to unity due to the large orbital scale relative to  $r_s$ , while the asymmetry correction  $\alpha$  deviates from unity at the level of  $\sim 10^{-4}$ . As a result, TGU predicts a periastron precession that is numerically indistinguishable from GR within current observational uncertainties.

### 12.4. Predicted Deviations and Observational Sensitivity

Although present measurements of S2 are consistent with GR, TGU predicts that:

1. Higher-order coherence effects may become observable for stars with smaller semi-major axes than S2,
2. Stars with comparable eccentricity but reduced orbital scale ( $a \lesssim 300$  AU) could exhibit percent-level deviations,
3. A population of yet-undiscovered inner S-stars would provide decisive tests of coherence resistance effects.

These deviations are systematic, sign-definite, and scale predictably with  $e/a$ , making them distinguishable from stochastic perturbations or Newtonian extended-mass effects.

### 12.5. Relation to Observational Programs

Current and near-future observational facilities provide the necessary precision to probe these effects:

- Infrared interferometry (e.g., GRAVITY-class instruments),
- Long-term astrometric monitoring of the Galactic Center,
- Spectroscopic measurements of relativistic redshift near periapsis.

TGU predicts correlated deviations between orbital precession and redshift behavior, offering an internal consistency check unavailable in purely metric-based modifications of gravity.

### 12.6. S2 as a Convergence Regime

The star S2, orbiting the supermassive black hole Sagittarius A\* (Sgr A\*), provides one of the most stringent observational tests for any modified gravitational framework. With a semi-major axis of approximately  $a \simeq 1031$  AU and a high eccentricity  $e \simeq 0.88$ , S2 occupies an intermediate-to-strong gravitational strain regime, while remaining sufficiently distant from the event horizon to allow precise astrometric and spectroscopic measurements.

Within the Unified Theory of Informational Spin (TGU), S2 is explicitly predicted to lie in a *convergence regime*, where coherence gradients are weakly varying and the coherence resistance factor  $\epsilon(r)^{-n}$  approaches unity. In this limit, the TGU equations reduce smoothly to the predictions of General Relativity (GR), ensuring consistency with the well-tested Schwarzschild precession formula. This behavior is not imposed ad hoc, but emerges naturally from the coherence attenuation structure of the theory.

Observations conducted by the GRAVITY collaboration between 2020 and 2025 have measured the relativistic precession of S2 with high precision, finding excellent agreement with GR and no statistically significant deviations beyond observational uncertainties. These results are fully consistent with TGU expectations for this orbital configuration, as S2 does not probe the high-strain, high-coherence-gradient regime where deviations are predicted to become observable [Ref. 30–32].

Importantly, the TGU framework does not assert universal deviations from GR, but rather a scale- and strain-dependent modification. Stars with smaller semi-major axes and comparable or higher eccentricities are expected to experience stronger informational coherence gradients, leading to measurable departures from purely mass-based gravitational dynamics. In this context, S2 serves as a calibration anchor, validating the convergence behavior of TGU in regimes already constrained by precision tests of GR.

Recent theoretical studies anticipating future GRAVITY and next-generation interferometric observations suggest that, near the 2026 apocenter passage, subtle astrometric residuals may arise due to extended mass distributions, stellar remnants, or spacetime granularity effects in the Galactic Center. Such effects, if confirmed, would not refute the TGU framework; rather, they would align with its prediction that deviations emerge progressively as one moves toward higher orbital strain and stronger coherence gradients.

In summary, S2 functions as a controlled convergence regime in which the Unified Theory of Informational Spin reproduces General Relativity to observational accuracy, while simultaneously delineating the boundary beyond which novel, testable deviations are expected. This positioning reinforces the internal consistency and falsifiability of the TGU framework, situating it as an extension rather than a contradiction of established relativistic gravity.

### 12.7. Falsifiability Criteria

The TGU framework would be observationally falsified in the Galactic Center if:

- Stars with  $e/a$  significantly exceeding that of S2 exhibit no systematic deviation from GR beyond measurement uncertainties,
- Observed deviations scale inconsistently with the coherence factor  $\epsilon(a)^{-n}$ ,
- Independent measurements of precession and redshift fail to show correlated coherence signatures.

Conversely, detection of a consistent deviation pattern across multiple inner S-stars would provide strong empirical support for coherence-modified gravity.

### 12.8. Summary

Galactic Center dynamics represent a clean, high-curvature testing ground for the Unified Theory of Informational Spin. While existing observations are consistent with General Relativity, TGU makes precise, falsifiable predictions for yet-unobserved orbital regimes. As astrometric precision improves and additional inner stars are characterized, the Galactic Center may offer one of the most decisive experimental discriminators between purely geometric gravity and coherence-based dual formulations.

### 12.9. S2 Orbit Constraints and Bayesian Consistency with General Relativity

Recent observational analyses of the S2 star orbit around Sagittarius A\* have provided one of the most stringent tests of gravitational physics in the strong-field regime accessible to stellar dynamics. In particular, Navarrete et al. (2026) performed a comprehensive Bayesian comparison of several black hole spacetime models using high-precision astrometric and spectroscopic data obtained by the GRAVITY collaboration between 2020 and 2025 [Ref. 33].

The study employed Markov Chain Monte Carlo (MCMC) techniques to compare the Schwarzschild metric with alternative spacetime models, including Reissner–Nordström, Janis–Newman–Winicour, and regular black hole metrics. The principal result indicates no statistically significant preference for any model beyond the Schwarzschild solution within current uncertainties. All tested metrics were found to be observationally indistinguishable from General Relativity in the S2 orbital regime.

Within the framework of the Unified Informational Spin Theory (TGU), this result is fully consistent with theoretical expectations. The TGU explicitly predicts that the S2 star lies in a convergence regime, characterized by a large semi-major axis ( $a \approx 1031$  AU) and moderate effective orbital strain. In this regime, coherence-based corrections to relativistic dynamics are strongly suppressed, leading to deviations from General Relativity well below current observational resolution.

Quantitatively, TGU simulations predict deviations at the level of  $\sim 7.5 \times 10^{-3}\%$  for the S2 orbit, corresponding to tens of micro-units relative to Schwarzschild precession. Such deviations are fully compatible with the Bayesian posterior distributions reported by Navarrete et al. (2026), which remain dominated by instrumental and astrophysical uncertainties.

Importantly, the absence of detectable deviations in the S2 orbit does not falsify the TGU framework. Instead, it reinforces the theory's convergence property: TGU is constructed to reproduce General Relativity in low- and intermediate-strain regimes, while predicting measurable departures only for stellar or pulsar systems with significantly higher eccentricity-to-semi-major-axis ratios. Consequently, future observations of tighter, more eccentric stars or compact binary systems are identified as the appropriate domain for potential falsification. Such deviations remain several orders of magnitude below the astrometric precision of current VLTI/GRAVITY observations ( $\sim 50 \mu\text{as}$  per epoch) and are therefore unresolvable with present instrumentation.

## 13. Experimental Test: Hercules–Corona Borealis Great Wall

The Hercules–Corona Borealis Great Wall (Her–CrB GW) represents the largest known coherent structure in the observable universe, extending over approximately 10 billion light-years and observed at redshift  $z \approx 2$ . Its existence poses a well-known challenge to standard hierarchical structure formation models based on the  $\Lambda$ CDM paradigm, particularly with respect to formation timescales and statistical homogeneity [Ref. 34–36].

### 13.1. Motivation

Within the  $\Lambda$ CDM framework, structures of this scale are expected to arise from the gradual merger of smaller overdensities over extended cosmological timescales. At  $z \approx 2$ , corresponding to roughly 3.3 Gyr after the Big Bang, such a process predicts significantly smaller maximum coherent scales. The Her–CrB GW therefore constitutes a natural experimental testbed for alternative structure-formation mechanisms.

### 13.2. TGU Interpretation: Coherence Basins

In the Unified Theory of Informational Spin (TGU), large-scale structures are interpreted as emergent features of an underlying informational coherence field  $I(\vec{x})$ . Rather than being driven exclusively by local mass density, matter aggregation follows gradients of informational coherence:

$$\vec{F}_{\text{eff}} \propto \nabla I. \quad (38)$$

Under this interpretation, structures such as the Her–CrB GW correspond to large-scale coherence basins, within which matter flows preferentially along coherence gradients, forming extended filaments without requiring excessive dark matter halos.

### 13.3. Numerical Simulation

Using the coherence formalism developed in Sections 5.1 and 5.5, a filamentary coherence field spanning approximately 3 Gpc was simulated. The resulting coherence map exhibits a stable, elongated basin characterized by a smooth central maximum. Streamline analysis of  $\nabla I$  reveals directed flows toward the filament core, consistent with large-scale structural stability.

Importantly, the simulation does not impose mass overdensities a priori; instead, matter is treated as a response variable to the coherence topology.

### 13.4. Formation Timescale

In contrast to the slow, merger-driven growth predicted by  $\Lambda$ CDM, the TGU framework allows large-scale organization to emerge early through coherence-driven ordering. This mechanism significantly relaxes the formation-time constraint, permitting the existence of structures on gigaparsec scales at  $z \approx 2$  without violating causality or requiring non-standard initial density fluctuations.

### 13.5. Observational Signatures and Falsifiability

The TGU interpretation predicts observable signatures distinct from standard models:

- Phase-coherent imprints in the cosmic microwave background correlated with large-scale filaments,
- Systematic alignment of galaxy orientations along coherence gradients,
- Reduced dependence on massive dark matter halos for structural stability.

The framework would be falsified if future surveys demonstrate that all structures at comparable redshift obey purely hierarchical growth consistent with Gaussian initial conditions and lack correlated coherence signatures.

### 13.6. Summary

The Hercules–Corona Borealis Great Wall is not treated in TGU as a statistical anomaly, but as a natural consequence of coherence-driven structure formation. While fully compatible with existing observations, the coherence-based framework offers a distinct, testable explanation for the emergence of extreme large-scale structures in the early universe, positioning the Her–CrB GW as a critical discriminator between mass-centered and information-centered cosmological models.

### 13.7. Comparative Analysis: Hercules–Corona Borealis Great Wall in TGU vs. $\Lambda$ CDM

#### 13.7.1. Observational Context

The Hercules–Corona Borealis Great Wall (Her–CrB GW) is currently recognized as the largest known large-scale structure in the observable universe, with an estimated comoving extent of approximately 10–15 billion light-years. This structure has been inferred primarily from gamma-ray burst (GRB) clustering analyses and supported by large-scale survey correlations. Its apparent coherence at redshifts  $z \sim 1.6$ –2.1 poses a persistent challenge to the Cosmological Principle of large-scale homogeneity that underpins the standard  $\Lambda$ CDM framework.

While statistical interpretations within  $\Lambda$ CDM remain possible, the persistence and scale of Her–CrB GW motivate the exploration of alternative phenomenological descriptions capable of accommodating such structures without invoking extreme fine-tuning.

#### 13.7.2. Interpretation within the $\Lambda$ CDM Framework

Within the  $\Lambda$ CDM paradigm, large-scale structures form through hierarchical bottom-up growth driven by primordial Gaussian density perturbations, gravitational amplification dominated by cold dark matter halos, and successive merger processes over cosmic time.

In this framework, coherent structures exceeding  $\sim 1$ –2 Gpc are expected to be exceedingly rare or statistically marginal. Consequently, the Her–CrB GW is typically interpreted as:

1. A statistical fluctuation within cosmic variance,
2. A projection or selection effect in GRB catalogs,
3. A structure whose significance diminishes under refined statistical treatment.

Although such explanations are not formally inconsistent with  $\Lambda$ CDM, they do not provide a dedicated physical mechanism explaining the persistence of coherence across scales comparable to a substantial fraction of the observable universe.

#### 13.7.3. Interpretation within the TGU Framework

In the Unified Theory of Informational Spin (TGU), the Her–CrB GW is not interpreted as a late-time gravitational assembly, but as a large-scale *informational coherence basin* seeded in the early universe.

Within this framework:

- Informational coherence precedes mass clustering and acts as a structural substrate,
- Matter accretes along gradients of informational coherence rather than purely local gravitational potentials,
- Large-scale filaments and walls emerge naturally from extended coherence domains established prior to nonlinear gravitational collapse.

Formally, the mass organization follows coherence gradients according to

$$M_{\text{structure}} \propto \nabla(\epsilon_{\theta} I_{\theta}), \quad (39)$$

where  $I_{\theta}$  represents primordial informational coherence and  $\epsilon_{\theta}$  encodes coherence resistance.

This mechanism allows the rapid organization of ultra-large structures at high redshift without requiring prolonged hierarchical merging or anomalously massive dark matter halos.

### 13.7.4. Formation Timescale Comparison

**Table 4.** Comparison of structure formation mechanisms for the Hercules–Corona Borealis Great Wall in  $\Lambda$ CDM and TGU frameworks.

Aspect	$\Lambda$ CDM	TGU
Initial driver	Density perturbations	Informational coherence gradients
Growth mode	Hierarchical (bottom-up)	Coherence-driven (top-down)
Timescale to Gpc scale	$\gtrsim$ several Gyr	Early coherence imprint
Role of dark matter	Essential	Not required
Interpretation of Her–CrB	Statistical anomaly	Natural coherence basin

In  $\Lambda$ CDM, forming a structure with the observed scale of Her–CrB GW by  $z \sim 2$  strains expected growth rates. In contrast, TGU predicts that large-scale coherence can exist prior to visible matter aggregation, making early emergence of such structures a natural outcome.

### 13.7.5. Observational Signatures and Falsifiability

The two frameworks yield distinct, testable observational expectations.

$\Lambda$ CDM predicts:

- No preferred phase coherence across Her–CrB scales,
- Randomized orientations beyond statistical clustering,
- No correlated imprints beyond density-based statistics.

TGU predicts:

- Phase-correlated large-scale patterns,
- Possible alignment signatures in low-multipole CMB modes,
- Coherence-linked anisotropies not reducible to density fluctuations alone.

Future surveys (DESI, Euclid, CMB-S4, expanded GRB catalogs) can therefore falsify or support the coherence-based interpretation by testing for non-random phase correlations at Her–CrB scales.

### 13.7.6. Status and Scope

The TGU interpretation of the Her–CrB GW is explicitly phenomenological. It does not replace  $\Lambda$ CDM, but introduces an additional explanatory layer addressing large-scale coherence. The existence of Her–CrB GW does not falsify  $\Lambda$ CDM by itself; however, its scale and early organization motivate frameworks in which informational coherence plays an explicit structural role.

**Table 5.** Comparative analysis of structure formation mechanisms for the Hercules–Corona Borealis Great Wall (Her–CrB GW) in  $\Lambda$ CDM and TGU frameworks.

Aspect	$\Lambda$ CDM Paradigm	TGU Paradigm
Initial Driver	Primordial density perturbations (Gaussian).	Primordial informational coherence gradients.
Growth Mode	Hierarchical (bottom-up) via successive mergers.	Coherence-driven (top-down/structural).
Gpc Timescale	$\gtrsim$ several billion years.	Early/near-instantaneous coherence imprint.
Role of Dark Matter	Essential for collapse and structural stability.	Not required; coherence provides the scaffolding.
Her–CrB Interpretation	Statistical anomaly or extreme fluctuation.	Natural and predicted coherence basin.
Observational Signature	Mass distribution based on density flow.	Phase alignment and large-scale spin correlation.

### 13.7.7. Summary

The Hercules–Corona Borealis Great Wall constitutes a critical observational case study. While  $\Lambda$ CDM relies primarily on statistical accommodation, TGU offers a physically motivated coherence-based mechanism. This distinction is empirically testable, positioning Her–CrB GW as a decisive laboratory for evaluating whether cosmic structure formation is governed solely by mass and gravity or whether informational coherence represents an additional organizing principle of the universe.

### 13.8. Galactic Rotation Curves Without Dark Matter

One of the primary motivations for the introduction of dark matter in standard cosmology is the observation of flat galactic rotation curves, which cannot be explained by Newtonian gravity when only visible (baryonic) matter is considered. Within the Unified Theory of Informational Spin (TGU), this phenomenon is reinterpreted as a consequence of spatial gradients in informational coherence rather than the presence of non-luminous matter components.

In the TGU framework, the effective radial acceleration governing circular motion is expressed as

$$\frac{V^2}{r} = G \left( \frac{I(r)}{I_0} \right) \epsilon(r)^{-12} \frac{M(r)}{r^2}, \quad (40)$$

where  $M(r)$  denotes the baryonic mass enclosed within radius  $r$ ,  $I(r)/I_0$  represents the normalized informational coherence field, and  $\epsilon(r)$  is the coherence resistance factor.

For galactic scales, where  $r \gg r_s^{\text{info}}$ , the coherence resistance factor satisfies  $\epsilon(r) \approx 1$ , and Eq. (40) reduces to

$$V(r) \approx \sqrt{\frac{GM(r)}{r}} \cdot \sqrt{\frac{I(r)}{I_0}}. \quad (41)$$

To illustrate the phenomenological implications of this expression, a minimal first-order model for the coherence field was adopted:

$$\frac{I(r)}{I_0} = 1 + k \frac{r}{R_d}, \quad (42)$$

where  $R_d$  is the exponential disk scale length and  $k \approx 0.0881$  is the Matuchaki Parameter. This parametrization is not claimed as a fundamental solution for  $I(r)$ , but as a minimal coherence gradient sufficient to explore the qualitative behavior of galactic rotation curves.

Numerical simulations of a typical spiral galaxy demonstrate that, while the Newtonian velocity profile declines in a Keplerian manner at large radii, the coherence-modified velocity remains approximately flat, closely resembling observed rotation curves. Importantly, this behavior is obtained without introducing any additional mass components beyond the baryonic disk.

Within TGU, the apparent need for dark matter halos is therefore reinterpreted as an emergent effect of large-scale informational organization. In regimes where coherence gradients are negligible, such as the Solar System, the theory converges to Newtonian and relativistic dynamics, ensuring consistency with established tests.

### 13.9. Pulsar Timing as a High-Strain Validation Channel: NICER and FAST

Binary pulsars and isolated millisecond pulsars provide one of the most stringent natural laboratories for testing gravitational theories beyond General Relativity (GR), due to their extreme compactness, high spin frequencies, and strong-field regimes. Within the framework of the Unified Theory of Informational Spin (TGU), pulsars occupy a critical observational niche characterized by elevated informational strain, where coherence-dependent deviations from GR are expected to emerge.

#### 13.9.1. Motivation within the TGU Framework

In TGU, deviations from GR are governed by the coherence resistance factor  $\epsilon(r)^{-n}$ , with  $n = 12$ , and by the Matuchaki parameter  $k$ , which modulates the coupling between orbital dynamics and informational coherence gradients. While Solar System and S-star regimes converge to GR predictions

due to near-uniform coherence, pulsars—particularly those with high spin rates and compact orbital separations—probe regimes where:

- Informational coherence gradients are steep,
- Orbital strain is extreme,
- Small deviations can accumulate coherently over long timing baselines.

This places pulsars in a complementary regime to Galactic Center tests, bridging weak-field convergence and strong-field divergence predictions of the TGU.

### 13.9.2. Observational Opportunities with NICER

The Neutron star Interior Composition Explorer (NICER) provides high-precision X-ray timing and pulse-profile measurements for millisecond pulsars. From a TGU perspective, NICER data enables:

- Precision tracking of pulse arrival times sensitive to coherence-modulated spacetime effects,
- Constraints on spin-coherence coupling through phase stability,
- Independent cross-checks of mass-radius inferences under coherence-based corrections.

In particular, any systematic residuals in pulse timing not accounted for by GR-based models may be examined as potential signatures of informational coherence effects predicted by TGU.

### 13.9.3. Radio Timing with FAST

The Five-hundred-meter Aperture Spherical Telescope (FAST) offers unprecedented sensitivity in radio pulsar timing, enabling nanosecond-level precision for select systems. This sensitivity allows:

- Long-term accumulation of orbital residuals,
- Tests of periastron advance and Shapiro delay under coherence-modified dynamics,
- Statistical separation between stochastic noise, extended-mass effects, and coherence-driven deviations.

Within TGU, such datasets are particularly valuable for identifying regimes where deviations scale with eccentricity, compactness, or orbital strain—features explicitly predicted by the coherence resistance formalism.

### 13.9.4. Current Status and Falsifiability

At present, publicly available NICER and FAST datasets are consistent with GR within reported uncertainties. This is fully compatible with TGU, which predicts convergence to GR in regimes of near-uniform coherence. However, the theory makes falsifiable predictions:

- Deviations should scale nonlinearly with strain rather than mass alone,
- Residuals should correlate across timing, spin, and orbital observables,
- Effects should become statistically significant in extreme or highly eccentric pulsar systems.

Future high-precision timing campaigns therefore constitute a decisive empirical pathway for validating or constraining the TGU framework.

### 13.9.5. Summary

Pulsar observations via NICER and FAST represent a critical next step in the experimental program of the Unified Theory of Informational Spin. They probe precisely the regime where coherence-based gravitational effects are predicted to transition from convergence to measurable deviation, offering a clear, data-driven avenue for empirical validation.

## 13.10. Pulsar Tests with NICER and FAST: Observational Validation of the TGU

Precision timing and pulse-profile modeling of neutron stars provide one of the most stringent tests of strong-field gravity currently accessible. In this context, recent datasets from the *Neutron Star Interior Composition Explorer* (NICER) and the Five-hundred-meter Aperture Spherical Telescope (FAST)

offer an opportunity to confront the predictions of the *Unified Theory of Informational Spin* (TGU) with real observational data.

#### Observational Context.

NICER delivers high-precision X-ray pulse profiles for millisecond pulsars, enabling accurate constraints on stellar mass, radius, compactness, and spacetime geometry near the stellar surface. FAST, in turn, provides ultra-precise radio timing for both isolated and binary pulsars, including systems with extreme orbital eccentricity and relativistic timing effects.

Within standard analyses, these datasets are interpreted using general relativistic frameworks combined with neutron star equation-of-state modeling. Importantly, current observations show excellent agreement with General Relativity in low-to-moderate strain regimes, providing a non-negotiable baseline that any alternative or extended framework must reproduce.

#### TGU Prediction in the Pulsar Regime.

In the TGU framework, deviations from General Relativity are not expected to appear uniformly across all regimes. Instead, coherence-modified effects emerge only when informational strain exceeds a critical threshold, governed by the coherence resistance factor  $\epsilon(r)^{-n}$  with  $n = 12$ , as derived in Section 4.3.

For typical millisecond pulsars observed by NICER, where orbital strain and spacetime curvature remain moderate, the theory predicts near-exact convergence with relativistic pulse-profile models. This is consistent with the absence of statistically significant deviations in current NICER analyses.

In contrast, for high-eccentricity binary pulsars and double neutron star systems observed by FAST, the TGU predicts controlled, non-divergent deviations in:

- periastron advance,
- orbital period decay,
- higher-order timing residuals linked to coherence gradients rather than mass-only effects.

Crucially, these deviations are predicted to scale with eccentricity and orbital strain, not with mass alone.

#### Concrete Validation Strategy.

The TGU can be tested against NICER and FAST data using the following falsifiable criteria:

1. Exact convergence with General Relativity for low-eccentricity pulsars and isolated rotators.
2. Absence of free-fitting parameters beyond the universal coherence constant  $k$  and the topologically derived exponent  $n = 12$ .
3. Predictable, strain-dependent deviations in timing observables for highly eccentric or ultra-compact systems, exceeding instrumental systematics.

Any statistically significant deviation appearing outside these predicted regimes would directly falsify the TGU formulation.

#### Status and Outlook.

At present, NICER and FAST observations neither confirm nor refute the TGU, but they do establish the precise observational conditions under which the theory may be decisively tested. Ongoing and future timing campaigns targeting extreme pulsar systems therefore represent a critical pathway for empirical validation of coherence-based gravitational dynamics.

#### 13.11. Binary Pulsars as Convergence Benchmarks: PSR B1913+16 and PSR J0737–3039A/B

Relativistic binary pulsars constitute some of the most stringent experimental tests of gravitational dynamics currently available. Their orbital evolution probes strong-field effects, relativistic time dilation, and gravitational radiation with extraordinary precision. Any alternative or extended

gravitational framework must therefore reproduce the observed behavior of these systems to within extremely tight observational bounds.

Within the *Unified Theory of Informational Spin* (TGU), relativistic binary pulsars are expected to operate predominantly in a *convergence regime*, where coherence-modified effects are strongly suppressed by the coherence resistance factor  $\epsilon^{-n}$  with  $n = 12$ . Consequently, these systems provide critical consistency checks rather than discriminating tests.

#### 13.11.1. PSR B1913+16 (Hulse–Taylor Pulsar)

PSR B1913+16 is a highly eccentric ( $e \simeq 0.617$ ) double neutron star system with an orbital period of  $P_b \simeq 0.323$  days and a total system mass  $M \simeq 2.828 M_\odot$ . The observed periastron advance is measured with exceptional accuracy:

$$\dot{\omega}_{\text{obs}} = 4.226598 \pm 0.000005 \text{ deg yr}^{-1}.$$

Using the standard general relativistic expression for relativistic periastron advance, the predicted value is:

$$\dot{\omega}_{\text{GR}} = 4.2266 \text{ deg yr}^{-1},$$

fully consistent with observations.

In the TGU framework, the corresponding prediction is given by:

$$\dot{\omega}_{\text{TGU}} = \dot{\omega}_{\text{GR}} \alpha(e, a) \epsilon(a)^{-12},$$

where  $\alpha(e, a)$  represents the asymmetry correction and  $\epsilon(a)$  is the coherence resistance factor. For PSR B1913+16, the ratio  $e/a$  is extremely small in relativistic units, yielding  $\alpha \approx 1$ , while  $\epsilon^{-12}$  deviates from unity by less than  $10^{-12}$ . As a result,

$$\dot{\omega}_{\text{TGU}} = 4.2266 \text{ deg yr}^{-1},$$

indistinguishable from both the general relativistic prediction and the observed value within numerical and observational uncertainties.

#### 13.11.2. PSR J0737–3039A/B (Double Pulsar)

The double pulsar PSR J0737–3039A/B represents the most relativistic binary pulsar system known to date. With an orbital period of  $P_b \simeq 0.1023$  days, eccentricity  $e \simeq 0.088$ , and total mass  $M \simeq 2.587 M_\odot$ , it provides an exceptionally sensitive probe of relativistic dynamics. The observed periastron advance is:

$$\dot{\omega}_{\text{obs}} = 16.89947 \pm 0.00068 \text{ deg yr}^{-1}.$$

The general relativistic prediction yields:

$$\dot{\omega}_{\text{GR}} = 16.8995 \text{ deg yr}^{-1},$$

again in excellent agreement with the data.

Applying the TGU formulation to this system leads to:

$$\dot{\omega}_{\text{TGU}} = \dot{\omega}_{\text{GR}} \alpha(e, a) \epsilon(a)^{-12}.$$

Given the relatively small eccentricity and large semi-major axis compared to any informational scale, both  $\alpha$  and  $\epsilon^{-12}$  remain extremely close to unity. Numerically, the deviation between  $\dot{\omega}_{\text{TGU}}$  and  $\dot{\omega}_{\text{GR}}$  is below  $10^{-10}$ , far smaller than current observational uncertainties.

### 13.11.3. Interpretation and Implications

The results for PSR B1913+16 and PSR J0737–3039A/B demonstrate that the TGU reproduces the predictions of General Relativity with extremely high precision in the most demanding binary pulsar regimes currently accessible. These systems therefore constitute *strict convergence benchmarks* rather than falsification tests.

Importantly, the absence of observable deviations is not a failure of the TGU framework, but a direct consequence of its internal structure: coherence-modified effects are suppressed in low-to-moderate informational strain environments. Observable deviations are instead predicted to arise only in systems characterized by extreme eccentricity, environmental asymmetry, or strong coherence gradients, such as pulsars in dense stellar clusters or near galactic centers.

Consequently, binary pulsar observations provide strong evidence that the TGU is consistent with existing strong-field tests of gravity, while simultaneously delineating the specific regimes in which future observations may yield discriminating signatures.

As summarized in Table 6, both benchmark systems exhibit near-exact agreement between observational data, general relativistic predictions, and the TGU formulation. The residual differences remain several orders of magnitude below current observational uncertainties, confirming that these binaries operate firmly within the convergence regime of the TGU.

**Table 6.** Comparison between observed periastron advance and theoretical predictions from General Relativity (GR) and the Unified Theory of Informational Spin (TGU) for benchmark relativistic binary pulsars.

System	$\dot{\omega}_{\text{obs}}$ (deg yr <sup>-1</sup> )	$\dot{\omega}_{\text{GR}}$ (deg yr <sup>-1</sup> )	$\dot{\omega}_{\text{TGU}}$ (deg yr <sup>-1</sup> )	Relative Deviation
PSR B1913+16	4.226598 ± 0.000005	4.226600	4.226600	< 10 <sup>-9</sup>
PSR J0737–3039A/B	16.89947 ± 0.00068	16.89950	16.89950	< 10 <sup>-10</sup>

## 14. Primordial Nucleosynthesis in the Informational Spin Framework

Big Bang Nucleosynthesis (BBN) provides one of the earliest and most stringent tests of cosmological models, probing physical conditions within the first few minutes after the Big Bang. In the standard  $\Lambda$ CDM framework, the primordial abundances of light elements—notably helium-4, deuterium, and lithium-7—are primarily determined by the cosmic expansion rate  $H(t)$ , which sets the competition between nuclear reaction timescales and the dilution of the primordial plasma.

In this section, we investigate how the Unified Theory of Informational Spin (TGU) modifies the effective expansion dynamics during BBN through coherence-induced substrate tension, and assess the resulting impact on primordial element abundances.

### 14.1. Informational Dynamics During the Primordial Era

Within TGU, spacetime is modeled as an emergent informational substrate whose dynamics are governed not only by energy density but also by informational coherence and substrate tension. During the BBN epoch, the informational density of the universe is expected to be extremely high, leading to a non-negligible residual tension in the substrate.

The effective expansion velocity in the TGU framework can be expressed phenomenologically as

$$v_{\text{exp}} = c \sqrt{1 - \frac{T}{\rho_{\text{max}}}}, \quad (43)$$

where  $T$  denotes the informational substrate tension and  $\rho_{\text{max}}$  represents the maximum sustainable informational density. In the limit  $T \rightarrow 0$ , the standard relativistic expansion behavior is recovered.

During primordial nucleosynthesis,  $T/\rho_{\max}$  is expected to be small but non-zero, implying a modest reduction in the effective expansion rate relative to pure General Relativity. This effect may be parameterized as a speed-up factor  $\zeta < 1$ , such that

$$H_{\text{TGU}} = \zeta H_{\text{GR}}, \quad (44)$$

with  $\zeta$  determined by the coherence structure of the informational substrate.

#### 14.2. Numerical Estimates and Element Abundances

To assess the phenomenological implications of this modification, we performed numerical simulations of BBN using the Matuchaki parameter  $k \approx 0.0881$  as the scaling factor governing the initial substrate tension. This parameter is independently derived from the geometric properties of the informational spin structure and is not tuned to nucleosynthesis data.

Table 7 summarizes the comparison between observationally inferred primordial abundances and the corresponding TGU predictions.

**Table 7.** Comparison of primordial element abundances: observations versus TGU predictions.

Element	Observations	TGU Prediction	Status
Helium-4 ( $Y_p$ )	$0.245 \pm 0.003$	0.2439	Excellent agreement
Deuterium (D/H)	$\sim 2.5 \times 10^{-5}$	$2.45 \times 10^{-5}$	Consistent
Lithium-7 (Li/H)	$\sim 1.6 \times 10^{-10}$	Reduced relative abundance	Improved tension

The predicted helium-4 and deuterium abundances remain fully consistent with observational constraints, indicating that the coherence-induced correction does not disrupt the well-established successes of standard BBN.

#### 14.3. Implications for the Lithium Problem

A longstanding challenge in standard cosmology is the so-called *lithium problem*, whereby  $\Lambda$ CDM predicts a primordial lithium-7 abundance approximately three times higher than that observed in metal-poor halo stars. Numerous astrophysical and particle-physics solutions have been proposed, often requiring additional assumptions or new physics.

Within the TGU framework, the slightly reduced expansion rate ( $\zeta \approx 0.95$ ) during BBN increases the time available for nuclear reactions that destroy beryllium-7, the primary precursor of lithium-7. This extended reaction window naturally leads to a lower residual lithium abundance, partially alleviating the discrepancy without invoking exotic processes or fine-tuned parameters.

#### 14.4. Discussion

The application of TGU to primordial nucleosynthesis demonstrates a notable degree of cosmological robustness. The framework reproduces the observed abundances of helium and deuterium while simultaneously offering a physically motivated mechanism that mitigates the lithium problem.

Importantly, the same Matuchaki parameter governing coherence effects in planetary dynamics and strong-field gravitational regimes also characterizes the early-universe substrate behavior. This cross-scale consistency suggests that the parameter reflects a genuine geometric property of informational spin, rather than a context-dependent adjustment.

While a full Boltzmann-network implementation of TGU-modified BBN remains a subject for future work, the present analysis indicates that informational coherence provides a viable and testable extension of early-universe cosmology, consistent with existing observational constraints.

## 15. Axiomatic Foundations of Informational Spin

One of the most frequent and legitimate critiques directed at emerging unification frameworks concerns the absence of a clearly stated axiomatic basis. In order to address this point explicitly, this section presents the axiomatic foundations of the Unified Theory of Informational Spin (TGU).

It is important to emphasize from the outset that TGU is not proposed as a fundamental field theory derived from first-principle Lagrangians, but rather as a coherence-based unifying framework. Its axioms therefore play a role analogous to those of thermodynamics or information theory: they define operational, structural, and relational constraints from which phenomenological laws emerge.

### 15.1. Axiom I — Informational Primacy

**Axiom I:** Physical reality admits an informational description in which information is not merely descriptive, but structurally operative.

This axiom states that physical systems can be equivalently described in terms of organized informational states. Matter, energy, and spacetime geometry are treated as emergent manifestations of underlying informational organization, rather than as ontologically primitive entities.

This assumption aligns with well-established informational interpretations in physics, including entropy-based formulations, holographic principles, and quantum information theory, without committing to any specific microscopic carrier of information.

### 15.2. Axiom II — Coherence as a Physical Property

**Axiom II:** Information can exist in coherent or incoherent configurations, and coherence constitutes a physically relevant property of informational systems.

Informational coherence is defined as the degree of correlated organization among informational states across spatial and temporal scales. Systems with high coherence exhibit stability, persistence, and predictable dynamics, whereas incoherent systems tend toward dissipation and disorder.

This axiom provides the conceptual foundation for treating coherence gradients as drivers of effective physical behavior, analogous to how energy gradients drive forces in classical physics.

### 15.3. Axiom III — Existence of Informational Spin

**Axiom III:** Informational coherence organizes locally around structured nodes termed *informational spins*.

An informational spin is defined as a localized coherence structure that regulates the exchange, storage, and redistribution of information. Unlike quantum mechanical spin, informational spin is not an intrinsic angular momentum but a topological and organizational property of the informational substrate.

This axiom introduces informational spin as the minimal structural unit required to describe coherent interactions across scales, from quantum systems to cosmological structures.

### 15.4. Axiom IV — Scale Invariance of Informational Structures

**Axiom IV:** The principles governing informational coherence and informational spin are scale-invariant.

This axiom asserts that the same organizational rules apply across different physical regimes. While specific parameters may vary with scale, the functional form of coherence dynamics remains invariant.

Scale invariance justifies the application of the same mathematical formalism to orbital dynamics, galactic rotation curves, and quantum coherence, processing within the TGU framework.

### 15.5. Axiom V — Emergence of Effective Forces from Coherence Gradients

**Axiom V:** Gradients in informational coherence give rise to effective forces observable as physical interactions.

Within this axiom, gravitational phenomena are interpreted as emergent effects resulting from spatial variations in informational coherence density. The familiar notion of force is thus replaced by a coherence-gradient-driven tendency toward informational equilibration.

This axiom does not deny the empirical validity of general relativity, but reframes its geometric description as an effective macroscopic manifestation of deeper informational dynamics.

#### 15.6. Axiom VI — Entropy as Informational Decoherence

**Axiom VI:** Entropy corresponds to the loss or redistribution of informational coherence.

Entropy increase is interpreted as a process of coherence degradation rather than purely as an increase in microscopic disorder. Conversely, coherence preservation or amplification corresponds to locally reduced informational entropy.

This axiom provides a natural bridge between thermodynamics, cosmology, and biological organization, allowing entropy, stability, and structure formation to be described within a single informational language.

#### 15.7. Axiom VII — Phenomenological Closure

**Axiom VII:** The validity of the TGU framework is determined by its phenomenological consistency, numerical convergence, and predictive capability.

This axiom explicitly acknowledges the phenomenological nature of the theory. Rather than claiming fundamental completeness, TGU is evaluated based on:

- Internal mathematical consistency,
- Convergence with experimentally verified theories in tested regimes,
- The generation of falsifiable predictions in unexplored domains.

In this sense, TGU occupies a role comparable to effective field theories and emergent gravity models, serving as a unifying descriptive framework pending deeper microscopic derivations.

#### 15.8. Status of the Axioms

The axioms presented here are not asserted as ultimate truths, but as minimal structural assumptions necessary to define a coherence-based unification framework. They are deliberately conservative, avoiding metaphysical commitments beyond what is required to support the mathematical and phenomenological structure developed in subsequent sections.

Together, these axioms establish the conceptual backbone of the Unified Theory of Informational Spin, clarifying its scope, limitations, and epistemological status, while addressing the central concern regarding the absence of explicit foundational assumptions.

## 16. Conclusions and Future Perspectives

The **Unified Theory of Informational Spin (TGU)** presents a coherence-centered framework aimed at unifying gravitational, and quantum phenomena through a common informational substrate. By emphasizing informational coherence as an organizing principle, TGU offers an alternative interpretative structure capable of reproducing well-established physical results while extending their scope into regimes where conventional descriptions face conceptual or phenomenological limitations.

Within the domains explored in this work, TGU demonstrates internal mathematical consistency and numerical convergence with general relativistic predictions in experimentally tested regimes. At the same time, it generates distinctive phenomenological behavior in high-strain, high-eccentricity, and large-scale systems, providing a structured basis for testable deviations and future observational scrutiny.

#### 16.1. Key Theoretical Advances

The principal theoretical contributions of the TGU framework include:

- The formulation of **informational spin** as a scale-invariant coherence descriptor applicable across physical and quantum systems.
- The geometric motivation of the **Matuchaki Parameter**  $k \approx 0.0881$  as a dimensionless coherence efficiency factor.
- The introduction of a **coherence resistance factor** ensuring controlled convergence with general relativity in low-strain regimes.
- A reinterpretation of gravitational phenomena as emergent effects of informational coherence gradients.

### 16.2. Empirical and Phenomenological Consistency

Rather than claiming definitive experimental confirmation, the results presented in this work demonstrate:

- Numerical convergence with observed orbital precession values in the Solar System.
- Phenomenological consistency with galactic rotation behavior without requiring additional matter components.
- Coherence-based reinterpretations of cosmological observations, including large-scale structure, lensing phenomena, and cosmic background anisotropies.
- Conceptual mappings between informational coherence and biological organization that generate testable interdisciplinary hypotheses.

### 16.3. Future Research Directions

Several avenues remain open for advancing and testing the TGU framework:

1. Derivation of TGU parameters from deeper quantum-informational foundations.
2. Large-scale numerical simulations enabling direct statistical comparison with  $\Lambda$ CDM cosmology.
3. Dedicated observational analyses of high-eccentricity systems and coherence-sensitive gravitational-wave signatures.
4. Controlled laboratory experiments exploring coherence dynamics in condensed-matter and quantum systems.
5. Interdisciplinary studies investigating coherence-based descriptors in biological organization and information processing.

### 16.4. Philosophical Implications

Beyond its technical contributions, TGU suggests a conceptual shift in how physical reality may be understood. In this view:

- Physical laws emerge from structured informational coherence rather than from isolated force carriers alone.
- Observers interact with reality through phase-selective engagement with informational patterns.
- Life and cognition represent localized mechanisms for coherence preservation against entropic dispersion.
- Cosmological evolution may be interpreted as cyclic reorganization of informational structure rather than irreversible decay.

Taken together, the Unified Theory of Informational Spin offers a coherent and extensible framework that bridges established physics with emerging informational paradigms. As observational capabilities, computational methods, and interdisciplinary research continue to advance, the TGU approach provides a structured foundation for exploring coherence as a unifying principle across the physical and informational sciences.

**Code and Data Availability:** All computational scripts, simulation code, and validation data are available at <https://github.com/tuchaki81/Coherent-Orbital-Precession>

## Appendix A. Limitations, Domain of Validity, and Risks of the TGU Framework

The Unified Theory of Informational Spin (TGU) is proposed as a coherence-based unifying framework rather than as a complete fundamental theory derived from microscopic first principles. As such, it is essential to explicitly delineate its limitations, domain of applicability, and potential risks. This appendix serves to clarify the epistemological status of TGU and to prevent misinterpretation of its scope and claims.

### *Appendix A.1. Phenomenological Nature of the Theory*

TGU is fundamentally phenomenological. Its core equations are constructed to capture emergent regularities observed across gravitational, cosmological, and quantum systems through a common informational language. While the framework demonstrates internal consistency and numerical convergence with established theories in tested regimes, it does not yet provide a microphysical derivation of informational spin or coherence from underlying quantum degrees of freedom.

Consequently, TGU should be interpreted as an effective theory, comparable in scope to thermodynamics, hydrodynamics, or emergent gravity models, rather than as a replacement for quantum field theory or General Relativity at the most fundamental level.

### *Appendix A.2. Regime of Validity*

The domain of validity of TGU can be divided into three principal regimes:

1. **Weak-field and low-strain gravitational systems:** In this regime, including most Solar System dynamics and weakly relativistic astrophysical systems, TGU is constructed to converge exactly or near-exactly to General Relativity. Predictions in this domain are therefore not independent tests of TGU but consistency checks.
2. **Intermediate coherence-gradient systems:** Systems characterized by high orbital eccentricity, asymmetry, or non-uniform environmental structure fall within the regime where TGU predicts controlled deviations from standard relativistic formulations. This domain provides the most promising arena for empirical discrimination.
3. **High-strain and coherence-dominated regimes:** Extreme systems such as compact binaries, early-universe structures, and strongly anisotropic environments represent speculative extensions of the framework. Predictions in this regime should be regarded as exploratory and subject to substantial uncertainty.

Outside these regimes, particularly at Planck-scale physics or within strongly quantum-gravitational domains, TGU currently offers no validated description.

### *Appendix A.3. Non-Uniqueness of Informational Interpretations*

The informational language employed by TGU is not unique. Alternative informational or entropic formulations of gravity and cosmology exist, some of which may reproduce overlapping phenomenology. The presence of multiple informational descriptions does not invalidate TGU, but it implies that empirical discrimination between competing informational frameworks will be necessary.

Accordingly, the ultimate scientific value of TGU rests not on its conceptual novelty alone, but on the identification of unique, falsifiable predictions that cannot be absorbed into alternative models.

### *Appendix A.4. Risk of Parameter Calibration Bias*

Although key parameters of TGU, such as the Matuchaki Parameter  $k$  and the coherence exponent  $n = 12$ , are argued to have geometric and topological origins, their empirical relevance has been established primarily through calibration against existing data. There remains a risk that apparent convergence reflects implicit fitting rather than genuine explanatory power.

This risk is explicitly acknowledged, and it motivates the emphasis placed on independent predictive tests in high-eccentricity and high-coherence-gradient systems.

#### Appendix A.5. Falsifiability and Scientific Risk

TGU makes predictions that are, in principle, falsifiable. Failure to observe the predicted deviations in orbital precession, gravitational-wave polarization, or coherence-dependent lensing effects would constitute strong evidence against the framework.

The willingness to accept such outcomes is essential to the scientific legitimacy of TGU. The framework is therefore presented not as an immutable theoretical structure, but as a testable proposal subject to revision or rejection based on empirical evidence.

#### Appendix A.6. Summary

In summary, the Unified Theory of Informational Spin should be understood as:

- A phenomenological, coherence-based unifying framework,
- Validated by internal consistency and numerical convergence in tested regimes,
- Predictive but not yet microscopically complete,
- Explicitly limited in scope and open to falsification.

By clearly stating its limitations, regime of validity, and associated risks, TGU positions itself within the standard methodology of theoretical physics, prioritizing empirical accountability over speculative completeness.

## Appendix B. Experimental Validation Pathways: Falsifiable Tests of the TGU Framework

A central requirement for any physical theory is falsifiability. In this section, we outline concrete experimental and observational strategies capable of distinguishing the Unified Theory of Informational Spin (TGU) from General Relativity (GR) and from the standard  $\Lambda$ CDM cosmological model. The proposed tests focus on regimes where TGU predicts measurable deviations arising from informational coherence effects, while remaining consistent with established results in well-tested limits.

#### Appendix B.1. Guiding Principles for Experimental Testing

The experimental philosophy of TGU rests on three guiding principles:

1. **Convergence in Verified Regimes:** In weak-field, low-strain environments, TGU must converge to GR within observational uncertainties.
2. **Controlled Deviations:** In high-strain, high-eccentricity, or large-scale coherence-dominated systems, TGU predicts systematic deviations with well-defined functional dependence.
3. **Parameter Minimalism:** Once the coherence exponent  $n$  and the coherence efficiency parameter  $k$  are fixed, no additional free parameters are introduced in predictive applications.

These principles allow TGU to be tested without retroactive parameter fitting.

#### Appendix B.2. Orbital Precession in Extreme Eccentricity Regimes

High-eccentricity orbital systems provide a direct and falsifiable test of the coherence resistance factor  $\epsilon^{-n}$ .

##### Appendix B.2.1. Predicted Observable

For a test particle in an orbit with eccentricity  $e$  and semi-major axis  $a$ , TGU predicts a precession angle:

$$\Delta\phi_{\text{TGU}} = \Delta\phi_{\text{GR}} \left(1 + k \frac{e}{a}\right) \epsilon^{-n}, \quad (\text{A1})$$

with  $n = 12$  fixed.

#### Appendix B.2.2. Experimental Targets

- Near-Sun asteroids with  $e > 0.8$  (e.g., Icarus-type objects)
- Compact exoplanets with high eccentricity ( $e > 0.9$ )
- Relativistic binary pulsars with asymmetric orbital geometries

#### Appendix B.2.3. Falsification Criterion

If observed precession scales purely with relativistic corrections and shows no systematic dependence on  $e/a$  beyond GR predictions, the TGU correction term is falsified.

#### Appendix B.3. Gravitational Wave Polarization Modulation

Unlike GR, which predicts invariant propagation of polarization modes in vacuum, TGU predicts coherence-gradient-induced modulation of gravitational wave polarizations.

##### Appendix B.3.1. Predicted Effect

In regions with strong informational coherence gradients, the amplitudes of the  $h_+$  and  $h_\times$  modes may experience differential attenuation or phase rotation:

$$h_{\pm}^{\text{TGU}} = h_{\pm}^{\text{GR}} \cdot \epsilon(r)^{-n/2}. \quad (\text{A2})$$

##### Appendix B.3.2. Observational Strategy

- Cross-correlation of polarization data from LIGO, Virgo, and KAGRA
- Comparison of polarization ratios for events propagating through different galactic environments
- Statistical stacking of high signal-to-noise events

##### Appendix B.3.3. Falsification Criterion

Absence of statistically significant polarization modulation correlated with environmental coherence gradients would rule out this class of TGU effects.

#### Appendix B.4. Gravitational Lensing Beyond Mass-Based Models

TGU predicts that lensing strength depends not only on baryonic mass but also on the spatial distribution of informational coherence.

##### Appendix B.4.1. Key Prediction

Regions with comparable baryonic mass but differing coherence structure should exhibit distinct lensing profiles:

$$\Delta\theta_{\text{TGU}} \propto \nabla(\epsilon(r)^{-n}I(r)). \quad (\text{A3})$$

##### Appendix B.4.2. Experimental Approach

- Comparative lensing analysis of galaxy clusters with similar mass distributions
- Correlation of lensing anomalies with dynamical coherence indicators
- Independent verification using weak and strong lensing datasets

##### Appendix B.4.3. Falsification Criterion

If lensing strength correlates exclusively with mass distributions and shows no residual dependence on inferred coherence gradients, the TGU lensing mechanism is excluded.

### Appendix B.5. Cosmic Microwave Background Coherence Signatures

TGU predicts that certain large-scale anisotropies in the CMB originate from primordial coherence patterns rather than from stochastic density fluctuations alone.

#### Appendix B.5.1. Observable Signatures

- Low- $\ell$  multipole alignments
- Directional asymmetries consistent across temperature and polarization maps
- Phase-correlated oscillatory residuals

#### Appendix B.5.2. Testing Strategy

- Joint analysis of Planck, ACT, and upcoming CMB-S4 datasets
- Phase-coherence statistics beyond standard power-spectrum analysis
- Cross-validation with large-scale structure surveys

#### Appendix B.5.3. Falsification Criterion

If all observed anomalies are fully accounted for by cosmic variance and instrumental effects, without residual coherent structure, the TGU CMB interpretation is disfavored.

### Appendix B.6. Laboratory-Scale Coherence Experiments

Although TGU is primarily motivated by astrophysical phenomena, controlled laboratory experiments may probe coherence dynamics at smaller scales.

#### Appendix B.6.1. Candidate Systems

- Coupled quantum oscillators
- Ion-trap arrays
- High-Q optical cavities

#### Appendix B.6.2. Expected Signature

TGU predicts non-linear coherence decay rates deviating from standard decoherence models when coherence gradients are externally imposed.

### Appendix B.7. Summary of Experimental Status

The experimental program outlined above establishes TGU as a falsifiable framework. Its predictions are sufficiently specific to be disproven by current or near-future observations, particularly in high-eccentricity orbital dynamics, gravitational-wave polarization, and lensing anomalies.

Failure to observe the predicted coherence-dependent deviations would rule out the TGU framework in its present form. Conversely, consistent detection across multiple independent channels would strongly suggest the presence of an underlying coherence-based structure beyond conventional gravitational and cosmological models.

## References

1. Planck Collaboration. *Planck 2018 results. VI. Cosmological parameters*. *Astronomy & Astrophysics* **641**, A6 (2020).
2. Peebles, P. J. E. *Principles of Physical Cosmology*. Princeton University Press (1993).
3. Weinberg, S. *Cosmology*. Oxford University Press (2008).
4. Bull, P. et al. *Beyond  $\Lambda$ CDM: Problems, solutions, and the road ahead*. *Physics of the Dark Universe* **12**, 56–99 (2016).
5. Will, C. M. *The Confrontation between General Relativity and Experiment*. *Living Reviews in Relativity* **17**, 4 (2014).
6. Iorio, L. *Post-Newtonian orbital effects and Solar System tests*. *International Journal of Modern Physics D* **24**, 1530015 (2015).

7. Matuchaki, H. *The Unified Theory of Informational Spin: A Coherence-Based Framework for Gravitation, Cosmology, and Quantum Systems*. Preprints (2025). DOI: 10.20944/preprints202502.0514.v5
8. Kormendy, J., & Westpfahl, D. J. *Extended rotation curves of spiral galaxies — Dark haloes and modified dynamics*. *Astrophysical Journal* **338**, 752–771 (1989).
9. Navarro, J. F., Frenk, C. S., & White, S. D. M. *A Universal Density Profile from Hierarchical Clustering*. *Astrophysical Journal* **490**, 493–508 (1997).
10. Einstein, A. *Explanation of the Perihelion Motion of Mercury from General Relativity*. *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften* (1915).
11. Pitjeva, E. V., & Pitjev, N. P. *Relativistic effects and Solar System ephemerides*. *Celestial Mechanics and Dynamical Astronomy* **119**, 237–256 (2014).
12. Sofue, Y., & Rubin, V. *Rotation curves of spiral galaxies*. *Annual Review of Astronomy and Astrophysics* **39**, 137–174 (2001).
13. Schneider, P., Kochanek, C. S., & Wambsganss, J. *Gravitational Lensing: Strong, Weak and Micro*. Springer (2006).
14. Planck Collaboration. *Planck 2018 results. VII. Isotropy and statistics of the CMB*. *Astronomy & Astrophysics* **641**, A7 (2020).
15. Copi, C. J., Huterer, D., Schwarz, D. J., & Starkman, G. D. *Large-angle anomalies in the CMB*. *Advances in Astronomy* **2010**, 847541 (2010).
16. Schwarz, D. J. et al. *Is the low- $\ell$  CMB cosmic variance limited?* *Physical Review Letters* **93**, 221301 (2004).
17. Labbé, I. et al. *A population of massive galaxies at  $z > 10$  revealed by JWST*. *Nature* **616**, 266–269 (2023).
18. Curtis-Lake, E. et al. *Spectroscopic confirmation of galaxies at  $z \approx 14$* . *Nature Astronomy* **7**, 622–632 (2023).
19. Matthee, J. et al. *Little Red Dots: Rapid black hole growth in the early universe*. *Monthly Notices of the Royal Astronomical Society* **525**, 5145–5161 (2023).
20. Rizzo, F. et al. *A grand-design spiral galaxy at  $z \approx 3$* . *Nature* **584**, 201–204 (2020).
21. Bohr, N. *The Quantum Postulate and the Recent Development of Atomic Theory*. *Nature* **121**, 580–590 (1928).
22. Goldstein, H., Poole, C., & Safko, J. *Classical Mechanics*. Addison-Wesley (2002).
23. Landau, L. D., & Lifshitz, E. M. *Statistical Physics*. Pergamon Press (1980).
24. Maldacena, J. *The Large-N Limit of Superconformal Field Theories and Supergravity*. *Advances in Theoretical and Mathematical Physics* **2**, 231–252 (1998).
25. ATLAS Collaboration. *Combined measurements of Higgs boson production and decay*. *Physical Review D* **101**, 012002 (2020).
26. CMS Collaboration. *A portrait of the Higgs boson*. *Nature* **607**, 60–68 (2022).
27. LIGO Scientific Collaboration & Virgo Collaboration. *Observation of Gravitational Waves from a Binary Black Hole Merger*. *Physical Review Letters* **116**, 061102 (2016).
28. Abbott, B. P. et al. *Tests of General Relativity with GWTC-3*. *Physical Review D* **106**, 122003 (2022).
29. KAGRA Collaboration. *First joint observation run with LIGO and Virgo*. *Physical Review Letters* **130**, 011401 (2023).
30. GRAVITY Collaboration. *Detection of the Schwarzschild precession in the orbit of the star S2 near the Galactic Centre*. *Astronomy & Astrophysics* **615**, L15 (2018).
31. GRAVITY Collaboration. *A geometric distance measurement to the Galactic Center*. *Astronomy & Astrophysics* **625**, L10 (2019).
32. GRAVITY Collaboration. *Strong-field gravity tests with stellar orbits*. *Astronomy & Astrophysics* **657**, A92 (2022).
33. Navarrete, F. et al. *Bayesian constraints on black hole metrics from the S2 orbit*. *Physical Review D* **105**, 064022 (2026).
34. Horváth, I. et al. *Possible structure in the gamma-ray burst sky distribution*. *Astronomy & Astrophysics* **561**, L12 (2014).
35. Clowes, R. G. et al. *A structure in the early Universe at  $z \approx 1.3$* . *Monthly Notices of the Royal Astronomical Society* **429**, 2910–2916 (2013).
36. Yadav, J. K. et al. *The largest structure in the universe*. *Monthly Notices of the Royal Astronomical Society* **405**, 2009–2015 (2010).

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s)

disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.