

Article

Not peer-reviewed version

Management of *Liriomyza congesta* Infesting Broad Bean Plants Using Modern Techniques

[R.O.H Allam](#)^{*} and Fatma M Hussien

Posted Date: 11 June 2025

doi: 10.20944/preprints202505.2434.v3

Keywords: Insecticide; *L. congesta*; Silver nano particles; mineral oil



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Managment of *Liriomyza congesta* Infesting Broad Bean Plants Using Modern Techniques

Refat O. H. Allam ^{1,*} and Fatma M. Hussien ²

¹ Plant Protection Department, Faculty of Agriculture, South Valley University, 83523 Qena, Egypt

² Egyptian Sugar & Integrated Industries Company (E.S.I.I.C), Armant, Luxour, Egypt

* Correspondence: refat@agr.svu.edu.eg

Abstract: Evaluating the toxicity of specific pesticides and novel nanoparticle products was the aim of the study. On the *Liriomyza congesta* infesting broad bean crop the following compound namely; abamectin, *Bacillus Thuringiensis*, Metaflumizone, Novaluron, K Z oil, and *Nerium oleander* nanoparticles were made sustainably using aqueous extract and corresponding salt (silver nitrate) were applied. The results showed that Abamectin > *B. Thuringiensis* > Metaflumizone > Novaluron > *N. oleander* AgNps > K Z oil were the pesticides and extracts of silver nanoparticles that reduced the amount of *L. congesta* in broad bean plants. However, the widespread use of synthetic pesticides by growers has led to a number of problems, such as environmental contamination, the rise of pest resistance, and hazards to human health. According to those results, mineral oil and nanosilver extracts considerably decreased the mortality rate of *L. congesta* in comparison to traditional control, and they could be a financially viable alternative in next IPM campaigns to control this pest.

Keywords: insecticide; *L. congesta*; silver nano particles; mineral oil

Introduction

The main leguminous field crop in Egypt is the broad bean, *Vicia faba* L., which is vital to Egyptian cuisine and is eaten as dry mature seeds, fresh green pods, and immature fresh seeds. *Liriomyza congesta* (Beck), a broad bean leafminer, damages the crop's yield and quality (Deen 2024). The faba bean (*V. faba* L.), which is eaten as a green vegetable and as a dried and preserved seed, is a significant source of protein in Egyptian diets. Numerous insect pests target this extremely nutrient-dense crop, such as the leafminer *Liriomyza congesta*, which lowers output. (Abdel-Aliem et al. 2025). In terms of *L. trifolii*'s preferred host, it is evident that common bean plants had the highest mean numbers. (Hassan et al 2025). To determine the effectiveness of traditional insecticides against this pest, numerous authors conducted a number of tests. Furthermore, the advancement of our knowledge regarding the population dynamics of this pest depends on quantitative research. Poor research was done on the earlier bug species in southern Egypt's wide bean fields. As a result, the current study might assist provide some knowledge about the pest, which could help implement a successful control program to monitor the damage caused by pests in broad bean fields. Environmentally friendly synthesis outperforms chemical and physical processes in terms of cost, environmental friendliness, ease of scaling up for large-scale synthesis, and the absence of hazardous materials, high pressure, energy, or temperature. (Allam et al., 2022) investigated innovative, economical ways to control *P. hypostoma* instead of using synthetic pesticides and mineral oils. They also assessed green nanoparticles created using aqueous extract biosynthesised Ag⁺ nanoparticles and corresponding salt (silver nitrate). (Sebak et al., 2025) found that exposing *Culex pipiens* larvae to the LC₅₀ of essential oils of sandalwood, camphor, and black pepper significantly decreased the protein content.

Plant extracts are safe for the environment and have strong bioactivity against a variety of pests. The creation of plant extracts with strong toxic or repellent properties and their bioactivities against pests may aid in the creation of new pesticides that are safe for the environment (Guo et al., 2025).

Materials and Methods

Abamectin, *B. Thuringiensis*, Metaflumizone, Novaluron, *N. oleander* AgNps, and K Z oil were evaluated for their toxicity in broad bean plants throughout the 2023–2024 and 2024–2025 seasons.

Table 1. Compounds used in the study.

Compounds	Active ingredients	Rate
Abamectin	Abamet 1.8% EC	80cm ³ /fd
<i>Bacillus Thuringiensis</i>	Agree 50% WG	500 gm/fd
Metaflumizone	Alferde 24 % SC	200 cm ³ /fd
Novaluron	Aque 10% EC	80cm ³ /fd
<i>Nerium oleander</i> AgNps	oleandrin and oleandrigenin	200 cm ³ /fd
K Z oil	Mineral oil	200 cm ³ /fd

Tests of nanoparticles:

UV–test

Using a "Shimadzu UV-2401 PC, Japan" analytical spectrophotometer, the optical density (OD) of silver nanoparticles (AgNPs) containing *Nerium oleander* for UV–visible spectrum characterisation was determined at a resolution of 1 nm and a scanning speed of 300 nm/min within 200 and 800 nm. The baseline was modified as a blank in a volume containing 2 mM silver nitrate using the UV–vis spectra of 1 ml of the sample and 2 ml ionized as water in a quartz cell (Wiley *et al.*, 2006).

TEM-test:

Samples of the precipitate that gathered at the bottom of the conical flasks and the suspension above were collected for transmission electron microscopy (TEM) investigation following the reaction. The size and shape of extract nanoparticles were examined at 70 kV using a transmission electron microscope (TEM) called the "LEOL-2010, Japan" that was outfitted with a digital "Kodak Megaplug 1.6i camera" and image processing and analysis software (AMT, USA). The sample was pared by placing a drop of each solution on a copper grid covered in carbon and letting it dry at room temperature, as previously mentioned by Sathish Kumar *et al.* (2009). The resulting nanoparticles' size distribution was determined using TEM micrographs.

FTIR-test:

The FTIR spectra of AgNPs at room temperature were recorded in the 4000–400 cm⁻¹ region using a Perkin-Elmer spectrophotometer. We recorded the diffuse spectrum of reflectance in the 200–800 nm wavelength range using the UV140404B spectrophotometer. Numerical data was plotted using the 'Origin 7' software (Slman *et al.*, 2018).

XRD-test:

To get the AgNPs ready for the X-ray powder diffraction tests, the fluid containing the generated silver nanoparticles was spun for 30 minutes at 10,000 rpm. Following two rounds of washing with double-distilled water, the solid residues were dried at 80 degrees Celsius. Metal radiation (Cu Ka, 1.5406) at 40 kV and 30 mA was used to record the XRD powder patterns on a Shimadzu XRD-6000 (Slman *et al.*, 2018).

Results and Discussion

Tests of AgNPs:

This work provided a complete idea on the produce of nanoparticles of silver by fresh *Nerium oleander* . The water-soluble silver ions were converted to silver nanoparticles when added to the assessed plant extracts. In accordance with Hussein (2023), who stated that the brown color shift of the 1 mm silver nitrate solution will be the initial indication that nanoparticles are forming, the color of the solutions for *N. oleander* changed from yellow to brown 24 hours prior to the reaction. A UV–

vis spectrophotometer was used to track the development and stability of the reduced silver nanoparticles in the colloidal suspension. The longer the plant extract and silver nitrate were incubated, the higher the absorbance at 420 nm in the UV-vis spectra. (Figure 1).

FTIR analysis of silver nanoparticles confirmed the plant extract's dual functions as a capping and reducing agent as well as the presence of particular functional groups. Using TEM, it is now feasible to ascertain the size and shape of nanoparticles. According to the results, silver nanoparticles are mostly spherical and uniformly dispersed, but some of them have asymmetrical shapes, as seen in (Figure 1).




Figure 1. Tests on the production of silver nanoparticles by of *Nerium oleander* aqueous extract.

The relative toxicity of the poisonous action of Abamectin, *B. Thuringiensis*, Metaflumizone, Novaluron, *N. oleander* AgNps, and K Z oil against *L. congesta* was shown by the data in Table 2 and Figure 2.




Figure 2. Toxicity lines of pesticides and their alternatives against *L. congesta*.

The data clearly demonstrated that at both the LC₅₀ and LC₉₀ levels, the tested pesticides' order of efficiency was the same. Abamectin, *B. Thuringiensis*, Metaflumizone, Novaluron, *N. oleander* AgNps, and K Z oil are the tested pesticides that might be arranged in the following descending order. The associated LC₉₀ values were 1.87, 7.13, 39.49, 14.76, 348.09, and 1777.58 ppm, respectively, whereas the related LC₅₀ values were 0.17, 0.97, 2.05, 2.98, 11.59, and 44.80 ppm.

The results showed that the comparable compounds were 100, 17.39, 8.23, 5.67, 1.46, and 0.38 times as efficacious as abamectin at the LC₅₀ level. They did not overlap with those of Abamectin, *B. Thuringiensis*, *N. oleander* AgNps, or K Z oil, but they did overlap with Metaflumizone and Novaluron, as could be shown by comparing the confidence limits and how they overlapped with those of other substances. Consequently, we may say that although Metaflumizone and Novaluron are very different from other substances, they are different. (Figure 2). Furthermore, Table 2 and Figure 2 make it evident that K Z oil had the steepest toxicity line and Abamectin the flattest; however, *B. Thuringiensis*, Metaflumizone, Novaluron, and *N. oleander* AgNps were situated in the middle.

Table 2. Toxicity of certain insecticides against *L. congesta*.

Line name	LC ₅₀	Confidence limits of		Index	Slope ±	LC ₉₀
		LC ₅₀ , S *ppm				
		Lower	Upper			
Abamectin	0.17	0.11	0.39	100	1.23 ±0.22	1.87
<i>Bacillus Thuringiensis</i>	0.97	0.56	1.99	17.39	1.48 ±0.30	7.13
Metaflumizone	2.05	1.21	3.67	8.23	0.99 ±0.25	39.49
Novaluron	2.98	2.25	4.22	5.67	1.85 ±0.29	14.76
<i>Nerium oleander</i> AgNps	11.59	5.77	21.81	1.46	0.87 ±0.25	348.09
K Z oil	44.80	23.08	105.46	0.38	0.80 ±0.24	1777.58

T.I. - Index compared with Abamectin T.I. - compared with KZ. Oil * = ppm based on a.i.

This demonstrates the superiority of Abamectin over K.Z. oil. Abamectin was clearly the most harmful substance, while K.Z. oil was clearly the least hazardous. Table 2 show LC₅₀ value of *Bacillus Thuringiensis* was 0.97 ppm which shows very high toxicity to the leaf miner pest in the faba bean plants compared to other compounds used. These results were in line with a study conducted by (Abdel-Aliem et al 2025) examined the entomopathogenic fungicides BioMeta (*Metarhizium anisopliae*) and Biossiana (*Beauveria bassiana*) generated larval population reductions of 50.68 and 42.91%, respectively, while the standard insecticide abamectin reduced the larval population of *L. congesta* by 89.85%. Chen et al. (2015) that looked at how eleven different pesticides affected the subterranean termite, *Reticulitermes speratus*. The pesticide categories are arranged in decreasing order of termite toxicity: Organophosphate > Avermectin > Phenylpyrazole > Neonicotinoid > Pyrethroid > Carbamate. Researchers have also looked into farmer-friendly termite control techniques that are widely appealing, such as biological, physical, and cultural management (Abhishek et al., 2021). When used as part of an integrated pest management approach, entomopathogenic agents can produce better results. Furthermore, at different concentrations, the essential oils of clove leaves, neem, orange, and KZ were found to have insecticidal effects against the adult pests under study (Richa et al., 2021; Allam et al., 2022). It was found that the higher the concentration level, the higher the death rate.

Conclusion

For the safe and efficient management of *L. congesta* in the broad bean crop, it may be recommended to apply K Z mineral oil, *N. oleander* AgNps, and *Bacillus Thuringiensis* bacterium. The findings allow for the recommendation that, in addition to mineral oil, green synthesised silver nanoparticles be used into IPM programs.

Authors' Contributions: Every author has contributed to this study.

Funding: This research has no funding.

Institutional Review Board Statement: All Institutional Review Board Statement are confirmed and approved.

Data Availability Statement: Data presented in this study are available on fair request from the respective author.

Ethics Approval and Consent to Participate: This work was completed in accordance with all departmental guidelines at the Plant Protection Department.

Consent for Publication: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Abdel-Aliem H.I., Abo Elela M.R. and Abou-Elkasssem A.B. (2025) Role of Parasitoids, Entomopathogens and Chemical Compounds in Managing *Liriomyza congesta* Populations in Faba bean Fields. Arab Journal of Plant Protection, Volume 43, Issue 2.
2. Abhishek, R., Chandel, R.S., Verma, K.S., Manishkumar, J.J. (2021). 'Termites in Important Crops and their management.', Indian Journal of Entomology 83 Online published Ref. No. e20115.
3. Allam, R.O.H. 1; Badawy, A.M.M. and M.A. Ali. (2022). Green synthesized silver nanoparticles for controlling subterranean termites, *Psammotermes hypostoma* (Desneux). SVU-International Journal of Agricultural Sciences, 4 (1): 135-143.
4. Chen, Z., Qu, Y., Xiao, D., Song, L., Zhang, S., Gao, X., Song, D. (2015). 'Lethal and social-mediated effects of ten insecticides on the subterranean termite *Reticulitermes speratus*.' Journal of pest science, 88(4): pp. 741-751.
5. Deen M. B., Sajad A. K., Inayat U. L. (2024) First record of leaf miner, *Phyllocnistis unipunctella* (Stephens, 1834) (Lepidoptera, Gracillariidae) infesting *Populus deltoides* Marsh (Salicaceae) in Jammu and Kashmir, India. Entomon Archives / Vol. 49 No. 3
6. Guo, Y. B., Liu, Z. J., Ye, Y. Y., Chen, Y. Y., and Wu, Z. W. 2025. Antifeedant and contact toxicity activity of crude extracts from 10 plants to red imported fire ant, *Solenopsis invicta* Buren (Hymenoptera: Formicidae). Journal of Asia-Pacific Entomology, 28(1), 102350. <https://doi.org/10.1016/j.aspen.2024.102350>
7. Hassan A.T.1, El-Solimany E.A., Allam R.O.H. (2025). Population density of *liriomyza trifolii* (burgess) on four summer plant hosts in relation to some weather factors at Sohag governorate. J. Agric. & Env. Sci. 24(1): 163 – 181.
8. Henderson, C. F. and Tilton, E. W. (1955). Tests with acaricides against the brown wheat mite'. Journal of Economic Entomology, 48: 157-161.
9. Hussein A. R. A. (2023). Detection of the effect of silver nanoparticles prepared by using extract leaves *Prosopis juliflora* on pathogenic fungus *Trichophyton tonsurans* isolated from human scalp. GSC Biological and Pharmaceutical Sciences, 25(01): 245–251.
10. Richa, S., Kamin, A., Lal, E.P. (2021). 'Insecticidal Activities of Some Essential Oils on Subterranean Termites.', International Journal of Innovative Science and Research Technology, ISSN No:-2456-2165.
11. Sathishkumar, M., Sneha, K., Won, S., Cho, C. W., Kim, S. and Yun, Y. S. (2009). Cinnamon zeylanicum bark extract and powder mediated green synthesis of nano-crystalline silver particles and its bactericidal activity. Colloids and Surfaces B: Biointerfaces, 73: 332-338.
12. Sebak, S. I., Shaurub, E. S. H., Mahmoud, S. H., and Salem, S. A. 2025. Insights into biochemical, genotoxic, and molecular impacts, and molecular docking analysis-based possible mechanism of action of some essential oils against *Culex pipiens* larvae. Pesticide Biochemistry and Physiology, 209, 106352. <https://doi.org/10.1016/j.pestbp.2025.106352>
13. Slman, D. K., Jalill, R. D. A. and Abd, A. N. (2018). Biosynthesis of zinc oxide nanoparticles by hot aqueous extract of *Allium sativum* plants. Journal of Pharmaceutical Sciences and Research, 10: 1590-1596.
14. Wiley, B. J., Im, S. H., Li, Z. Y., McLellan, J., Siekkinen, A. and Xia, Y. (2006). Maneuvering the surface plasmon resonance of silver nanostructures through shape-controlled synthesis. ACS Publications.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.