

Short Note

Not peer-reviewed version

Non-Archimedean Massera-Schaffer-Maligranda-Pecaric-Rajic Inequality

[K. Mahesh Krishna](#) *

Posted Date: 15 January 2026

doi: 10.20944/preprints202601.0908.v2

Keywords: normed linear space; triangle inequality; non-Archimedean linear space; ultra-norm



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Short Note

Non-Archimedean Massera-Schaffer-Maligranda-Pecaric-Rajic Inequality

K. Mahesh Krishna

School of Mathematics and Natural Sciences, Chanakya University Global Campus, NH-648, Haralur Village, Devanahalli Taluk, Bengaluru North District, Karnataka State, 562 110, India; kmaheshak@gmail.com

Abstract

Massera and Schaffer [Ann. Math. (2), 1958] derived a breakthrough upper bound for the Clarkson angle between two nonzero vectors in a normed linear space, which was later improved by Maligranda [Am. Math. Mon., 2006]. Pecaric and Rajic [Math. Inequal. Appl., 2007] extended Maligranda's inequality to finitely many nonzero vectors. We derive a non-Archimedean version of Massera-Schaffer-Maligranda-Pecaric-Rajic inequality.

Keywords: normed linear space; triangle inequality; non-archimedean linear space; ultra-norm

MSC: 12J25; 46S10

1. Introduction

Let $\mathcal{X}$ be a normed linear space (NLS). Recall that for $x, y \in \mathcal{X} \setminus \{0\}$, the **Clarkson angle** [1] between x and y is defined as

$$\alpha[x, y] := \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\|.$$

Important Massera-Schaffer inequality [2] says that

$$\alpha[x, y] = \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\| \leq \frac{2\|x - y\|}{\max\{\|x\|, \|y\|\}}, \quad \forall x, y \in \mathcal{X} \setminus \{0\}. \tag{1}$$

Maligranda improved Inequality (1) in 2006 [3].

Theorem 1. [3,4] (*Massera-Schaffer-Maligranda Inequality*) Let $\mathcal{X}$ be a NLS. Then for all $x, y \in \mathcal{X} \setminus \{0\}$,

$$\alpha[x, y] = \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\| \leq \frac{\|x - y\| + \left| \|x\| - \|y\| \right|}{\max\{\|x\|, \|y\|\}} \leq \frac{2\|x - y\|}{\max\{\|x\|, \|y\|\}}. \tag{2}$$

We note that, in 1964 Dunkl and Williams [5] independently showed that

$$\left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\| \leq \frac{4\|x - y\|}{\|x\| + \|y\|}, \quad \forall x, y \in \mathcal{X} \setminus \{0\}. \tag{3}$$

Note that

$$\frac{2\|x - y\|}{\max\{\|x\|, \|y\|\}} \leq \frac{4\|x - y\|}{\|x\| + \|y\|}, \quad \forall x, y \in \mathcal{X} \setminus \{0\}.$$

Therefore Inequality (1) improves Inequality (3). Inequality (3) is famously known as Dunkl-Williams inequality in the literature. In 2007, Pecaric and Rajic extended Inequality (2) to finitely many nonzero elements [6].

Theorem 2. [6] (*Massera-Schaffer-Maligranda-Pecaric-Rajic Inequality*) Let $\mathcal{X}$ be a NLS and $n \in \mathbb{N}$. Then for all $x_1, \dots, x_n \in \mathcal{X} \setminus \{0\}$,

$$\left\| \sum_{j=1}^n \frac{x_j}{\|x_j\|} \right\| \leq \min_{1 \leq k \leq n} \left\{ \frac{1}{\|x_k\|} \left(\left\| \sum_{j=1}^n x_j \right\| + \sum_{j=1}^n \left| \|x_j\| - \|x_k\| \right| \right) \right\}.$$

It is important to ask what are non-Archimedean versions of Theorems 1 and 2? We answer the question by deriving non-Archimedean Massera-Schaffer-Maligranda-Pecaric-Rajic Inequality (Theorem 4).

2. Non-Archimedean Massera-Schaffer-Maligranda-Pecaric-Rajic Inequality

Let $\mathbb{K}$ be a field. A map $|\cdot| : \mathbb{K} \rightarrow [0, \infty)$ is said to be a non-Archimedean valuation if following conditions holds.

- (i) If $\lambda \in \mathbb{K}$ is such that $|\lambda| = 0$, then $\lambda = 0$.
- (ii) $|\lambda\mu| = |\lambda||\mu|$ for all $\lambda, \mu \in \mathbb{K}$.
- (iii) (Ultra-triangle inequality) $|\lambda + \mu| \leq \max\{|\lambda|, |\mu|\}$ for all $\lambda, \mu \in \mathbb{K}$.

In this case, $\mathbb{K}$ is called as non-Archimedean valued field [7]. Let $\mathcal{X}$ be a vector space over a non-Archimedean valued field $\mathbb{K}$ with valuation $|\cdot|$. A map $\|\cdot\| : \mathcal{X} \rightarrow [0, \infty)$ is said to be a non-Archimedean norm if following conditions holds.

- (i) If $x \in \mathcal{X}$ is such that $\|x\| = 0$, then $x = 0$.
- (ii) $\|\lambda x\| = |\lambda|\|x\|$ for all $\lambda \in \mathbb{K}$, for all $x \in \mathcal{X}$.
- (iii) (Ultra-norm inequality) $\|x + y\| \leq \max\{\|x\|, \|y\|\}$ for all $x, y \in \mathcal{X}$.

In this case, $\mathcal{X}$ is called as non-Archimedean linear space (NALS) [8]. Let $\mathcal{X}$ be a NALS. Let $x, y \in \mathcal{X} \setminus \{0\}$ with $\|x\|, \|y\| \in \mathcal{X}$. We define the **non-Archimedean Clarkson angle** between x and y as

$$\alpha[x, y] := \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\|.$$

Non-Archimedean version of Theorem 1 now reads as follows.

Theorem 3. (*Non-Archimedean Massera-Schaffer Inequality*) Let $\mathcal{X}$ be a NALS. Then for all $x, y \in \mathcal{X} \setminus \{0\}$ with $\|x\|, \|y\| \in \mathcal{X}$ it holds

$$\alpha[x, y] = \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\| \leq \frac{\max\{\|x - y\|, |\|x\| - \|y\||\}}{\max\{\|x\|, \|y\|\}}.$$

Proof. Let $x, y \in \mathcal{X} \setminus \{0\}$ with $\|x\|, \|y\| \in \mathcal{X}$. Then

$$\begin{aligned} \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\| &= \left\| \frac{x - y}{\|x\|} + \left(\frac{1}{\|x\|} - \frac{1}{\|y\|} \right) y \right\| \\ &\leq \max \left\{ \left\| \frac{x - y}{\|x\|} \right\|, \left\| \left(\frac{1}{\|x\|} - \frac{1}{\|y\|} \right) y \right\| \right\} \\ &= \max \left\{ \frac{\|x - y\|}{\|x\|}, \frac{|\|y\| - \|x\||}{\|x\|} \right\} \end{aligned}$$

and

$$\begin{aligned}\left\|\frac{x}{\|x\|} - \frac{y}{\|y\|}\right\| &= \left\|\left(\frac{1}{\|x\|} - \frac{1}{\|y\|}\right)x + \frac{x-y}{\|y\|}\right\| \\ &\leq \max\left\{\left\|\left(\frac{1}{\|x\|} - \frac{1}{\|y\|}\right)x\right\|, \left\|\frac{x-y}{\|y\|}\right\|\right\} \\ &= \max\left\{\frac{|\|y\| - \|x\||}{\|y\|}, \frac{\|x-y\|}{\|y\|}\right\}.\end{aligned}$$

Therefore

$$\left\|\frac{x}{\|x\|} - \frac{y}{\|y\|}\right\| \leq \frac{1}{\|x\|} \max\{\|x-y\|, |\|x\| - \|y\||\} \quad (4)$$

and

$$\left\|\frac{x}{\|x\|} - \frac{y}{\|y\|}\right\| \leq \frac{1}{\|y\|} \max\{\|x-y\|, |\|x\| - \|y\||\}. \quad (5)$$

Inequalities (4) and (5) give

$$\begin{aligned}\left\|\frac{x}{\|x\|} - \frac{y}{\|y\|}\right\| &\leq \left(\min\left\{\frac{1}{\|x\|}, \frac{1}{\|y\|}\right\}\right) \max\{\|x-y\|, |\|x\| - \|y\||\} \\ &= \frac{\max\{\|x-y\|, |\|x\| - \|y\||\}}{\max\{\|x\|, \|y\|\}}.\end{aligned}$$

□

Note the additional assumption $\|x\|, \|y\| \in \mathcal{X}$ in the previous theorem. This is necessary because, since the norm is a real number, we generally cannot guarantee that it belongs to the given non-Archimedean field. We now derive the non-Archimedean version of Theorem 2.

Theorem 4. (Non-Archimedean Massera-Schaffer-Maligranda-Pecaric-Rajic Inequality) Let $\mathcal{X}$ be a NALS and $n \in \mathbb{N}$. Then for all $x_1, \dots, x_n \in \mathcal{X} \setminus \{0\}$ with $\|x_1\|, \dots, \|x_n\| \in \mathcal{X}$ it holds

$$\left\|\sum_{j=1}^n \frac{x_j}{\|x_j\|}\right\| \leq \min_{1 \leq k \leq n} \left\{ \frac{1}{\|x_k\|} \max\left\{ \left\|\sum_{j=1}^n x_j\right\|, \max_{1 \leq j \leq n} |\|x_j\| - \|x_k\|| \right\} \right\}.$$

Proof. Let $x_1, \dots, x_n \in \mathcal{X} \setminus \{0\}$ with $\|x_1\|, \dots, \|x_n\| \in \mathcal{X}$. Let $1 \leq k \leq n$ be fixed. Then

$$\begin{aligned}
 \left\| \sum_{j=1}^n \frac{x_j}{\|x_j\|} \right\| &= \left\| \frac{x_k}{\|x_k\|} + \sum_{j=1, j \neq k}^n \frac{x_j}{\|x_j\|} \right\| \\
 &= \left\| \sum_{j=1}^n \frac{x_j}{\|x_k\|} - \sum_{j=1, j \neq k}^n \frac{x_j}{\|x_k\|} + \sum_{j=1, j \neq k}^n \frac{x_j}{\|x_j\|} \right\| \\
 &= \left\| \sum_{j=1}^n \frac{x_j}{\|x_k\|} - \sum_{j=1, j \neq k}^n \left(\frac{1}{\|x_k\|} - \frac{1}{\|x_j\|} \right) x_j \right\| \\
 &= \left\| \sum_{j=1}^n \frac{x_j}{\|x_k\|} - \sum_{j=1}^n \left(\frac{1}{\|x_k\|} - \frac{1}{\|x_j\|} \right) x_j \right\| \\
 &\leq \max \left\{ \left\| \sum_{j=1}^n \frac{x_j}{\|x_k\|} \right\|, \left\| \sum_{j=1}^n \left(\frac{1}{\|x_k\|} - \frac{1}{\|x_j\|} \right) x_j \right\| \right\} \\
 &= \max \left\{ \frac{1}{\|x_k\|} \left\| \sum_{j=1}^n x_j \right\|, \left\| \sum_{j=1}^n \left(\frac{1}{\|x_k\|} - \frac{1}{\|x_j\|} \right) x_j \right\| \right\} \\
 &\leq \max \left\{ \frac{1}{\|x_k\|} \left\| \sum_{j=1}^n x_j \right\|, \max_{1 \leq j \leq n} \left\| \left(\frac{1}{\|x_k\|} - \frac{1}{\|x_j\|} \right) x_j \right\| \right\} \\
 &= \frac{1}{\|x_k\|} \max \left\{ \left\| \sum_{j=1}^n x_j \right\|, \max_{1 \leq j \leq n} \left| \|x_j\| - \|x_k\| \right| \right\}.
 \end{aligned}$$

By varying k and taking minimum in the right side of previous inequality gives

$$\left\| \sum_{j=1}^n \frac{x_j}{\|x_j\|} \right\| \leq \min_{1 \leq k \leq n} \left\{ \frac{1}{\|x_k\|} \max \left\{ \left\| \sum_{j=1}^n x_j \right\|, \max_{1 \leq j \leq n} \left| \|x_j\| - \|x_k\| \right| \right\} \right\}.$$

□

3. Conclusions

- (1) In 1958, Massera and Schaffer derived a surprising upper bound for the Clarkson angle between two nonzero elements in a normed linear space [2].
- (2) In 2006, Maligranda improved Massera-Schaffer inequality [3].
- (3) In 2007, Pecaric and Rajic extended Maligranda inequality for finitely many nonzero elements [6].
- (4) In this article, we introduced the notion of non-Archimedean Clarkson angle and derived non-Archimedean version of Massera-Schaffer-Maligranda-Pecaric-Rajic inequality.

References

1. Clarkson, J.A. Uniformly convex spaces. *Trans. Am. Math. Soc.* **1936**, *40*, 396–414. <https://doi.org/10.2307/1989630>.
2. Massera, J.L.; Schäffer, J.J. Linear differential equations and functional analysis. I. *Ann. Math.* (2) **1958**, *67*, 517–573. <https://doi.org/10.2307/1969871>.
3. Maligranda, L. Simple norm inequalities. *Am. Math. Mon.* **2006**, *113*, 256–260. <https://doi.org/10.2307/27641893>.
4. Mercer, P.R. A refined Cauchy-Schwarz inequality. *International Journal of Mathematical Education in Science and Technology* **2007**, *38*, 839–843. <https://doi.org/10.1080/00207390701359370>.
5. Dunkl, C.F.; Williams, K.S. A simple norm inequality. *Am. Math. Mon.* **1964**, *71*, 53–54. <https://doi.org/10.2307/2311304>.
6. Pečarić, J.; Rajić, R. The Dunkl-Williams inequality with n elements in normed linear spaces. *Math. Inequal. Appl.* **2007**, *10*, 461–470. <https://doi.org/10.7153/mia-10-44>.

7. Schikhof, W.H. *Ultrametric calculus. An introduction to p -adic analysis*; Vol. 4, *Camb. Stud. Adv. Math.*, Cambridge: Cambridge University Press, 2006.
8. Perez-Garcia, C.; Schikhof, W.H. *Locally convex spaces over non-Archimedean valued fields*; Vol. 119, *Camb. Stud. Adv. Math.*, Cambridge: Cambridge University Press, 2010. <https://doi.org/10.1017/CBO9780511729959>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.