

Article

Not peer-reviewed version

# Bioactive Factors Isolated and Purified from Bovine Colostrum Can Restore Extracellular Matrix under Degradation by Metalloproteinases

[Federica Coppa](#) , [Graziella Giuffrida](#) , [Giulia Iannello](#) , [Stefania Pennisi](#) , [Greta Ferruggia](#) ,  
[Maria Violetta Brundo](#) \*

Posted Date: 20 August 2024

doi: 10.20944/preprints202408.1271.v1

Keywords: mechanical signals; wound healing; collagen; integrin; mechanotransducer



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Article*

# Bioactive Factors Isolated and Purified from Bovine Colostrum Can Restore Extracellular Matrix Under Degradation by Metalloproteinases

Federica Coppa, Graziella Giuffrida, Giulia Iannello, Stefania Pennisi, Greta Ferruggia <sup>1</sup> and Maria Violetta Brundo \*

Department of Biological, Geological and Environmental Sciences, University of Catania, Catania, Italy; federica.coppa@phd.unict.it (F.C.); graziella.giuffrida@phd.unict.it (G.G.); giulia.iannello@phd.unict.it (G.I.); stefania.pennisi@phd.unict.it (S.P.); greta.ferruggia@phd.unict.it (G.F.); mariavioletta.brundo@unict.it (M.V.B.)

\* Correspondence: mariavioletta.brundo@unict.it

**Abstract:** The ECM is composed of a considerable number of biochemically and structurally diverse constituents. ECM is a highly dynamic system that constantly receives and sends biological, chemical and mechanical signals. Several studies suggest that mechanical signals derived from the extracellular microenvironment regulate skin regeneration and wound healing. We evaluated effects of a mixture of biomolecules extracted and purified from bovine colostrum on restoring the mechanical properties of ECM. Test measuring collagen contraction showed a significant difference in contraction activation in samples treated with the 2% colostrum derivative mixture compared to the control. The analysis of the supernatant showed an inhibition of metalloproteinase-2 expression and an increase in collagen secretion by fibroblasts in treatment samples. Our hypothesis is that the molecules extracted and purified from bovine colostrum restore the ECM environment qualitative and quantitative characteristics, thus guaranteeing through a mechanical action the restoration of the wound due to the transduction of the signal activated by the integrins.

**Keywords:** mechanical signals; wound healing; collagen; integrin; mechanotransducer

## 1. Introduction

The cells within tissues are inserted in a highly structured microenvironment and are very sensitive to the geometric and mechanical constraints of the latter [1]. The cellular microenvironment, formed by the extracellular matrix (ECM) and neighboring cells, influences not only cellular architecture and mechanics, but also the polarity and functions of the cell [1,2]. Mechanical properties of the ECM depend on a complex protein network that forms the scaffolding to which cells adhere and acts as a reservoir for growth factors, cytokines and proteolytic enzymes [3]. ECM provides a fibrous 3D scaffold whose structural components are collagen fibers, proteoglycans, and glycosaminoglycans [4]. The diameter and organization of collagen fibers are adapted to the biomechanical function of each tissue [4]. The ECM also contains nonstructural components that modulate cell-ECM interactions. By moving under the influence of forces, ECM proteins can also act as mechanotransducers by exposing cryptic sites and growth factors [5].

The ECM is a dynamic structure, constantly undergoing a remodeling process where ECM components are deposited, degraded and modified [6]. ECM remodeling is an important mechanism whereby cell differentiation can be regulated wound repair. In physiological conditions, the balance between processes of destruction and regeneration of the constituents of the extracellular matrix is regulated by specific tissue inhibitors of metalloproteinases (MMPs) [7]. An alteration of this balance, as in the case of wounds, is related to a catalytic hyperactivity of MMPs that play a critical role in wound healing. Their main function is degradation with removal of the damaged ECM during the

inflammatory phase; breakdown of the basement membrane of the capillaries for angiogenesis and cell migration during the proliferation phase, and tissue contraction and remodeling in the remodeling phase. For effective wound healing, the presence of these enzymes is necessary, but they play a harmful role at high concentrations, causing excessive tissue degradation and slow wound healing [7].

Knowledge of the cellular response to mechanical stimuli coming from the cellular microenvironment can be fundamental for applications in regenerative medicine to de-sign new and more effective scaffolds or biomaterials [8].

Scaffolds should mimic the properties of the target tissue, ensure the restoring of anisotropy (one of the fundamental characteristics of most tissues) [9–11] and through mechanotransduction activate integrins and induce the cascade of growth factors (TGF-beta 1, CTGF, IGF-1, etc.) necessities, among other things, for the production of new collagen by the fibroblast [12]. The response of integrins to force involves three mechanochemical steps. Integrins must first bind to extracellular matrix molecules. Second, these forces must be transmitted into the cell where they are converted into biochemical signals (mechanotransduction). Finally, integrins connect to the cytoskeleton to transmit forces throughout the cell and strengthen adhesions to resist forces. Mechanical signals that are transmitted through structural components of the cytoskeleton play a key role in events that regulate cell migration, polarity, and proliferation [13].

The aim of our research is to demonstrate that bioactive factors purified from bovine colostrum are like molecules normally present in the extracellular matrix and that if supplied to stressed tissues (such as in chronic wounds), they can provide the scaffolding necessary to activate the signal transduction mechanisms that activate the response of integrins. Bovine colostrum is a nutrient milk secretion containing bioactive compounds that support calf nutrition and immune development. Colostrum is profuse in bioactive compounds like immunoglobulins, growth factors, lysozymes, lactoferrin, lactoperoxidase, but also possesses elevated levels of fats, proteins, minerals and vitamins. Bovine colostrum is also rich in extracellular nanovesicles, such as exosomes, which protect bioactive components from degradation [14]. These bioactive molecules are also components of the ECM, and their mechanical role is fundamental in its remodeling [15–17]. In order to demonstrate our hypothesis, we carried out also a test to evaluate collagen contraction and measured the expression of collagen production and metalloproteinases-2, crucial elements for wound healing [7,18,19].

## 2. Materials and Methods

### 2.1. Colostrum Derivative Mixture Preparation

The colostrum derivative mixture (CDM) preparation was processed according to the procedure described by [20]. Bovine colostrum was collected from Holstein cows from 1 up to 6 h after parturition. This methodology, through micro and nano filtration, allows the elimination of casein, fats and other non-functional macromolecules including bovine immunoglobulins from the colostrum, and essentially isolates growth factors and cytokines. In all experiments, we used 2% colostrum derivative mixture because, our unpublished data, we have seen that this corresponds to concentration growth factors and cytokines normally present in the extracellular matrix.

### 2.2. Cell Lines

The Human Dermal Fibroblasts (HDF) (106-05A, Merck) were grown in Dulbecco's Modified Eagle Medium (DMEM). Medium were supplemented with 10% fetal bovine serum. The medium was replaced every 2 days, starting when the cells reached 80% confluence. Cultures were maintained at 37°C in a humidified environment containing 5% CO<sub>2</sub>.

### 2.3. Collagen Contraction

To assess matrix contraction free-floating collagen lattice models were used according to [18]. The lattices were detached before addition of test substances. The reduction of the lattice area due to contraction was evaluated at 24 hours intervals for up to 96 hours.

#### 2.4. Collagen Production

Matrix synthesis was determined over 7 days in the presence or absence of colostrum derivative mixture 2%. Secreted C-terminal propeptide of collagen type I (CICP) was measured in culture supernatant using an enzyme immunoassay kit (Metra Biosystem, Quidel Corporation).

#### 2.5. Determination of MMP-2 Concentration

MMP-2 expression, constitutively secreted from fibroblasts, was determined with a commercially available quantitative ELISA test (Amersham Pharmacia Biotech). All procedures were performed according to the instructions of the manufacturer.


#### 2.6. Statistical Analysis

Data were analyzed using the GraphPad Prism 5.0 program (GraphPad Software, La Jolla, CA, USA), using the analysis of variance (ANOVA) test and Tukey's test. A  $p$  value  $\leq 0.05$  was used to identify statistically significant differences.


### 3. Results and Discussions

Tissues are not only made up of cells, but a significant part of their volume is formed by the extracellular space, occupied by an intricate network of macromolecules, whose three-dimensional organization is represented by the extracellular matrix. Biochemical analysis of the ECM reveals that it is composed of several proteins and polysaccharides, which aggregate in a compactly organized network connected to the surface of the cells that produced it and the surrounding ones [6,15,21]. ECM forms the bio-scaffold of our body and adapts to the environmental variables. The biomechanical properties of ECM are negatively affected by aging, but also during wound healing or restore skin in burning injuries [22]. Many computational models and experimental studies have revealed the important effects of cell-generated mechanical forces, forces acting upon cells, and mechanical characteristics of the extracellular matrix on cell morphology and function [22]. The microenvironment surrounding cells play a large role in directing cell behavior and the mechanical functions are important for both understanding cell behavior and excellent scaffolds designed. Cell can actively sense the mechanical properties of their surroundings by exerting contractile force, which can transmit between cell-matrix or cell-cell. Cells produce and can modify the organization of this ECM, which can vary widely in both composition and mechanical proprieties. Thus, these mechanical properties are a direct result of cellular activity, leading to the principle of dynamic reciprocity between the cell and its environment. But on cells can act mechanical forces exerted from ECM if as tissues are deformed, such as in wounds [23–25]. Our hypothesis is that the molecules extracted from colostrum restore chemical/physical ECM environment, thus guaranteeing through a mechanical action the restoration of the wound due to the transduction of the signal activated by the integrins. Our hypothesis is supported by the results obtained with collagen contraction test that demonstrated exposure of fibroblast-populated collagen lattices to concentrations of colostrum derivative mixture significantly has activated contraction by day 7, compared with the control (Figure 1). These results are of great interest especially if correlated to the simultaneous results of ELISA test made during collagen contraction. Tests demonstrated that fibroblasts actively have secreted CICP during lattice contraction up to 7 days (Figure 2) and inhibited MMP-2 expression (Figure 3) in the samples treated with the 2% colostrum derivative mixture. Contraction and remodeling of the extracellular matrix are essential processes during wound healing [26,27]. At the center of these two phenomena are fibroblasts, which not only produce and secrete extracellular matrix proteins but can also reorganize them through mechanical interactions [28,29]. Collagen is the predominant structural protein in the ECM providing not only tensile strength but also play a role in cells adhesion and


migration [30]. Matrix metalloproteinases (MMPs) are a family of proteolytic enzymes capable of degrading various structural components of the matrix [31–36]. They therefore play a fundamental role in various physiological and pathological processes including wound healing [16,27]. It is now known that the biomechanical properties of the ECM contribute to physiological development of cells and tissues [23].



**Figure 1.** Graphical representation of collagen contraction. A. Collagen gel contraction over 96 h at 24 h time intervals. B. Fold change in collagen gel contraction at 96 h. Asterisk denotes the degree of significance between results: \* $p < 0.01$ .



**Figure 2.** Graphical representation of collagen production during collagen contraction. Fibroblasts actively has secreted C1CP during lattice contraction up to 7 days. In control samples, secretion of C1CP has increased up to 3 days and after was reduced. Asterisk denotes the degree of significance between results: \*\* $p < 0.001$ .





**Figure 3.** Graphical representation of proenzyme MMP-2 production during collagen contraction. Asterisk denotes the degree of significance between results: \*\*\* $p < 0.0001$ .

The elasticity of the ECM, for example, helps determine how a cell feels and perceives external forces and, therefore, provides an important environmental input signal that determines the cell's response. Indeed, the focal adhesion complex, which consists of integrins and a multicomplex of adapters and signaling proteins, can be seen as a mechanosensor that connects the actomyosin cytoskeleton with the ECM [37]. Many of the focal adhesion components undergo conformational changes that determine functional consequences in the response to applied force. Together with the cytoskeleton and nuclear matrices, nuclear envelope, and chromatin, they constitute a sophisticated mechanosensing machinery that determines how cells react to forces transmitted by the ECM, which regulates various essential cellular behaviors, including cell fate determination, differentiation, and cell function [37–41]. The different properties of the ECM are not independent of each other but influence each other. Therefore, when the ECM increases in stiffness, as, for example, in pathological conditions, its biomechanical properties change, and cells respond by exerting markedly different types of force. Furthermore, the stiffening of the matrix also determines a change in the other physical properties of the ECM and, consequently, directly modifies the cellular ability to migrate. Second, the ECM is highly dynamic, constantly undergoing restructuring in different tissues at various embryonic and postnatal stages [41]. ECM dynamics can arise from changes in the absolute amount or composition of the ECM, for example as result of altered synthesis or degradation of one or more components of the ECM. Alternatively, ECM dynamics may show no changes in the composition of its components, but instead involve only the transformation of its components, into spatially organized structures, thanks to covalent and non-covalent bonds [6,21].

Finally, one of the most important features of cell-ECM interactions is reciprocity. On the one hand, cells constantly make, break down, or otherwise rearrange and realign components of the ECM to modify one or more properties; on the other hand, since the ECM regulates the behavior of cells, any change in it, as result of cellular activities, will in turn influence adjacent cells and modify their behaviors. This feedback regulatory mechanism between cells and the ECM allows cells and tissues to rapidly adapt to the environment.

#### 4. Conclusions

Wounds heal through coordinated action of fibroblast-mediated extracellular matrix (ECM) deposition, ECM remodeling, and wound contraction. For these processes, are fundamental the mechanical signals between ECM and cells. In this paper, we demonstrated that biomolecules extracted and purified of bovine colostrum can restore chemical/physical ECM environment, reestablish the mechanical properties of ECM and the anisotropy characteristics of the damaged tissue. This process activates transduction of the signal by the integrins and reactivates the ability of the fibroblast to synthesize new collagen, inducing the autologous mechanisms of repair and remodeling of damaged connective tissue. Fibroblasts, in fact, can be able to generate tensile forces as well as receive them; these contraction forces of the fibroblast are indispensable for wound resolution processes.

**Author Contributions:** Conceptualization, M.V.B.; methodology, M.V.B.; software, G.F.; formal analysis, F.C., G.G., G.I., and S.P.; investigation, F.C., G.G., G.I., and S.P.; data curation, M.V.B.; writing—original draft preparation, M.V.B.; writing—review and editing, G.F. and M.V.B.; supervision, M.V.B.; funding acquisition, M.V.B. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Acknowledgments:** G.F. thanks the Ph.D. program FSE Notice 1/2021. F.C., G.G., G.I. S.P. thanks the Ph.D. program PNRR—D.M. 117/2023.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. Mierke, C.T. Bidirectional Mechanical Response Between Cells and Their Microenvironment. *Front Phys* **2021**, *9*, 749830.
2. Xie, N.; Xiao, C.; Shu, Q.; Cheng, B.; Wang, Z.; Xue, R.; Wen, Z.; Wang, J.; Shi, H.; Fan, D.; Liu, N.; Xu, F. Cell response to mechanical microenvironment cues via Rho signaling: From mechanobiology to mechanomedicine. *Acta Biomater* **2023**, *159*, 1-20.
3. Boyd, D.F.; Thomas, P.G. Towards integrating extracellular matrix and immunological pathways. *Cytokine* **2017**, *98*, 79-86.
4. Jansen KA, Donato DM, Balcioglu HE, Schmidt T, Danen EH, Koenderink GH. A guide to mechanobiology: Where biology and physics meet. *Biochim Biophys Acta*. 2015 Nov;1853(11 Pt B):3043-52.
5. Di, X.; Gao, X.; Peng, L.; Ai, J.; Jin, X.; Qi, S.; Li H.; Wang K.; Luo D. Cellular mechanotransduction in health and diseases: from molecular mechanism to therapeutic targets. *Sig Transduct Target Ther* **2023**, *8*, 282.
6. Lu, P.; Takai, K.; Weaver, V.M.; Werb, Z. Extracellular matrix degradation and remodeling in development and disease. *Cold Spring Harb Perspect Biol* **2011**, *3*(12), a005058.
7. Caley, M.P.; Martins, V.L.; O'Toole, E.A. Metalloproteinases and Wound Healing. *Adv Wound Care* (New Rochelle) **2015**, *4*(4), 225-234.
8. Zhao, X.; Li, Q.; Guo, Z.; Li, Z. Constructing a cell microenvironment with biomaterial scaffolds for stem cell therapy. *Stem Cell Res Ther* **2021**, *12*(1), 583.
9. Caliar, S.R.; Harley, B.A. The effect of anisotropic collagen-GAG scaffolds and growth factor supplementation on tendon cell recruitment, alignment, and metabolic activity. *Biomaterials* **2011**, *32*(23), 5330-40.
10. Harmansa, S.; Erlich, A.; Eloy, C.; Zurlo, G.; Lecuit, T. Growth anisotropy of the extracellular matrix shapes a developing organ. *Nat Commun* **2023**, *14*, 1220.
11. Datta, P.; Vyas, V.; Dhara, S.; Chowdhury, A.; Barui, A. Anisotropy Properties of Tissues: A Basis for Fabrication of Biomimetic Anisotropic Scaffolds for Tissue Engineering. *J Bionic Eng* **2019**, *16*, 842-868.
12. Reid, J.A.; Dwyer, K.D.; Schmitt, P.R.; Soepriatna, A.H.; Coulombe, K.L.; Callanan, A. Architected fibrous scaffolds for engineering anisotropic tissues. *Biofabrication* **2021**, *13*(4), 10.1088/1758-5090/ac0fc9.
13. Ross, T.D.; Coon, B.G.; Yun, S.; Baeyens, N.; Tanaka, K.; Ouyang, M.; Schwartz, M.A. Integrins in mechanotransduction. *Curr Opin Cell Biol* **2013**, *25*(5), 613-8.
14. Ross, M.; Atalla, H.; Karrow, N.; Mallard, B.A. The bioactivity of colostrum and milk exosomes of high, average, and low immune responder cows on human intestinal epithelial cells. *J Dairy Sci* **2021**, *104*(3), 2499-2510.
15. Yue, B. Biology of the extracellular matrix: an overview. *J Glaucoma* **2014**, *23*(8 Suppl 1), S20-3.
16. Diller, R.B.; Tabor, A.J. The Role of the Extracellular Matrix (ECM) in Wound Healing: A Review. *Biomimetics* (Basel) **2022**, *7*(3), 87.
17. Kuehlmann, B.; Bonham, C.A.; Zucal, I.; Prantl, L.; Gurtner, G.C. Mechanotransduction in Wound Healing and Fibrosis. *J Clin Med* **2020**, *9*(5), 1423.
18. Daniels, J.T.; Cambrey, A.D.; Ocleston, N.L.; Garrett, Q.; Tarnuzzer, R.W.; Schultz, G.S.; Khaw, P.T. Matrix metalloproteinase inhibition modulates fibroblast-mediated matrix contraction and collagen production *in vitro*. *Invest Ophthalmol Vis Sci* **2003**, *44*(3), 1104-10.
19. Singh, D.; Rai, V.; Agrawal, D.K. Regulation of Collagen I and Collagen III in Tissue Injury and Regeneration. *Cardiol Cardiovasc Med* **2023**, *7*(1), 5-16.
20. Sacerdote, P.; Mussano, F.; Franchi, S.; Panerai, A.E.; Bussolati, G.; Carossa, S.; Bartorelli, A.; Bussolati, B. Biological components in a standardized derivative of bovine colostrum. *J Dairy Sci* **2013**, *96*, 1745-1754.
21. Hu, M.; Ling, Z.; Ren, X. Extracellular matrix dynamics: tracking in biological systems and their implications. *J Biol Eng* **2022**, *16*, 13.
22. Franchi, M.; Piperigkou, Z.; Mastronikolis, N.S.; Karamanos, N. Extracellular matrix biomechanical roles and adaptation in health and disease. *FEBS J* **2024**, *291*(3), 430-440.
23. Holle, A.W.; Young, J.L.; Van Vliet, K.J.; Kamm, R.D.; Discher, D.; Janmey, P.; Spatz, J.P.; Saif, T. Cell-Extracellular Matrix Mechanobiology: Forceful Tools and Emerging Needs for Basic and Translational Research. *Nano Lett* **2018**, *18*(1), 1-8.
24. Arnold, M.; Cavalcanti-Adam, E.A.; Glass, R.; Blümmel, J.; Eck, W.; Kantelechner, M.; Kessler, H.; Spatz, J.P. Activation of integrin function by nanopatterned adhesive interfaces. *Chemphyschem* **2004**, *5*(3), 383-8.
25. Xu, R.; Boudreau, A.; Bissell, M.J. Tissue architecture and function: dynamic reciprocity via extra- and intra-cellular matrices. *Cancer Metastasis Rev* **2009**, *28*(1-2), 167-76.
26. Olczyk, P.; Mencner, Ł.; Komosinska-Vassev, K. The role of the extracellular matrix components in cutaneous wound healing. *Biomed Res Int* **2014**, *2014*, 747584.
27. Potekaev, N.N.; Borzykh, O.B.; Medvedev, G.V.; Pushkin, D.V.; Petrova, M.M.; Petrov, A.V.; Dmitrenko, D.V.; Karpova, E.I.; Demina, O.M.; Shnayder, N.A. The Role of Extracellular Matrix in Skin Wound Healing. *J Clin Med* **2021**, *10*(24), 5947.

28. Tracy, L.E.; Minasian, R.A.; Caterson, E.J. Extracellular Matrix and Dermal Fibroblast Function in the Healing Wound. *Adv Wound Care* (New Rochelle) **2016**, 5(3), 119-136.
29. Kular, J.K.; Basu, S.; Sharma, R.I. The extracellular matrix: Structure, composition, age-related differences, tools for analysis and applications for tissue engineering. *J Tissue Eng* **2014**, 5, 2041731414557112.
30. Sun, B. The mechanics of fibrillar collagen extracellular matrix. *Cell Rep Phys Sci* **2021**, 2(8), 100515.
31. Gill, S.E.; Parks, W.C. Metalloproteinases and their inhibitors: regulators of wound healing. *Int J Biochem Cell Biol* **2008**, 40(6-7), 1334-47.
32. Lukes, A.; Mun-Bryce, S.; Lukes, M.; Rosenberg, G.A. Extracellular matrix degradation by metalloproteinases and central nervous system diseases. *Mol Neurobiol* **1999**, 19(3), 267-84.
33. Cabral-Pacheco, G.A.; Garza-Veloz, I.; Castruita-De la Rosa, C.; Ramirez-Acuña, J.M.; Perez-Romero, B.A.; Guerrero-Rodriguez, J.F.; Martinez-Avila, N.; Martinez-Fierro, M.L. The Roles of Matrix Metalloproteinases and Their Inhibitors in Human Diseases. *Int J Mol Sci* **2020**, 21(24), 9739.
34. Burbridge, M.F.; Cogé, F.; Galizzi, J.P.; Boutin, J.A.; West, D.C.; Tucker, G.C. The role of the matrix metalloproteinases during in vitro vessel formation. *Angiogenesis* **2002**, 5(3), 215-26.
35. Lombard, C.; Saulnier, J.; Wallach, J. Assays of matrix metalloproteinases (MMPs) activities: a review. *Biochimie* **2005**, 87(3-4), 265-72.
36. Agren, M.S.; Mirastschijski, U.; Karlsmark, T.; Saarialho-Kere, U.K. Topical synthetic inhibitor of matrix metalloproteinases delays epidermal regeneration of human wounds. *Exp Dermatol* **2001**, 10(5), 337-48.
37. Wong, V.W.; Akaishi, S.; Longaker, M.T.; Gurtner, G.C. Pushing back: wound mechanotransduction in repair and regeneration. *J Invest Dermatol* **2011**, 131(11), 2186-96.
38. DuFort, C.C.; Paszek, M.J.; Weaver, V.M. Balancing forces: architectural control of mechanotransduction. *Nat Rev Mol Cell Biol* **2011**, 12(5), 308-19.
39. Ricca, B.L.; Venugopalan, G.; Fletcher, D.A. To pull or be pulled: parsing the multiple modes of mechanotransduction. *Curr Opin Cell Biol* **2013**, 25(5), 558-64.
40. Saraswathibhatla, A.; Indana, D.; Chaudhuri, O. Cell-extracellular matrix mechanotransduction in 3D. *Nat Rev Mol Cell Biol* **2023**, 24(7), 495-516.
41. Xie, W.; Wei, X.; Kang, H.; Jiang, H.; Chu, Z.; Lin, Y.; Hou, Y.; Wei, Q. Static and Dynamic: Evolving Biomaterial Mechanical Properties to Control Cellular Mechanotransduction. *Adv Sci* (Weinh) **2023**, 10(9), e2204594.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.