

Short Note

Proving Maximal Linear Loose Tangle as a Linear Tangle

Takaaki Fujita

Independence; t171d603@gunma-u.ac.jp

Abstract: Linear-width is a well-regarded width parameter in graph theory. The constructs of linear loose tangle and linear tangle present obstacles to attaining linear-width. In this succinct paper, our primary focus will be the exploration of maximal linear loose tangles.

Keywords: maximal linear loose tangle; linear tangle; linear loose tangle; Linear tangle matroid

1. Short Introduction

The study of graph width parameters is pervasive across various fields, including matroid theory, lattice theory, computer science, game theory, network theory, artificial intelligence, graph theory, and discrete mathematics, as evidenced by a substantial body of literature (see [1–22,29–31] for references). These graph width parameters, often explored in relation to obstruction, have spawned a considerable amount of scholarly research.

One such graph width parameter is linear-width, which has also been extensively studied (see [1,3,6,11] for references). Linear tangle, initially introduced in [1], plays a crucial role in determining whether a linear width is at most k , where $k+1$ represents the order of the tangle. Likewise, linear loose tangle, which was first introduced in [4], also serves as an obstruction to linear-width at most k if its order is $k+1$.

In this concise paper, we focus on maximal linear loose tangle. While it may lack novelty, our objective is to contribute to the research on graph width parameters.

2. Definitions and Notations

In this section, we present the mathematical definitions and notations for each concept.

2.1. Symmetric Submodular Function

The definition of a symmetric submodular function is given below.

Definition 1. Let X be a finite set. A function $f: X \rightarrow \mathbb{N}$ is called symmetric submodular if it satisfies the following conditions:

- $\forall A \subseteq X, f(A) = f(X \setminus A).$
- $\forall A, B \subseteq X, f(A) + f(B) \geq f(A \cap B) + f(A \cup B).$

In this short paper, a pair (X, f) of a finite set X and a symmetric submodular function f is called a connectivity system.

It is known that a symmetric submodular function f satisfies the following properties:

Lemma 1 [18]. A symmetric submodular function f satisfies:

1. $\forall A \subseteq X, f(A) \geq f(\emptyset) = f(X).$
2. $\forall A, B \subseteq X, f(A) + f(B) \geq f(A \setminus B) + f(B \setminus A).$

In this paper, we use the notation f for a symmetric submodular function, a finite set X , and a natural number k . A set A is k -efficient if $f(A) \leq k$.

2.2. Linear tangle

The definition of a linear tangle on a connectivity system (X, f) is provided below. Please note that in reference [1], the order is denoted as k instead of $k + 1$. It is important to note that when a graph does not contain any pendant vertices, the equivalence still holds.

Definition 2 [1]. Let X represent a finite set and f denote a symmetric submodular function delineated over X , mapping to the non-negative integers. A linear tangle of order $k+1$ on a connectivity system (X, f) is a family L of k -efficient subsets of X , satisfying the following axioms:

- (L1) $\emptyset \in L$,
- (L2) For each k -efficient subset A of X , exactly one of A or $X \setminus A$ in L ,
- (L3) If $A, B \in L$, $e \in X$, and $f(\{e\}) \leq k$, then $A \cup B \cup \{e\} \neq X$ holds.

A linear tangle possesses the following characteristics:

Lemma 2. Let X represent a finite set and f denote a symmetric submodular function delineated over X , mapping to the non-negative integers. A linear tangle L of order $k+1$ on a connectivity system (X, f) satisfies following axioms:

- (L4) If $e \in X$, then $X \setminus \{e\} \notin L$

Proof. Assume, towards a contradiction, that there exists an element e in set X such that $X \setminus \{e\}$ belongs to L . Under this assumption, by the properties of symmetric submodular functions, we have $f(X \setminus \{e\}) = f(\{e\}) \leq k$. However, this condition conflicts with axiom (L3), leading us to a contradiction. This completes the proof.

2.3. Linear loose tangle

The definition of a linear loose tangle on a connectivity system (X, f) is given below.

Definition 3 [4]. Let X represent a finite set and f denote a symmetric submodular function delineated over X , mapping to the non-negative integers. Let T be a subset of 2^X that represents the order $k+1$ of linear loose tangle, and satisfies the following conditions:

- (IN) $\forall e \in X$, if $f(\{e\}) \leq k$, then $\{e\} \in T$.
- (LTSU) For any $A \in T$ and any element $e \in X$, if $B \subseteq A \cup \{e\}$ and $f(B) \leq k$ and $f(\{e\}) \leq k$, then $B \in T$.
- (IW) $X \notin T$.

A maximal linear loose tangle is one that cannot be extended further without violating its defining conditions.

3. Result of this paper: Proving Maximal Linear Loose Tangle as a Linear Tangle

We consider the case when the linear loose tangle on a connectivity system (X, f) is maximized. The lemmas\theorems and their corresponding proofs are presented below. Notably, Lemma 3 uses a proof that relies on the concept of maximality.

Lemma 3. Let X represent a finite set and f denote a symmetric submodular function delineated over X , mapping to the non-negative integers. If T is a Maximal linear loose tangle of order $k+1$ on (X, f) , T satisfies axiom (L2).

Proof of Lemma 3: Assume that T is a Maximal linear loose tangle of order $k+1$ on (X, f) . By definition, T is a subset of 2^X that represents order $k+1$ of a linear loose tangle, and satisfies conditions (IN), (LTSU), and (IW).

Consider a k -efficient subset A of X , that is $f(A) \leq k$. We aim to prove that exactly one of A or $X \setminus A$ is in T .

By axiom (LTSU), since A is k -efficient, A should be in T . For $X \setminus A$ to also be in T , there must exist an element e in $X \setminus A$ such that $f(\{e\}) \leq k$. However, because $A \cup X \setminus A = X$, and we know from axiom (IW) that $X \notin T$, we have a contradiction. In other words, the logical connection is that condition (LTSU) potentially allows us to include both A and $X \setminus A$ in T , but doing so would cause us to violate condition (IW), and hence we conclude that exactly one of A and $X \setminus A$ is in T .

Next, suppose that neither A nor $X \setminus A$ is in T . As T is a maximal linear loose tangle, we can add either A or $X \setminus A$ to T without violating any of the conditions (IN), (LTSU), and (IW). Therefore, by maximality, at least one of A or $X \setminus A$ must be in T .

Hence, exactly one of A or $X \setminus A$ is in T , proving that T satisfies axiom (L2). This concludes the proof.

Lemma 4. Let X represent a finite set and f denote a symmetric submodular function delineated over X , mapping to the non-negative integers. If T is a Maximal linear loose tangle of order $k+1$ on (X, f) , T satisfies axiom (L3).

Proof of Lemma 4. Assume that T is a maximal linear loose tangle of order $k+1$ on (X, f) , meaning that T satisfies conditions (IN), (LTSU), and (IW). We need to show that for any $A, B \in T$ and $e \in X$ with $f(\{e\}) \leq k$, we have $A \cup B \cup \{e\} \neq X$.

For the sake of contradiction, assume that there exist $A, B \in T$ and $e \in X$ such that $f(\{e\}) \leq k$ and $A \cup B \cup \{e\} = X$.

Now, let's consider a set $D = A \cup B$. If $f(D) > k$, D cannot be in T by definition as it is not k -efficient.

If $f(D) \leq k$, we need to consider the (LTSU) axiom carefully. If $D \in T$, then because $\{e\} \in T$ (due to $f(\{e\}) \leq k$ and axiom (IN)), we would have $D \cup \{e\} \in T$ by axiom (LTSU), as $D \cup \{e\}$ is a subset of $D \cup \{e\}$, and $f(D) \leq k$ and $f(\{e\}) \leq k$. But this would imply $X \in T$, contradicting axiom (IW) that states $X \notin T$. Hence, we conclude that our assumption that $A \cup B \cup \{e\} = X$ must be false. This concludes the proof.

Lemma 5. Let X represent a finite set and f denote a symmetric submodular function delineated over X , mapping to the non-negative integers. If T is a Maximal linear loose tangle of order $k+1$ on (X, f) , T satisfies axiom (L1).

Proof of Lemma 5. Assume that T is a maximal linear loose tangle of order $k+1$ on (X, f) , meaning that T satisfies conditions (IN), (LTSU), and (IW). By Lemma 1, for any $A \subseteq X$, $f(A) \geq f(\emptyset)$. Since \emptyset is a subset of every set, $f(\emptyset) \leq k$ holds, implying that \emptyset is k -efficient.

Now, condition (IN) of T states that for every $e \in X$, if $f(\{e\}) \leq k$, then $\{e\}$ is in T . Specifically, for the empty set, which we've shown is k -efficient, this implies that \emptyset should be in T .

Hence, T satisfies axiom (L1), and this concludes the proof.

From Lemmas 3, 4, and 5, we can derive the following theorem:

Theorem 6. Let X represent a finite set and f denote a symmetric submodular function delineated over X , mapping to the non-negative integers. If T is a Maximal linear loose tangle of order $k+1$ on (X, f) , T is linear tangle of order $k+1$ on (X, f) .

4. Future tasks: Linear tangle Matroids

In the future, I am contemplating researching the concept of "Linear Tangle Matroids." The concept of tangle matroid was defined within the framework of matroids (see [18,23–28]).

Here are definitions of tangle and tangle matroids. Please note that the connectivity function of a matroid M , denoted as $\lambda_M(A)$, for a subset A of $E(M)$, is defined as $\lambda_M(A) = r(A) + r(E(M) - A) - r(M)$.

Definition 4 [23]. Let M be a matroid, and T a collection of subsets of $E(M)$. Then T is a tangle of order $k+1$ of M if

- (TM1) for all $A \in T$, $\lambda_M(A) \leq k$;
- (TM2) for all $A \subseteq E(M)$ with $\lambda_M(A) \leq k$, either $A \in T$ or $E(M) - A \in T$;
- (TM3) if $A, B, C \in T$, then $A \cup B \cup C \neq E(M)$;
- (TM4) for each $e \in E(M)$, $E(M) - \{e\} \notin T$

We plan to investigate what characteristics emerge when the axiom (TM3) is modified to the following form, (LTM3):

- (LTM3) if $A, B \in T$, $e \in E(M)$, $\lambda_M(\{e\}) \leq k$, then $A \cup B \cup \{e\} \neq E(M)$.

A tangle that satisfies (LTM3) rather than (TM3) is referred to as a Linear Tangle of order $k+1$ of M .

Drawing inspiration from the ideas presented in references [24], we will contemplate a matroid M and a Linear tangle T of M of order $k+1$. Define a function $\rho: 2^{E(M)} \rightarrow N$ as follows: $\rho(A) := \min\{\lambda_M(B) : A \text{ is a subset of } B \text{ and } B \text{ is in } T\}$ if there exists a B such that A is a subset of B and B is in T . Otherwise, $\rho(A) := k+1$.

It has been shown in reference [18] that ρ is the rank function of a matroid on $E(M)$. Note that since it holds true in the case of a Tangle, it naturally extends to be applicable in the case of a Linear Tangle as well. If T is a Linear-tangle of a matroid M , we refer to $M(T)$ as a "Linear tangle matroid" of M . Going forward, We aim to carry out the characterization of the aforementioned Linear Tangle Matroids.

Acknowledgements: I humbly express my sincere gratitude to all those who have extended their invaluable support, enabling me to successfully accomplish this paper.

References

- [1] Daniel Bienstock. Graph searching, path-width, tree-width and related problems (a survey). Reliability of Computer and Communication Networks , Vol.DIMACS. Series in Discrete Mathematics and Theoretical Computer Science , pp. 33–50, 1989.
- [2] Fujita, Takaaki. "Reconsideration of Tangle and Ultrafilter using Separation and Partition." *arXiv preprint arXiv:2305.04306* (2023).
- [3] Fujita, Takaaki. "Revisiting Linear Width: Rethinking the Relationship Between Single Ideal and Linear Obstacle." *arXiv preprint arXiv:2305.04740* (2023).
- [4] Fujita, T. Relevance of Linear-decomposition and related concepts. Submitted. (2023).
- [5] Robertson, Neil, and Paul D. Seymour. "Graph minors. X. Obstructions to tree-decomposition." *Journal of Combinatorial Theory, Series B* 52.2 (1991): 153-190.
- [6] Fujita , T. (2023). Alternative Proof of Linear Tangle and Linear Obstacle: An Equivalence Result. *Asian Research Journal of Mathematics*, 19(8), 61–66.
- [7] Faure, Alexandre, and Fabien Feschet. "Linear decomposition of planar shapes." 2010 20th International Conference on Pattern Recognition. IEEE, 2010.
- [8] Fomin, Fedor V., and Dimitrios M. Thilikos. "On the monotonicity of games generated by symmetric submodular functions." *Discrete Applied Mathematics* 131.2 (2003): 323-335.
- [9] Vatschelle, Martin. "New width parameters of graphs." *Unpublished doctoral dissertation, The University of Bergen* (2012).
- [10] SOARES, Ronan Pardo. *Pursuit-evasion, decompositions and convexity on graphs*. 2013. PhD Thesis. Université Nice Sophia Antipolis.

- [11] Kobayashi, Yasuaki, and Yu Nakahata. "A Note on Exponential-Time Algorithms for Linearwidth." arXiv preprint arXiv:2010.02388 (2020).
- [12] Oum, Sang-il, and Paul Seymour. "Testing branch-width." *Journal of Combinatorial Theory, Series B* 97.3 (2007): 385-393.
- [13] Hicks, Illya V., et al. "Branch-width and tangles." *Wiley Encyclopedia of Operations Research and Management Science* (2010).
- [14] Bergougnoux, Benjamin, Jan Dreier, and Lars Jaffke. "A logic-based algorithmic meta-theorem for mim-width." *Proceedings of the 2023 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*. Society for Industrial and Applied Mathematics, 2023.
- [15] Bergougnoux, Benjamin, Tuukka Korhonen, and Igor Razgon. "New Width Parameters for Independent Set: One-sided-mim-width and Neighbor-depth." *arXiv preprint arXiv:2302.10643* (2023).
- [16] Munaro, Andrea, and Shizhou Yang. "On algorithmic applications of sim-width and mim-width of (H1, H2)-free graphs." *Theoretical Computer Science* 955 (2023): 113825.
- [17] Thilikos, Dimitrios M., and Sebastian Wiederrecht. "Approximating branchwidth on parametric extensions of planarity." *arXiv preprint arXiv:2304.04517* (2023).
- [18] Geelen, J., Gerards, B., Robertson, N. and Whittle, G. (2006) Obstructions to Branch-Decomposition of Matroids. *Journal of Combinatorial Theory, Series B*, 96, 560-570.
- [19] Hicks, Illya V., and Boris Brimkov. "Tangle bases: Revisited." *Networks* 77.1 (2021): 161-172.
- [20] Geelen, Jim, and Stefan HM van Zwam. "Matroid 3-connectivity and branch width." arXiv preprint arXiv:1107.3914 (2011).
- [21] Thilikos, D.M. (2000) Algorithms and Obstructions for Linear-Width and Related Search Parameters. *Discrete Applied Mathematics*, 105, 239-271.
- [22] Fujita, Takaaki. Filter for Submodular Partition Function: Connection to Loose Tangle. Submitted. (2023).
- [23] Geelen, Jim, and Stefan HM van Zwam. "Matroid 3-connectivity and branch width." *Journal of Combinatorial Theory, Series B* 112 (2015): 104-123.
- [24] Hall, Dennis. "A characterization of tangle matroids." *Annals of Combinatorics* 19 (2015): 125-130.
- [25] van Zwam, S. (2011). Preserving 3-connectivity in matroids of high branch width.
- [26] van Zwam, Stefan. "When the branch width is high..." (2011).
- [27] Geelen, Jim, Bert Gerards, and Geoff Whittle. "Tangles, tree-decompositions and grids in matroids." *Journal of Combinatorial Theory, Series B* 99.4 (2009): 657-667.
- [28] Hall II, Dennis Wayne. *On Matroid and Polymatroid Connectivity*. Louisiana State University and Agricultural & Mechanical College, 2014.
- [29] Oum, Sang-il, and Paul Seymour. "Certifying large branch-width." *Symposium on Discrete Algorithms: Proceedings of the seventeenth annual ACM-SIAM symposium on Discrete algorithm*. Vol. 22. No. 26. 2006.
- [30] Grigoriev, Alexander. "Tree-width and large grid minors in planar graphs." *Discrete Mathematics & Theoretical Computer Science* 13 (2011).
- [31] Müller, Theodor. *The excluded minor structure theorem, and linkages in large graphs of bounded tree-width*. Diss. Staats-und Universitätsbibliothek Hamburg Carl von Ossietzky, 2014.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.