

Article

Not peer-reviewed version

In Vitro Evaluation of the Antimicrobial Activity of *Origanum vulgare* Essential Oil Against Esbl-Producing Strains of *Escherichia coli* and *Klebsiella pneumoniae*

[Andrea González](#)*, Mónica Espadero, [Inés Malo](#), [Ricardo Alejandro](#)

Posted Date: 2 March 2026

doi: 10.20944/preprints202603.0101.v1

Keywords: essential oil; antimicrobial activity; ESBL; *Klebsiella pneumoniae*; *Escherichia coli*; cefepime; synergism



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

In vitro Evaluation of the Antimicrobial Activity of *Origanum vulgare* Essential Oil Against ESBL-Producing Strains of *Escherichia coli* and *Klebsiella pneumoniae*

Andrea González ^{1*}, Mónica Espadero ¹, Inés Malo ¹ and Ricardo Alejandro ²

¹ Biotechnology degree, Universidad Politécnica Salesiana, Cuenca 010102, Ecuador

² National Institute of Public Health Research (INSPI), Center of reference antimicrobial resistance (CR-RAM), Cuenca 010107, Ecuador

* Correspondence: angy2003gc@outlook.com

Abstract

Background/Objectives: One of the greatest threats to global public health is antimicrobial resistance (AMR), due to the increasing number of infections caused by extended-spectrum β -lactamase (ESBL)-producing Enterobacteriaceae. *Escherichia coli* and *Klebsiella pneumoniae* ESBL-producing bacteria possess resistance mechanisms that inactivate β -lactam antibiotics by hydrolyzing their β -lactam ring, thereby limiting conventional therapeutic options. In response to this problem, the objective in this exploratory *in vitro* study was to evaluate the antimicrobial activity of *Origanum vulgare* (oregano) essential oil and its interaction with the antibiotic cefepime using *in vitro* methods. **Methods:** Antimicrobial susceptibility tests were performed, including determination of the minimum inhibitory concentration by the microdilution method with statistical analysis, and evaluation of the fractional inhibitory concentration index using the checkerboard method. In addition, advanced methods such as bacterial identification by MALDI-TOF mass spectrometry and PCR were employed for the identification of resistance genes. **Results:** The studied strains exhibited both phenotypic and genotypic resistance. The MIC of the essential oil was 1024 $\mu\text{g/mL}$ for ESBL-producing *E. coli* and 2048 $\mu\text{g/mL}$ for ESBL-producing *K. pneumoniae*, whereas the ATCC strains showed higher susceptibility. The FICI values indicated synergism in *E. coli* (FICI = 0.188) and an additive effect in *K. pneumoniae* (FICI = 0.563). **Conclusions:** Oregano essential oil exhibits antimicrobial activity and the ability to potentiate the effect of cefepime, suggesting its potential as a therapeutic adjuvant. Additional studies are required, including a larger number of strains, cytotoxicity analyses, and clinical validation.

Keywords: essential oil; antimicrobial activity; ESBL; *Klebsiella pneumoniae*; *Escherichia coli*; cefepime; synergism

1. Introduction

Antimicrobial resistance (AMR) is one of the main challenges for public health because it compromises the effectiveness of antibiotic treatments [1], particularly due to the increasing prevalence of extended-spectrum β -lactamase (ESBL)-producing Enterobacteriaceae, which inactivate β -lactam antibiotics such as penicillins and cephalosporins. These β -lactamase enzymes are encoded by acquired genes or variants such as: *bla*_{CTX-M}, *bla*_{TEM} and *bla*_{SHV}, which confer resistance to β -lactam antibiotics [2], thereby limiting therapeutic options and increasing morbidity and mortality associated with infections. The World Health Organization (WHO) classifies ESBL-producing bacteria as priority pathogens for the development of new treatments, and reports indicate

that 60% of nosocomial infections in Latin America are caused by β -lactam-resistant Enterobacteriaceae [3].

Escherichia coli is an opportunistic bacterium associated with cases of cystitis, sepsis, and pneumonia due to the acquisition of resistance genes through mobile genetic elements (plasmids) [4,5]. *Klebsiella pneumoniae* also acquires ESBL production and is a nosocomial pathogen associated with approximately 50% of bacteremia cases, as well as a high capacity for resistance to multiple antibiotics [6].

Medicinal plants have been valued for their therapeutic properties, playing a fundamental role in traditional medicine. This practice has been integrated into modern medicine through the isolation and characterization of phytochemicals or bioactive compounds such as terpenoids, phenolic compounds, glycosides, and alkaloids, which exhibit anti-inflammatory, antimicrobial, and antifungal effects [7,8].

Among the methods used for the extraction of bioactive compounds are non-conventional techniques such as ultrasound-assisted extraction and supercritical fluid extraction, as well as conventional methods including maceration, percolation, decoction, solvent extraction, and steam distillation [9], the latter is particularly useful because it yields essential oils containing bioactive components with therapeutic potential [10].

The essential oil of *Origanum vulgare* (oregano), belonging to the Lamiaceae family, is rich in phenolic monoterpenes and phenylpropanoids, with carvacrol and thymol being the main compounds responsible for its antimicrobial activity [11]. Studies by Nurzyńska demonstrate that essential oils exert inhibitory effects on the growth of both Gram-positive and Gram-negative bacteria, with particular emphasis on *Escherichia coli* and *Klebsiella pneumoniae*, due to the presence of metabolites in oregano essential oil that act through multiple mechanisms, including destabilization of the outer membrane, suppression of resistance gene expression, and inhibition of protein synthesis. In light of this emergency, oregano essential oil has been considered a natural adjuvant [12].

Carvacrol has demonstrated synergistic effects with β -lactam antibiotics, reversing resistance to cephalosporins in resistant *E. coli* strains [13]. Si et al. reported minimum inhibitory concentrations (MIC) of *Origanum vulgare* essential oil of 0.5 $\mu\text{g/mL}$ and 0.8 $\mu\text{g/mL}$ against ESBL-producing *E. coli* strains, in addition to a synergistic effect with fluoroquinolones or doxycycline, which reduced the standard dose of these antibiotics by up to eightfold [14].

2. Materials and Methods

2.1. Bacterial Strains

Four bacterial strains were used, provided by the National Institute of Public Health Research (INSPI) due to their epidemiological relevance regarding resistance mechanisms: *E. coli* ATCC 25922 and *K. pneumoniae* ATCC 700603, employed as quality control strains, and two ESBL-producing bacterial isolates: *E. coli* (ID 323) and *K. pneumoniae* (ID 132). The strains were preserved at $-20\text{ }^{\circ}\text{C}$ in cryovials with glycerol and reactivated in BHI. For each assay, fresh cultures ($\leq 24\text{ h}$) were used to ensure viability.

2.2. Identification by MALDI-TOF MS

Taxonomic identification of the study strains was performed by MALDI-TOF mass spectrometry at the MEDICULT laboratory (Cuenca, Ecuador). Fresh colonies (18 h) were taken from blood agar plates and placed onto a polished steel target plate for identification using the Microflex LT system (Bruker Daltonics, Bremen, Germany). The generated protein spectrum was compared with a reference database, providing score values, where scores ≥ 2.0 indicated reliable identification at the genus and species levels.

2.3. Susceptibility Analysis for Phenotypic Detection

Phenotypic detection of ESBL production was carried out using the double-disk synergy test (Jarlier et al., 1988), employing the antibiotics amoxicillin/clavulanic acid (20/10 µg), aztreonam (30 µg), cefotaxime (30 µg), ceftazidime (30 µg), and cefepime (30 µg).

Mueller–Hinton agar plates were inoculated with a bacterial suspension adjusted to the 0.5 McFarland standard. The disks were placed at a distance of 25 mm and incubated at 37 °C for 18–24 h. The appearance of deformation or enlargement of the inhibition zone toward the amoxicillin/clavulanic acid disk was considered indicative of ESBL production. Inhibition zone diameters were measured and interpreted according to CLSI breakpoints (2025).

2.4. Molecular Identification by PCR

Five to six fresh colonies from each strain were collected and placed in 1 mL of TAE buffer, then incubated in a Thermomixer (Senacyt) at 95 °C with constant agitation for 10 minutes, followed by centrifugation to mechanically extract the genetic material. PCR reactions were prepared in a final volume of 24 µL containing 10 µL of molecular-grade water, 12 µL of GoTaq® Master Mix, 0.5 µL of forward primer, 0.5 µL of reverse primer, and 1 µL of DNA. Amplification was performed for two hours over 36 cycles in a Bio-Rad thermocycler. The samples were loaded onto a 1% agarose gel prepared with 1% TAE buffer and SafeView Classic® as the staining agent, and electrophoresis was carried out at 80 V for 40 minutes. After electrophoresis, the gel was visualized using a Bio-Rad gel documentation system for gene detection.

Table 1. PCR conditions for DNA amplification.

Gene	Primer sequence		Size	Denaturation	Hybridization	Extension	Final extension
bla _{SHV}	SHV-F	ATGCGTTATATTCGCCTGTG	449	94° for 30 seconds	52.8° for 40 seconds	72° for 50 seconds	72° for 5 minutes
	SHV-R	TGCTTTGTTATTCGGGCCAA					
bla _{TEM}	TEM-F	AAACGCTGCTGAAAGTA	516	94° for 30 seconds	52.8° for 40 seconds	72° for 50 seconds	72° for 5 minutes
	TEM-R	AGCGATCTGTCTAT					
bla _{CTX-M}	CTX-M-F	TTTGCGATGTGCAGTACCAGTA	172	94° for 30 seconds	52.8° for 40 seconds	72° for 50 seconds	72° for 5 minutes
	CTXM-R	CGATATCGTTGGTGGTGCCATA					

2.5. *Origanum vulgare* Essential Oil

The essential oil was acquired from the company KHIWA NATURAL (Quito, Ecuador), obtained by steam distillation, and accompanied by a certificate reporting the purity of the essential oil, density (0.935 g/mL), and the quantified content of secondary metabolites detailed in Table 2.

Table 2. Secondary metabolites of *Origanum vulgare* (oregano) essential oil.

Bioactives	Percentage
α-pinene	1.03%
α-thuyene	1.28%
β-myrcene	1.82%
α-terpinene	1.25%
δ-terpinene	6.8%
q-cymene	10.63%
Linalool	2.1%
β-caryophyllene	2.01%
Thymol	5.44%
Carvacrol	86.74%

Due to its hydrophobic nature, the essential oil was solubilized using 1% DMSO at a 1:1 ratio.

2.6. *Determination of the Minimum Inhibitory Concentration (MIC)*

The MIC of oregano essential oil and the antibiotic cefepime was determined using the broth microdilution method in 96-well plates, in accordance with CLSI guidelines (2025).

Ten serial dilutions of the essential oil were prepared with a bacterial inoculum of 5 × 10⁵ CFU/mL in Mueller–Hinton broth. Plates were incubated at 37 °C for 24 h, and absorbance was measured using an mcr® microplate reader at an OD of 630 nm. The MIC was defined as the lowest concentration with no visible growth. Each assay was performed in triplicate to ensure data consistency, including a DMSO control with no significant differences. For statistical analysis, data normality was initially assessed using the Shapiro–Wilk test. Since the data did not follow a normal distribution and corresponded to ordered concentrations in a dose–response design with a small sample size (n = 3), the non-parametric Jonckheere–Terpstra test was applied to evaluate the monotonic trend of the inhibitory effect of the essential oil as a function of concentration. Given the discrete dilution design and limited number of replicates, regression-based approaches were not applied.

The MIC was reported in an exploratory manner as the concentration associated with maximum inhibition (minimum median response), considering this criterion conservative and consistent with the microbiological interpretation of the assay, based on experimentally observed inhibition patterns rather than exclusively on statistical criteria. Statistical analyses were performed using RStudio software.

2.7. *Determination of the Fractional Inhibitory Concentration*

A 6 × 6 well matrix was prepared, in which the antibiotic was placed in serial horizontal dilutions across columns, and the essential oil was placed in serial vertical dilutions across rows. For this method, it is necessary to start from the MIC observed for each strain. Interpretation was performed according to the following criteria: FICI ≤ 0.5, synergy; 0.5 < FICI ≤ 1.0, additive effect; 1.0 < FICI ≤ 4.0, indifference; FICI > 4.0, antagonism. The interaction between the essential oil and cefepime was evaluated by calculating the FICI using the following formula:

$$FICI = \frac{MIC\ of\ the\ oil\ in\ combination}{MIC\ of\ the\ oil\ alone} + \frac{MIC\ of\ antibiotic\ in\ combination}{MIC\ of\ antibiótico\ alone}$$

Equation 1. Calculation of the FICI

3. Results

3.1. Susceptibility Testing

The study strains showed resistance to all evaluated β -lactam antibiotics, confirming their ESBL phenotype. The ATCC strains behaved according to the QC values established by CLSI (2025) and were identified by MALDI-TOF mass spectrometry. The MALDI-TOF results are detailed in Table 1, the inhibition zone diameters in Table 2, and the genotypes identified by PCR in Table 3.




Figure 1. Protein spectra of the study strains (a) *Escherichia coli* (b) *Klebsiella pneumoniae*.

Table 3. Identification of bacterial strains.

Sample ID	Score	Corresponding genus and species
323	2.02	<i>Escherichia coli</i>
132	2.03	<i>Klebsiella pneumoniae</i>




Figure 2. Mueller–Hinton agar plates: (a) antibiogram of the *E. coli* study strain; (b) antibiogram of the *E. coli* ATCC 25922 control strain; (c) antibiogram of the *K. pneumoniae* study strain; (d) antibiogram of the *K. pneumoniae* ATCC 700603 control strain.

Table 4. Results and interpretation of inhibition zone diameters in the antibiograms.

Antibiotic	<i>E. coli</i> 323	<i>E. coli</i> ATCC 25922	<i>K. pneumoniae</i> 132	<i>K. pneumoniae</i> ATCC 700603
AMC	6 mm (R)	30 mm (S)	13 mm (R)	16 mm (I)
ATM	10 mm (R)	33 mm (S)	13 mm (R)	14 mm (R)
FEP	18 mm (R)	37 mm (S)	17mm (R)	30 mm (S)
CTX	10 mm (R)	34 mm (S)	17 mm (R)	23 mm (I)
CAZ	6 mm (R)	33 mm (S)	14 mm (R)	17 mm (R)

Note: amoxicillin/clavulanic acid (AMC), aztreonam (ATM), cefepime (FEP), cefotaxime (CTX), ceftazidime (CAZ), R (resistant), I (intermediate), S (susceptible). Interpretation was performed according to CLSI (2025).




Figure 3. Agarose gel electrophoresis evaluating, in order: lanes 1 to 6, *K. pneumoniae* study strain, *E. coli* study strain, *K. pneumoniae* ATCC, *E. coli* ATCC, positive control, and negative PCR control for the *bla*_SHV; lanes 8 to 13, *K. pneumoniae* study strain, *E. coli* study strain, *K. pneumoniae* ATCC, *E. coli* ATCC, positive control, and negative PCR control for the *bla*_TEM; finally, lanes 14 to 19, *K. pneumoniae* study strain, *E. coli* study strain, *K. pneumoniae* ATCC, *E. coli* ATCC, positive control, and negative PCR control for the *bla*_CTX-M. Lanes 7 and 20 corresponded to the molecular weight marker.

Table 5. PCR results and interpretation.

Bacterial strains	<i>bla</i> _SHV	<i>bla</i> _TEM	<i>bla</i> _CTX-M
<i>K. pneumoniae</i> study strain	Positive	Positive	Positive
<i>E. coli</i> study strain	Positive	Positive	Positive
<i>K. pneumoniae</i> ATCC 700603	Positive	Negative	Negative
<i>E. coli</i> ATCC 25922	Negative	Negative	Negative

3.2. Determination of the Minimum Inhibitory Concentration

All figures and tables should be cited in the main text as Figure 1, Table 1, etc.
Analysis for MIC determination was performed using the broth microdilution method. Cefepime was used as the antibiotic, and ten concentrations were tested to determine the MIC.

3.2.1. Experimental MIC Results of Oregano Essential Oil and Cefepime.

The essential oil showed an inhibitory effect against all strains, with greater potency against *E. coli* ATCC 25922 (256 µg/mL) and lower potency against the *K. pneumoniae* study strain (2048 µg/mL). These values are attributed to differences in permeability and efflux pumps. The results are shown in Table 6.

Table 6. MIC results.

Bacteria	<i>E. coli</i>	<i>E. coli</i> ATCC 25922	<i>K. pneumoniae</i>	<i>K. pneumoniae</i> ATCC 700603
Concentration µg/mL	1024	256	2048	1024
OEO				
Concentration µg/mL	64	0.12	64	1
Cefepime				

3.2.2. Statistical Analysis of the MIC of *Origanum vulgare* Essential Oil

The minimum inhibitory concentration (MIC) values of *Origanum vulgare* essential oil against the *E. coli* and *K. pneumoniae* study strains and ATCC control strains are shown in Table 7. The Shapiro–Wilk test indicated that the data did not follow a normal distribution ($p < 0.05$). Considering the small sample size ($n = 3$) and the natural ordering of the evaluated concentrations, the non-parametric Jonckheere–Terpstra test was applied to evaluate the dose–response trend.

Table 7. Results of the Shapiro–Wilk and Jonckheere–Terpstra analyses for the exploratory MIC of *Origanum vulgare* essential oil.

Bacterial strains	S-W	P values in S-W	Interpretation of normality	P values in Jonckheere-Terpstra	Exploratory MIC (experimental criterion)
<i>E. coli</i> study	0.78935	4.219e-05	Not normal	7e-04	1024 µg/mL

<i>E. coli</i> ATCC 25922	0.77984	2.874e-05	Not normal	0.0012	512 µg/mL
<i>K. pneumoniae</i> study	0.87583	0.002265	Not normal	1e-04	1024 µg/mL
<i>K. pneumoniae</i> ATCC 700603	0.64886	3.048e-07	Not normal	0.0156	1024 µg/mL




Figure 4. Diagrams for the exploratory determination of the MIC of *Origanum vulgare* essential oil against: (a) *E. coli* study strain; (b) *E. coli* ATCC 25922 control strain; (c) *K. pneumoniae* study strain; (d) *K. pneumoniae* ATCC 700603 control strain. The central line represents the median, the box represents the interquartile range, and the individual points represent experimental replicates (n = 3). The vertical dashed line indicates the exploratory MIC, defined as the concentration associated with maximum inhibition.

The Jonckheere–Terpstra test showed a significant concentration-dependent trend in all evaluated strains ($p < 0.05$), indicating that the inhibitory effect of the essential oil increased as the

concentration increased. Based on this trend and the inhibition observed experimentally, the exploratory MIC was defined as the concentration associated with maximum inhibition for each strain. Consequently, in some strains the exploratory MIC differed from the visually observed MIC, reflecting a conservative statistical interpretation rather than a discrepancy in the experimental results. For example, while the observed MIC for the *E. coli* ATCC 25922 strain was 256 µg/mL, the exploratory MIC was defined as 512 µg/mL based on the statistical trend analysis. Similarly, for the *K. pneumoniae* study strain, the observed MIC was 2048 µg/mL, whereas the exploratory MIC was conservatively established at 1024 µg/mL.

Given the limited number of clinical isolates analyzed (n = 2), the main objective of the statistical analysis was to support the existence of a dose–response trend. In this sense, the MIC values reported in the present study should be interpreted as exploratory and primarily supported by the experimental evidence observed during the broth microdilution assays.

3.3. Determination of the Fractional Inhibitory Concentration Index (FICI)

Based on the FICI results of oregano essential oil against the bacterial strains, a complete checkerboard assay was performed using a 6 × 6 matrix, and all combinations were evaluated. Six combinations showing the greatest inhibition, measured by optical density, were selected, and the most representative result is detailed in Table 8.

Table 8. FICI results showing synergy between the essential oil and the antibiotic.

Bacterial strains	Concentration	Concentration	FICI
	oregano essential oil µg/mL	Cefepime µg/mL	
<i>E. coli</i> study	64	8	0.188 Synergy
<i>E. coli</i> ATCC 25922	8	0.0375	0.0625 Synergy
<i>K. pneumoniae</i> study	128	512	0.563 Addition
<i>K. pneumoniae</i> ATCC 700603	64	0.06	0.1225 Synergy

The results indicated a predominantly synergistic effect of the combination of oregano essential oil and cefepime against the *E. coli* study strain, *E. coli* ATCC, and *K. pneumoniae* ATCC, with a higher frequency of synergy observed in the *E. coli* study strain and a tendency toward additivity in the *K. pneumoniae* study strain, which required significantly higher concentrations.

4. Discussion

Phenotypic resistance profile

The results obtained from the disk diffusion test (Kirby–Bauer) for the *E. coli* study strain showed resistance to AMC (6 mm), ATM (10 mm), FEP (18 mm), CTX (10 mm), and CAZ (6 mm), similar to the study by Kumar et al. (2014), in which 65% of clinical isolates exhibited complete resistance to cephalosporins (ceftazidime, cefotaxime, and cefepime). Likewise, Rehman et al. reported resistance rates of up to 100% to cefepime and cefotaxime, with inhibition zones < 16 mm, confirming the multidrug-resistant profile of the strain evaluated in the present study [15], In contrast, the control strain *E. coli* ATCC 25922 showed susceptibility to all evaluated antibiotics, consistent with the

findings reported by Hudzicki, who indicated susceptibility to monobactams with inhibition zones equal to or greater than 30 mm in diameter [16], ATCC strains are essential as reference controls to confirm the reliability of the assay [17].

For the *K. pneumoniae* study strain, reduced inhibition zones were observed: AMC (13 mm), ATM (13 mm), FEP (17 mm), CTX (17 mm), and CAZ (14 mm), in agreement with Smith et al., who established that inhibition zones below 18 mm for FEP, AMC, and ATM indicate the presence of ESBL production [18]. Similarly, Husna described that resistance to monobactams and combinations such as AMC/ATM in ESBL-producing *K. pneumoniae* strains may exceed 90% of cases, which is consistent with the results obtained in this study [19]. Regarding the control strain *K. pneumoniae* ATCC 700603, the inhibition zones were within the quality control ranges established by EUCAST/CLSI, demonstrating a typical ESBL phenotype and validating the disk diffusion method. This strain is used as an ESBL control because it carries natural resistance associated with the *bla*_SHV-18 gene [20].

Antimicrobial activity of the essential oil (MIC)

The MIC of oregano essential oil for the *E. coli* study strain was 1024 µg/mL. This value is consistent with, although slightly higher than, that reported by Burt, who obtained an average MIC of 500 µg/mL for oregano essential oil against ESBL-producing *E. coli* strains [21]. This variability is attributed to differences in the chemical composition of the essential oil or to the study strains used, which may present distinct resistance profiles, resulting in greater natural or acquired susceptibility to natural compounds. For *E. coli* ATCC 25922, the obtained concentration (256 µg/mL) is consistent with the study by Khan et al., who reported an MIC of 225 µg/mL for the compound carvacrol [22]. Similarly, Stamova et al. reported a concentration of 250 µg/mL [23], supporting the validity of the results obtained in this study, as the essential oil used contains 86.74% carvacrol, providing high efficacy and confirming its antimicrobial activity through damage to the cytoplasmic membrane and alteration of cellular metabolism [21].

For the *K. pneumoniae* study strain, the MIC was 2048 µg/mL. Several studies have also evaluated oregano essential oil against clinical *K. pneumoniae* strains, including those with ESBL mechanisms, and have reported MIC values ranging from 128 to 256 µg/mL [24]. Another study conducted in Bulgaria using clinical isolates reported a very broad MIC range of 390 to 12,500 µg/mL; therefore, the strain used in the present study falls within the range observed for resistant strains. Finally, for the control strain *K. pneumoniae* ATCC 700603, the observed concentration of 1024 µg/mL represented significant inhibition due to its resistant nature. In comparison, a study that also evaluated the antimicrobial activity of oregano essential oil reported an MIC of 512 µg/mL [25], while Bouhaddouda evaluated the same strain and reported MIC values of 1560 µg/mL, indicating a variable range [26].

It should be noted that, although the statistical trend analysis supported a concentration-dependent inhibitory effect starting at 1024 µg/mL, the MIC for the *K. pneumoniae* study strain was conservatively reported as 2048 µg/mL. This decision was based on the microbiological criterion of complete and consistent inhibition observed across replicates, rather than solely on statistical significance. Given the exploratory nature of the study and the intrinsic resistance of *K. pneumoniae*, this conservative approach was adopted to ensure coherence with experimental inhibition patterns.

Antimicrobial activity of cefepime (MIC)

The *E. coli* study strain exhibited a cefepime MIC of 64 µg/mL. This value is consistent with the study by Kumar et al. (2014), in which more than 60% of clinical isolates presented a cefepime MIC of 64 µg/mL, providing clear evidence of the epidemiology of cefepime resistance in ESBL-producing strains. In addition, CLSI and EUCAST [27] indicate that MIC values above 16 µg/mL are associated with CTX-M-type enzymes and porin loss mechanisms, indicating a high resistance profile. In contrast, the cefepime MIC for *E. coli* ATCC 25922 was 0.12 µg/mL, which falls within the quality control (QC) range established by the CLSI 2025 standard, defined as 0.016 to 0.12 µg/mL; therefore, the obtained values do not exceed the established threshold.

For the *K. pneumoniae* study strain, the results showed a cefepime MIC of 64 µg/mL, confirming its resistant condition. Liang et al. evaluated 110 clinical ESBL-producing *K. pneumoniae* isolates and

reported that 47% of these strains exhibited resistance greater than 16 µg/mL, suggesting that such resistance levels are common in strains producing CTX-M or SHV-5 β-lactamases [27]. Patel reported that ESBL-producing *K. pneumoniae* with MIC values ranging from 2 to 8 µg/mL are associated with poor clinical response to cefepime [28]. In the case of the control strain *K. pneumoniae* ATCC 700603, the observed MIC was 1 µg/mL, which is within the QC ranges established by CLSI 2025 (0.25–2 µg/mL).

Antimicrobial interaction (FICI) between *Origanum vulgare* essential oil and cefepime

The combination of oregano essential oil with cefepime exhibited a synergistic effect against *E. coli*, indicating that carvacrol and thymol enhance the action of the antibiotic by increasing outer membrane permeability and facilitating the entry of the β-lactam, a mechanism previously described in the literature [13]. In addition, recent studies have reported that essential oils rich in carvacrol can reduce the expression of β-lactamases through gene suppression and protein inhibition, thereby decreasing the MIC of antibiotics in resistant Enterobacteriaceae, which is consistent with the marked MIC reduction observed in this study [12,29].

In *K. pneumoniae* the observed effect was additive, which is consistent with investigations indicating that this species exhibits lower susceptibility to hydrophobic compounds due to its polysaccharide capsule and reduced membrane permeability [30]. Despite this, the partial reduction of the antibiotic MIC in combination suggests an adjuvant effect of the essential oil. These results, together with findings reported in recent reviews, reinforce the potential use of essential oils as enhancers of antimicrobial therapies against multidrug-resistant bacteria.

5. Conclusions

This study demonstrates that the evaluated clinical strains of *E. coli* and *K. pneumoniae* exhibit a high resistance phenotype to β-lactam antibiotics, consistent with the expected profiles of ESBL-producing bacteria. The use of ATCC reference strains allowed validation of the applied methodologies and reinforced the reliability of the experimental results.

Origanum vulgare essential oil exhibited antimicrobial activity against all evaluated strains, although differences in susceptibility between species were observed. *E. coli* showed greater sensitivity to the essential oil, whereas *K. pneumoniae* required higher concentrations to achieve inhibition, reflecting its greater intrinsic resistance. Statistical analysis supported the observed trend, indicating a concentration-dependent inhibitory trend.

The evaluation of the fractional inhibitory concentration index (FICI) revealed synergistic interactions between the essential oil and cefepime in *E. coli*, as well as additive effects in *K. pneumoniae*. These findings suggest that the essential oil, particularly due to its high content of carvacrol and thymol, may act as an adjuvant in combination approaches against resistant Enterobacteriaceae.

Overall, the findings support the antimicrobial potential of *O. vulgare* essential oil and its ability to enhance the activity of β-lactam antibiotics against ESBL-producing bacteria. However, as this was a pilot study, further investigations involving a larger number of strains, cytotoxicity assessments, and evaluation in in vivo models are required to determine safety, efficacy, and potential clinical relevance.

Author Contributions: Conceptualization, A.G.; methodology, A.G. and R.A.; software, A.G.; validation, A.G. and M.E.; formal analysis, A.G., M.E., and R.A.; investigation, A.G.; resources, A.G. and M.E.; data curation, A.G. and M.E.; writing—original draft preparation, A.G. and M.E.; writing—review and editing, A.G., M.E., I.M., and R.A.; visualization, M.E. and R.A.; supervision, M.E. and R.A.; project administration, I.M. and M.E.; funding acquisition, I.M. and R.A. All authors have read and agreed to the published version of the manuscript.

Funding: this research received no external funding.

Institutional Review Board Statement: not applicable, as the study used only bacterial strains.

Informed Consent Statement: not applicable.

Data Availability Statement: the data presented in this study are available upon request from the corresponding author.

Acknowledgments: The authors thank Universidad Politécnica Salesiana for providing access to academic resources, training, and facilities used for the execution of this study. The authors also acknowledge INSPI for their valuable collaboration in the development of this work. The authors reviewed and edited the manuscript and assume full responsibility for the content of this publication.

Conflicts of Interest: the authors declare no conflict of interest.

References

1. K. Tang, B. Millar and J. Moore, «Antimicrobial Resistance (AMR),» British Journal of Biomedical Science, p. 80, 2023.
2. J. Munita and C. Arias, «Mechanisms of Antibiotic Resistance,» Microbiology Spectrum, p. 4 (2), 2016.
3. WHO, «Antimicrobial Resistance ,» World Health Organization, 17 11 2021.
4. M. Mueller and C. Tainter, «Escherichia coli Infection. StatPearls Publishing,» NCBI, 2023.
5. H. Namikawa, W. Imoto, K. Yamada, Y. Tochino, Y. Kaneko, H. Takeya and T. Shuto, «Predictors of mortality from extended-spectrum beta-lactamase-producing Enterobacteriaceae bacteremia,» Emerging Microbes & Infections, p. 12 (1), 2023.
6. A. Fostervold, N. Raffelsberger, M. Hetland, R. Bakksjø, E. Bernhoff, O. Samuelsen, A. Sundsfjord, J. Afset, C. Berntsen, R. Bævre-Jensen, M. Ebbesen, K. Gammelsrud, A. Guleng and I. Löhr, «Risk of death in Klebsiella pneumoniae bloodstream infections is associated with specific phylogenetic lineages,» Journal of Infection, pp. 88(5), 106155, 2024.
7. M. Contaldo, F. D'Ambrosio, G. Ferraro, D. di Stasio, M. di Palo, R. Serpico and M. Simeone, «A Narrative Review of the Evidence beyond the Myth. International Journal of Environmental Research and Public Health,» Antibiotics in Dentistry, pp. 20(11), 6025, 2023.
8. B. Petrovska, «Historical review of medicinal plants' usage,» Pharmacognosy Reviews, pp. 6(11), 1, 2012.
9. Q. Zhang, L. Lin and W. Ye, «Techniques for extraction and isolation of natural products: a comprehensive review,» Chinese Medicine, pp. 13(1), 20, 2018.
10. J. Hamuel, «Phytochemicals: Extraction Methods, Basic Structures and Mode of Action as Potential Chemotherapeutic Agents. In Phytochemicals,» A Global Perspective of Their Role in Nutrition and Health, 2012.
11. R. Nurzyńska-Wierdak and M. Walasek-Janusz, «Chemical Composition, Biological Activity, and Potential Uses of Oregano (*Origanum vulgare* L.) and Oregano Essential Oil,» Pharmaceuticals, pp. 18(2), 267, 2025.
12. N. Wijesundara, S. Lee, D. Langelaan, A. Lefsay and H. Rupasinghe, «Carvacrol alters the membrane phospholipids in erythromycin-resistant *Streptococcus pyogenes*,» Phytomedicine Plus, pp. 4(3), 100614, 2024.
13. Bassolé and H. Juliani, «Essential Oils in Combination and Their Antimicrobial Properties,» Molecules, pp. 17(4), 3989-4006, 2012.
14. H. Si, J. Hu, Z. Liu and Z. Zeng, «Antibacterial effect of oregano essential oil alone and in combination with antibiotics against extended-spectrum β -lactamase-producing *Escherichia coli*: Table 1,» FEMS Immunology & Medical Microbiology, pp. 53(2), 190-194, 2008.
15. N. Rehman, S. Azam, A. Ali, I. Khan, M. Asghar, M. Ali, M. Waqas, F. Ullah and G. Sehra, «Molecular epidemiology of antibiotic-resistant genes and potent inhibitors against TEM, CTX-M-14, CTX-M-15, and SHV-1 proteins of *Escherichia coli* in district Peshawar, Pakistan. Saudi Journal of Biological Sciences,» Saudi Journal of Biological Sciences, pp. 28(11), 6568-6581, 2021.
16. Hudzicki, «Kirby-Bauer disk diffusion susceptibility test protocol,» American society for microbiology, pp. 15(1), 1-23, 2009.
17. R. Miller, R. Walker, A. Baya, K. Clemens, M. Coles, M. Papapetropoulou and R. Reimschuessel, «Antimicrobial Susceptibility Testing of Aquatic Bacteria: Quality Control Disk Diffusion Ranges for *Escherichia coli* ATCC 25922 and *Aeromonas salmonicida* subsp. *salmonicida* ATCC 33658 at 22 and 28°C,» Journal of Clinical Microbiology, pp. 41(9), 4318-4323, 2003.

18. Smith, T. Brennan-Krohn, S. Weir and J. Kirby, «Improved Accuracy of Cefepime Susceptibility Testing for Extended-Spectrum-Beta-Lactamase-Producing Enterobacteriaceae with an On-Demand Digital Dispensing Method,» *Journal of Clinical Microbiology*, pp. 55(2), 470–478, 2017.
19. A. Husna, M. Rahman, A. Badruzzaman, M. Sikder, M. Islam, T. Rahman, J. Alam and H. Ashour, «Extended-Spectrum β -Lactamases (ESBL): Challenges and Opportunities,» *Biomedicines*, pp. 11(11), 2937, 2023.
20. EUCAST, «European Committee on Antimicrobial Susceptibility Testing,» EUCAST QC Tables v, 2025.
21. S. Burt, «Essential oils: their antibacterial properties and potential applications in foods—a review,» *International Journal of Food Microbiology*, pp. 94(3), 223–253, 2004.
22. Khan, A. Bahuguna, P. Kumar, V. Bajpai and S. Kang, «Antimicrobial Potential of Carvacrol against Uropathogenic *Escherichia coli* via Membrane Disruption, Depolarization, and Reactive Oxygen Species Generation,» *Frontiers in Microbiology*, p. 8, 2017.
23. S. Stamova, S. Mihaylova, N. Ermenlieva and G. Emilia, «IN VITRO STUDY OF ANTIMICROBIAL ACTIVITY OF ESSENTIAL OILS OF ORIGANUM VULGARE AGAINST STAPHYLOCOCCUS AUREUS ATCC 29213 AND ESCHERICHIA COLI ATCC 25922,» *Scripta Scientifica Pharmaceutica*, 8(2), 34-37. doi: HYPERLINK "http://dx.doi.org/10.14748/ssp.v8i2.8357" http://dx.doi.org/10.14748/ssp.v8i2.8357 , 2021.
24. S. Silva, F. Araújo, P. Silva, E. Silva, M. Bezerra, A. Diniz, D. Oliveira, H. Jesus, J. B. Nascimento, L. Medeiros and A. Oliveira Filho, «Evaluation of the antimicrobial effect of the *Origanum vulgare* L essential oil on strains of *Klebsiella pneumoniae*,» *Brazilian Journal of Biology*, p. 83, 2023.
25. Farhadi, E. Rajabi, H. Varpaei, M. Iranzadasl, S. Khodaparast and M. Salehi, «Thymol and carvacrol against *Klebsiella*: anti-bacterial, anti-biofilm, and synergistic activities—a systematic review,» *Frontiers in Pharmacology*, p. 15, 2024.
26. N. Bouhaddouda, S. Aouadi and R. Labiod, «Evaluation of Chemical Composition and Biological Activities of Essential Oil and Methanolic Extract of *Origanum vulgare* L. ssp. *glandulosum* (Desf.) Ietswaart from Algeria,» *International Journal of Pharmacognosy and Phytochemical Research*, pp. 8(1), 104–112, 2016.
27. W. Liang, M. Pfaller, P. Winokur and R. Jones, «Cefepime MIC as a Predictor of the Extended-Spectrum β -Lactamase Type in *Klebsiella pneumoniae*, Taiwan,» *China Medical College Hospital*, pp. 8,26, 2002.
28. H. Patel, «Cefepime use for ESBL infections: Usable or Inducible?,» *University of the Incarnate Word Feik School of Pharmacy*, 2014.
29. Muzammil, M. Ijaz, M. Saleem and A. Ali, «Drug repurposing strategy: An emerging approach to identify potential therapeutics for treatment of bovine mastitis,» *Microbial Pathogenesis*, pp. 171, 105691, 2022.
30. A. Karaś, A. Turska-Szewczuk, I. Komaniecka and B. Łotocka, «Lipidomics Analysis of Multilamellar Bodies Produced by *Amoeba Acanthamoeba castellanii* in Co-Culture with *Klebsiella aerogenes*,» *Pathogens*, pp. 12(3), 411, 2023.
31. A. Balley, «Development of Broth Microdilution MIC and Disk Diffusion Antimicrobial Susceptibility Test Quality Control Ranges for the Combination of Cefepime and the Novel β -Lactamase Inhibitor Enmetazobactam,» *Journal of Clinical Microbiology*, 26 7 2019.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.