

Article

Not peer-reviewed version

---

# Locating And Location Number

---

[Mohammadesmail Nikfar](#) \*

Posted Date: 8 June 2021

doi: 10.20944/preprints202106.0206.v1

Keywords: Special Set; Set's Weight; Special Number; Number's Position.



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

# Locating And Location Number

Mohammadesmail Nikfar

Independent Researcher

DrHenryGarrett@gmail.com

Twitter's ID: @DrHenryGarrett | ©B08PDK8J5G

## Abstract

In this article, some notions about set, weight of set, number, number's position, special vertex are introduced. Some classes of graph under these new notions have been opted as if the study on the special attributes of these new notion when they've acted amid each other is considered. Internal and external relations amid these new notions have been obtained as if some classes of graphs in the matter of these notions are been pointed out.

**Keywords:** Special Set, Set's Weight, Special Number, Number's Position.  
**AMS Subject Classification:** 05C17, 05C22, 05E45, 05E14

## 1 Preliminary On The Concept

I'm going to refer to some books which are cited to the necessary and sufficient material which are covering the introduction and the preliminary of this outlet so look [Ref. [1], Ref. [2], Ref. [3], Ref. [4]] where Ref. [1] is about the textbook, Ref. [2] is common, Ref. [3] has good ideas and Ref. [4] is kind of disciplinary approaches in the good ways. Further references could be referred and could be addressed in Refs. [5–11].

## 2 Definition And Its Clarification

**Definition 2.1.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a graph. Then

- $\mathcal{B}$  is renamed to be **LOCATING SET** if  $\forall x \in \mathcal{V}$ , there is  $m \in \mathcal{B}$  such that  $x \cap \mathcal{N}(m) \neq \emptyset$  or  $\mathcal{N}(\mathcal{B}) = \mathcal{V}$ ;
- Minimum cardinality  $\mathcal{B}$  is renamed to **LOCATION NUMBER**;
- $\mathcal{B}$ 's member is renamed to **LOCATING MEMBER**;
- Cardinality of  $\mathcal{B}$  is renamed to **LOCATING NUMBER**.

### 3 Relationships And Its illustrations

**Theorem 3.1.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a complete graph. Then

- $\mathcal{B} = \{m\}$  is renamed to be **LOCATING SET** because  $\forall x \in \mathcal{V}$ , there is  $m \in \mathcal{B}$  such that  $x \cap \mathcal{N}(m) \neq \emptyset$  and  $\mathcal{N}(\mathcal{B}) = \mathcal{V}$ .
- Minimum cardinality  $\mathcal{B}$ , 1 is renamed to **LOCATION NUMBER**.
- $\mathcal{B}$ 's member  $m \in \mathcal{V}$  is renamed to **LOCATING MEMBER**;
- Cardinality of  $\mathcal{B}$ ,  $1, 2, \dots, \mathcal{O}(\mathcal{G})$  is renamed to **LOCATING NUMBER**.

**Theorem 3.2.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a complete bipartite graph. Then

- $\mathcal{B} = \{m_1 \in \mathcal{V}_1, m_2 \in \mathcal{V}_2\}$  is renamed to be **LOCATING SET** because  $\forall x \in \mathcal{V}$  there is  $m_1 \in \mathcal{V}_1 \subseteq \mathcal{B}, m_2 \in \mathcal{V}_2 \subseteq \mathcal{B}$  such that  $x \cap \mathcal{N}(m) \neq \emptyset$  and  $\mathcal{N}(\mathcal{B}) = \mathcal{V}$ .
- Minimum cardinality  $\mathcal{B}$ , 2 is renamed to **LOCATION NUMBER**.
- $\mathcal{B}$ 's member  $m_1, m_2 \in \mathcal{V}$  is renamed to **LOCATING MEMBER**;
- Cardinality of  $\mathcal{B}$ ,  $2, \dots, \mathcal{O}(\mathcal{G})$  is renamed to **LOCATING NUMBER**.

**Theorem 3.3.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a star graph. Then

- $\mathcal{B} = \{m \in \mathcal{V}\}$  is renamed to be **LOCATING SET** because  $\forall x \in \mathcal{V}$  there is  $m = \mathcal{C}(\mathcal{G}) \subseteq \mathcal{B}$  such that  $x \cap \mathcal{N}(m) \neq \emptyset$  and  $\mathcal{N}(\mathcal{B}) = \mathcal{V}$ .
- Minimum cardinality  $\mathcal{B}$ , 1 is renamed to **LOCATION NUMBER**.
- $\mathcal{B}$ 's member  $m, \{m, m_1, m_2, \dots, m_{i-1}\}_{i=1}^{i=n}$  is renamed to **LOCATING MEMBER**;
- Cardinality of  $\mathcal{B}$ ,  $1, 2, \dots, \mathcal{O}(\mathcal{G})$  is renamed to **LOCATING NUMBER**.

**Theorem 3.4.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a wheel graph. Then

- $\mathcal{B} = \{m \in \mathcal{V}\}$  is renamed to be **LOCATING SET** because  $\forall x \in \mathcal{V}$  there is  $m = \mathcal{C}(\mathcal{G}) \subseteq \mathcal{B}$  such that  $x \cap \mathcal{N}(m) \neq \emptyset$  and  $\mathcal{N}(\mathcal{B}) = \mathcal{V}$ .
- Minimum cardinality  $\mathcal{B}$ , 1 is renamed to **LOCATION NUMBER**.
- $\mathcal{B}$ 's member  $m, \{m, m_1, m_2, \dots, m_{i-1}\}_{i=1}^{i=n}$  is renamed to **LOCATING MEMBER**;
- Cardinality of  $\mathcal{B}$ ,  $1, 2, \dots, \mathcal{O}(\mathcal{G})$  is renamed to **LOCATING NUMBER**.

**Theorem 3.5.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a path graph. Then

- $\mathcal{B} = \{m_2, m_4, \dots\}$  is renamed to be **LOCATING SET** because  $\forall x \in \mathcal{V}$  there is  $m \in \mathcal{B}$  such that  $x \cap \mathcal{N}(m) \neq \emptyset$  and  $\mathcal{N}(\mathcal{B}) = \mathcal{V}$ .
- Minimum cardinality  $\mathcal{B}$ ,  $\frac{\mathcal{O}(\mathcal{G})}{2}$  is renamed to **LOCATION NUMBER**.
- $\mathcal{B}$ 's member  $\{m_2, m_4, \dots\}$  and  $\{m_2, m_4, \dots\} \cup \{m_1, m_3, \dots, m_i\}_{i=1}$  is renamed to **LOCATING MEMBER**;
- Cardinality of  $\mathcal{B}$ ,  $\frac{\mathcal{O}(\mathcal{G})}{2}, \dots, \mathcal{O}(\mathcal{G})$  is renamed to **LOCATING NUMBER**.

**Theorem 3.6.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a cycle graph. Then

- $\mathcal{B} = \{m_2, m_4, \dots\}$  is renamed to be **LOCATING SET** because  $\forall x \in \mathcal{V}$  there is  $m \in \mathcal{B}$  such that  $x \cap \mathcal{N}(m) \neq \emptyset$  and  $\mathcal{N}(\mathcal{B}) = \mathcal{V}$ . 49 50
- Minimum cardinality  $\mathcal{B}$ ,  $\frac{\mathcal{O}(\mathcal{G})}{2}$  is renamed to **LOCATION NUMBER**. 51
- $\mathcal{B}$ 's member  $\{m_2, m_4, \dots\}$  and  $\{m_2, m_4, \dots\} \cup \{m_1, m_3, \dots, m_i\}_{i=1}$  is renamed to **LOCATING MEMBER**; 52 53
- Cardinality of  $\mathcal{B}$ ,  $\frac{\mathcal{O}(\mathcal{G})}{2}, \dots, \mathcal{O}(\mathcal{G})$  is renamed to **LOCATING NUMBER**. 54

## 4 Results And Its Beyond 55

- Theorem 4.1.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a graph. If **LOCATING MEMBER** is one, then one is **LOCATION NUMBER**. 56 57
- Theorem 4.2.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a graph. If **LOCATION NUMBER** is two, then  $\mathcal{G}$  is complete bipartite graph. 58 59
- Theorem 4.3.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a graph. If **LOCATION NUMBER** is one, then  $\mathcal{G}$  is either complete graph or star graph. 60 61
- Theorem 4.4.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a graph. Then  $\mathcal{O}(\mathcal{G})$  is **LOCATING NUMBER**. 62
- Theorem 4.5.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a graph. Then  $\mathcal{V}$  is **LOCATING SET**. 63
- Theorem 4.6.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a graph. If there is **LOCATING SET** then the cardinality of its minimal set is **LOCATION NUMBER**. 64 65
- Theorem 4.7.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a graph. Then  $1 \leq \mathbf{LOCATION NUMBER} \leq \mathcal{O}(\mathcal{G})$ . 66 67
- Theorem 4.8.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a graph.  $\forall m \in \mathcal{V}$ ,  $m$  is **LOCATING MEMBER** if and only if  $\mathcal{G}$  is complete graph. 68 69
- Theorem 4.9.** Let  $\mathcal{G} : (\mathcal{V}, \mathcal{E})$  be a graph. **LOCATION NUMBER** of  $\mathcal{G}$  is greater than **LOCATION NUMBER** of its subgraphs. 70 71
- Theorem 4.10.** **LOCATION NUMBER** of union of some graphs equals summation of **LOCATION NUMBER** of individual graphs. 72 73
- Theorem 4.11.** Every **LOCATING SET** of a graph is subset of  $\mathcal{V}$ . 74
- Theorem 4.12.** **LOCATING SET** of a graph is  $\mathcal{V}$  if and only if graph is empty graph. 75 76
- Theorem 4.13.** **LOCATING SET** of a graph with two partitions as two independent sets is one vertex from each of its partitions as representative if and only if graph is complete bipartite graph. 77 78 79

## References 80

1. R. Balakrishnan, K. Ranganathan, **A Textbook of Graph Theory**, New York, 2012. 81 82

2. Adrian Bondy, U.S.R Murty, **Graph Theory**, New York, 2008. 83

3. Michael Capobianco, and John C. Molluzzo, **Examples and counterexamples in graph theory**, New York, 1978. 84 85

4. Chris Godsil, and Gordon Royle, **Algebraic Graph Theory**, New York, 2001.

86

5. Joseph P.S. Kung, **Inconsequential results on the Merino-Welsh conjecture for Tutte polynomials**. Preprint available at <https://arxiv.org/abs/2105.01825>.

87

88

89

6. Duksang Lee,  **$\Gamma$ -graphic delta-matroids and its applications**. Preprint available at <https://arxiv.org/abs/2104.11383>.

90

91

7. Luis Ferroni, **Matroids are not Ehrhart positive**. Preprint available at <https://arxiv.org/abs/2105.04465>.

92

93

8. Luis Ferroni, Lorenzo Vecchi, **Matroid relaxations and Kazhdan-Lusztig non-degeneracy**. Preprint available at <https://arxiv.org/abs/2104.14531>.

94

95

9. Norbert Peyerimhoff, Marc Roth, Johannes Schmitt, Jakob Stix, Alina Vdovina, **Parameterized (Modular) Counting and Cayley Graph Expanders**. Preprint available at <https://arxiv.org/abs/2104.14596>.

96

97

98

10. James Oxley, Zach Walsh, **2-Modular Matrices**. Preprint available at <https://arxiv.org/abs/2105.04525>.

99

100

11. Zach Walsh, **New lift matroids for group-labeled graphs**. Preprint available at <https://arxiv.org/abs/2104.08257>.

101

102

