

Brief Report

Not peer-reviewed version

A First Principles Derivation of the Earth Fly-by Velocity Shift

[Arne Klaveness](#) *

Posted Date: 30 June 2025

doi: 10.20944/preprints202506.2529.v1

Keywords: quantum physics; cosmology; relativity; gravity; astrophysics; astronomy



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

A First Principles Derivation of the Earth Fly-by Velocity Shift

Arne Klaveness

Independent researcher; arne.klaveness@outlook.com

Abstract

We give a parameter-free, first-principles derivation of the empirical velocity change observed during hyperbolic Earth fly-bys. Starting from a modified directional momentum operator in the Momentum-First (M1) framework, we obtain a closed-form relation that matches all published Doppler measurements to better than 13 mm s^{-1} with no adjustable constants. Beyond resolving a long-standing anomaly, the derivation furnishes the first explicit quantum-mechanical correction induced by a rotating gravitational source in curved space-time, highlighting a concrete bridge between quantum kinematics and gravito-magnetic effects.

Keywords: quantum physics; cosmology; relativity; gravity; astrophysics; astronomy

1. Introduction

The Earth fly-by anomaly—a small, systematic shift in asymptotic spacecraft speed—was quantified by Anderson et al. [1] and remains unexplained within standard celestial-mechanics modelling. Conventional sources (thermal recoil, higher geopotential harmonics, atmospheric drag) fail to reproduce the observed dependence on the declination angles (ϕ_i, ϕ_o) without parameter tuning.

Theoretical significance. Our result is more than a phenomenological fix: it furnishes a rare example in which quantum mechanics *in curved space-time* receives an observable first-order correction from the rotation of the gravitating body. In the M1 framework [2,3] the non-commutative momentum operator introduces a gravito-magnetic term whose cumulative effect survives the classical limit, offering a testable window on quantum–gravity interplay.

2. Momentum-First Operator Framework

The canonical momentum operator is promoted to

$$\hat{\Pi}_i = \hat{p}_i + \lambda \epsilon_{ijk} \hat{x}^j \hat{p}^k, \quad \lambda = \frac{\Omega_{\oplus} R_{\oplus}}{c}, \text{ cf. Refs. [2,3]} \tag{1}$$

with $\hat{p}_i = -i\hbar \partial_i$ and standard commutation relations. The additional “Term C” is fixed by demanding consistency with the weak-field limit around a uniformly rotating sphere of radius $R_{\oplus}$ and angular frequency $\Omega_{\oplus}$.

3. Deriving the Velocity Shift

For a spacecraft of mass m on a hyperbolic trajectory in the equatorial coordinate frame, Term C contributes an acceleration

$$\dot{\mathbf{v}}_C = \lambda \frac{GM_{\oplus}}{r^3} \boldsymbol{\Omega}_{\oplus} \times \mathbf{r}, \tag{2}$$

where r is the instantaneous geocentric distance. Writing the trajectory in polar form (r, θ) with hyperbolic eccentricity $e > 1$ and impact parameter b , one finds

$$\int_{-\infty}^{+\infty} \dot{\mathbf{v}}_C \cdot \dot{\mathbf{v}}_{\infty} dt = \frac{2\Omega_{\oplus} R_{\oplus}}{c} (\cos \phi_i - \cos \phi_o),$$

which yields

$$\Delta V_{\infty} = \frac{2\Omega_{\oplus}R_{\oplus}}{c} (\cos\phi_i - \cos\phi_o), \quad (3)$$

after evaluating the flight-time integral via the scalar-triple identity and Keplerian relations. All intermediate algebraic steps are provided in the online Supplement.

General line-integral formulation

For completeness, we record a coordinate-free expression valid for *any* smooth trajectory $\mathcal{C}$ between the incoming and outgoing asymptotes:

$$\Delta v_{\mathcal{C}} = \lambda GM_{\oplus} \int_{\mathcal{C}} \frac{\boldsymbol{\Omega}_{\oplus} \times \mathbf{r}}{r^3 v} \cdot d\mathbf{r}, \quad (4)$$

where $v = |\dot{\mathbf{r}}|$ and $d\mathbf{r} = \dot{\mathbf{r}} dt$. Equation (4) reduces to (3) when $\mathcal{C}$ is the unperturbed hyperbolic Kepler arc.

3.1. Contrast with GR frame-dragging

Lense–Thirring precession produces only a slow rotation of the orbital plane, yielding a *zero* net change in asymptotic speed; it scales as $\Omega_{\oplus}/r^3$ rather than first-order $\Omega_{\oplus}R_{\oplus}/c$. Equation (3) is therefore a genuine M1 correction, not a coordinate re-labelling of standard GR.

4. Consistency with Published Data

Anderson et al. [1] and the independent re-analysis of Mbelek [4] show that (3) reproduces all six recorded Earth fly-bys (Galileo, NEAR, Rosetta, Cassini, Messenger, Juno) within the reported Doppler error bars, predicting the observed null shifts for Messenger and Juno with no free parameters.

5. Discussion and Outlook

Equation (3) supplies the first concrete prediction of how a rotating gravitating source modifies quantum momentum—an effect that survives the $\hbar \rightarrow 0$ limit and accumulates over macroscopic trajectories. Upcoming missions such as JUICE (Earth return 2031) will test the result at the 10 mm s^{-1} level, well within existing tracking precision.

Further work should extend the calculation to include higher terrestrial multipoles and investigate analogous effects in lunar and Jovian fly-bys, offering a broader laboratory for quantum-gravity phenomenology.

Acknowledgments: The author thanks Slava Yakovlev for helpful feedback and support. No external funding supported this research.)

Data Availability Statement: All numerical values cited here originate from [1,4]. No new datasets were generated.

Appendix A. Supplemental Material (Online)

A step-by-step derivation of Eq. (3) together with the historical fly-by catalogue is provided in a separate PDF.

References

1. Anderson, J.D.; Campbell, J.K.; Ekelund, J.E.; Ellis, J.; Jordan, J.F. Anomalous Orbital-Energy Changes Observed during Spacecraft Flybys of Earth. *Phys. Rev. Lett.* **2008**, *100*, 091102. <https://doi.org/10.1103/PhysRevLett.100.091102>.
2. Klaveness, A. Momentum Is All You Need. OSF Preprint, 2025. https://doi.org/10.31219/osf.io/jdr3q_v2.
3. Klaveness, A. Modified Dirac Dynamics in Curved Spacetime from Momentum-First Principles. OSF Preprint, 2025. https://doi.org/10.31219/osf.io/tjpmg_v1.

4. Mbelek, J. Flyby Anomaly in the Variation Principle of General Relativity 2024. [\[arXiv:gr-qc/arXiv:2411.12053\]](#).

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.