

SOME INEQUALITIES OF THE BELL POLYNOMIALS

FENG QI

Institute of Mathematics, Henan Polytechnic University, Jiaozuo City, Henan Province, 454010, China; Department of Mathematics, College of Science, Tianjin Polytechnic University, Tianjin City, 300387, China

ABSTRACT. In the paper, the author

- (1) presents an explicit formula and its inversion formula for higher order derivatives of generating functions of the Bell polynomials, with the help of the Faà di Bruno formula, properties of the Bell polynomials of the second kind, and the inversion theorem for the Stirling numbers of the first and second kinds;
- (2) recovers an explicit formula and its inversion formula for the Bell polynomials in terms of the Stirling numbers of the first and second kinds, with the aid of the above explicit formula and its inversion formula for higher order derivatives of generating functions of the Bell polynomials;
- (3) derives the (logarithmically) absolute and complete monotonicity of generating functions of the Bell polynomials;
- (4) constructs some determinantal and product inequalities and deduces the logarithmic convexity of the Bell polynomials, with the assistance of the complete monotonicity of generating functions of the Bell polynomials. These inequalities are main results of the paper.

CONTENTS

1. Introduction	2
2. Higher order derivatives for generating functions	3
3. An explicit formula and its inversion formula	4
4. Monotonicity of generating functions	5
5. Inequalities for the Bell polynomials	6
References	9

E-mail address: qifeng618@gmail.com, qifeng618@hotmail.com.

2010 *Mathematics Subject Classification.* Primary 11B83; Secondary 11A25, 11B73, 11C08, 11C20, 15A15, 26A24, 26A48, 26C05, 26D05, 33B10, 34A05.

Key words and phrases. Bell polynomial; Bell number; Bell polynomial of the second kind; higher order derivative; generating function; Faà di Bruno formula; inversion theorem; Stirling number of the first kind; Stirling number of the second kind; explicit formula; inversion formula; logarithmically absolute monotonicity; logarithmically complete monotonicity; determinantal inequality; product inequality.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

1. INTRODUCTION

In combinatorics, the Bell numbers, usually denoted by B_k for $k \in \{0\} \cup \mathbb{N}$, where \mathbb{N} denotes the set of all positive integers, count the number of ways a set with k elements can be partitioned into disjoint and nonempty subsets. These numbers have been studied by mathematicians since the 19th century, and their roots go back to medieval Japan, but they are named after Eric Temple Bell, who wrote about them in the 1930s. The Bell numbers B_k for $k \geq 0$ can be generated by

$$e^{e^t-1} = \sum_{k=0}^{\infty} B_k \frac{t^k}{k!} = 1 + t + t^2 + \frac{5}{6}t^3 + \frac{5}{8}t^4 + \frac{13}{30}t^5 + \frac{203}{720}t^6 + \frac{877}{5040}t^7 + \dots$$

and the first ten Bell numbers B_n are

$$\begin{array}{lllll} B_0 = 1, & B_1 = 1, & B_2 = 2, & B_3 = 5, & B_4 = 15, \\ B_5 = 52, & B_6 = 203, & B_7 = 877, & B_8 = 4140, & B_9 = 21147. \end{array}$$

In [1, Section 3], Asai, Kubo, and Kuo mentioned applications of the Bell numbers B_n and its some generalizations to white noise distribution theory. For more information on the Bell numbers B_k , please refer to [1, 4, 5, 6, 8, 18, 19, 25] and plenty of references therein.

As a well-known generalization of the Bell numbers B_k for $k \geq 0$, the Bell polynomials $B_k(x)$ for $k \geq 0$ can be generated by

$$\begin{aligned} e^{x(e^t-1)} &= \sum_{k=0}^{\infty} B_k(x) \frac{t^k}{k!} = 1 + xt + \frac{1}{2}x(x+1)t^2 + \frac{1}{6}x(x^2+3x+1)t^3 \\ &+ \frac{1}{24}x(x^3+6x^2+7x+1)t^4 + \frac{1}{120}x(x^4+10x^3+25x^2+15x+1)t^5 + \dots \end{aligned}$$

and the first six Bell polynomials $B_k(x)$ are

$$\begin{aligned} &1, \quad x, \quad x(x+1), \quad x(x^2+3x+1), \quad x(x^3+6x^2+7x+1), \\ &x(x^4+10x^3+25x^2+15x+1), \quad x(x^5+15x^4+65x^3+90x^2+31x+1). \end{aligned}$$

In [22] it was pointed out that there have been studies on interesting applications of the Bell polynomials $B_k(x)$ in soliton theory, including links with bilinear and trilinear forms of nonlinear differential equations which possess soliton solutions. See, for example, [11, 12, 13]. Therefore, applications of the Bell polynomials $B_k(x)$ to integrable nonlinear equations are greatly expected and any amendment on multilinear forms of soliton equations, even on exact solutions, would be beneficial to interested audiences in the research community. For more information about the Bell polynomials $B_k(x)$, please refer to [9, 10, 22, 24] and closely related references therein.

In this paper, continuing the article [19], we present an explicit formula and its inversion formula for higher order derivatives with respect to t of generating functions $e^{xe^{\pm t}}$ for the Bell polynomials $B_k(x)$ with the help of the Faà di Bruno formula, properties of the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$, and the inversion theorem for the Stirling numbers of the first and second kinds $s(n, k)$ and $S(n, k)$, recover an explicit formula and its inversion formula for the Bell polynomials $B_k(x)$ in terms of the Stirling numbers $s(n, k)$ and $S(n, k)$, derive the (logarithmically) absolute and complete monotonicity of generating functions $e^{xe^{\pm t}}$

for the Bell polynomials $B_k(x)$, construct some determinantal and product inequalities for the Bell polynomials $B_k(x)$, and deduce the logarithmic convexity of the Bell polynomials $B_k(x)$.

2. HIGHER ORDER DERIVATIVES FOR GENERATING FUNCTIONS

In this section, by the Faà di Bruno formula, properties of the Bell polynomials of the second kind $B_{n,k}$, and the inversion theorem for the Stirling numbers $s(n, k)$ and $S(n, k)$, we present an explicit formula and its inversion formula for higher order derivatives with respect to t of the generating function $e^{xe^{\pm t}}$.

Theorem 2.1. *For $n \geq 0$, the n th derivative of the generating function $e^{xe^{\pm t}}$ with respect to t can be computed by*

$$\frac{\partial^n e^{xe^{\pm t}}}{\partial t^n} = (\pm 1)^n e^{xe^{\pm t}} \sum_{k=0}^n S(n, k) (xe^{\pm t})^k \quad (2.1)$$

and the generating function $e^{xe^{\pm t}}$ satisfies a family of nonlinear ordinary differential equations

$$\sum_{k=0}^n (\pm 1)^k s(n, k) f^{(k)}(t) = f(t) [\ln f(t)]^n, \quad (2.2)$$

where $x \in \mathbb{C}$, $S(n, k)$ for $n \geq k \geq 0$, which can be generated by

$$\frac{(e^x - 1)^k}{k!} = \sum_{n=k}^{\infty} S(n, k) \frac{x^n}{n!},$$

represent the Stirling numbers of the second kind, and $s(n, k)$ for $n \geq k \geq 0$, which can be generated by

$$\frac{[\ln(1+x)]^k}{k!} = \sum_{n=k}^{\infty} s(n, k) \frac{x^n}{n!}, \quad |x| < 1,$$

stand for the Stirling numbers of the first kind.

Proof. In combinatorics, the Bell polynomials of the second kind $B_{n,k}$ are defined by

$$B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{\ell_1, \dots, \ell_n \in \{0\} \cup \mathbb{N} \\ \sum_{i=1}^n i \ell_i = n \\ \sum_{i=1}^n \ell_i = k}} \frac{n!}{\prod_{i=1}^{n-k+1} \ell_i!} \prod_{i=1}^{n-k+1} \left(\frac{x_i}{i!} \right)^{\ell_i}$$

for $n \geq k \geq 0$, see [5, p. 134, Theorem A], and satisfy identities

$$B_{n,k}(abx_1, ab^2x_2, \dots, ab^{n-k+1}x_{n-k+1}) = a^k b^n B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) \quad (2.3)$$

and

$$B_{n,k}(1, 1, \dots, 1) = S(n, k), \quad (2.4)$$

see [5, p. 135], where a and b are any complex numbers. The Faà di Bruno formula for computing higher order derivatives of composite functions can be described in terms of the Bell polynomials of the second kind $B_{n,k}$ by

$$\frac{d^n}{dt^n} f \circ g(x) = \sum_{k=0}^n f^{(k)}(g(x)) B_{n,k}(g'(x), g''(x), \dots, g^{(n-k+1)}(x)), \quad (2.5)$$

see [5, p. 139, Theorem C]. Applying $f(u) = e^{xu}$ and $u = g(t) = e^t$ to (2.5) and making use of identities (2.3) and (2.4) yield

$$\begin{aligned} \frac{\partial^n e^{xe^t}}{\partial t^n} &= \sum_{k=0}^n \frac{\partial^k e^{xu}}{\partial u^k} B_{n,k}(e^t, e^t, \dots, e^t) \\ &= \sum_{k=0}^n x^k e^{xu} e^{kt} B_{n,k}(1, 1, \dots, 1) = e^{xe^t} \sum_{k=0}^n x^k e^{kt} S(n, k). \end{aligned}$$

The explicit formula (2.1) for the plus sign case is thus proved.

In [27, p. 171, Theorem 12.1], it is stated that, if b_α and a_k are a collection of constants independent of n , then

$$a_n = \sum_{\alpha=0}^n S(n, \alpha) b_\alpha \quad \text{if and only if} \quad b_n = \sum_{k=0}^n s(n, k) a_k. \quad (2.6)$$

Combining this inversion theorem for the Stirling numbers with (2.1) arrives at equations

$$\sum_{k=0}^n s(n, k) \frac{\partial^k e^{xe^t}}{\partial t^k} = e^{xe^t} (xe^t)^n$$

which is equivalent to that the generating function e^{xe^t} satisfies the family of non-linear ordinary differential equations in (2.2) for the plus sign case.

The equation (2.12) in the second proof of [19, Theorem 2.2] reads that

$$B_{n,n}(\alpha) = \alpha^n \quad \text{and} \quad B_{n+k+1,k}(\alpha, 0, \dots, 0) = 0, \quad (2.7)$$

where $\alpha \in \mathbb{C}$ and $k, n \in \{0\} \cup \mathbb{N}$. Applying $f(u) = e^{xe^u}$ and $u = g(t) = -t$ in (2.5), taking $\alpha = -1$ in (2.7), and utilizing (2.1) lead to

$$\begin{aligned} \frac{\partial^n e^{xe^{-t}}}{\partial t^n} &= \sum_{k=0}^n \frac{\partial^k e^{xe^u}}{\partial u^k} B_{n,k}(-1, 0, \dots, 0) = \frac{\partial^n e^{xe^u}}{\partial u^n} B_{n,n}(-1) \\ &= (-1)^n e^{xe^u} \sum_{k=0}^n S(n, k) (xe^u)^k = (-1)^n e^{xe^{-t}} \sum_{k=0}^n S(n, k) (xe^{-t})^k. \end{aligned}$$

The formula (2.1) for the minus sign case follows immediately.

Employing the inversion theorem for Stirling numbers, demonstrated in (2.6), to consider the formula (2.1) for the minus sign case give

$$\sum_{k=0}^n s(n, k) (-1)^k \frac{\partial^k e^{xe^{-t}}}{\partial t^k} = e^{xe^{-t}} (xe^{-t})^n$$

which implies that the generating function $e^{xe^{-t}}$ satisfies the family of nonlinear ordinary differential equations in (2.2) for the minus sign case. The proof of Theorem 2.1 is thus complete. \square

3. AN EXPLICIT FORMULA AND ITS INVERSION FORMULA

In this section, with the aid of Theorem 2.1, we recover an explicit formula and its inversion formula for the Bell polynomials $B_n(x)$ in terms of the Stirling numbers $s(n, k)$ and $S(n, k)$.

Theorem 3.1. For $n \geq 0$, the Bell polynomials $B_n(x)$ can be computed by

$$B_n(x) = \sum_{k=0}^n S(n, k)x^k \quad (3.1)$$

and satisfy

$$\sum_{k=0}^n s(n, k)B_k(x) = x^n. \quad (3.2)$$

Proof. In light of the theory of series, it is easy to see that

$$B_n(x) = (\pm 1)^n \lim_{t \rightarrow 0} \frac{\partial^n e^{x(e^{\pm t} - 1)}}{\partial t^n} = (\pm 1)^n e^{-x} \lim_{t \rightarrow 0} \frac{\partial^n e^{xe^{\pm t}}}{\partial t^n}.$$

Combining this with (2.1) gives

$$B_n(x) = (\pm 1)^n e^{-x} \lim_{t \rightarrow 0} (\pm 1)^n e^{xe^{\pm t}} \sum_{k=0}^n S(n, k)(xe^{\pm t})^k = \sum_{k=0}^n S(n, k)x^k.$$

The formula (3.1) follows.

Substituting $f(t) = e^{xe^{\pm t}}$ into (2.2) and taking $t \rightarrow 0$ result in

$$\sum_{k=0}^n (\pm 1)^k s(n, k)(\pm 1)^k e^x B_k(x) = e^x x^n$$

which can be simplified as (3.2). The proof of Theorem 3.1 is complete. \square

4. MONOTONICITY OF GENERATING FUNCTIONS

Recall from [16, 31] that a function f is said to be absolutely monotonic on an interval I if it has derivatives of all orders and $f^{(k-1)}(t) \geq 0$ for $t \in I$ and $k \in \mathbb{N}$.

Recall from [7, p. 23, Definition 1] that a positive function f is said to be logarithmically absolutely monotonic on an interval I if it has derivatives of all orders and $[\ln f(t)]^{(k)} \geq 0$ for $t \in I$ and $k \in \mathbb{N}$.

Recall from [16, Chapter XIII], [28, Chapter 1], and [31, Chapter IV] that an infinitely differentiable function f is said to be completely monotonic on an interval I if it satisfies $(-1)^k f^{(k)}(x) \geq 0$ on I for all $k \geq 0$.

Recall from [2, 3, 20], [21, Remark 8], [23, pp. 2315–2316], [26, Remark 4.7], and [28, p. 67] that an infinitely differentiable function f is said to be logarithmically completely monotonic on an interval I if $(-1)^k [\ln f(x)]^{(k)} \geq 0$ on I for all $k \geq 1$.

We are now in a position to derive the (logarithmically) absolute and complete monotonicity of generating functions $e^{xe^{\pm t}}$ for the Bell polynomials $B_k(x)$. These results are seemingly simple, however, they are our basis of constructing determinantal and product inequalities for the Bell polynomials $B_k(x)$ in next section.

Theorem 4.1. When $x > 0$, the generating function e^{xe^t} is absolutely monotonic and, strongly, logarithmically absolutely monotonic with respect to $t \in \mathbb{R}$.

When $x < 0$, the generating function e^{-xe^t} is logarithmically absolutely monotonic and, consequently, absolutely monotonic with respect to $t \in \mathbb{R}$.

When $x > 0$, the generating function $e^{xe^{-t}}$ is logarithmically completely monotonic on \mathbb{R} and, consequently, completely monotonic with respect to $t \in \mathbb{R}$.

When $x < 0$, the generating function $e^{-xe^{-t}}$ is logarithmically completely monotonic on \mathbb{R} and, consequently, completely monotonic with respect to $t \in \mathbb{R}$.

Proof. By definition, the formula (2.1), and the positivity of the Stirling numbers of the second kind $S(n, k)$, it follows immediately that, when $x > 0$, the generating function e^{xe^t} is absolutely monotonic with respect to $t \in \mathbb{R}$.

Since $\ln e^{xe^t} = xe^t$, by definition, it follows readily that, when $x > 0$, the generating function e^{xe^t} is logarithmically absolutely monotonic with respect to $t \in \mathbb{R}$.

The absolute monotonicity of e^{xe^t} can also be derived from [7, p. 23, Theorem 1] which reads that a logarithmically absolutely monotonic function on an interval I is also absolutely monotonic on I , but not conversely.

The rest proofs are also straightforward, as did in the above argument. The proof of Theorem 4.1 is complete. \square

5. INEQUALITIES FOR THE BELL POLYNOMIALS

In light of complete monotonicity of generating functions $e^{xe^{\pm t}}$ in Theorem 4.1 and with the assistance of properties of completely monotonic functions, we now start out to construct some determinantal and product inequalities for the Bell polynomials $B_n(x)$. From these inequalities, we can derive the logarithmic convexity of the sequence of the Bell polynomials $B_n(x)$. These inequalities are our main results in this paper.

Theorem 5.1. *Let $m \geq 1$ be a positive integer and let $|a_{ij}|_m$ denote a determinant of order m with elements a_{ij} .*

- (1) *If a_i for $1 \leq i \leq m$ are non-negative integers, then*

$$|B_{a_i+a_j}(x)|_m \geq 0, \quad x > 0 \quad (5.1)$$

and

$$|(-1)^{a_i+a_j} B_{a_i+a_j}(x)|_m \geq 0, \quad x > 0. \quad (5.2)$$

- (2) *If $a = (a_1, a_2, \dots, a_n)$ and $b = (b_1, b_2, \dots, b_n)$ are non-increasing n -tuples of non-negative integers such that $\sum_{i=1}^k a_i \geq \sum_{i=1}^k b_i$ for $1 \leq k \leq n-1$ and $\sum_{i=1}^n a_i = \sum_{i=1}^n b_i$, then*

$$\prod_{i=1}^n B_{a_i}(x) \geq \prod_{i=1}^n B_{b_i}(x), \quad x > 0. \quad (5.3)$$

Proof. In [15] and [16, p. 367], it was obtained that if f is completely monotonic on $[0, \infty)$, then

$$|f^{(a_i+a_j)}(t)|_m \geq 0 \quad (5.4)$$

and

$$|(-1)^{a_i+a_j} f^{(a_i+a_j)}(t)|_m \geq 0. \quad (5.5)$$

Applying $f(t)$ to the function $e^{xe^{-t}}$ in (5.4) and (5.5) and taking the limit $t \rightarrow 0^+$ give

$$\lim_{t \rightarrow 0^+} |(e^{xe^{-t}})^{(a_i+a_j)}|_m = |(-1)^{a_i+a_j} e^x B_{a_i+a_j}(x)|_m \geq 0$$

and

$$\lim_{t \rightarrow 0^+} |(-1)^{a_i+a_j} (e^{xe^{-t}})^{(a_i+a_j)}|_m = |(-1)^{a_i+a_j} (-1)^{a_i+a_j} e^x B_{a_i+a_j}(x)|_m \geq 0.$$

The determinant inequalities (5.1) and (5.2) follow.

In [16, p. 367, Theorem 2], it was stated that if f is a completely monotonic function on $[0, \infty)$, then

$$\prod_{i=1}^n [(-1)^{a_i} f^{(a_i)}(t)] \geq \prod_{i=1}^n [(-1)^{b_i} f^{(b_i)}(t)]. \quad (5.6)$$

Applying $f(t)$ to the function $e^{xe^{-t}}$ in (5.6) and taking the limit $t \rightarrow 0^+$ give

$$\begin{aligned} \lim_{t \rightarrow 0^+} \prod_{i=1}^n [(-1)^{a_i} (e^{xe^{-t}})^{(a_i)}] &= \prod_{i=1}^n [e^x B_{a_i}(x)] \\ &\geq \lim_{t \rightarrow 0^+} \prod_{i=1}^n [(-1)^{b_i} (e^{xe^{-t}})^{(b_i)}] = \prod_{i=1}^n [e^x B_{b_i}(x)]. \end{aligned}$$

The product inequality (5.3) follows. The proof of Theorem 5.1 is complete. \square

Corollary 5.1. *The sequence $\{B_n(x)\}_{n \geq 0}$ for $x > 0$ is logarithmically convex.*

Proof. In [16, p. 369] and [17, p. 429, Remark], it was stated that if $f(t)$ is a completely monotonic function such that $f^{(k)}(t) \neq 0$ for $k \geq 0$, then the sequence

$$\ln[(-1)^{k-1} f^{(k-1)}(t)], \quad k \geq 1 \quad (5.7)$$

is convex. Applying this result to the function $e^{xe^{-t}}$ for $x > 0$ figures out that the sequence

$$\ln[(-1)^{k-1} (e^{xe^{-t}})^{(k-1)}] \rightarrow x + \ln B_{k-1}(x), \quad t \rightarrow 0^+$$

for $k \geq 1$ is convex. Hence, the sequence $\{B_n(x)\}_{n \geq 0}$ is logarithmically convex.

Alternatively, letting

$$\ell \geq 1, \quad n = 2, \quad a_1 = \ell + 2, \quad a_2 = \ell, \quad \text{and} \quad b_1 = b_2 = \ell + 1$$

in the inequality (5.3) leads to $B_\ell(x)B_{\ell+2}(x) \geq B_{\ell+1}^2(x)$ which means that the sequence $\{B_k(x)\}_{k \in \mathbb{N}}$ is logarithmically convex. The proof of Corollary 5.1 is complete. \square

Corollary 5.2. *If $\ell \geq 0$ and $n \geq k \geq 0$, then*

$$[B_{n+\ell}(x)]^k [B_\ell(x)]^{n-k} \geq [B_{k+\ell}(x)]^n, \quad x > 0.$$

Proof. This follows from taking

$$a = (\overbrace{n+\ell, \dots, n+\ell}^k, \overbrace{\ell, \dots, \ell}^{n-k}) \quad \text{and} \quad b = (k+\ell, k+\ell, \dots, k+\ell)$$

in the inequality (5.3). The proof of Corollary 5.2 is complete. \square

Theorem 5.2. *If $\ell \geq 0$, $n \geq k \geq m$, $2k \geq n$, and $2m \geq n$, then*

$$B_{k+\ell}(x)B_{n-k+\ell}(x) \geq B_{m+\ell}(x)B_{n-m+\ell}(x), \quad x > 0. \quad (5.8)$$

Proof. In [29, p. 397, Theorem D], it was recovered that if $f(t)$ is completely monotonic on $(0, \infty)$ and if $n \geq k \geq m$, $k \geq n-k$, and $m \geq n-m$, then

$$(-1)^n f^{(k)}(t) f^{(n-k)}(t) \geq (-1)^n f^{(m)}(t) f^{(n-m)}(t).$$

Replacing $f(t)$ by the function $(-1)^\ell (e^{xe^{-t}})^{(\ell)}$ in the above inequality leads to

$$(-1)^n (e^{xe^{-t}})^{(k+\ell)} (e^{xe^{-t}})^{(n-k+\ell)} \geq (-1)^n (e^{xe^{-t}})^{(m+\ell)} (e^{xe^{-t}})^{(n-m+\ell)}.$$

Further taking $t \rightarrow 0^+$ finds the inequality (5.8). The proof of Theorem 5.2 is complete. \square

Theorem 5.3. For $x, \ell \geq 0$ and $m, n \in \mathbb{N}$, let

$$\begin{aligned}\mathcal{G}_{\ell,m,n} &= B_{\ell+2m+n}(x)[B_{\ell}(x)]^2 - B_{\ell+m+n}(x)B_{\ell+m}(x)B_{\ell}(x) \\ &\quad - B_{\ell+n}(x)B_{\ell+2m}(x)B_{\ell}(x) + B_{\ell+n}(x)[B_{\ell+m}(x)]^2, \\ \mathcal{H}_{\ell,m,n} &= B_{\ell+2m+n}(x)[B_{\ell}(x)]^2 - 2B_{\ell+m+n}(x)B_{\ell+m}(x)B_{\ell}(x) \\ &\quad + B_{\ell+n}(x)[B_{\ell+m}(x)]^2, \\ \mathcal{I}_{\ell,m,n} &= B_{\ell+2m+n}(x)[B_{\ell}(x)]^2 - 2B_{\ell+n}(x)B_{\ell+2m}(x)B_{\ell}(x) \\ &\quad + B_{\ell+n}(x)[B_{\ell+m}(x)]^2.\end{aligned}$$

Then

$$\begin{aligned}\mathcal{G}_{\ell,m,n} &\geq 0, \quad \mathcal{H}_{\ell,m,n} \geq 0, \\ \mathcal{H}_{\ell,m,n} &\leq \mathcal{G}_{\ell,m,n} \quad \text{when } m \leq n, \\ \mathcal{I}_{\ell,m,n} &\geq \mathcal{G}_{\ell,m,n} \geq 0 \quad \text{when } n \geq m.\end{aligned}\tag{5.9}$$

Proof. In [30, Theorem 1 and Remark 2], it was obtained that if f is completely monotonic on $(0, \infty)$ and

$$\begin{aligned}G_{m,n} &= (-1)^n \{ f^{(n+2m)} f^2 - f^{(n+m)} f^{(m)} f - f^{(n)} f^{(2m)} f + f^{(n)} [f^{(m)}]^2 \}, \\ H_{m,n} &= (-1)^n \{ f^{(n+2m)} f^2 - 2f^{(n+m)} f^{(m)} f + f^{(n)} [f^{(m)}]^2 \}, \\ I_{m,n} &= (-1)^n \{ f^{(n+2m)} f^2 - 2f^{(n)} f^{(2m)} f + f^{(n)} [f^{(m)}]^2 \}\end{aligned}$$

for $n, m \in \mathbb{N}$, then

$$\begin{aligned}G_{m,n} &\geq 0, \quad H_{m,n} \geq 0, \\ H_{m,n} &\leq G_{m,n} \quad \text{when } m \leq n, \\ I_{m,n} &\geq G_{m,n} \geq 0 \quad \text{when } n \geq m.\end{aligned}\tag{5.10}$$

Replacing $f(t)$ by $(-1)^\ell (e^{xe^{-t}})^{(\ell)}$ in $G_{m,n}$, $H_{m,n}$, and $I_{m,n}$ and simplifying produce

$$\begin{aligned}G_{m,n} &= (-1)^{\ell+n} \left\{ (e^{xe^{-t}})^{(\ell+2m+n)} \left[(e^{xe^{-t}})^{(\ell)} \right]^2 \right. \\ &\quad - (e^{xe^{-t}})^{(\ell+m+n)} (e^{xe^{-t}})^{(\ell+m)} (e^{xe^{-t}})^{(\ell)} \\ &\quad \left. - (e^{xe^{-t}})^{(\ell+n)} (e^{xe^{-t}})^{(\ell+2m)} (e^{xe^{-t}})^{(\ell)} + (e^{xe^{-t}})^{(\ell+n)} \left[(e^{xe^{-t}})^{(\ell+m)} \right]^2 \right\}, \\ H_{m,n} &= (-1)^{\ell+n} \left\{ (e^{xe^{-t}})^{(\ell+2m+n)} \left[(e^{xe^{-t}})^{(\ell)} \right]^2 \right. \\ &\quad \left. - 2(e^{xe^{-t}})^{(\ell+m+n)} (e^{xe^{-t}})^{(\ell+m)} (e^{xe^{-t}})^{(\ell)} + (e^{xe^{-t}})^{(\ell+n)} \left[(e^{xe^{-t}})^{(\ell+m)} \right]^2 \right\}, \\ I_{m,n} &= (-1)^{\ell+n} \left\{ (e^{xe^{-t}})^{(\ell+2m+n)} \left[(e^{xe^{-t}})^{(\ell)} \right]^2 \right. \\ &\quad \left. - 2(e^{xe^{-t}})^{(\ell+n)} (e^{xe^{-t}})^{(\ell+2m)} (e^{xe^{-t}})^{(\ell)} + (e^{xe^{-t}})^{(\ell+n)} \left[(e^{xe^{-t}})^{(\ell+m)} \right]^2 \right\}.\end{aligned}$$

Further taking $t \rightarrow 0^+$ reveals

$$\lim_{t \rightarrow 0^+} G_{m,n} = e^3 \mathcal{G}_{\ell,m,n}, \quad \lim_{t \rightarrow 0^+} H_{m,n} = e^3 \mathcal{H}_{\ell,m,n}, \quad \text{and} \quad \lim_{t \rightarrow 0^+} I_{m,n} = e^3 \mathcal{I}_{\ell,m,n}.$$

Substituting these quantities into (5.10) and simplifying bring about inequalities in (5.9). The proof of Theorem 5.3 is complete. \square

Theorem 5.4. For $x, k \geq 0$ and $n \in \mathbb{N}$, we have

$$\left[\prod_{\ell=0}^n B_{k+2\ell}(x) \right]^{1/(n+1)} \geq \left[\prod_{\ell=0}^{n-1} B_{k+2\ell+1}(x) \right]^{1/n}. \quad (5.11)$$

Proof. If f is a completely monotonic function on $(0, \infty)$, then, by the convexity of the sequence (5.7) and Nanson's inequality listed in [14, p. 205, 3.2.27],

$$\left[\prod_{\ell=0}^n (-1)^{k+2\ell+1} f^{(k+2\ell+1)}(t) \right]^{1/(n+1)} \geq \left[\prod_{\ell=1}^n (-1)^{k+2\ell} f^{(k+2\ell)}(t) \right]^{1/n}$$

for $k \geq 0$. Replacing $f(t)$ by $e^{xe^{-t}}$ in the above inequality results in

$$\left[\prod_{\ell=0}^n (-1)^{k+2\ell+1} (e^{xe^{-t}})^{(k+2\ell+1)} \right]^{1/(n+1)} \geq \left[\prod_{\ell=1}^n (-1)^{k+2\ell} (e^{xe^{-t}})^{(k+2\ell)} \right]^{1/n}$$

for $k \geq 0$. Letting $t \rightarrow 0^+$ in the above inequality leads to (5.11). The proof of Theorem 5.4 is complete. \square

REFERENCES

- [1] N. Asai, I. Kubo, and H.-H. Kuo, *Bell numbers, log-concavity, and log-convexity*, Acta Appl. Math. **63** (2000), no. 1-3, 79–87; Available online at <https://doi.org/10.1023/A:1010738827855>.
- [2] R. D. Atanassov and U. V. Tsoukrovski, *Some properties of a class of logarithmically completely monotonic functions*, C. R. Acad. Bulgare Sci. **41** (1988), no. 2, 21–23.
- [3] C. Berg, *Integral representation of some functions related to the gamma function*, Mediterr. J. Math. **1** (2004), no. 4, 433–439; Available online at <https://doi.org/10.1007/s00009-004-0022-6>.
- [4] E. R. Canfield, *Engel's inequality for Bell numbers*, J. Combin. Theory Ser. A **72** (1995), no. 1, 184–187; Available online at [https://doi.org/10.1016/0097-3165\(95\)90033-0](https://doi.org/10.1016/0097-3165(95)90033-0).
- [5] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Revised and Enlarged Edition, D. Reidel Publishing Co., Dordrecht and Boston, 1974.
- [6] K. Engel, *On the average rank of an element in a filter of the partition lattice*, J. Combin. Theory Ser. A **65** (1994), no. 1, 67–78; Available online at [https://doi.org/10.1016/0097-3165\(94\)90038-8](https://doi.org/10.1016/0097-3165(94)90038-8).
- [7] B.-N. Guo and F. Qi, *A property of logarithmically absolutely monotonic functions and the logarithmically complete monotonicity of a power-exponential function*, Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys. **72** (2010), no. 2, 21–30.
- [8] B.-N. Guo and F. Qi, *An explicit formula for Bell numbers in terms of Stirling numbers and hypergeometric functions*, Glob. J. Math. Anal. **2** (2014), no. 4, 243–248; Available online at <https://doi.org/10.14419/gjma.v2i4.3310>.
- [9] F. T. Howard, *A special class of Bell polynomials*, Math. Comp. **35** (1980), no. 151, 977–989; Available online at <https://doi.org/10.2307/2006208>.
- [10] T. Kim, D. V. Dolgy, D. S. Kim, H. I. Kwon, and J. J. Seo, *Differential equations arising from certain Sheffer sequence*, J. Comput. Anal. Appl. **23** (2017), no. 8, 1359–1367.
- [11] W.-X. Ma, *Bilinear equations, Bell polynomials and linear superposition principle*, J. Phys. Conf. Ser. **411** (2013), no. 1, Article ID 012021, 11 pages; Available online at <https://doi.org/10.1088/1742-6596/411/1/012021>.
- [12] W.-X. Ma, *Bilinear equations and resonant solutions characterized by Bell polynomials*, Rep. Math. Phys. **72** (2013), no. 1, 41–56; Available online at [https://doi.org/10.1016/S0034-4877\(14\)60003-3](https://doi.org/10.1016/S0034-4877(14)60003-3).

- [13] W.-X. Ma, *Trilinear equations, Bell polynomials, and resonant solutions*, Front. Math. China **8** (2013), no. 5, 1139–1156; Available online at <https://doi.org/10.1007/s11464-013-0319-5>.
- [14] D. S. Mitrinović, *Analytic Inequalities*, Springer-Verlag, 1970.
- [15] D. S. Mitrinović and J. E. Pečarić, *On two-place completely monotonic functions*, Anzeiger Öster. Akad. Wiss. Math.-Naturwiss. Kl. **126** (1989), 85–88.
- [16] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, 1993; Available online at <https://doi.org/10.1007/978-94-017-1043-5>.
- [17] J. E. Pečarić, *Remarks on some inequalities of A. M. Fink*, J. Math. Anal. Appl. **104** (1984), no. 2, 428–431; Available online at [https://doi.org/10.1016/0022-247X\(84\)90006-4](https://doi.org/10.1016/0022-247X(84)90006-4).
- [18] F. Qi, *An explicit formula for the Bell numbers in terms of the Lah and Stirling numbers*, Mediterr. J. Math. **13** (2016), no. 5, 2795–2800; Available online at <https://doi.org/10.1007/s00009-015-0655-7>.
- [19] Feng Qi, *Some inequalities for the Bell numbers*, Proc. Indian Acad. Sci. Math. Sci. **127** (2017), no. 4, in press; Available online at <https://doi.org/10.1007/s12044-017-0355-2>.
- [20] F. Qi and C.-P. Chen, *A complete monotonicity property of the gamma function*, J. Math. Anal. Appl. **296** (2004), no. 2, 603–607; Available online at <https://doi.org/10.1016/j.jmaa.2004.04.026>.
- [21] F. Qi, S. Guo, and B.-N. Guo, *Complete monotonicity of some functions involving polygamma functions*, J. Comput. Appl. Math. **233** (2010), no. 9, 2149–2160; Available online at <https://doi.org/10.1016/j.cam.2009.09.044>.
- [22] F. Qi, D. Lim, and B.-N. Guo, *Explicit formulas and identities for the Bell polynomials and a sequence of polynomials applied to differential equations*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM (2018), in press; Available online at <https://doi.org/10.1007/s13398-017-0427-2>.
- [23] F. Qi, Q.-M. Luo, and B.-N. Guo, *Complete monotonicity of a function involving the divided difference of digamma functions*, Sci. China Math. **56** (2013), no. 11, 2315–2325; Available online at <https://doi.org/10.1007/s11425-012-4562-0>.
- [24] F. Qi, D.-W. Niu, and B.-N. Guo, *Some identities for a sequence of unnamed polynomials connected with the Bell polynomials*, Preprints **2017**, 2017080044, 10 pages; Available online at <https://doi.org/10.20944/preprints201708.0044.v1>.
- [25] F. Qi, X.-T. Shi, and F.-F. Liu, *Expansions of the exponential and the logarithm of power series and applications*, Arab. J. Math. (Springer) **6** (2017), no. 2, 95–108; Available online at <https://doi.org/10.1007/s40065-017-0166-4>.
- [26] F. Qi, C.-F. Wei, and B.-N. Guo, *Complete monotonicity of a function involving the ratio of gamma functions and applications*, Banach J. Math. Anal. **6** (2012), no. 1, 35–44; Available online at <https://doi.org/10.15352/bjma/1337014663>.
- [27] J. Quaintance and H. W. Gould, *Combinatorial Identities for Stirling Numbers*. The unpublished notes of H. W. Gould. With a foreword by George E. Andrews. World Scientific Publishing Co. Pte. Ltd., Singapore, 2016.
- [28] R. L. Schilling, R. Song, and Z. Vondraček, *Bernstein Functions—Theory and Applications*, 2nd ed., de Gruyter Studies in Mathematics **37**, Walter de Gruyter, Berlin, Germany, 2012; Available online at <https://doi.org/10.1515/9783110269338>.
- [29] H. van Haeringen, *Completely monotonic and related functions*, J. Math. Anal. Appl. **204** (1996), no. 2, 389–408; Available online at <https://doi.org/10.1006/jmaa.1996.0443>.
- [30] H. van Haeringen, *Inequalities for real powers of completely monotonic functions*, J. Math. Anal. Appl. **210** (1997), no. 1, 102–113; Available online at <https://doi.org/10.1006/jmaa.1997.5376>.
- [31] D. V. Widder, *The Laplace Transform*, Princeton University Press, Princeton, 1946.