

Article

Predictors of Speed and Agility in Youth Male Basketball Players

Denis Čaušević¹, Nedim Čović¹, Ensar Abazović¹, Babina Rani², Gabriel Marian Manolache^{3*}, Cătălin Vasile Ciocan^{4*}, Gabriel Zaharia⁵, Dan Iulian Alexe⁴

¹ Faculty of Sport and Physical Education, University of Sarajevo, 71000 Sarajevo, Bosnia and Herzegovina

² Department of Physical Rehabilitation & Medicine (Physiotherapy), Post Graduate Institute of Medical Education and Research, 160012 Chandigarh, India

³ Department of Sports Games and Physical of Education Faculty of Physical Education and Sport "Dunarea de Jos" University of Galati, 800008, Romania

⁴ Department of Physical Education and Sports Performance, Faculty of Movement, Sports and Health, Sciences, "Vasile Alecsandri", University of Bacău, 600115 Bacău, Romania

⁵ National University of Physical Education and Sport Bucharest, 060057, Romania

* Correspondence: gabriel.manolache@ugal.ro (G.M.M); ciocan.catalin@ub.ro (C.C)

Abstract: Player's performance in an intense sport such as basketball is known to be related to attributes like speed, agility, and power. This study presents a comparative analysis of associations between anthropometric assessment and physical performance in different age-group elite youth basketball players, while simultaneously identifying the predictors for speed and agility in these players. U14 (n=44), U15 (n=45) and U16 (n=51) players were tested for anthropometry, lower body power, speed, and agility. U16 players were found to be taller, heavier, more muscular than U14 and U15 players. Also, the U16 group showed better performance in all performance tests. Age had a significant positive correlation with countermovement (CMJ) and drop jump (DJ) performance in U14 players, and a significant negative correlation with 10m and 20m sprint times in U15 group. CMJ and DJ emerged as the most significant predictors for sprint and agility variables, respectively. Body fat percentage was found to be a significant predictor for the speed and agility tests in all age groups, but a negative lower-body power predictor. Therefore, besides all sport-specific and fitness tests, it is essential to place emphasis on the percentage of body fat when designing players' individualized training programs, and during team selection.

Keywords: team sports; explosive strength; body composition; CMJ

1. Introduction

Basketball is a dynamic and complex aerobic game that involves frequent interchange of high-intensity anaerobic activities such as jumping, accelerations and decelerations, changes of direction (COD), and sprinting [1] (Besides the player's technical skills, performance and success are related to lower body physical components of multidirectional running, power, vertical jump, and agility [2,3,4]).

Special research interest is directed towards physical fitness and its relation with high performance in young basketball players. In fact, speed, agility, and power are the most performance-related, evaluated, and tested abilities in basketball that tend to peak with attaining body height peak velocity (roughly around age 14) in male basketball players [5].

A well-known fact is that significant relations exist between jump performance, running speed, and agility in basketball players [3,4,6,7,8]. The aforementioned physical components have been shown to vary by gender and age [9], especially from 12-14 years of age [10]. Buchanan and Vardaxis (2003) reported that speed tends to be similar in the 11-13 age group but disparate for the 15-17-year-olds [11]. Additionally, 14-year-olds have better speed and agility performance when compared to 12-year-olds [10]. How speed and agility in young players are affected by violent changes in the

morphological structure, occurring from 12 to 16 years [12] and which differences among age groups are present still remains unclear. Such differences probably emerge due to morphological diversity in the muscle mass as well as muscle structure and differences in muscle fiber types. Variation in speed and agility is also affected by puberty, adolescent growth spurt, relative age, and maturation level [13,14,15].

Selection of the well-talented basketball players is usually performed by anthropometric screening, emphasizing body height. Even though the height is genetically predetermined, general and sports-specific fitness tests such as 5,10- and 20-meter sprint, Countermovement jump (CMJ), Drop Jump (DJ), Squat jump, and T-test and Lane agility are most commonly used for the assessment of players' physical potential related to basketball success [10,16] independent of training status and experience. As suggested by Cui et al. (2019), drafted NBA players outperformed undrafted players in body height, arm span, vertical jumps, and agility [17]. This is of special importance in relation to the player's position. Body dimensions such as height, limb length, body fat percentage and BMI make a consequential influence on running speed, explosive strength, and agility [3,4,18,19,20,21]. Body fat percentage has been negatively associated with explosive actions, change of direction and vertical jumps [22,23].

Available literature focusing on correlations between anthropometric assessment and physical performance in elite youth basketball players is highly limited. Moreover, it is important to comparatively analyze these relations in different levels of basketball players, in order to accurately predict the lacking component to be improved in a player, thus enhancing the sport performance as well as facilitating early return to sports in case of an injury. Therefore, the aim of this research was to identify the anthropometric factors predicting the speed and agility profiling in under-14 (U14), under-15 (U15), and under-16 (U16) basketball players. In addition, we aimed to investigate the correlation of physical performance tests with the players' anthropometric profile.

2. Materials and Methods

2.1. Participants

One hundred and forty male basketball players from under 14 (U14), under 15 (U15), and under 16 (U16) age groups participated in this study. Forty-four players were U14 (age = 13.41 ± 0.54 years, height = 175.14 ± 8.58 cm, body mass = 63.46 ± 13.70 kg); 45 players were U15 (age = 14.71 ± 0.29 years, height = 183.60 ± 7.91 cm, body mass = 75.20 ± 11.87 kg); and 51 players were U16 (age = 15.64 ± 0.32 years, height = 185.99 ± 5.83 cm, body mass = 76.25 ± 10.29 kg). All participants were enrolled in a regional competition in Bosnia and Herzegovina.

2.2. Measurement Procedure

All the tests/measurements were performed by experienced personnel at Institute of Sport at Faculty of Sport and Physical Education, University of Sarajevo. Specifically for this study, an appropriate testing cycle lasting two days (in a row) was planned and designed in a specific order. The testing sessions took place between 09:00 and 12:00 in the morning. On the first day, body height, body mass and body fat percentage (PBF) measurements were performed. Body height was measured to one decimal place (0.1 cm) by digital stadiometer (InBody BSM 370; Biospace Co., Ltd., Seoul, Korea). Body mass and PBF were estimated using a direct segmental high-frequency bioelectrical impedance scale (InBody 720; Biospace Co., Ltd., Seoul, Korea). Prior to testing of lower-body power (CMJ and DJ), players were instructed to complete a standard warm-up consisting of three minutes of jogging, three minutes of dynamic stretching, and three minutes of acceleration-deceleration. The following day, all speed and agility tests were performed, including 15 m and 20 m sprint tests, and T-test and lane agility test. Prior to the tests, all players performed standard warm-up protocol. All tests were conducted on a wooden basketball court. The study was approved by the Ethics Committee (No: 01-2603/22; 01.07.2022.) of the Faculty of Sports and Physical Education, University of Sarajevo and was carried out in accordance with the Declaration of Helsinki. The

informed consent was obtained from the participants' parents or legal guardians, prior to enrolment into the study.

2.3. Lower-Body Power

For assessment of the lower-body power, countermovement jump (CMJ) and drop jump (DJ) from a height of 40 cm were performed [24] (Young et al. 1995). Both protocols included two trials that were measured by using Optojump Next system (Microgate, Bolzano, Italy). For CMJ, players started from an upright standing position (with feet shoulder-width apart and hands fixed on hips throughout the jump to eliminate the use of arm swing), made a preliminary downward countermovement position to a self-selected depth by flexing the hips and knees. Players then immediately extend their hips and knees to make a vertical jump. The player returned to the starting position after the jump. The best of the two trials was used for analysis. The trial was considered invalid in case of knee flexion at landing, or if arm swing was detected.

The DJ was performed from a 40 cm wooden box, as recommended by previous study [25]. The DJ protocol began with the athlete standing on the box with hands fixed on the hips for the entire protocol to eliminate the arm swing influence. They were then instructed to step off the box one foot at the time (self-chosen order of limbs) and then after contact with the ground, to jump as fast and high as possible. The jump was invalid if hands were removed from the hips at any point or if athlete jumped off the box. After each jump, the athlete returned to the starting position, and the procedure was repeated two times. The best of the two trials was used for analysis.

2.4. Sprint tests

As part of the running speed testing, the participants ran a 20 m maximal sprint starting from a stationary standing position, with their lead foot 20 cm behind the photocell. Three photocells (Microgate, Bolzano, Italy) were used to measure 15 m and 20 m sprint times with precision of 0.01 s). Two trials were performed by each player with a 3-minute rest in between, and the best of the two was used for analysis.

2.5. Agility

Agility was evaluated by using the T-test and lane agility test, as recommended in previous studies [26,27]. For the T-test, athletes started the test from the standing position with their lead foot placed 20 cm behind the first photocell. Players were asked to sprint forwards 9.14 m to the center cone and touch it with right hand tip, then shuffle 4.57 m to the left to touch the second cone, then 9.50 m to the right to the third cone and then shuffle back 4.75 m to touch the center cone with left hand before they finally ran backwards to the starting point. Test time was recorded in seconds (to 0.01 s) with a photocell placed at the starting line (Microgate, Bolzano, Italy). Two trials were performed for the T-test, with a rest period of 3 minutes in-between.

For the lane agility test, athletes started the test from a standing position with their lead foot placed 20 cm behind first photocell. Players sprinted 5.79m forward towards the cone on the top right, then side shuffled 4.87m to the cone on the top left, backstep 5.79m to the cone on the bottom left, then go to the side shuffle 4.87m to the cone on the bottom right. Then, athletes returned to the starting point, around the cones in a reverse order (shuffle left, run forward, shuffle left, run backward). Two trials were completed, with three-minute rest intervals between trials. Test time was recorded in seconds with photocells placed at the starting line (Microgate, Bolzano, Italy).

2.5. Statistical Analysis

Descriptive statistics for was expressed as means \pm standard deviations. Two-way analysis of variance (ANOVA) was carried out to investigate the differences in, body composition and physical performance test results between three age-groups (U14, U15, U16). Pairwise comparisons were done by using Bonferroni post-hoc test. The relationship between physical performance and body composition variables was assessed using Pearson's r product-moment correlation coefficient. The

size of correlations was evaluated using the following criteria [28,29]: trivial (0.0), small (0.1), medium (0.3), large (0.5), very large (0.7), nearly perfect (0.9), and perfect (1.0). A hierarchical model of multiple regression analysis was conducted to investigate the amount of variance in speed and agility tests explained by CMJ and DJ (Step 3), after adjusting for CA (Step 1), and body composition variables (Step 2). The regression analysis was performed to investigate whether the superior performance in lower body power tasks (CMJ, DJ) would correspond to the lower sprinting and agility time scores. All analyses were performed using IBM SPSS Statistics software 22.0 (SPSS Inc., Chicago, IL, USA). The significance level was set at $p \leq 0.05$.

3. Results

Table 1 summarizes the body composition, speed and agility characteristics for three age categories of basketball players. The U16 players were significantly taller and heavier than their peers from other age groups. Overall, the U16 group showed better performance in all physical fitness tests.

Table 1. Descriptive statistics and post-hoc comparisons for body composition and physical fitness tests

Variables	U14		U15		U16		ANOVA		
	Mean	SD	Mean	SD	Mean	SD	η^2	F	p-value
CA (years)	13.41	0.54	14.71	0.29	15.64	0.32	/	304.879	≤ 0.01
Height (cm)	175.14 [€]	8.58	183.60	7.91	185.99 [€]	5.83	.28	26.888	≤ 0.01
Body mass (kg)	63.46 [€]	13.70	75.20	11.87	76.25 [€]	10.29	.19	16.044	≤ 0.01
PBF (%)	12.88	5.98	13.92	5.62	12.92	5.74	.07	0.477	0.62
CMJ (cm)	28.83	4.36	30.39	5.62	31.44 [€]	4.88	.04	3.255	0.04
DJ (cm)	27.68	4.63	29.89	5.28	31.68 [€]	5.49	.09	7.039	≤ 0.01
15m sprint (s)	2.69	0.19	2.64 [§]	0.18	2.53 [€]	0.14	.12	9.345	≤ 0.01
20m sprint (s)	3.40	0.24	3.35 [§]	0.23	3.20 [€]	0.19	.13	10.254	≤ 0.01
T-test (s)	11.45	0.87	11.21 [§]	0.78	10.77 [€]	0.85	.10	8.129	≤ 0.01
Lane agility (s)	13.48	0.98	13.13	0.85	12.77 [€]	0.89	.09	7.215	≤ 0.01

SD = standard deviation); CA = chronological age; PBF = body fat percentage; CMJ = countermovement jump; DJ = drop jump; η^2 = partial eta squared; [€] = sig. difference U14 vs U15 $p < 0.01$; [§] = sig. difference U15 vs U16 $p < 0.01$; [€] = sig. difference U14 vs U16 $p < 0.01$

Results of Pearson product-moment correlations according to age groups are presented in Table 2. Among the U14 group, CA presented significant positive correlations with CMJ ($r = .50$, $p \leq 0.01$) and DJ ($r = .43$, $p \leq 0.01$). In the U15 group, CA showed significant correlation with 10m speed ($r = -.30$, $p \leq 0.05$) and 20 m speed ($r = -.34$, $p \leq 0.05$). In the U16 group, CA did not correlate with any of analysed variables. PBF correlated significantly and positively with speed and agility tests in all age groups, but in contrast, it was negative predictor of lower-body power in all groups (CMJ from $r = -.39$ to $r = -.63$, $p \leq 0.01$; DJ from $r = -.42$ to $r = -.56$, $p \leq 0.01$). Generally, speed and agility tests were strongly and negatively correlated with lower-body power across all groups. A significant and positive correlation was observed between sprint time and agility performance in all age groups.

Table 3 summarizes the results (whole sample) of a hierarchical multiple regression conducted on CA (Step 1), body composition (Step 2) and lower-body power (Step 3) on speed and agility performance. CA and body composition variables explained 49% and 53% of variance observed in 10 and 20m sprint time. By adding lower-body power variables, an additional 11% and 13% of variance were explained. Along with PBF ($\beta = 0.39$; $p \leq 0.01$), DJ ($\beta = -0.33$; $p \leq 0.05$) remained the most powerful predictor in predictive model. Regarding the agility tests, CA and body composition variables explained 42% (T-test) and 36% (Lane agility) of variance observed. By adding lower-body power variables, an additional 13% and 10% of variance were explained. The DJ ($\beta = -0.51$; $p \leq 0.01$) remained the most significant predictor of the whole agility model.

Table 2. Descriptive statistics and post-hoc comparisons for body composition and physical fitness tests

	Variable	1.	2.	3.	4.	5.	6.	7.	8.	9.	10.
U14	1. CA (age)	1	.284	.178	-.203	.495**	.434**	-.459**	-.536**	-.571**	-.370*
	2. Height (cm)		1	.769**	-.001	.074	-.050	-.348*	-.343*	-.264	-.250
	3. Body mass (kg)			1	.515**	-.082	-.147	-.022	-.017	-.130	-.163
	4. PBF (%)				1	-.493**	-.421**	.632**	.661**	.405**	.301*
	5. CMJ (cm)					1	.898**	-.711**	-.739**	-.719**	-.486**
	6. DJ (cm)						1	-.635**	-.650**	-.690**	-.530**
	7. 15m sprint (s)							1	.962**	.702**	.480**
	8. 20m sprint (s)								1	.726**	.509**
	9. T-test (s)									1	.772**
	10. Lane agility (s)										1
U15	1. CA (age)	1	.276	.096	-.121	.180	.209	-.349*	-.343*	-.194	-.231
	2. Height (cm)		1	.564**	-.208	.350*	.285	-.278	-.349*	-.308*	-.159
	3. Body mass (kg)			1	.400**	.011	-.009	-.058	-.068	.012	.241
	4. PBF (%)				1	-.629**	-.560**	.506**	.566**	.573**	.551**
	5. CMJ (cm)					1	.898**	-.652**	-.719**	-.629**	-.555**
	6. DJ (cm)						1	-.549**	-.621**	-.587**	-.550**
	7. 15m sprint (s)							1	.969**	.588**	.543**
	8. 20m sprint (s)								1	.662**	.600**
	9. T-test (s)									1	.844**
	10. Lane agility (s)										1
U16	1. CA (age)	1	-.229	-.059	.017	-.079	-.050	-.119	-.126	-.181	-.036
	2. Height (cm)		1	.433**	-.329*	.163	.057	-.079	-.120	-.021	-.066
	3. Body mass (kg)			1	.259	.053	.031	.032	.031	.039	.063
	4. PBF (%)				1	-.394**	-.446**	.544**	.531**	.469**	.426**
	5. CMJ (cm)					1	.861**	-.659**	-.658**	-.516**	-.475**
	6. DJ (cm)						1	-.718**	-.702**	-.634**	-.595**
	7. 15m sprint (s)							1	.967**	.690**	.658**
	8. 20m sprint (s)								1	.679**	.693**
	9. T-test (s)									1	.873**
	10. Lane agility (s)										1

CA = chronological age; PBF = body fat percentage; CMJ = countermovement jump; DJ = drop jump; * = $p \leq 0.05$; ** = $p \leq 0.01$

Table 3. Multiple Regression analysis

	15m sprint			20m sprint			T-test			Lane agility		
	Model I	Model II	Model III	Model I	Model II	Model III	Model I	Model II	Model III	Model I	Model II	Model III
	β	β	β	β	β	β	β	B	β	β	β	β
1. CA (age)	-0.43**	-0.34**	-0.23**	-0.46**	-0.35**	-0.26**	-0.43**	-0.36**	-0.23*	-0.37**	-0.30**	-0.16
2. Height (cm)		0.11	0.06		0.08	0.01		0.19	0.08		0.03	-0.09
3. Body mass (kg)		-0.33*	-0.19		0.30*	-0.16		-0.36*	-0.19		-0.17*	-0.05
4. PBF (%)		0.65**	0.39**		0.66**	0.38**		0.59**	0.32**		0.44**	0.18
5. CMJ (cm)			-0.35*			-0.38**			-0.10			0.07
6. DJ (cm)			-0.09			-0.07			-0.36*			-0.51**
R ²	0.19	0.49	0.61	0.21	0.53	0.66	0.18	0.42	0.55	0.14	0.36	0.46
F for change in R ²	31.939**	32.764**	34.564**	36.010**	38.614**	43.502**	30.988**	23.984**	26.646**	21.519**	13.661**	15.921**

Model I: CA, Model II: CA, Height, Body mass and PBF; Model III: CA, Height, Body mass, PBF, CMJ height and DJ height; CA = chronological age; PBF = body fat percentage; CMJ = countermovement jump; DJ = drop jump; * = $p \leq 0.05$; ** = $p \leq 0.01$

4. Discussion

This research aimed to identify the anthropometric factors predicting the speed and agility profiling in under-14 (U14), under-15 (U15), and under-16 (U16) basketball players. In addition, we aimed to investigate the correlation of physical performance tests with the players' anthropometric profile.

Besides obvious differences in anthropometric variables, we found significant between age-group differences in all performance variables. Interestingly, no PBF differences were found between age groups. Further on, although the U14 and U16 groups differed in all of the measured variables, the U14 and U15 groups differed in only two variable (height and body mass), and the U15 and U16 groups differed in three variables (10 and 20m sprints and T-test). This is understandable considering the fact that great biological and morphological transformations and variations occur in adolescence [30]. In fact, similar inconsistencies have been found in numerous earlier studies. For example, Jakovljević et al. (2012) [10] found similar trends in mixed young basketball and football players aged 12-15 years. In their study, they divided participants according to their chronological age (12, 13, 14, and 15 years). No differences in speed and agility test results were found between 12 and 13, and 14 and 15, whilst 13 and 14 differed in all performance variables.

Within-age-group correlation showed that CA correlated with all performance variables in U14 and sprint performance in the U15 group. No significant correlations were found in the U16 group implying that the phenomenon known as relative age effect [31] is present in U15 and younger age groups. This implies that a specific biological development stabilization takes place, and it is crucial to take this stabilization into account during testing and scouting.

Further on, body height and weight correlation with performance tasks was found to be weak and insignificant (< 0.35), whereas PBF showed large and constant, correlation with power and sprint tests. In each age group, sprint and power performance exhibited a large to nearly perfect correlation with correlation coefficients ranging 0.475 - 0.969 (irrespective of the direction). Similar results were found previously in young [10,32] and professional basketball players [6].

The results indicate that CMJ and DJ are significantly correlated with sprint and agility performance in all groups and, while CMJ showed a stronger correlation with these results than DJ in U14 and U15, it was opposite in the U16 age category.

It is possible that the findings could be influenced by several factors, including the age and maturity of the participants, as well as the specific training methods used. It was previously reported that power performance may be mediated by biological maturation [33], agility performance tends to improve naturally with age, but there are significant increases from childhood to early adolescence, followed by a near-plateau in mid-adolescence [34]. Further on, training regimens designed to increase lower-body power and strength which have the potential to improve agility performance [35,36] are commonly introduced after peak height velocity age (13-15 years for boys) [12] because strength training is generally less effective before the growth spurt [37]. Therefore, training adaptations (in agility performance) are not solely attributable to the impacts of exposed training stimuli but also to the natural developmental processes of the young athlete [38,39]. Additionally, DJ is a measure of reactive muscular strength and may be more closely associated with agility, which requires rapid changes of direction, compared to CMJ, which provides information on lower limb power development during both the eccentric and concentric phase under the influence of the stretch-shortening cycle (SSC) [40,41].

The multiple regression analysis supported the aforementioned findings. CMJ and DJ both emerged as the significant predictors for the sprint and agility variables, respectively, showing lower limb power and SSC utilization ability are crucial components of high-quality sports performance. Additionally, PBF with its moderate, but constant, correlation coefficients with performance tasks showed to be a significant predictor for the majority speed and agility capacities. This is

understandable since excess body fat can negatively affect the ability to move quickly and efficiently. Additionally, carrying excess body fat can increase the energy cost of movement [42] thus leading to decreased speed and endurance. Therefore, athletes with lower percentages of body fat are typically able to move more quickly and efficiently, leading to improved speed performance.

Given that talent identification refers to the process of recognizing current participants with the potential to become elite players [43], it is reasonable to assume that these tests could be utilized for appropriate talent selection and evaluation of both, player development and training programs in young basketball players.

Possible limitations could be non-inclusion of specific basketball ball-dribbling tests and between-playing-position comparisons since players specialize in their positions and develop more specific skills related to those positions.

5. Conclusions

Significant age-group differences were found in all performance variables, with no differences in percentage body fat between age groups. Within-age-group correlation showed interesting results, with the relative age effect present in U15 and younger age groups. Lower-body explosive strength was linked to essential attributes of high-quality sports performance such as speed and agility, with CMJ and DJ emerging as the most significant predictors for sprint and agility variables, respectively. Additionally, PBF showed moderate but constant correlation coefficients with performance tasks and emerged as a significant predictor for most speed and agility capacities, highlighting the importance of maintaining lower percentages of body fat for improved performance. When interpreting the results, it is crucial to consider potential variations in age, biological maturation, and training specificity among basketball players.

Author Contributions: Conceptualization: D.Č., N.Č., G.Z and D.I.A.; methodology, N.Č., and D.Č.; software, B.R.; validation, E.A., D.I.A., and C.V.C.; formal analysis, D.Č. and B.R.; investigation, D.Č., N.Č., and E.A.; resources, G.Z., G.M.M and D.I.A; data curation, D.Č.; writing—original draft preparation, D.Č., N.Č., B.R., and E.A.; writing—review and editing, D.I.A. and D.Č.; visualization, N.Č., D.I.A., and E.A; supervision, D.Č. and D.I.A.; project administration, N.Č.; funding acquisition, G.Z., G.M.M and C.V.C. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding

Informed Consent Statement: Informed consent was obtained from participants' parents or legal guardians.

Data Availability Statement: The data presented in this study are available on reasonable request from the corresponding author.

Acknowledgments: The work of Dan Iulian Alexe and Cătălin Vasile Ciocan was supported by "Vasile Alecsandri" University of Bacau, Romania. The work of Gabriel Zaharia was supported by the project "Proinvent", Contract no. 62487/03.06.2022 - POCU/993/6/13 - Code 153299, by The Human Capital Operational Programme 2014-2020 POCU, Romania.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Jeličić M, Ivančev V, Čular D, Čović N, Stojanović E, Scanlan AT, Milanović Z. The 30-15 intermittent fitness test: A reliable, valid, and useful tool to assess aerobic capacity in female basketball players. *Research Quarterly for Exercise and Sport*. 2020 Jan 2;91(1):83-91. <https://doi.org/10.1080/02701367.2019.1648743>
2. Deletrat A, Cohen D. Physiological testing of basketball players: toward a standard evaluation of anaerobic fitness. *The Journal of Strength & Conditioning Research*. 2008 Jul 1;22(4):1066-72. <https://doi.org/10.1519/JSC.0b013e3181739d9b>
3. Čaušević D, Mašić S, Doder I, Matulaitis K, Spicer S. Speed, agility and power potential of young basketball players. *Baltic journal of sport & health sciences*. 2022;4(127):29-34. <https://doi.org/10.33607/bjshs.v127i4.1297>

4. Čaušević D, Abazović E, Mašić S, Hodžić A, Ormanović Š, Doder I, Čović N, Lakota R. Agility, sprint and vertical jump performance relationship in young basketball players. *Acta Kinesiologica*. 2021;1:133-7. <https://doi.org/10.51371/issn.1840-2976.2021.15.1.16>
5. Philippaerts RM, Vaeyens R, Janssens M, Van Renterghem B, Matthys D, Craen R, Bourgois J, Vrijens J, Beunen G, Malina RM. The relationship between peak height velocity and physical performance in youth soccer players. *Journal of sports sciences*. 2006 Mar 1;24(3):221-30. <https://doi.org/10.1080/02640410500189371>
6. Alemdaroğlu U. The relationship between muscle strength, anaerobic performance, agility, sprint ability and vertical jump performance in professional basketball players. *Journal of human kinetics*. 2012 Mar 1;31(2012):149-58. <https://doi.org/10.2478/v10078-012-0016-6>
7. Chaouachi A, Brughelli M, Chamari K, Levin GT, Abdelkrim NB, Laurencelle L, Castagna C. Lower limb maximal dynamic strength and agility determinants in elite basketball players. *The Journal of Strength & Conditioning Research*. 2009 Aug 1;23(5):1570-7. <https://doi.org/10.1519/JSC.0b013e3181a4e7f0>
8. Shalfawi SA, Sabbah A, Kailani G, Tønnessen E, Enoksen E. The relationship between running speed and measures of vertical jump in professional basketball players: a field-test approach. *The Journal of Strength & Conditioning Research*. 2011 Nov 1;25(11):3088-92. <https://doi.org/10.1519/JSC.0b013e318212db0e>
9. Mancha-Triguero D, García-Rubio J, Gamonales JM, Ibáñez SJ. Strength and speed profiles based on age and sex differences in young basketball players. *International Journal of Environmental Research and Public Health*. 2021 Jan;18(2):643. <https://doi.org/10.3390/ijerph18020643>
10. Jakovljevic ST, Karalejic MS, Pajic ZB, Macura MM, Erculj FF. Speed and agility of 12-and 14-year-old elite male basketball players. *The Journal of Strength & Conditioning Research*. 2012 Sep 1;26(9):2453-9. <https://doi.org/10.1519/JSC.0b013e31823f2b22>
11. Buchanan PA, Vardaxis VG. Sex-related and age-related differences in knee strength of basketball players ages 11–17 years. *Journal of athletic training*. 2003 Jul;38(3):231.
12. Malina RM, Bouchard C, Bar-Or O. Growth, maturation, and physical activity. *Human kinetics*; 2004.
13. Arede J, Fernandes J, Moran J, Norris J, Leite N. Maturity timing and performance in a youth national basketball team: Do early-maturing players dominate? *International Journal of Sports Science & Coaching*. 2021 Jun;16(3):722-30. <https://doi.org/10.1177/1747954120980712>
14. Toselli S, Campa F, Maietta Latessa P, Greco G, Loi A, Grigoletto A, Zaccagni L. Differences in maturity and anthropometric and morphological characteristics among young male basketball and soccer players and non-players. *International Journal of Environmental Research and Public Health*. 2021 Apr 8;18(8):3902. <https://doi.org/10.3390/ijerph18083902>
15. Towlson C, Cogley S, Parkin G, Lovell R. When does the influence of maturation on anthropometric and physical fitness characteristics increase and subside?. *Scandinavian journal of medicine & science in sports*. 2018 Aug;28(8):1946-55. <https://doi.org/10.1111/sms.13198>
16. Sansone P, Makivic B, Csapo R, Hume P, Martínez-Rodríguez A, Bauer P. Body fat of basketball players: A systematic review and meta-analysis. *Sports Medicine-Open*. 2022 Dec;8(1):26. <https://doi.org/10.1186/s40798-022-00418-x>
17. Cui Y, Liu F, Bao D, Liu H, Zhang S, Gómez MÁ. Key anthropometric and physical determinants for different playing positions during National Basketball Association draft combine test. *Frontiers in Psychology*. 2019 Oct 22;10:2359. <https://doi.org/10.3389/fpsyg.2019.02359>
18. Duncan MJ, Woodfield L, Al-Nakeeb Y. Anthropometric and physiological characteristics of junior elite volleyball players. *British journal of sports medicine*. 2006 Jul 1;40(7):649-51. <http://dx.doi.org/10.1136/bjism.2005.021998>
19. Mancha-Triguero D, Garcia-Rubio J, Calleja-Gonzalez J, Ibanez SJ. Physical fitness in basketball players: A systematic review. *J. Sports Med. Phys. Fit*. 2019 Sep 1;59:1513-25. <https://doi.org/10.23736/S0022-4707.19.09180-1>
20. Mancha-Triguero D, García-Rubio J, Gamonales JM, Ibáñez SJ. Strength and speed profiles based on age and sex differences in young basketball players. *International Journal of Environmental Research and Public Health*. 2021 Jan;18(2):643. <https://doi.org/10.3390/ijerph18020643>
21. Popowczak, M.; Horička, P.; Šimonek, J.; Domaradzki, J. The Functional Form of the Relationship between Body Height, Body Mass Index and Change of Direction Speed, Agility in Elite Female Basketball and Handball Players. *Int. J. Environ. Res. Public Health* 2022, 19, 15038. <https://doi.org/10.3390/ijerph192215038>
22. Ribeiro BG, Mota HR, Sampaio-Jorge F, Morales AP, Leite TC. Correlation between body composition and the performance of vertical jumps in basketball players. *J. Exerc. Physiol. Online*. 2015 Oct 1;18:69-79.
23. Spiteri T, Newton RU, Binetti M, Hart NH, Sheppard JM, Nimphius S. Mechanical determinants of faster change of direction and agility performance in female basketball athletes. *The Journal of Strength & Conditioning Research*. 2015 Aug 1;29(8):2205-14. <https://doi.org/10.1519/JSC.0000000000000876>
24. Young WB, Pryor JF, Wilson GJ. Effect of instructions on characteristics of countermovement and drop jump performance. *The Journal of Strength & Conditioning Research*. 1995 Nov 1;9(4):232-6.

25. Zhang M, Liang X, Huang W, Ding S, Li G, Zhang W, Li C, Zhou Y, Sun J, Li D. The effects of velocity-based versus percentage-based resistance training on athletic performances in sport-collegiate female basketball players. *Frontiers in Physiology*. 2023 Jan 10;13:2739. <https://doi.org/10.3389/fphys.2022.992655>
26. Gál-Pottyondy, A.; Petró, B.; Czétényi, A.; Négyesi, J.; Nagatomi, R.; Kiss, R.M. Collection and Advice on Basketball Field Tests—A Literature Review. *Appl. Sci.* 2021, 11, 8855. <https://doi.org/10.3390/app11198855>
27. Bae, J.-Y. Positional Differences in Physique, Physical Strength, and Lower Extremity Stability in Korean Male Elite High School Basketball Athletes. *Int. J. Environ. Res. Public Health* 2022, 19, 3416. <https://doi.org/10.3390/ijerph19063416>
28. Silva, A.F.; Alvrdu, S.; Akyildiz, Z.; Clemente, F.M. Relationships of Final Velocity at 30-15 Intermittent Fitness Test and Anaerobic Speed Reserve with Body Composition, Sprinting, Change-of-Direction and Vertical Jumping Performances: A Cross-Sectional Study in Youth Soccer Players. *Biology* 2022, 11, 197. <https://doi.org/10.3390/biology11020197>
29. Hopkins, W. Spreadsheets for Analysis of Validity and Reliability. *Sports Sci.* 2015, 19, 36–42.
30. Vaeyens R, Lenoir M, Williams AM, Philippaerts RM. Talent identification and development programmes in sport: Current models and future directions. *Sports medicine*. 2008 Sep;38:703-14. <https://doi.org/10.2165/00007256-200838090-00001>
31. Musch J, Grondin S. Unequal competition as an impediment to personal development: A review of the relative age effect in sport. *Developmental review*. 2001 Jun 1;21(2):147-67. <https://doi.org/10.1006/drev.2000.0516>
32. Asadi A. Relationship between jumping ability, agility and sprint performance of elite young basketball players: A field-test approach. *Revista Brasileira de Cineantropometria & Desempenho Humano*. 2016 Mar;18:177-86. <https://doi.org/10.5007/1980-0037.2016v18n2p177>
33. Moran JJ, Sandercock GR, Ramírez-Campillo R, Meylan CM, Collison JA, Parry DA. Age-related variation in male youth athletes' countermovement jump after plyometric training: a meta-analysis of controlled trials. *The Journal of Strength & Conditioning Research*. 2017 Feb 1;31(2):552-65. <https://doi.org/10.1519/JSC.0000000000001444>
34. Zemková, E.; Hamar, D. Age-related changes in agility time in children and adolescents. *Int. J. Sci. Res.* 2014, 3, 280–285
35. Kovaleski JE, Heitman RJ, Andrew DP, Gurchiek LR, Pearsall AW. Relationship between closed-linear-kinetic-and open-kinetic-chain isokinetic strength and lower extremity functional performance. *Journal of Sport Rehabilitation*. 2001 Aug 1;10(3):196-204. <https://doi.org/10.1123/jsr.10.3.196>
36. Young, W.B.; James, R.; Montgomery, I. Is muscle power related to running speed with changes of direction? *J. Sports Med. Phys. Fit.* 2002, 42, 282–288.
37. Meylan CM, Cronin JB, Oliver JL, Hopkins WG, Contreras B. The effect of maturation on adaptations to strength training and detraining in 11–15-year-olds. *Scandinavian journal of medicine & science in sports*. 2014 Jun;24(3):e156-64. <https://doi.org/10.1111/sms.12128>
38. Harrison CB, Eisenmann J, Knight CJ. 17 Creating a holistic environment for young athletes. *Strength and Conditioning for Young Athletes: Science and Application*. 2019 Aug 16:362.
39. Thieschäfer L, Büsch D. Development and trainability of agility in youth: A systematic scoping review. *Frontiers in Sports and Active Living*. 2022:340. <https://doi.org/10.3389/fspor.2022.952779>
40. Bosco C. Valoraciones funcionales de la fuerza dinámica, de la fuerza explosiva y de la potencia anaeróbica aláctica con los test de Bosco. *Apunts Medicina de l' Esport (Castellano)*. 1987 Sep 1;24(093):151-6.
41. Bishop, C.; Read, P.; Chavda, S.; Jarvis, P.; Turner, A. Using Unilateral Strength, Power and Reactive Strength Tests to Detect the Magnitude and Direction of Asymmetry: A Test-Retest Design. *Sports* 2019, 7, 58. <https://doi.org/10.3390/sports7030058>
42. Hills AP, Hennig EM, Byrne NM, Steele JR. The biomechanics of adiposity—structural and functional limitations of obesity and implications for movement. *Obesity reviews*. 2002 Feb;3(1):35-43. <https://doi.org/10.1046/j.1467-789X.2002.00054.x>
43. Ramos S, Volossovitch A, Ferreira AP, Fragoso I, Massaça L. Differences in maturity, morphological and physical attributes between players selected to the primary and secondary teams of a Portuguese Basketball elite academy. *Journal of sports sciences*. 2019 Aug 3;37(15):1681-9. <https://doi.org/10.1080/02640414.2019.1585410>