

Influence of Harvesting Stage on the Phytochemical Composition, Antioxidant, and Antidiabetic Activity of Immature *Ceratonia siliqua* L. Pulp from Béni-Mellal-Khenifra Region, Morocco: In Silico, In Vitro, and In Vivo Approaches.

[Salah Laaraj](#)^{*}, Hanane Choubbane, [Aziz Tikent](#), [Amal Elrherabi](#), [Ayoub Farihi](#), Meriem Laaroussi, [Mohamed Bouhrim](#), [Abdelaaty A. Shahat](#), [Younes Noutfia](#), [Rashed N. Herqash](#), [Fatiha Chigr](#), [Souad Salmaoui](#), [Kaoutar Elfazazi](#)^{*}

Posted Date: 26 August 2024

doi: 10.20944/preprints202408.1777.v1

Keywords: immature carob pulp; phytochemical compounds; antioxidant activity; in silico, in vitro, and in vivo studies; antidiabetic agent.



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Influence of Harvesting Stage on the Phytochemical Composition, Antioxidant, and Antidiabetic Activity of Immature *Ceratonia siliqua* L. Pulp from Béni-Mellal-Khenifra Region, Morocco: *In Silico*, *In Vitro*, and *In Vivo* Approaches

Salah Laaraj ^{1,2,*}, Hanane choubbane ³, Aziz Tikent ⁴, Amal Elrherabi ⁵, Ayoub Farihi ⁶, Meriem Laaroussi ⁷, Mohamed Bouhrim ⁷, Abdelaaty A. Shahat ⁸, Younes Noutfia ⁹, Rashed N. Herqash ⁸, Fatiha chigr ⁷, Souad Salmaoui ² and Kaoutar Elfazazi ^{1,*}

- ¹ Agri-food Technology and Quality Laboratory, Regional Centre of Agricultural Research of Tadla, National Institute of Agricultural Research (INRA); avenue Ennasr, Bp 415 Rabat principal, Rabat 10090, Morocco.
 - ² Environmental; Ecological, and Agro-Industrial Engineering Laboratory, LGEEAI, Sultan Moulay Slimane University (USMS), Faculty of Science and Technology (FST), Beni Mellal, Morocco.
 - ³ University Cadi Ayyad of Marrakech, FSTG, Laboratory of Sustainable Development and Health, Marrakech, Morocco.
 - ⁴ Laboratoire d'Amélioration des Productions agricoles, Biotechnologie & Environnement (LAPABE), Faculté des Sciences, Université Mohammed Premier, BP: 717, 60000, Oujda, Morocco.
 - ⁵ Laboratory of Bioresources, Biotechnology, Ethnopharmacology, and Health, Faculty of Sciences, Mohammed First University, Oujda, B.P. 717, Morocco.
 - ⁶ Oriental Center for Water and Environmental Sciences and Technologies (COSTE), Mohammed Premier University, Oujda 60000, Morocco.
 - ⁷ Biological Engineering Laboratory, Faculty of Sciences and Techniques, Sultan Moulay Slimane University, Beni Mellal, Morocco.
 - ⁸ Department of Pharmacognosy, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia.
 - ⁹ Fruit and Vegetable Storage and Processing Department, The National Institute of Horticultural Research, Konstytucji 3 Maja 1/3, 96-100 Skierniewice, Poland.
- * Correspondence: salah.laaraj@usms.ma / kaoutar.elfazazi@inra.ma.

Abstract: *Ceratonia siliqua* L. is a medicinal plant that has long been used in traditional Moroccan medicine to treat many diseases. This study aimed to assess the impact of the stages of the immature phase of carob pulp (M1, M2, M3, M4, and M5) on phytochemical composition, antioxidant activity, and antidiabetic activity of *Ceratonia siliqua* L. The identification of the phenolic profile by HPLC-UV/MS-MS and the study of the antidiabetic effect by *in silico*, *in vitro*, and *in vivo* studies were carried out for extracts with a high content of phenolic compounds from immature wild carob pulp from the communes of Timoulit (TM), Bin Elouidane (AW), and Ouaouizerth (TG) in the province of Azilal in the Beni Mellal-Khenifra region. The results revealed a gradual increase in total sugar content over the pulp's ripening period, reaching a value of 2134 ± 56.23 mg GE/100 g fresh matter (FM) for TG. The three locations showed peak values for total polyphenol content (TPC), total flavonoid content (TFC), and total condensed tannin (TCT) at the M2 stage. AW has the highest concentrations of TPC (3819 ± 226.4 mg GAE/100 g FM), TFC (1034 ± 57.08 mg QE/100 g FM), and TCT (1472 ± 28.46 mg CE/100 g FM). The DPPH assay (7892 ± 296.1 mg TE/100 g FM) and the FRAP assay (278.2 ± 7.85 mg TE/100 g FM) both demonstrate that the TG zone is a highly potent antioxidant zone. In contrast, the AW site exhibits a markedly elevated value of 725.4 ± 103.6 mg TE/100 g FM in the ABTS assay. HPLC-UV-MS/MS analysis showed that the methanolic extracts of immature carob pulp (MEICP) from the three areas contained several different chemical compounds. The most prevalent were 3-O-p-coumaroyl-5-O-caffeoylquinic acid, quercetin 3-methyl ether, gallic acid, and galloylquinic acid. Immature carob pulp extract (ICPE) from AW showed the strongest *in vitro* inhibition of pancreatic α -amylase ($IC_{50} = 0.405 \mu\text{g/mL}$) and TG extracts were most potent against intestinal α -glucosidase ($IC_{50} = 0.063 \mu\text{g/mL}$). *In vivo*, AW, TG, and TM extracts significantly reduced postprandial glycemia in rats, with AW having the greatest effect. These results highlight the

antidiabetic potential of ICPE. The 3-O-p-Coumaroyl-5-O-caffeoylquinic acid showed a better affinity for α -amylase compared to acarbose and interacted significantly with several amino acid residues of the enzyme. Similarly, this molecule and 3,4-Dicaffeoylquinic acid demonstrated a strong affinity for α -glucosidase, suggesting their potential as natural inhibitors of enzymes involved in carbohydrate metabolism. Most of the compounds are not substrates of P-glycoprotein and exhibit high intestinal absorption. Furthermore, the majority of these compounds do not act as inhibitors or substrates of CYP450 enzymes, reinforcing their suitability for development as oral medications. These results underscore the potential of immature carob pulp as a promising antidiabetic agent.

Keywords: immature carob pulp; phytochemical compounds; antioxidant activity; *in silico*; *in vitro*; and *in vivo* studies; antidiabetic agent

1. Introduction

Ceratonia siliqua L. is a native tree found in the Mediterranean and Middle Eastern regions. It belongs to the Leguminosae family and is commonly called the Carob tree. This plant is classified as an evergreen and can thrive in its natural habitat or when intentionally cultivated. It has a high level of tolerance towards various soil types. This species is essential for preventing soil erosion, degradation, and desertification due to its deep roots, evergreen leaves, and unique agroecological properties [1]. The carob tree's fruit has recently attracted significant attention, evident from the publication of numerous research publications in the fields of food science and technology that specifically examine food products generated from the carob fruit. These possess exceptional characteristics, including their low-fat composition, abundant dietary fiber, minerals, and a significant quantity of polyphenols [2–4]. On average, the pulp and seeds make up 90% and 10% of the weight of a carob tree fruit, respectively. The chemical composition of the pulp exhibits significant variations based on factors such as the cultivar or species, climate, place of origin, and the stage of ripeness during harvest [5,6]. The pulp and seeds are frequently differentiated based on their high concentration of polyphenolic compounds, which have advantageous impacts on human well-being [7]. The high sugar content of the pulp makes it a valuable product in the culinary industry, where it is utilized to produce syrups and as a chocolate alternative in cookies and ice cream formulations [8–10]. The presence of dietary fiber in it confers numerous advantageous benefits on blood sugar, cholesterol levels, and cancer, making it suitable for use in the pharmaceutical industry [11]. The seed contains a substantial proportion of galactomannan (80–85%) [12], a widely used food additive (E410), that plays a vital role in food technology as it can be utilized as a stabilizer, thickener, and gelling agent. In addition, it finds application in the manufacturing of cosmetics, paint, textiles, and pharmaceuticals [13–15]. The carob fruit is a rich source of natural antioxidants that effectively protect against reactive oxygen species, including both free radicals and non-free radicals. Oxidative stress, resulting from oxidation processes affecting proteins, lipids, and nucleic acids, has been linked to several health problems, including cardiovascular disease, renal disease, and degenerative disease [16]. Polyphenols present in the carob pulp have been demonstrated to exert a range of physiological effects, including improvements in digestive function [17] and reductions in both blood cholesterol [18] and glucose levels [19]. An increase in glucose levels leads to oxidative stress, which in turn increases the non-enzymatic glycation of proteins and glucose oxidation [20]. The efficacy of medications such as sulfonylurea and biguanides in managing diabetes and mitigating its long-term complications has been inadequate [21]. Various medicinal plants and their extracts have shown efficacy in treating diabetes, including the carob fruit. The carob fruit stands out due to its abundant concentration of antidiabetic compounds and antioxidants, specifically flavonoids and gallotannins [22]. According to previous investigations, unripe carob fruits have higher levels of phenolic acids, flavonoids, and tannins compared to ripe fruits. They also exhibit higher levels of *in vitro* antioxidant activity [23–27]. Regarding the carbohydrate content, unripe fruit had lower levels of sugars, whereas ripe fruits had higher amounts of sucrose [28]. Rtibi et al. (2017) found that immature carob pulp aqueous extract exhibits antidiabetic properties. Their findings indicate that the extract inhibits

intestinal glucose transport, thereby reducing blood glucose levels [29]. Based on this information, we proceeded to study the effect of immature stages on the phytochemical composition of unripe carob pulp to consider their use in the pharmaceutical field. The objective of this study was to ascertain the levels of total sugars, total polyphenol content, total flavonoid content, and total condensed tannins in immature carob pulp from the Béni-Mellal-Khnéfra region in Morocco and to evaluate its antioxidant activity. Secondly, the elevated polyphenol content prompted the identification of the respective phenolic profiles and an investigation into the anti-diabetic effects of immature carob pulp during this stage. This was achieved by utilizing *in silico*, *in vitro*, and *in vivo* studies.

2. Materials and Methods

2.1. Plant Materials Harvest

Immature carob pods were collected from three different areas in the Beni Mellal-Khenifra region of Morocco. Timoulilte commune (collection zone: Tanzight, abbreviated as TN, coordinates: 32°11'52.9"N 6°19'52.2"W), Bin Elouidane commune (collection zone: Ait-Waada, abbreviated as AW, coordinates: 32°08'41.1"N 6°30'04.8"W), and Ouaouizerth commune (collection zone: Tizi Ghnim, abbreviated as TG, coordinates: 32°12'24.9"N 6°23'30.8"W). Sampling was carried out between April and June 2022 to track the immature stage of the carob pods. Based on pod dimensions, five different stages (M1, M2, M3, M4, and M5) determined the time of harvest at the immature stage. Thus, ten trees from each area were selected and a total of 50 carob pods were systematically obtained from each tree, by collecting an equivalent number of samples from all four cardinal directions (north, south, east, west) and within the tree itself, with a careful selection process to ensure that only ripe and disease-free carob pods were sampled.

2.2. Preparation of the Immature Carob Juice

On arrival at the laboratory, the collected fruits were separated from the seeds to extract only the pulp, which was then crushed using an electric juice extractor. The collected juice was centrifuged at 4500 rpm for 15 minutes. The supernatant obtained was stored in opaque containers in the refrigerator to avoid oxidation. The carob juice was used for compositional analysis.

2.3. Quantification of Total Sugar Content

The Dubois et al. (1956) method was used to quantify the amounts of sugars in the juice [76]. A mixture was prepared by adding 1 mL of diluted juice (1/100), 5 mL of sulphuric acid, and 1 mL of 5% phenol. After shaking, the mixture was left to stay in the dark environment for almost 45 min. The optical density was then measured at 488 nm using a Shimadzu UV-2401 PC UV spectrophotometer.

2.4. Quantification of Total Phenolic Content

A volume of 500 µL of the diluted juice was added to a test tube. Subsequently, 2.5 mL of Folin-Ciocalteu reagent (10%) and 2 mL of sodium carbonate, 7.5% solution were added at a two-minute interval. The samples were then maintained in a dark environment at a standard laboratory temperature for approximately 90 min before measuring their absorbance at a wavelength of 760 nm using a Shimadzu UV-2401 PC spectrophotometer. The amount of total phenolic contents was calculated by using a standard curve and was expressed as mg gallic acid per 100 g FM [77].

2.5. Quantification of Flavonoid Content

The flavonoid content was assessed using the AlCl₃ procedure [78]. A mixture of 1 mL of carob juice and 1 mL of 2% AlCl₃ was prepared and stored in a dark environment at lab temperature for almost 10 min. Then the absorbance was carefully measured at 430 nm using a Shimadzu-UV-2401 PC spectrophotometer. Following a comparison with the standard curve, the flavonoid content was calculated and expressed as mg catechin per 100 g FM.

2.6. Quantification of Total Condensed Tannins

To determine condensed tannins, present in the carob juice, we adapted the method developed by Broadhurst and Jones (1979) [79]. A solution of 400 μL of diluted juice (1/10) was prepared, with a maximum of 3 mL of 4% vanillin methanol solution added. Subsequently, 1.5 mL of 99% concentrated hydrochloric acid was added. The mixture was then left to stand in the dark at room temperature for approximately 15 minutes, after which the final absorbance was measured at 500 nm using a Shimadzu UV-2401 PC UV spectrophotometer with methanol as the blank.

2.7. Antioxidant Activity Evaluation Assays

2.7.1. 2,2-Diphenyl 1-Picrylhydrazyle Assay

The antioxidant capacity of immature carob pulp juice was determined using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay, by the methodology previously established by Szabo et al. (2019) [80]. Specifically, 0.1 mL of juice was combined with ethanol/water (6:4) at a ratio of 1/1000 and 2 mL of a 0.1 mM DPPH solution. The mixture was shaken and then incubated in complete darkness for approximately 45 minutes. The absorbance of DPPH was then measured at a specified wavelength of 517 nm using a Shimadzu UV-2410 PC spectrophotometer. For comparison, a standard of Trolox (1-5 mg/mL) was employed ($R^2 = 0.9706$), and the final values of free radical scavenging activity of juice samples were expressed as milligrams of Trolox equivalents per 100 g FM.

2.7.2. ABTS Scavenging Assay

The antioxidant capacity was determined by performing the 2,2'-azino-bis (3-éthylbenzothiazoline-6-sulfonate) (ABTS) assay by the protocol established by Uysal et al. (2016) [81]. Initially, a 7 mM ABTS solution was combined with 2.45 mM potassium persulfate and incubated in the dark at room temperature for 12–16 hours. Before the commencement of the experiment, the solution was diluted with ethanol to achieve an absorbance of 0.700 ± 0.02 at 734 nm. To ascertain the ABTS scavenging activity in milligrams of Trolox equivalents per 100 g of FM, 1 mL of extract was combined with 2 mL of ABTS solution and incubated at room temperature for 30 minutes. The absorbance was then measured at 734 nm. To ensure accuracy, triplicate measurements were conducted.

2.7.3. Ferric Reducing Antioxidant Power Assay

In their study, Benzie and Strain (1996) describe the preparation of the FRAP reagent, which was created by combining 0.3 M acetate buffer (pH 3.6), 10 mmol/L TPTZ, and 40 mmol/L FeCl_3 in a 10:1:1 ratio [82]. The juice was combined with 3 mL of the freshly prepared FRAP reagent, incubated for 30 minutes, and vortexed. The absorbance was measured at 593 nm, and the results were expressed in milligrams of Trolox equivalents per 100 mL of FM, using Trolox as a standard from 1 to 5 mg/mL ($R^2 = 0.9706$).

2.8. Identification of Unripe Carob Pulp Juice Phytochemical Compounds

2.8.1. Solubilization of Polyphenols of Unripe Carob Pulp Juice

The polyphenols were extracted using the process of dissolving them in methanol [83]. The juices extracted from unripe carob pulp were subjected to vacuum dehydration and then transformed into powder using a vacuum oven (Memmert UM 400, made in Germany). 5 g of immature carob pulp powder in 50 mL of aqueous methanol (70%, v/v) was prepared and left for 24 hours to obtain the methanolic extracts of immature carob pulp (MEICP).

2.8.2. HPLC- UV-MS/MS Analysis

Extracts with a high phenolic compound content from TM, AW, and TG unripe carob pulps were analyzed qualitatively using a liquid chromatography system coupled with a triple quadrupole mass

spectrometer (Thermo Fisher Scientific, San Jose, CA, USA). A Kinetex C18 reversed-phase column (100 x 4.6 mm, 2.6 μ m particles) was employed, with a gradient separation of solvent A (0.1% formic acid aqueous solution) and solvent B (methanol), as described in the study of Bouaouda et al. (2023) [58]: 0–3 min, linear gradient from 5% to 25% B; 3–6 min, at 25% B; 6–9 min, from 25% to 37% B; 9–13 min, at 37% B; 13–18 min, from 37% to 54% B; 18–22 min, at 54% B; 22–26 min, from 54% to 95% B; 26–29 min, at 95% B; 29–29.15 min, returning to initial conditions at 5% B, and re-stabilizing the column from 29.15 to 31 min to the initial gradient 5% B. A concentration of 1 mg/mL of each standard were analyzed with the same conditions. The identification of phenolic compounds relied on comparing retention time and spectra with nine standards, along with referencing the NIST-MS/MS library.

2.9. Animals

This study was conducted on albino mice (male and female). The animals were raised in the animal facility of the Department of Biology at the Faculty of Science and Technology of Béni Mellal, with a photoperiod of 12 hours of light/12 hours of darkness and a temperature of 22 ± 2 °C. The animals were kept in favorable breeding conditions with free access to water and food. All rats were cared for in compliance with the internationally accepted Guide for the Care and Use of Laboratory Animals, published by the US National Institutes of Health, 2011 [84].

2.10. Acute Toxicity Test

For each sample, two groups of fasting albino mice (25-35 g, 14h) were divided into four groups (n=6; $\sigma/\varnothing=1$): a control group gavaged with distilled water (10 ml/kg) and three treated groups gavaged with plant extract at doses of 2 g/kg, 1 g/kg, or 0.5 g/kg. At the beginning of the experiment, the mice were weighed and then gavaged with a single dose of the plant extract. They were monitored continuously for 10 hours to detect any obvious signs of toxicity, and then daily for 14 days to observe any additional clinical or behavioral signs of toxicity.

2.11. Evaluation of the Inhibitory Effect of Unrip Carob Pulp on Pancreatic α -Amylase Activity

2.11.1. In Vitro Assay

To investigate the in vitro inhibition of pancreatic α -amylase, we conducted antidiabetic tests on digestive enzymes using extracts from the stages that yielded the greatest concentrations of polyphenols and flavonoids, along with the most substantial antioxidant effects. This approach was based on an adapted version of the procedure outlined by Elrherabi et al. (2023) [85]. The test mixtures were prepared by combining 200 μ L of a 0.02 M phosphate buffer solution (pH 6.9), and 200 μ L of α -amylase enzyme solution, and incubating the mixture at 37 °C for 10 minutes. Subsequently, 200 μ L of P1, P2, P3, P4, and P5 solutions (at concentrations of 0.062, 0.125, 0.25, 0.5, and 1 mg/mL) were added to the respective tubes. Following this, 200 μ L of a 1% starch solution was added to each tube, and the mixtures were incubated at 37 °C for 15 minutes. To terminate the enzymatic reaction, 600 μ L of 3,5-dinitrosalicylic acid (DNSA) reagent was added, and the tubes were heated in an incubator at 100 °C for 8 minutes. After this thermal treatment, the tubes were cooled in an ice-water bath, and 1 mL of distilled water was added to dilute the samples. The absorbance of the samples was recorded at 540 nm with a spectrophotometer. To measure 100% enzyme activity, a blank was prepared by replacing the extract with 200 μ L of phosphate buffer. Additionally, another blank was set up by applying the plant extract at various doses without adding the enzyme solution. Acarbose was employed as a positive control at comparable concentrations, with the reaction process mirroring that of the plant extract. The inhibition ratio was then computed using the formula provided below:

$$\% \text{ of Inhibition} = \frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \times 100$$

A_{Control} : Absorbance of enzymatic effect in the absence of an inhibitor.

A_{Sample} : Absorbance of enzymatic effect in the presence of extracts or acarbose.

The IC₅₀ (concentration of samples that inhibit 50% of α -amylase enzyme activity) was calculated graphically using the formula: percentage of inhibition = f (log (sample concentration)).

2.11.2. In Vivo Assay

The oral starch tolerance test was utilized to evaluate the antihyperglycemic effect of the unripe carob pulp, *in vivo*. Normal mice were divided into 3 groups (n=6; σ/φ =1). Control Group: Normal mice administered with distilled water (10 mL/kg), Test Group: Normal mice administered with the plant aqueous extract (400 mg/kg), and Medication Group: Normal mice administered with acarbose (10 mg/mL). The test was conducted as follows: a blood glucose measurement was taken at t₀. Immediately afterward, the test product (ED, the aqueous extract, or acarbose) was administered. Thirty minutes later, a second blood glucose measurement was taken, and then the mice were orally loaded with starch (3 g/kg). Subsequently, the variation in blood glucose levels was monitored for 2 hours at 60, 90, and 120 minutes.

2.12. Assessment of the Inhibitory Effect of Unripe Carob Pulp on Intestinal α -Glucosidase Activity

2.12.1. In Vitro Assay

The methodology for studying the inhibition of α -glucosidase activity adhered to the method described by Elrherabi et al. (2023b) [85]. The assay mixtures consisted of 1 mL of phosphate buffer (pH = 7.5), 0.1 mL of α -glucosidase enzyme solution (10 IU), and 200 μ L of E.E, AG, or M (at concentrations of 0.031, 0.062, 0.125, 0.25, 0.5, and 1 mg/mL), with a corresponding volume of distilled water. Acarbose (at concentrations of 0.041, 0.082, 0.165, 0.328, and 0.656 mg/mL) served as the control, including appropriate negative and positive controls. The mixtures underwent pre-incubation for 20 minutes at 37 °C. Subsequently, 0.1 mL of sucrose was introduced to each tube, and the reaction was halted by incubating the tubes at 100 °C for 5 minutes. Following this, 1 mL of GOD-POD (Glucose oxidase-peroxidase) was added, and the tubes were further incubated at 37 °C for 10 minutes. The absorbance was recorded at 500 nm using a spectrophotometer. The inhibition ratio was calculated using the formula:

$$\% \text{ of Inhibition} = \frac{A \text{ control} - A \text{ sample}}{A \text{ control}} \times 100$$

Where:

A control: is the absorbance of enzymatic activity without the inhibitor.

A sample: is the absorbance of enzymatic activity in the presence of the extract or acarbose.

The concentration of samples required to inhibit 50% (IC₅₀) of α -amylase enzyme activity was determined graphically using the formula: inhibition percentage = f (log sample concentration).

2.12.2. In Vivo Assay

The oral sucrose tolerance test was employed to evaluate the antihyperglycemic impact of the immature carob pulp, *in vivo*. Normal mice were segregated into 3 groups (n=6; σ/φ =1). Control Group: Normal mice administered with distilled water (10 mL/kg), Test Group: Normal mice administered with the plant aqueous extract (400 mg/kg), and Medication Group: Normal mice administered with acarbose (10 mg/mL). The test was conducted as follows: a blood glucose measurement was taken at t₀. Immediately afterward, the test product (ED, the plant aqueous extract, or acarbose) was given. After thirty minutes, a second blood glucose measurement was taken, and then the mice were orally loaded with sucrose (2 g/kg). Subsequently, the variation in blood glucose levels was monitored for 2 hours at 60, 90, and 120 minutes.

2.13. Molecular Docking Analysis

2.13.1. Ligand Preparation

The phytochemicals detected by HPLC-UV-MS/MS in methanolic extracts of immature carob pulp (ICPME), as well as the drug Acarbose (CID: 41774) (a standard inhibitor), were sourced in 3D

SDF format from the PubChem database (available at <https://pubchem.ncbi.nlm.nih.gov/>). However, the 3D model of 3-O-p-coumaroyl-5-O-caffeoylquinic acid was created using Avogadro 1.2 software. To assess the potential of these ligands as inhibitors of intestinal α -glucosidase and pancreatic α -amylase, the ligands were first converted to pdb format using PyMOL Molecular Graphics System (version 2.5.3) [86], and then further processed to pdb format using AutoDockTools (ADT; version 1.5.7, The Scripps Research Institute) [87]. Visualization of the molecular interactions between the ligands and receptors, including their binding affinities compared to the standard inhibitor, was achieved with Discovery Studio Visualizer (Biovia, 2021), aiding in the generation of the graphics [86].

2.13.2. Protein Preparation

The crystal structures of human α -glucosidase (PDB ID: 5NN8) and α -amylase (PDB ID: 1B2Y) were obtained from the Protein Data Bank (PDB) (accessible at www.rcsb.org) [88]. Each crystal structure was individually processed by removing water molecules and adding polar hydrogens and Kollman charges using AutoDockTools (ADT; version 1.5.7). For molecular docking, a grid with a point spacing of 0.375 Å and dimensions of 40 × 40 × 40 was created [89]. This grid was positioned to cover both the peripheral regions and active sites of the proteins, based on the x, y, and z coordinates. The processed macromolecules were then saved in PDB format for further molecular analysis.

2.14. ADME Studies

Understanding pharmacokinetic properties, such as absorption, distribution, metabolism, and excretion (ADME), is vital for determining how a substance behaves in the body [90]. These stages describe the path a substance takes from its absorption to its elimination. Computational tools have become indispensable for predicting the ADME characteristics of molecules. They allow for the evaluation of a molecule's ability to cross cellular barriers, interact with key transporters and enzymes involved in absorption and excretion, and assess its metabolic stability [91]. In our molecule evaluation process, we opted to use the SwissADME platform. This platform allows for a comprehensive analysis of the physicochemical properties of phytochemicals identified in methanolic extract of unripe carob pods, their potential as therapeutic agents, and their pharmacokinetic characteristics, providing a thorough understanding of their ADME profile [91].

2.15. Statistical Analysis

The analyses were conducted in duplicate to get three results, which were then reported as the means and standard deviations. The statistical analysis was performed using the one-way ANOVA method in GraphPad PRISM 8.0.1 software. Using Duncan's multiple-range test, the mean values were separated with a level of significance as $P < 0.05$.

3. Results

3.1. Total Sugar Content

Table 1 presents the total sugar content at different stages of maturity harvested in different areas of the province of Azilal, Morocco. During the immature stage, the sugar content of all sources increased, with the highest amounts observed in TG and AW at M5 (2134 ± 56.23 mg GE/100g FM and 2098 ± 34.57 mg GE/100g FM, respectively). Nevertheless, M3 exhibited the highest TM value (1987 ± 71.89 mg GE/100g FM). The overall sugar analysis revealed no statistically significant differences ($p > 0.05$) between TG, TM, and AW, except for the M3 stage, where TG differed from TM and AW. However, during the immature stage, we observed significant differences among the zones themselves, the variation in total sugar content across the three regions may be attributed to variations in their geographical provenance, meteorological factors, and soil properties [30]. There is a limited amount of research that has specifically investigated the overall sugar content of unripe carob pulp. The total sugar content in TG, TM, and AW was found to be lower than that reported by

Benchikh et al. (2016), with values ranging from 15.98 ± 0.04 to 18.93 ± 0.07 g/100 g DW [24]. Vekiari et al. (2012) conducted research that demonstrated a substantial increase in the sugar content of two different types of maturing Greek carob pods [31]. As stated by Prasanna et al. (2007), the rise in sugar content and sweetness is a natural occurrence during the maturation of numerous fruits. The fruit softens due to the enzymatic breakdown of cell wall polysaccharides through hydrolysis [32]. The low sugar content of fruit can be beneficial for blood sugar management, which is essential for people with diabetes. Furthermore, a study conducted on the polysaccharides in baobab fruit demonstrated that they possess notable anti-diabetic properties by inhibiting enzymes such as α -amylase and α -glucosidase, which are involved in the breakdown of carbohydrates. This inhibition results in a slower release of glucose into the bloodstream, facilitating the regulation of blood sugar levels [33].

Table 1. Change in total sugar content of unrip carob pulp during ripening stages.

Total sugar mg GE/100 g FM			
	TG	TM	AW
M1	1720±76.22 ^{dA}	1709±72.63 ^{cA}	1889 ± 124.8 ^{bA}
M2	1814±84.36 ^{cdA}	1721±91.97 ^{bcA}	1636 ± 36.63 ^{cA}
M3	1880±7.31 ^{bCB}	1987±71.89 ^{aA}	1982 ±55.23 ^{abA}
M4	2032±22.35 ^{bA}	1880±25.34 ^{abA}	1940±27.93 ^{abA}
M5	2134±56.23 ^{aA}	1904 ± 20.8 ^{aA}	2098 ±34.57 ^{aA}

The results are shown as the mean value ±SD (n=3). Distinct lowercase letters indicate statistically significant variations (p<0.05) across unripening phases, while distinct capital letters indicate statistically significant variations (p<0.05) between provenances as determined by the Tukey test. TG., Tizi-Ghnm site; TM., Timouliit site; AW., Ait Waarda site; (M1, M2, M3, M4, and M5), stages of the immature phase of carob pulp; GE., Glucose equivalent; FM., Fresh Matter.

3.2. Total Phenolics, Flavonoids, and Condensed Tannins Contents

Table 2 presents the results for bioactive compounds in immature carob pulp, with a focus on total polyphenol content (TPC), total flavonoid content (TFC), and total condensed tannins (TCT) for the three zones (TG, TN, and AW) in the five immature stages. The TPC for TG, TN, and AW ranged from 2532 ± 48.3 to 3475 ± 189.6 mg GAE/100 g FM, 2120 ± 150.9 to 3819 ± 226.4 mg GAE/100 g FM, and 1947 ± 55.37 to 2765 ± 144.3 mg GAE/100 g FM, respectively. The M2 stage demonstrated the greatest TPC value among the three zones tested, whereas the M1, M5, M3, and M4 stages exhibited TPC values in a declining order.

Table 2. Change in total phenolic content, total flavonoid content, and total condensed tannins of unrip carob pulp during ripening stages.

	TPC mg GAE/100g FM			TFC mg QE/100g FM			TCT mg CE/100g FM		
	TG	TN	AW	TG	TN	AW	TG	TN	AW
M1	2813 ± 58.02 ^{bA}	3014 ± 59.13 ^{bA}	2734 ± 99.24 ^{aA}	826.2 ± 40.77 ^{bB}	298.7 ± 20.1 ^{cC}	1423 ± 78.47 ^{aA}	613.9 ± 74.18 ^{bA}	231.2 ± 69.86 ^{cB}	567.9 ± 21.35 ^{bA}
M2	3475 ± 189.6 ^{aB}	3819 ± 226.4 ^{aA}	2765 ± 144.3 ^{aC}	925.4 ± 13.27 ^{aAB}	1034 ± 57.08 ^{aA}	830.1 ± 87.62 ^{bB}	848.1 ± 18.91 ^{aB}	1472 ± 28.46 ^{aA}	756.1 ± 92.54 ^{aC}

M3	2750	2368	2209	900.6	827.5	848.4	628.5	633.7	539.6
	±	±	±	±	±	±	±	±	±
	98.37 ^{bcA}	43.12 ^{cB}	86.94 ^{bc}	65.7 ^{a^bA}	49.6 ^{bA}	52.7 ^{bA}	93.07 ^{bA}	61.17 ^{bA}	35.26 ^{bB}
M4	2532	2120	1947	173.2	196.2	161.2	724.7	706.9	565.8
	±	±	±	±	±	±	±	±	±
	48.3 ^{cA}	150.9 ^{cB}	55.37 ^{cB}	11.25 ^{dA}	10.96 ^{dA}	13.73 ^{cA}	22.69 ^{abA}	37.82 ^{bAB}	18.38 ^{bB}
M5	2744	2391	2595	294	249.6	305.6	75.71	71.42	77.17
	±	±	±	±	±	±	±	±	±
	39.47 ^{bcA}	76.58 ^{cC}	91.21 ^{aB}	12.06 ^{cA}	16.76 ^{cdA}	11.95 ^{cA}	5.52 ^{cA}	3.32 ^{dA}	6.11 ^{cA}

The results are shown as the mean value ±SD (n=3). Distinct lowercase letters indicate statistically significant variations (p<0.05) across unripening phases, while distinct capital letters indicate statistically significant variations (p<0.05) between provenances as determined by the Tukey test. TG, Tizi-Ghnm site; TM., Timouliit site; AW, Ait Waarda site; (M1, M2, M3, M4, and M5), stages of the immature phase of carob pulp; GAE., Gallic acid equivalent; QE., Quercetin equivalent; CE., Catechin equivalent; FM., Fresh Matter.

The levels of TPC observed in the areas studied in this investigation were markedly lower than those reported by Benchikh et al. (2014) (19.82 ± 0.48 g GAE/100 g DW) [23], and almost similar to those found by Moumou et al. (2023) (41.5 ± 1.57 mg/g) [34], whereas our results are higher than those found by Ouzounidou et al.(2012) and Sebai et al. (2013) for the respondent populations in Greece (between 13.6 and 24.8 mg GAE/100g DW) and cultures in Tunisia (28.07 ± 0.99 mg/kg), respectively [35,36].

The statistical analysis revealed a significant decrease (p < 0.05) in the total phenolic content of TG, TN, and AW from M1 to M3. However, there was a slight increase in AW from M4 to M5. It is important to mention that there was considerable variation between zones for each stage, except for the M1 stage. Gull et al. [37] found that the decrease in polyphenolic content in carob fruit is due to the oxidation of phenolic compounds by polyphenol oxidase as the fruit approaches maturity. This leads to the conversion of soluble phenols into insoluble phenolic compounds that bind to the cell walls' polysaccharides. The TFC results demonstrated that the TG values ranged from 173.2 ± 11.25 to 925.4 ± 13.27 mg QE/100 g FM, while the TN values exhibited a broader range, from 196.2 ± 10.96 to 1034 ± 57.08 mg QE/100 g FM. AW values exhibited a similar range, spanning from 161.2 ± 13.73 to 1423 ± 78.47 mg QE/100 g FM. The results obtained in this study are significantly higher than those reported by Sebai et al. (2013), who reported a mean value of 6.14 ± 1.03 mg/kg [36]. Moreover, the results presented include the findings reported by Benchikh et al. (2014), which span a range of 263.81 to 320.24 mg QE/100g DW [23], as well as those reported by Ben Othmen et al. (2019a) (ranging from 275.91± 2.54 to 562.26 ± 8.20 mg RE/ 100g DW) [38], and Moumou et al. (2023) reported a value of 410 ± 140 mg QE/100 g DW [34]. The results indicate a notable decline in flavonoid levels from M1 to M4, followed by a slight increase at M5 in all three studied zones. Furthermore, in the initial three stages, there is a significant difference in flavonoid levels among the three zones. The role of flavonoids in protecting the photosynthetic apparatus may account for their greater abundance during the early stages of the immature phase compared to the final stage. According to Harborne and Williams (2000), flavonoids play a protective role in photosynthetic tissues against damage. When chlorophylls break down, the photosynthetic process ceases, leading to a decrease in TFC in the carob pod during ripening [39]. Flavonoids are also renowned for their antioxidant properties, which may assist in reducing the oxidative stress associated with diabetes. This diminished oxidative stress may facilitate enhanced insulin sensitivity and glucose metabolism. Furthermore, flavonoids may facilitate an increase in insulin secretion from pancreatic cells, thereby providing additional support for regulating blood sugar levels [33].


The Total condensed tannins (TCT) analysis reveals that the juices extracted from the immature carob pulp of TG, TN, and AW have the highest concentrations at the M2 stage, with values of 848.1 ± 18.91, 1472 ± 28.46, and 756.1 ± 92.54 mg CE/100 g FM, respectively. On the other hand, the M5 stage yields the lowest concentrations, measuring 75.71 ± 5.52, 71.42 ± 3.32, and 77.17 ± 6.11mg CE/100 g

FM, respectively. The results show that TN has the highest concentration of condensed tannins, which indicates its potential role in a range of biological activities. Only Ben Othmen et al. (2019) and Saci et al. (2020) have investigated the changes in tannin levels in maturing carob pulp to our knowledge. The condensed tannin levels reported in those studies (ranging from 1.18 ± 0.03 to 6.67 ± 0.05 g CE/100 g DW and 3.84 to 16.24 g CE/g DM, respectively) were similar to the range observed in our study [28,40]. The concentration of condensed tannins decreased significantly as the carob fruit matured from the M1 to M4 stages, and there was a significant decrease at the M5 stage. Substantial polymerization and condensation processes are responsible for the decrease in tannin content during carob ripening [41]. The obtained results suggest that factors such as ecological conditions during harvesting, genotype, growth conditions, genetic diversity, and fruit maturity stage may link to the variations in TPC, TFC, and TCT in immature carob pulp [41–43]. The high levels of polyphenols, flavonoids, and gallotannins in carob have demonstrated that immature carob can inhibit intestinal glucose absorption in addition to exerting other beneficial effects, including a potent antioxidant action [44].


3.3. Antioxidant Activities

The main role of bioactive compounds in plant extracts is to prevent diseases and promote human health. These compounds have gained significant attention in recent years because of their ability to act as antioxidants. The results shown in Figure 1 demonstrate that the unripe carob pulp is proficient in efficiently trapping free radicals across all phases of analysis.


The highest DPPH antioxidant capacity of immature carob pulp for the three zones (TG, AW, and TM) (Figure 1a) was exhibited by Stages M2 (7892 ± 296.1 , 7458 ± 178.4 , 5986 ± 205.9 mg TE/100g FM), and M5 (7844 ± 249.1 , 7786 ± 64.59 , and 6772 ± 99.37 mg TE/100g FM). The results exceeded those reported by Benchikh et al. (2014) (30.37 ± 0.03 g AAE/100 g DW) [23], and Ben Othmen et al. (2019) (20.98 ± 0.15 – 32.78 ± 0.78 g TE/ 100g DW) [38]. The obtained values demonstrate a statistically significant difference ($p < 0.05$) primarily between the two zones TG, AW, and TN. The majority of the analyzed internships show a high degree of similarity between TG and TN. The capacity to capture free radicals may decrease in stages M1, M3, and M4 due to the declining levels of phenolic compounds caused by fruit growth. There is a lack of published data on the influence of the ripening stage on carob's antioxidant activity. We also evaluated the plant extracts for their ability to scavenge free radicals using the ABTS radical decolorization assay. This assay is based on the reduction of ABTS+• radicals by the antioxidants present in the tested plant extracts. As depicted in Figure 1b, the outcomes were largely comparable to the DPPH activity, except TG, where the activity exhibited a decline after M3. Stage M2 had the highest amount of ABTS for TG, AW, and TM, with values of 705.7 ± 37.27 , 725.4 ± 103.6 , and 733.2 ± 35.76 mg TE /100g FM, respectively. The lowest levels of ABTS were observed in M5 for TG (624.0 ± 15.94 mg TE/100g FM), M3 for AW (607.4 ± 78.94 mg TE/100g FM), and M4 for TM (598.1 ± 55.78 mg TE/100g FM). In acidic conditions, the FRAP assay checks how well antioxidants work at changing Fe³⁺ to Fe²⁺ [45]. We acquired the results using the Trolox calibration curve. The antioxidant capacity of TG ranged from 113 ± 14.28 to 278.2 ± 7.85 mg TE /100g FM, AW varied between 79.52 ± 10.92 and 189.5 ± 18.62 mg TE /100 g FM, while the TM ranged from 56.32 ± 11.70 to 133.6 ± 16.22 mg TE /100 g of FM (Figure 1c). The TG, AW, and TM showed the highest reducing ability with values of 278.2 ± 7.85 mg TE /100g FM, 189.5 ± 18.62 mg TE /100g FM, and 133.6 ± 16.22 mg TE /100g FM, respectively. Comparable to the results observed in DPPH assays. According to Ferreira-Santos et al. (2024), the variations in antioxidant capacity observed in the three zones studied can be attributed to the chemical structure of the radical molecules and the specific phenolic compounds present in each residue [46]. In addition, variables such as residue diversity, and ripening stage can also have an impact on antioxidant capacity [47]. Our results indicate that immature carob pulp from the three study sites has high antioxidant activity. Consequently, it may be a valuable source for the prevention or mitigation of diseases related to oxidative stress. The pronounced antioxidant activity of carob may be attributed to its high concentration of polyphenolic compounds. These findings confirm that carob has the potential to serve as a rich source of natural antioxidants, which may play a key role in maintaining health and preventing disease [44].



(a)



(b)



(c)

Figure 2. The antioxidant activity of unripe carob pulp during ripening stages using DPPH (a), ABTS (b), and FRAP (c) tests. The results are shown as the mean value \pm SD ($n=3$). Distinct lowercase letters indicate statistically significant variations ($p<0.05$) across unripening phases, while distinct capital letters indicate statistically significant variations ($p<0.05$) between provenances as determined by the Tukey test. TG, Tizi-Ghnm site; TM., Timoulit site; AW, Ait Waarda site; (M1, M2, M3, M4, and M5), stages of the immature phase of carob pulp; TE., Trolox equivalent; FM., Fresh Matter.

3.4. HPLC- UV-MS/MS Analysis

The methanolic extracts of immature carob pulp (MEICP) from the M2 stage were selected for study because of their high polyphenolic content. The HPLC-UV-MS/MS method was used to determine the phenolic profile and identify the phenolic compounds present in the MEICP from the three zones.

A total of seven peaks were identified through HPLC analysis as part of our investigation. 3-O-p-coumaroyl 5-O-caffeoylquinic acid, quercetin 3'-methyl ether, gallic acid, galloylquinic acid, and 3,4-dicaffeoylquinic acid were the main chemicals found in MEICP from all three areas (Table 3, Figure 2). When examining the individual components, 3-O-p-coumaroyl 5-O-caffeoylquinic acid has almost identical percentages in different locations. Specifically, TM and TG have percentages of 53.40% and 55.84%, respectively, whereas AW has a greater proportion, up to 63.62% of this acid. Quercetin 3-methyl ether is present at similar percentages in TM (13.87%), AW (12.91%), and TG (13.83%). Conversely, galloylquinic acid is only present in TM (12.55%). Gallic acid is particularly high in TM (11.24%), whereas it is relatively low in AW and TG (5.17% and 2.35% respectively). All areas contain 3,4-Dicaffeoylquinic acid, with TG exhibiting comparatively greater proportions than TM and AW. In their study, Can et al. (2020) investigated the protective effects of *Daphne gnidioides* Jaub and Spach extracts on DNA, their antibacterial properties, their potential in combating cancer, and their ability to inhibit enzyme activity. They concluded that significant amounts of vicenin-2 and 3-O-p-coumaroyl-5-o-caffeoylquinic acid in these extracts could be associated with these biological activities [48]. To our knowledge, there has been no research into the potential antidiabetic properties of this particular molecule. Several plants such as *Allagopappus viscosissimus* [49], *Opuntia ficus-indica* var. *saboten* [50], *Lychnophora staavioides* [51], and *Rhamnus* species contain the flavonoid quercetin-3-methyl ether, which, to the best of our knowledge, has not previously been reported in carob pod. Studies have shown that quercetin-3-methyl ether has anti-inflammatory [52], antioxidant [53], significant anticancer-promoting properties [54], antitrypanocidal, antimutagenic, tracheal relaxing, radical scavenging, and xanthine oxidase inhibitory properties [51]. The research conducted by Motta et al. (2017) discovered that galloylquinic acid, which was detected in zone TM (12.55%), was

primarily responsible for the gastroprotective effect [55]. Furthermore, galloylquinic acid exhibited a considerable level of cytotoxicity against stomach cancer cells when tested in isolation [55]. TM extract contains substantial amounts of gallic acid, while TG and AW only contain minor amounts. Multiple studies have demonstrated that this compound, which is prevalent in various plant compounds, possesses antitumor properties [56], and acts as an antioxidant, safeguarding against a variety of diseases. This includes the prevention of neurodegeneration resulting from oxidative stress and other neuroprotective mechanisms. It is thought that gallic acid could be a useful first compound for treating several neurodegenerative diseases, such as brain tumors, glioblastoma, cerebral ischemia/reperfusion, anxiety, memory loss, depression, sedation, psychosis, neuropathic pain, Alzheimer’s disease, Parkinson’s disease, and neuroinflammation [57]. The variation in compound percentages indicates potential regional factors that may affect the chemical makeup of carob pods. In some areas, compounds such as galloylquinic acid and ferulic acid are noticeably absent or found in very small quantities, suggesting the specificity of their distribution. These findings demonstrate the significance of considering geographical variables when examining the chemical composition of natural extracts. They also give us useful information for future research on the bioactivity and possible uses of carob pods from various geographical origins [28].

Table 3. HPLC-UV MS/MS phenolic profile of unripe carob jus from various locations (TG, TM, and AW).

N°	Compounds		Chemical formula	RT (min)	%AIR			Ref
					TM	AW	TG	
1	3-O-p-coumaroyl caffeoylquinic acid	5-O-	C ₂₅ H ₂₄ O ₁₁	4.28	53.40	63.62	55.84	N
2	Quercetin 3-methyl ether		C ₁₆ H ₁₂ O ₇	5.26	13.87	12.91	13.83	N
3	Gallic acid		C ₇ H ₆ O ₅	6.60	11.24	5.17	2.35	S
4	3,4-Dicaffeoylquinic acid		C ₂₅ H ₂₄ O ₁₂	8.43	3.04	2.85	5.06	N
5	Galloylquinic acid		C ₁₄ H ₁₆ O ₁₀	10.95	12.55	n.d	n.d	(1)
6	Ferulic acid		C ₁₀ H ₁₀ O ₄	13.52	0.34	0.37	0.60	S
7	Catechine		C ₁₅ H ₁₄ O ₆	17.84	n.d	0.37	n.d	S

RT.: Retention time; ND.: Not detected; ref.: reference, N.: Nist/MSMS; S.: Standard; TG.: Tizi-Ghnm site; TM.: Timoulilt site; AW.: Ait Waarda site; (1): [58].




Figure 2. Phenolic profile of the TN, AW, and TG unripe carob pods sites. RT., Retention time; TG., Tizi-Ghnm site; TM., Timoulilt site; AW., Ait Waarda site.

3.5. Acute Safety

The results of the acute toxicity evaluation show that none of the three extracts demonstrated any toxicity, even at a dose of 2 g/kg. No toxic effects such as diarrhea, vomiting, or abnormal mobility were observed, and no deaths were recorded during the entire observation period (Table 4). A study by Moumou et al. (2023) investigated the acute toxicity of a single dose of immature locust bean pod extract at 2 and 5 g/kg body weight in mice. The results indicated that immature carob pod extract is non-toxic with an LD50 greater than 5 g/kg body weight [34].

Table 4. A comprehensive overview of additional pre-clinical or behavioral signs of acute toxicity.

	2 g/kg	1 g/kg	0.5 g/kg
Doses			
Symptoms			
Locomotion and mobility	Normal	Normal	Normal
Hair loss or erection	-	-	-
Diarrhea	-	-	-
Abnormal agitation	-	-	-
Anorexia	-	-	-
Spontaneous startle	-	-	-
Isolation in the corner	-	-	-
Mortality	-	-	-
Body weight gain	Normal	Normal	Normal

-: the absence or non-occurrence of the sign.

3.6. *In Vitro*, Inhibitory Effect of the Plant Extract on the Pancreatic α -Amylase and Intestinal α -Glucosidase Activities

3.6.1. Pancreatic α -Amylase

Among the many digestive enzymes in humans, pancreatic α -amylase (EC 3.2.1.1) is particularly crucial. It drives the hydrolysis of α -1,4 glycosidic bonds in starch, facilitating its digestion. Since large starch molecules cannot cross the blood-brain barrier, α -amylase breaks them down into smaller sugar fragments that can enter the bloodstream and subsequently reach the brain. If starch is converted into sugars excessively, blood sugar levels rise, prompting insulin to instruct cells to metabolize and store the excess sugar as glycogen. This process maintains equilibrium in a healthy individual. However, excessive amylase activity coupled with insulin deficiency or resistance can lead to elevated blood glucose levels, potentially causing hyperglycemia. Inhibiting this enzyme can slow down the digestion of dietary polysaccharides, this process decreases the uptake of carbohydrates in the intestines and reduces post-meal blood sugar levels. Consequently, research into inhibiting amylase activity is ongoing to manage and control hyperglycemia [59].

The inhibitory effect on α -amylase activity observed in vitro by immature carob pulp extract (ICPE) from the three regions TM, AW, and TG shown in Table 5 and Figure 5 notable differences. The ICPE from AW demonstrated the most potent inhibition with an IC_{50} of $0.405 \pm 0.247 \mu\text{g/mL}$, followed closely by TM with an IC_{50} of $0.450 \pm 0.239 \mu\text{g/mL}$, and TG with an IC_{50} of $0.591 \pm 0.104 \mu\text{g/mL}$. Luísa Custódio et al. (2015) revealed that at a concentration of 1 mg/mL, decoctions from the leaves (74%) and stem bark (65%) of *C. siliqua* L. had potent α -amylase inhibition, surpassed the standard α -amylase inhibitor (Acarbose) [60]. This is the first time we have assessed the impact of the stages of the immature phase of carob pulp (M1, M2, M3, M4, and M5) on digestive enzymes. We concentrated on the stages that produced the greatest quantities of polyphenols and flavonoids, along with the most significant antioxidant activity.

Table 5. Inhibition of α -amylase activity by TG, TM, AW, and acarbose.

Extraits	IC ₅₀ $\mu\text{g/mL}$
TG	0,591 \pm 0.104a
TM	0,450 \pm 0.239 ab
AW	0,405 \pm 0.247 ab
Acarbose (control)	0,098 \pm 0.004 b

Values are presented as mean \pm SEM (n = 3). Data were analyzed using one-way ANOVA with post hoc Tukey's test at a 5% significance level. Means within the same column that have different letters are significantly different (p < 0.05). TG., Tizi-Ghnm site; TM., Timoulilt site; AW., Ait Waarda site.

Germ flour exhibited moderate inhibition (40% at 1 mg/mL, 42% at 10 mg/mL), while locust bean gum and pulps had low or no activity [60]. A separate study reported that the aqueous extract of *C. siliqua* L. demonstrated a significant inhibitory effect on all digestive enzymes [61]. These findings are consistent with Custódio et al. (2015), who observed that both the leaves and stem bark of the carob tree exhibit strong inhibition of α -amylase and α -glucosidase activities, along with notable antioxidant properties [60]. Diabetes mellitus is a metabolic disorder often linked to increased cellular oxidative stress and diminished antioxidant activity. Natural antidiabetic compounds such as polyphenols and flavonoids mitigate free radical formation, decrease oxidative stress, and inhibit digestive enzymes, which in turn helps lower post-meal glucose levels [62]. The superior inhibitory activity observed in the AW and TM regions can be ascribed to their higher levels of total secondary metabolites, including catechin for AW, a known potent α - α -amylase inhibitor. Additionally, recent research often highlights the α -glucosidase inhibitory effects of basic polyphenols, including phenolic acids and flavonoids.

We conducted this test to find traditional alternatives for treating diabetes because the initial phase of acarbose treatment may be linked with side effects such as bloating, gas, and diarrhea [63].




Figure 3. Inhibitory effect of TG, TM, AW, and acarbose on pancreatic α -amylase activity of human pancreatic cells in vitro. The results are shown as the mean value \pm SD (n=3). TG, Tizi-Ghnm site; TM., Timoulilt site; AW, Ait Waarda site; (M1, M2, M3, M4, and M5), stages of the immature phase of carob pulp.

3.6.2. Intestinal α -Glucosidase

Intestinal α -glucosidase performs an essential function in the digestion of carbohydrates by breaking down starch and disaccharides into glucose, facilitating their absorption in the intestines. Inhibitors of α -glucosidase are considered key targets for developing new treatments for diabetes, as they can help manage blood glucose levels [64]. Polyphenol-rich plant foods have been documented to mimic insulin's effects on glucose metabolism and serve as effective inhibitors of crucial enzymes like α -amylase and α -glucosidase, which are linked to type 2 diabetes and tissue lipid peroxidation [65]. In our study, we investigated the inhibitory effect of ICPE from three different regions (TG, TM, and AW) on α -glucosidase activity. The IC_{50} values demonstrated a strong inhibitory effect, with TG showing the most potent activity at $0.063 \pm 0.002 \mu\text{g/mL}$, followed by AW at $0.065 \pm 0.004 \mu\text{g/mL}$, and TM at $0.086 \pm 0.014 \mu\text{g/mL}$ (Table 6, Figure 4). These results are particularly significant relative to the standard inhibitor, acarbose, which has an IC_{50} of $0.089 \pm 0.004 \mu\text{g/mL}$. The extracts from TG and AW regions exhibit superior inhibitory effects on α -glucosidase, indicating a higher potential for managing postprandial blood glucose levels. This enhanced inhibition is likely due to the rich polyphenolic content, including compounds such as 3-O-p-coumaroyl 5-O-caffeoylquinic acid, quercetin 3-methyl ether, galloylquinic acid, and gallic acid, known for their enzyme inhibitory and antioxidant properties, as demonstrated in this study. A previous study demonstrated that the leaf and stem bark samples of the plant exhibited a potent inhibitory activity on α -glucosidase, with both samples showing 99% inhibition at a concentration of 10 mg/mL. This inhibition was significantly higher than that of acarbose, which showed an 85% inhibition [60]. It has been documented that hydrolyzable tannins have been shown to inhibit α -glucosidase activity. Moreover, *C. siliqua* germ is known to be rich in glycosylated derivatives of (iso)schaftoside. Apigenin 6,8-C-di-glycosides have also been identified as effective α -glucosidase inhibitors [66].




Figure 4. Inhibitory effect of TG, TM, AW, and acarbose on intestinal α -glucosidase activity of human pancreatic cells in vitro. The results are shown as the mean value \pm SD (n=3). TG, Tizi-Ghnm site; TM., Timoulilt site; AW, Ait Waarda site; (M1, M2, M3, M4, and M5), stages of the immature phase of carob pulp.

Table 6. Inhibitory activity of α -glucosidase by TG, TM, AW, and acarbose.

Extractions	IC ₅₀ $\mu\text{g/mL}$
TG	0.063 ± 0.002^b
TM	0.086 ± 0.014^a

AW	0,065 ± 0.004 ^b
Acarbose (control)	0,089 ± 0.004 ^a

Values are presented as mean ± SEM (n = 3). Statistical analysis was performed using one-way ANOVA with Tukey’s post hoc test at a 5% significance level. Means within the same column marked with different letters are significantly different (p < 0.05). TG., Tizi-Ghnm site; TM., Timoulilt site; AW., Ait Waarda site.

3.7. In Vivo, the Plant Extract’s Effect on Inhibiting Pancreatic α-Amylase and Intestinal α-Glucosidase Activities

3.7.1. Pancreatic α-Amylase

We evaluated the inhibitory effect of ICPE on α-amylase *in vitro*, which demonstrated significant enzyme inhibition. To validate and confirm these findings, it is crucial to examine the effects *in vivo*, as *in vitro* studies alone cannot fully replicate the complex interactions and metabolic processes occurring within a living organism. *In vivo* studies offer a deeper insight into the bioavailability, efficacy, and potential side effects of the extracts in a physiological context. Therefore, we proceeded to test the extracts’ inhibitory effects on α-amylase activity *in vivo*. After administering starch (2 g/kg), glycemia peaked at 2.18 ± 0.02 g/L at 60 minutes, as shown in Figure 3. It then gradually decreased to 1.79 ± 0.01 g/L at 90 minutes, 1.0 ± 0.005 g/L at 150 minutes, and stabilized around 0.94 ± 0.001 g/L. However, with the administration of AW (400 mg/kg), postprandial glycemia significantly decreased (P < 0.001 at 60 and 90 minutes, and P < 0.01 at 150 and 210 minutes), reaching 1.05 ± 0.06 g/L at 60 minutes, 0.89 ± 0.004 g/L at 90 minutes, 0.72 ± 0.005 g/L at 150 minutes, and 0.70 ± 0.001 g/L at 210 minutes relative to the control. Similarly, the presence of TG (400 mg/kg) significantly reduced postprandial glycemia (P < 0.001 at 60 and 90 minutes), resulting in values of 1.22 ± 0.008 g/L at 60 minutes and 1.21 ± 0.006 g/L at 90 minutes. Additionally, TM (400 mg/kg) significantly decreased postprandial glycemia (P < 0.001 at 60 and 90 minutes), with levels reaching 1.35 ± 0.008 g/L at 60 minutes and 1.05 ± 0.001 g/L at 90 minutes compared to the control. Acarbose administration (10 mg/kg) significantly reduced starch-induced hyperglycemia (P < 0.001) at 150 and 210 minutes, with values of 0.72 ± 0.005 g/L at 120 minutes and 0.57 ± 0.001 g/L at 210 minutes. After a starch overload, the area under the curve (AUC) in the control group was 81.29 ± 0.01 g/L/h. Pretreatment with AW (400 mg/kg) significantly inhibited the glycemic load (P < 0.001), resulting in an AUC of 50.03 ± 0.003 g/L/h. TG (400 mg/kg) inhibited the glycemic load (P < 0.05) with an AUC of 67.89 ± 0.001 g/L/h. TM (400 mg/kg) also showed significant inhibition (P < 0.01) with an AUC of 64.93 ± 0.003 g/L/h. Acarbose (10 mg/kg) demonstrated inhibitory activity (P < 0.05) with an AUC of 67.35 ± 0.006 g/L/h. These results indicate that AW, TG, and TM significantly inhibit α-amylase activity *in vivo*, resulting in lower post-meal blood glucose levels. The effectiveness of AW was particularly notable, showing the most substantial decrease in glycemia and glycemic load. This suggests that the increased concentration of total secondary metabolites in these extracts is vital for the inhibition of α-amylase, thereby controlling postprandial hyperglycemia effectively. Previous *in vivo* research indicated that *C. siliqua* extracts significantly inhibited α-amylase activity and improved blood glucose levels in type 2 diabetic rats. High-dose carob-treated (CS1000) and glibenclamide-treated (GD) groups had higher total protein and serum amylase levels compared to untreated diabetics (D0). The treated groups also had lower serum fasting blood glucose (FBG) and higher insulin levels. Additionally, the HOMA-B index improved significantly in the treated groups, indicating better beta-cell function. These results highlight the potential of *C. siliqua* extracts in managing diabetes [67].




Figure 3. Impact of AW, TG, TM, and acarbose on glycemia following starch overload (2g/kg) in normal rats (n=6). B: Area under the curve (AUC) for rats subjected to starch overload in the presence of AW, TG, TM, or acarbose. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. TG., Tizi-Ghnm site; TM., Timoulilt site; AW., Ait Waarda site.

3.7.2. Intestinal α -Glucosidase

We began by evaluating the inhibitory effects of ICPE on α -glucosidase *in vitro*, where we observed substantial enzyme inhibition. However, to validate these findings, *in vivo* studies are necessary. This step is crucial as *in vitro* experiments cannot fully replicate the intricate interactions and metabolic processes that occur within a living organism. Following sucrose administration (2 g/kg), glycemia peaked at 1.58 ± 0.003 g/L at 60 minutes (Figure 4), then gradually declined to 1.23 ± 0.001 g/L at 90 minutes, 1.0 ± 0.001 g/L at 150 minutes, and stabilized around 0.66 ± 0.01 g/L. However, with the administration of AW (400 mg/kg), postprandial glycemia significantly decreased ($P < 0.01$ at 90 minutes and $P < 0.001$ at 210 minutes), reaching 0.72 ± 0.001 g/L at 90 minutes and 0.45 ± 0.01 g/L at 210 minutes compared to the control. Similarly, TG (400 mg/kg) significantly reduced postprandial glycemia ($P < 0.05$ at 150 minutes), resulting in values of 0.76 ± 0.005 g/L at 150 minutes. Additionally, TM (400 mg/kg) and Acarbose (10 mg/kg) did not exhibit significant inhibitory activity on α -glucosidase *in vivo*. After a sucrose overload, the area under the curve (AUC) in the control group was 62.61 ± 0.003 g/L/h. Pretreatment with AW (400 mg/kg) significantly reduced the glycemic load ($P < 0.001$), resulting in an AUC of 41.60 ± 0.001 g/L/h. TG (400 mg/kg) inhibited the glycemic load ($P < 0.05$) with an AUC of 50.32 ± 0.01 g/L/h, while TM (400 mg/kg) also showed significant inhibition ($P < 0.01$) with an AUC of 50.32 ± 0.001 g/L/h. Acarbose (10 mg/kg) demonstrated an inhibitory effect ($P < 0.05$) with an AUC of 50.32 ± 0.01 g/L/h. These findings suggest that AW, TG, and TM significantly inhibit α -glucosidase activity *in vivo*, leading to reduced postprandial blood

glucose levels. The effectiveness of AW was particularly notable, showing the most substantial decrease in glycemia and glycemic load. This indicates that the greater amount of total secondary metabolites in these extracts are keys to inhibiting α -glucosidase, effectively controlling postprandial hyperglycemia. This study is pioneering in its exploration of the inhibitory activity of α -glucosidase in vivo using ICPE. While previous research has primarily focused on in vitro analyses, our work is the first to extend these investigations to a living organism, providing a more comprehensive understanding of the extract's potential antidiabetic effects. Even though other studies have already demonstrated the antidiabetic effect of *C. siliqua* in vivo, they have done so using different methods [22].

This study offers a valuable understanding of the potential of using immature carob pulp at different stages of maturity (M1, M2, M3, M4, and M5) as a traditional alternative for treating diabetes. By demonstrating significant inhibitory effects on digestive enzymes and highlighting the stage with the highest polyphenol and flavonoid levels, along with antioxidant properties, our findings underscore the therapeutic promise of carob pulp. Given the gastrointestinal side effects associated with conventional treatments like acarbose, incorporating immature carob pulp could offer a beneficial, natural alternative for diabetes management.




Figure 4. Impact of AW, TG, TM, and acarbose on glycemia following sucrose overload (2g/kg) in normal rats (n = 6). B: Area under the curve (AUC) for rats subjected to sucrose overload in the presence of AW, TG, TM, or acarbose. * P < 0.05; ** P < 0.01; *** P < 0.001. TG., Tizi-Ghnm site; TM., Timoulilt site; AW., Ait Waarda site.

3.8. Molecular Docking: Targeting α -Glucosidase and α -Amylase

Pancreatic α -Amylase is a crucial enzyme in carbohydrate metabolism. It is also studied as a therapeutic target for diabetes treatment, as modulating its activity can help manage postprandial blood glucose levels. Therefore, α -amylase is important not only for digestion but also for its medical and therapeutic applications in diabetes management [68]. In this study, results obtained from molecular docking of ligands to human α -amylase (PDB: 1B2Y) revealed that the 3-O-p-Coumaroyl-5-O-caffeoylquinic acid molecule exhibited a higher binding affinity higher (-9.2 kcal/mol) for α -

amylase than acarbose (-8.2 kcal/mol), which is a standard inhibitor of α -amylase. Additionally, quercetin 3-methyl ether exhibited a similar binding affinity to acarbose (-8.2 kcal/mol) (Table 7). The 3-O-p-Coumaroyl-5-O-caffeoylquinic acid molecule stood out for its good affinity towards α -amylase, interacting effectively with several amino acid residues of the enzyme. Specifically, it demonstrated hydrogen bond interactions with Tyr 6, Arg 252, Trp 280, Pro 332, Arg 398, and Arg 421. Moreover, it exhibited a pi-alkyl interaction with Pro 4 and a pi-pi T-shaped interaction with His 331, offering deeper insight into its binding with α -amylase (Figure 5). α -Glucosidase is a crucial enzyme in carbohydrate metabolism. It plays a crucial role in the digestive process by catalyzing the hydrolysis of α -glucose residues at the non-reducing ends of carbohydrates, such as starch and disaccharides. Due to its central role in carbohydrate digestion and absorption, α -glucosidase has become an important target in the treatment of diabetes. Inhibition of this enzyme can effectively delay glucose absorption, leading to a more gradual rise in blood sugar levels after meals [69]. In this study, the results obtained from molecular docking of ligands to human α -glucosidase (PDB:5NN8) revealed that the 3-O-p-Coumaroyl-5-O-caffeoylquinic acid and 3,4-Dicaffeoylquinic acid molecules exhibited higher binding affinities, with -8.2 kcal/mol and -7.9 kcal/mol (Table 7). The 3-O-p-Coumaroyl-5-O-caffeoylquinic acid molecule stood out for its good affinity towards α -amylase, interacting effectively with several amino acid residues of the enzyme. Specifically, it demonstrated hydrogen bond interactions with Lys 348, Arg 725, Ile 823, and Glu 856, pi-cation interaction with His 708, and pi-alkyl interaction with Ala 749. Additionally, the 3,4-Dicaffeoylquinic acid molecule interacted effectively with several amino acid residues of the enzyme. It demonstrated hydrogen bond interactions with Arg 585, Tyr 609, and His 717, thus providing further insight into its interaction with α -glucosidase (Figure 5). These findings indicate that the compounds in the methanolic extract of immature carob pods exhibit significant affinity for α -amylase and α -glucosidase enzymes. This affinity aligns with the inhibitory effects observed in our in vitro and in vivo studies, further supporting the potential of this extract as a natural inhibitor of enzymes involved in carbohydrate metabolism.

Table 7. Summary of molecular docking studies on phytochemicals identified in the methanolic extract of unripe carob pods, targeting human α -glucosidase and α -amylase.

Compound name	α -amylase (PDB:1B2Y)		α -glucosidase (PDB: 5NN8)	
	Affinity (Kcal/mol)	Interaction Site	Affinity (Kcal/mol)	Interaction Site
Acarbose (Standard inhibitor)	-8.2	Tyr 2, Ser 3, Pro 4, Thr 6, Gln 8, Phe 222, Arg 252, Pro 332, Gly 334, Arg 398.	-7.2	Asp 356, Val 357, Arg 608, His 584, His 717, Leu 865, Glu 866.
3-O-p-Coumaroyl-5-O-caffeoylquinic acid	-9.2	Pro 4, Tyr 6, Arg 252, Trp 280, His 331, Pro 332, Arg 398, Arg 421	-8.2	Lys 348, His 708, Arg 725, Ala 749, Ile 823 Glu 856
Quercetin 3-methyl ether	-8.2	Arg 267, Arg 303, Thr 314, Arg 346, Asp 356	-7.1	Met 363, Arg 585, Arg 594
Gallic acid	-5.9	Val 129, Gly 181, Tyr 182	-5.8	His 708, Gln 715, Glu 748, Ala 749, Ile 823, Glu 856.
3,4-Dicaffeoylquinic acid	-8.2	Pro4, Gln 8, Gly 9, Pro 332, Phe 335, Thy 336,	-7.9	Met 363, Arg 585, Arg 594, Tyr 609, His 717, Glu 866,

		Arg 398, Asp 402, Arg 421	
Galloylquinic acid	-7.2	Arg 10, Arg 252, Ser 289, Gly 334, Asp 402, Gly 403	Leu 195, Arg 585, Tyr 609
Ferulic acid	-6.3	Arg 552, Ser 289, Pro 332, Phe 335, Arg 424	His 708, Glu 721, Arg 725, Tyr 822, Gly 855, Glu 856
Catechin	-7.8	Asn 301, Ile 312, Thr 314, Asp 317, Arg 346, Asn 352	Pro 194, Leu 195, Phe 490, Thr 491, Leu 577, Ile 581, Tyr 609




Figure 5. The 2D and 3D binding interactions of phytochemicals identified in the methanolic extract of unripe carob pods and acarbose (standard inhibitor) against the human α -glucosidase (PDB: 5NN8) and human α -amylase enzyme (PDB: 1B2Y).

3.9. ADME Analysis

In silico ADME studies are crucial for advancing pharmaceutical development, as they offer an affordable approach to forecasting a drug’s actions within the human body [70]. Leveraging computational models for early pharmacokinetic evaluation allows for the rapid identification of promising drug candidates, optimizing the development process. This approach reduces the risk of adverse side effects, lowers the probability of drug development failures, and enhances the likelihood of clinical success, making it a crucial asset in contemporary drug discovery and development [71]. To adhere to Lipinski’s rule of five and Veber’s rule, compounds that are considered appropriate for oral drug development should typically not violate more than one of the following criteria: (1) a maximum of 10 hydrogen bond acceptors (nitrogen or oxygen atoms), (2) an octanol-water partition coefficient log P (MLogP) of less than 5, (3) a molecular weight of less than 500 daltons, and (4) no more than 5 hydrogen bond donors [72]. In our study, all compounds tested conformed to Lipinski’s guidelines, suggesting their potential for oral drug development, except 3-O-p-Coumaroyl-5-O-caffeoylquinic acid and 3,4-Dicaffeoylquinic acid. The blood-brain barrier (BBB) acts as a vital barrier between the systemic circulation and the central nervous system, safeguarding the brain through biochemical processes such as enzyme reactions and physical mechanisms like active efflux systems [73]. Our study found that only ferulic acid can cross the BBB, as indicated by the yellow region in Figure 6. Additionally, most of the compounds analyzed were identified as P-glycoprotein non-substrates (PGP-), except for 3-O-p-coumaroyl-5-O-caffeoylquinic acid and catechin, which were classified as PGP+ (Figure 6). A molecule that is efficiently absorbed by the intestine presents significant benefits in terms of bioavailability, efficacy, convenience, and tolerance, making it a strong candidate for oral drug development [74]. Notably, all the phytochemicals analyzed in our study exhibit high intestinal absorption, except Galloylquinic acid, 3-O-p-Coumaroyl-5-O-caffeoylquinic acid, and 3,4-Dicaffeoylquinic acid. Cytochrome P450, a key enzyme for detoxification, is primarily found in the liver [75]. Our analysis showed that most compounds are neither inhibitors nor substrates of CYP450 enzymes, particularly CYP1A2, except for Quercetin-3-methyl ether, which was identified as a CYP1A2 inhibitor (Table 8). This suggests a lower likelihood of interference with drug metabolism, enhancing the safety profile of the phytochemicals.

Table 8. Evaluation of the pharmacokinetic properties (ADME) of phytochemicals identified in methanolic extract of unripe carob pods (*in silico*).

Compound	Physicochemical Properties					Lipophilicity		Druglikeness		Pharmacokinetics		
	MW	HB	HB	TPS	RO	Mlo	Wlo	Lipins	Verbe	GI	BBB	CYP1
s	g/mol	A	D	A	Å ²	gP	gP	ki’s	r’s	absorption	Permeation	A2 Inhibitor
3-O-p-Coumaroyl-5-O-caffeoylquinic acid	500.4	11	6	191.0	9	0.1	1.1	3	1	Low	No	No
Quercetin 3-methyl ether	316.2	7	4	120.3	2	-0.3	2.2	0	0	High	No	No

Gallic acid	170.	5	4	97.9	1	-0.1	0.5	0	0	High	No	No
	1											
3,4-Dicaffeoylquinic acid	516.	12	7	211.	9	-0.3	0.8	3	1	Low	No	No
	4			2								
Galloylquinic acid	344.	10	8	195.	3	-2.7	-2.1	1	1	Low	No	No
	2			8								
Ferulic acid	194.	4	2	66.7	3	1.0	1.3	0	0	High	Yes	No
	1											
Catechin	290.	6	5	110.	1	0.2	1.2	0	0	High	No	No
	2			3								

MW: molecular weight; ROTB: rotatable bonds; HBD: Hydrogen-Bond Donors; HBA: Hydrogen-Bond Acceptors; WLogP: Lipophilicity; TPSA: Topological polar surface area; MLogP: Octanol/water partition coefficient; GI: Gastrointestinal; BBB: Blood-Brain Barrier.




Figure 6. BOILED-Egg Model of the GI absorption and BBB permeability of phytochemicals identified in methanolic extract of unripe carob pods. (1) Ferulic acid, (2) Gallic acid, (3) Catechin, (4) Quercetin-3-methylether, (5) 3-O-p-coumaroyl-5-O-caffeoylquinic acid, (6) Galloylquinic acid. PGP+: P-glycoprotein substrate, PGP-: non-substrate of P-glycoprotein.

4. Conclusions

Maturity can be used to predict the potential content of secondary metabolites with antioxidant capacity, such as polyphenols and flavonoids. The phytochemical composition and antioxidant capacity of immature carob pulp exhibited variation concerning the maturity stage, as evidenced by this study. The stage containing the highest amount of polyphenols demonstrated a notable anti-diabetic effect, suggesting that these compounds may play a role in the prevention and management of diabetes. It can be posited that this activity is responsible for the presence of a significant amount of 3-O-p-coumaryol-5-O-caffeoylquinique in the carob pulp extracts. In light of these new findings, further rigorous studies must be conducted on how this compound functions as a medicinal agent in a variety of animal models to gain a comprehensive understanding of its mechanism of action

concerning blood sugar control. Our study's findings indicate that immature carob pulp extracts contain active principles. The *in silico* analysis reveals that these compounds exhibit varied affinities with key enzymes involved in carbohydrate metabolism and also possess favorable properties for the development of oral medications. In particular, immature carob pulp extracts offer a valuable opportunity for the development of new natural therapeutic strategies against diabetes. Their potential as effective alternatives to established synthetic inhibitors underscores the importance of continuing research to explore and leverage these compounds in clinical applications.

Author Contributions: Conceptualization, S.L., M.B., A.T. and K.E.; Methodology, S.L., M.B., A.E., and K.E.; Formal Analysis, S.L., A.T., M.B., H.C. and A.F.; Data Curation, M.L., Y.N. and F.C.; Writing- original draft, S.L., M.B., A.T. and A.E.; Writing- Review and editing, S.L., M.B., A.E., K.E., and Y.N.; Investigation, H.C., R.N.H. and F.C.; Validation, S.S., F.C., Y.N. and K.E.; Visualization, Y.N., A.A.S. and S.S; Resources, M.L. and A.F.; Software, S.L., A.F., A.T., and H.C.; Supervision, S.S., Y.N. and K.E.; funding acquisition, A.A.S. and R.N.H. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Researchers Supporting Project number (RSPD2024R1057) at King Saud University, Riyadh, Saudi. This research was also funded by PROMEDLIFE (Novel food products for Promotion of Mediterranean LIFE style and healthy diet), grant number 2132 (PRIMA program supported by the European Union).

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Institutional Review Board of the Faculty of Sciences and Technology, Beni Mellal, Morocco (11/19-LGB-08 and 05/06/2024).

Data Availability Statement:

Acknowledgments: The authors are thankful to the Researchers Supporting Project number (RSPD2024R1057) at King Saud University, Riyadh, Saudi Arabia. Special thanks are extended to the staff of the Food Technology and Quality Laboratory at the "Qualipôle alimentation de Béni-Mellal," affiliated with the National Institute for Agricultural Research (INRA). The invaluable assistance and unwavering support provided by the team at the Biological Engineering Laboratory, Faculty of Sciences and Techniques, Sultan Moulay Slimane University, are deeply appreciated. The authors also acknowledge the financial support provided by the PRIMA Program on Carob Research through the PROMEDLIFE Project (Section I), Agreement No. 2132.

Conflicts of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. Kassout J, Hmimsa Y, Fatehi S El, Kadaoui K, Houssni M, Chakkour S, et al. Aridity Gradients Shape Intraspecific Variability of Morphological Traits in Native *Ceratonia siliqua* L. of Morocco. *Plants* 2023;12. <https://doi.org/10.3390/plants12193447>.
2. Frühbauerová M, Červenka L, Hájek T, Pouzar M, Palarčík J. Bioaccessibility of phenolics from carob (*Ceratonia siliqua* L.) pod powder prepared by cryogenic and vibratory grinding. *Food Chem* 2022;377. <https://doi.org/10.1016/j.foodchem.2021.131968>.
3. Papaefstathiou E, Agapiou A, Giannopoulos S, Kokkinofa R. Nutritional characterization of carobs and traditional carob products. *Food Sci Nutr* 2018;6:2151–61. <https://doi.org/10.1002/fsn3.776>.
4. Roseiro LB, Tavares CS, Roseiro JC, Rauter AP. Antioxidants from aqueous decoction of carob pods biomass (*Ceratonia siliqua* L.): Optimisation using response surface methodology and phenolic profile by capillary electrophoresis. *Ind Crops Prod* 2013;44:119–26. <https://doi.org/10.1016/j.indcrop.2012.11.006>.
5. Kyriacou MC, Antoniou C, Roupheal Y, Graziani G, Kyrtatzis A. Mapping the primary and secondary metabolomes of carob (*Ceratonia siliqua* L.) fruit and its postharvest antioxidant potential at critical stages of ripening. *Antioxidants* 2021;10:1–21. <https://doi.org/10.3390/antiox10010057>.
6. Owen RW, Haubner R, Hull WE, Erben G, Spiegelhalder B, Bartsch H, et al. Isolation and structure elucidation of the major individual polyphenols in carob fibre. *Food and Chemical Toxicology* 2003;41:1727–38. [https://doi.org/10.1016/S0278-6915\(03\)00200-X](https://doi.org/10.1016/S0278-6915(03)00200-X).
7. Ortega N, Macià A, Romero MP, Trullols E, Morello JR, Anglès N, et al. Rapid determination of phenolic compounds and alkaloids of carob flour by improved liquid chromatography tandem mass spectrometry. *J Agric Food Chem* 2009;57:7239–44. <https://doi.org/10.1021/jf901635s>.
8. Bengoechea C, Romero A, Villanueva A, Moreno G, B MA, Milla F, et al. Composition and structure of carob (*Ceratonia siliqua* L.) germ proteins. *Food Chem* 2008;107:675–83. <https://doi.org/10.1016/j.foodchem.2007.08.069>.

9. Achchoub M, Azzouzi H, Elhajji L, Benbati M, Elfazazi K, Salmaoui S. Evaluation of Physicochemical, Functional and Sensory Properties of Carob Pulp Beverage (*Ceratonia Siliqua* L). *Biosci Biotechnol Res Asia* 2021;18:611–8. <https://doi.org/10.13005/bbra/2945>.
10. Elfazazi K, Harrak H, Achchoub M, Benbati M. Physicochemical criteria, bioactive compounds and sensory quality of Moroccan traditional carob drink. *Mater Today Proc* 2020;27:3249–53. <https://doi.org/10.1016/j.matpr.2020.04.868>.
11. Goulas V, Stylos E, Chatziathanasiadou M V., Mavromoustakos T, Tzakos AG. Functional Components of Carob Fruit: Linking the Chemical and Biological Space. *International Journal of Molecular Sciences* 2016, Vol 17, Page 1875 2016;17:1875. <https://doi.org/10.3390/IJMS17111875>.
12. Maier H, Anderson M, Karl C, Magnuson K, Whistler RL. Guar, Locust Bean, Tara, and Fenugreek Gums. *Industrial Gums: Polysaccharides and Their Derivatives: Third Edition* 2013:181–226. <https://doi.org/10.1016/B978-0-08-092654-4.50012-7>.
13. Patmore J V., Goff HD, Fernandes S. Cryo-gelation of galactomannans in ice cream model systems. *Food Hydrocoll* 2003;17:161–9. [https://doi.org/10.1016/S0268-005X\(02\)00048-6](https://doi.org/10.1016/S0268-005X(02)00048-6).
14. Laaraj S, Hussain A, Mouhaddach A, Noutfia Y, Gorski FI, Yaqub S, et al. Nutritional Benefits and Antihyperglycemic Potential of Carob Fruit (*Ceratonia siliqua* L.): An Overview. *Ecological Engineering and Environmental Technology* 2024;25:124–32. <https://doi.org/10.12912/27197050/178456>.
15. Laaraj S, Salmaoui S, Addi M, El-Rhouttaï C, Tikent A, Elbouzidi A, et al. Carob (*Ceratonia siliqua* L.) Seed Constituents: A Comprehensive Review of Composition, Chemical Profile, and Diverse Applications. *J Food Qual* 2023;2023. <https://doi.org/10.1155/2023/3438179>.
16. Palipoch S, Punsawad C, Suwannalert P. *Thunbergia laurifolia*, a new choice of natural antioxidant to prevent oxidative stress-related pathology: A review 2013;7:698–701. <https://doi.org/10.5897/JMPR012x.003>.
17. Slavin J. Fiber and prebiotics: Mechanisms and health benefits. *Nutrients* 2013;5:1417–35. <https://doi.org/10.3390/nu5041417>.
18. Raninen K, Lappi J, Mykkänen H, Poutanen K. Dietary fiber type reflects physiological functionality: comparison of grain fiber, inulin, and polydextrose. *Nutr Rev* 2011;69:9–21. <https://doi.org/10.1111/J.1753-4887.2010.00358.X>.
19. de Bock M, Derraik JGB, Cutfield WS. Polyphenols and glucose homeostasis in humans. *J Acad Nutr Diet* 2012;112:808–15. <https://doi.org/10.1016/j.jand.2012.01.018>.
20. Dewanjee S, Das AK, Sahu R, Gangopadhyay M. Antidiabetic activity of *Diospyros peregrina* fruit: Effect on hyperglycemia, hyperlipidemia and augmented oxidative stress in experimental type 2 diabetes. *Food and Chemical Toxicology* 2009;47:2679–85. <https://doi.org/10.1016/j.fct.2009.07.038>.
21. Kumar S, Kumar V, Prakash O, Kumar V. Antidiabetic, hypolipidemic and histopathological analysis of *Dillenia indica* (L.) leaves extract on alloxan induced diabetic rats *Asian Pacific Journal of Tropical Medicine* Antidiabetic *Dillenia indica* Hypolipidemic Rat Serum cholesterol. 2011.
22. Rîbi K, Selmi S, Grami D, Sebai H, Marzouki L. Cronicon In Vitro α -Amylase/ α -Glucosidase Inhibitory Activities and In Vivo Improving Glucose Tolerance and Hypoglycemic Effect of *Ceratonia siliqua* Leaves Aqueous Extract *Corresponding. vol. 13. 2018.
23. Benchikh Y, Louaileche H, George B, Merlin A. Changes in bioactive phytochemical content and in vitro antioxidant activity of carob (*Ceratonia siliqua* L.) as influenced by fruit ripening. *Ind Crops Prod* 2014;60:298–303. <https://doi.org/10.1016/j.indcrop.2014.05.048>.
24. Benchikh Y, Paris C, Louaileche H, Charbonnel C, Ghoul M, Chebil L. Comparative characterization of green and ripe carob (*Ceratonia siliqua* L.): physicochemical attributes and phenolic profile. 2016.
25. Benchikh Y, Louaileche H. Effects of extraction conditions on the recovery of phenolic compounds and in vitro antioxidant activity of carob (*Ceratonia siliqua* L.) pulp. *Acta Botanica Gallica* 2014;161:175–81. <https://doi.org/10.1080/12538078.2014.909325>.
26. Farag MA, El-Kersh DM, Ehrlich A, Choucri MA, El-Seedi H, Frolov A, et al. Variation in *Ceratonia siliqua* pod metabolome in context of its different geographical origin, ripening stage and roasting process. *Food Chem* 2019;283:675–87. <https://doi.org/10.1016/j.foodchem.2018.12.118>.
27. Ydjedd S, Chaalal M, Richard G, Kati DE, López-Nicolás R, Fauconnier ML, et al. Assessment of antioxidant potential of phenolic compounds fractions of Algerian *Ceratonia siliqua* L. pods during ripening stages. *Int Food Res J* 2017;24:2041–9.
28. Ben Othmen K, Elfalleh W, Lachiheb B, Haddad M. Evolution of phytochemical and antioxidant activity of Tunisian carob (*Ceratonia siliqua* L.) pods during maturation. *Eurobiotech Journal* 2019;3:135–42. <https://doi.org/10.2478/ebtj-2019-0016>.
29. Rîbi K, Selmi S, Grami D, Saidani K, Sebai H, Amri M, et al. *Ceratonia siliqua* L. (immature carob bean) inhibits intestinal glucose absorption, improves glucose tolerance and protects against alloxan-induced diabetes in rat. *J Sci Food Agric* 2017;97:2664–70. <https://doi.org/10.1002/jsfa.8091>.
30. Tikent A, Laaraj S, Marhri A, Taïbi M, Elbouzidi A, Khalid I, et al. The Antioxidant and Antimicrobial Activities of Two Sun-Dried Fig Varieties (*Ficus carica* L.) Produced in Eastern Morocco and the

- Investigation of Pomological, Colorimetric, and Phytochemical Characteristics for Improved Valorization. *International Journal of Plant Biology* 2023;14:845–63. <https://doi.org/10.3390/ijpb14030063>.
31. Vekiari AS, Ouzounidou G, Gork G, Ozturk M, Asfi M. Compositional changes of major chemical compounds in Greek carob pods during development. *Bull Chem Soc Ethiop* 2012;26:343–51. <https://doi.org/10.4314/BCSE.V26I3.3>.
 32. Prasanna V, Prabha TN, Tharanathan RN. Fruit Ripening Phenomena—An Overview. *Crit Rev Food Sci Nutr* 2007;47:1–19. <https://doi.org/10.1080/10408390600976841>.
 33. Song S, Abubaker MA, Akhtar M, Elimam AM, Zhu X, Zhang J. Chemical Characterization Analysis, Antioxidants, and Anti-Diabetic Activity of Two Novel Acidic Water-Soluble Polysaccharides Isolated from Baobab Fruits. *Foods* 2024;13. <https://doi.org/10.3390/foods13060912>.
 34. Moumou M, Mokhtari I, Tayebi A, Milenkovic D, Amrani S, Harnafi H. Immature carob pods extract and its fractions prevent lipid metabolism disorders and lipoprotein-rich plasma oxidation in mice: A phytochemical and pharmacological study. *J Ethnopharmacol* 2023;117557. <https://doi.org/10.1016/j.jep.2023.117557>.
 35. Ouzounidou G, Vekiari S, Asfi M, Gork MG, Sakcali MS, Ozturk M. PHOTOSYNTHETIC CHARACTERISTICS OF CAROB TREE (*CERATONIA SILIQUA* L.) AND CHEMICAL COMPOSITION OF ITS FRUIT ON DIURNAL AND SEASONAL BASIS. *Pak J Bot* 2012;44:1689–95.
 36. Sebai H, Souli A, Chehimi L, Rtibi K, Amri M, El-Benna J, et al. In vitro and in vivo antioxidant properties of Tunisian carob (*Ceratonia siliqua* L.). *Journal of Medicinal Plants Research* 2013;7:85–90. <https://doi.org/10.5897/JMPR12.915>.
 37. Gull J, Sultana B, Anwar F, Naseer R, Ashraf M, Ashrafuzzaman M. Variation in Antioxidant Attributes at Three Ripening Stages of Guava (*Psidium guajava* L.) Fruit from Different Geographical Regions of Pakistan. *Molecules* 2012, Vol 17, Pages 3165-3180 2012;17:3165–80. <https://doi.org/10.3390/MOLECULES17033165>.
 38. Ben Othmen K, Elfalleh W, Lachiheb B, Haddad M. Evolution of phytochemical and antioxidant activity of Tunisian carob (*Ceratonia siliqua* L.) pods during maturation. *Eurobiotech J* 2019;3:135–42. <https://doi.org/10.2478/ebtj-2019-0016>.
 39. Harborne JB, Williams CA. Advances in flavonoid research since 1992. *Phytochemistry* 2000;55:481–504. [https://doi.org/10.1016/S0031-9422\(00\)00235-1](https://doi.org/10.1016/S0031-9422(00)00235-1).
 40. Saci F, Bachir bey M, Louaileche H, Gali L, Bensouici C. Changes in anticholinesterase, antioxidant activities and related bioactive compounds of carob pulp (*Ceratonia siliqua* L.) during ripening stages. *Journal of Food Measurement and Characterization* 2020;14:937–45. <https://doi.org/10.1007/s11694-019-00344-9>.
 41. Saci F, Bachir bey M, Louaileche H, Gali L, Bensouici C. Changes in anticholinesterase, antioxidant activities and related bioactive compounds of carob pulp (*Ceratonia siliqua* L.) during ripening stages. *Journal of Food Measurement and Characterization* 2020;14:937–45. <https://doi.org/10.1007/s11694-019-00344-9>.
 42. Kim I, Lee J. Variations in anthocyanin profiles and antioxidant activity of 12 genotypes of mulberry (*Morus* spp.) fruits and their changes during processing. *Antioxidants* 2020;9. <https://doi.org/10.3390/antiox9030242>.
 43. Saensouk S, Senavongse R, Papayrata C, Chumroenphat T. Evaluation of Color, Phytochemical Compounds and Antioxidant Activities of Mulberry Fruit (*Morus alba* L.) during Ripening. *Horticulturae* 2022;8. <https://doi.org/10.3390/horticulturae8121146>.
 44. Dahmani W, Elaouni N, Abousalim A, Akissi ZLE, Legssyer A, Ziyat A, et al. Exploring Carob (*Ceratonia siliqua* L.): A Comprehensive Assessment of Its Characteristics, Ethnomedicinal Uses, Phytochemical Aspects, and Pharmacological Activities. *Plants* 2023, Vol 12, Page 3303 2023;12:3303. <https://doi.org/10.3390/PLANTS12183303>.
 45. Jafri SAA, Khalid ZM, Khan MZ, Jomezai NU. Evaluation of phytochemical and antioxidant potential of various extracts from traditionally used medicinal plants of Pakistan. *Open Chem* 2022;20:1337–56. <https://doi.org/10.1515/chem-2022-0242>.
 46. Ferreira-Santos P, Nobre C, Rodrigues RM, Genisheva Z, Botelho C, Teixeira JA. Extraction of phenolic compounds from grape pomace using ohmic heating: Chemical composition, bioactivity and bioaccessibility. *Food Chem* 2024;436:137780. <https://doi.org/10.1016/J.FOODCHEM.2023.137780>.
 47. Inga M, Betalleluz-Pallardel I, Puma-Isuiza G, Cumpa-Arias L, Osorio C, Valdez-Arana J-D-C, et al. Chemical analysis and bioactive compounds from agrifood by-products of peruvian crops. *Front Sustain Food Syst* 2024;8:1341895. <https://doi.org/10.3389/FSUFS.2024.1341895/BIBTEX>.
 48. Can-Cauich CA, Sauri-Duch E, Betancur-Ancona D, Chel-Guerrero L, González-Aguilar GA, Cuevas-Glory LF, et al. Tropical fruit peel powders as functional ingredients: Evaluation of their bioactive compounds and antioxidant activity. *J Funct Foods* 2017;37:501–6. <https://doi.org/10.1016/J.JFF.2017.08.028>.
 49. Rubio S, Quintana J, Eiroa JL, Triana J, Estévez F. Acetyl derivative of quercetin 3-methyl ether-induced cell death in human leukemia cells is amplified by the inhibition of ERK. *Carcinogenesis* 2007;28:2105–13. <https://doi.org/10.1093/CARCIN/BGM131>.

50. Lee EH, Kim HJ, Song YS, Jin C, Lee KT, Cho J, et al. Constituents of the stems and fruits of *Opuntia ficus-indica* var. *saboten*. *Arch Pharm Res* 2003;26:1018–23. <https://doi.org/10.1007/BF02994752/METRICS>.
51. Takeara R, Albuquerque S, Lopes NP, Callegari Lopes JL. Trypanocidal activity of *Lychnophora staavioides* Mart. (Vernonieae, Asteraceae). *Phytomedicine* 2003;10:490–3. <https://doi.org/10.1078/094471103322331430>.
52. Wei BL, Lu CM, Tsao LT, Wang JP, Lin CN. In vitro anti-inflammatory effects of quercetin 3-O-methyl ether and other constituents from *Rhamnus* species. *Planta Med* 2001;67:745–7. <https://doi.org/10.1055/S-2001-18339/ID/4/BIB>.
53. Kim SH, Naveen Kumar C, Kim HJ, Kim DH, Cho J, Jin C, et al. Glucose-containing flavones—their synthesis and antioxidant and neuroprotective activities. *Bioorg Med Chem Lett* 2009;19:6009–13. <https://doi.org/10.1016/J.BMCL.2009.09.062>.
54. Li J, Mottamal M, Li H, Liu K, Zhu F, Cho Y-Y, et al. Quercetin-3-methyl ether suppresses proliferation of mouse epidermal JB6 P1 cells by targeting ERKs. *Carcinogenesis* 2012;33:459–65. <https://doi.org/10.1093/carcin/bgr281>.
55. Motta EVS, Lemos M, Costa JC, Banderó-Filho VC, Sasse A, Sheridan H, et al. Galloylquinic acid derivatives from *Copaifera langsdorffii* leaves display gastroprotective activity. *Chem Biol Interact* 2017;261:145–55. <https://doi.org/10.1016/j.cbi.2016.11.028>.
56. Hassani S, Ghanbari F, Lotfi M, Alam W, Aschner M, Popović-Djordjević J, et al. How gallic acid regulates molecular signaling: role in cancer drug resistance. *Medical Oncology* 2023 40:11 2023;40:1–18. <https://doi.org/10.1007/S12032-023-02178-4>.
57. Bhuia MS, Rahaman MM, Islam T, Bappi MH, Sikder MI, Hossain KN, et al. Neurobiological effects of gallic acid: current perspectives. *Chinese Medicine* 2023 18:1 2023;18:1–19. <https://doi.org/10.1186/S13020-023-00735-7>.
58. Bouaouda K, Elagdi C, El Hachlafi N, Mohtadi K, Hsaine M, Kettani A, et al. HPLC-UV-MS/MS Profiling of Phenolics from *Euphorbia nicaeensis* (All.) Leaf and Stem and Its Antioxidant and Anti-Protein Denaturation Activities. *Progress In Microbes & Molecular Biology* 2023;6. <https://doi.org/10.36877/PMMB.A0000331>.
59. Agarwal P, Gupta R. Alpha-amylase inhibition can treat diabetes mellitus. *Research and Reviews Journal of Medical and Health Sciences RRJMHS* 2016;5.
60. Custódio L, Patarra J, Alberício F, Neng NR, Nogueira JMF, Romano A. In vitro antioxidant and inhibitory activity of water decoctions of carob tree (*Ceratonía siliqua* L.) on cholinesterases, α -amylase and α -glucosidase. *Nat Prod Res* 2015;29:2155–9. <https://doi.org/10.1080/14786419.2014.996147>.
61. Darwish WS, Khadr AES, Kamel MAEN, Abd Eldaim MA, El Sayed IET, Abdel-Bary HM, et al. Phytochemical Characterization and Evaluation of Biological Activities of Egyptian Carob Pods (*Ceratonía siliqua* L.) Aqueous Extract: In Vitro Study. *Plants* 2021, Vol 10, Page 2626 2021;10:2626. <https://doi.org/10.3390/PLANTS10122626>.
62. Kamtekar S, Keer V, Patil V. Estimation of Phenolic content, Flavonoid content, Antioxidant and Alpha amylase Inhibitory Activity of Marketed Polyherbal Formulation. *J Appl Pharm Sci* 2014;4:061–5. <https://doi.org/10.7324/JAPS.2014.40911>.
63. Hanefeld M. The Role of Acarbose in the Treatment of Non-Insulin-Dependent Diabetes Mellitus. *J Diabetes Complications* 1998;12:228–37. [https://doi.org/10.1016/S1056-8727\(97\)00123-2](https://doi.org/10.1016/S1056-8727(97)00123-2).
64. Subramanian R, Asmawi MZ, Sadikun A. In vitro α -glucosidase and α -amylase enzyme inhibitory effects of *Andrographis paniculata* extract and andrographolide. 2008.
65. Reddy NVLS, Anarthe SJ, Raghavendra NM. In Vitro Antioxidant and Antidiabetic activity of *Asystasia gangetica* (Chinese Violet) Linn. (Acanthaceae). *International Journal of Research in Pharmaceutical and Biomedical Sciences* 2010;1.
66. Picariello G, Sciammaro L, Siano F, Volpe MG, Puppo MC, Mamone G. Comparative analysis of C-glycosidic flavonoids from *Prosopis* spp. and *Ceratonía siliqua* seed germ flour. *Food Research International* 2017;99:730–8. <https://doi.org/10.1016/j.foodres.2017.06.058>.
67. Qasem MA, Noordin MI, Arya A, Alsalahi A, Jayash SN. Evaluation of the glycemic effect of *Ceratonía siliqua* pods (Carob) on a streptozotocin-nicotinamide induced diabetic rat model. *PeerJ* 2018;2018:e4788. <https://doi.org/10.7717/PEERJ.4788/SUPP-7>.
68. Zheng Y, Tian J, Yang W, Chen S, Liu D, Fang H, et al. Inhibition mechanism of ferulic acid against α -amylase and α -glucosidase. *Food Chem* 2020;317:126346. <https://doi.org/10.1016/j.foodchem.2020.126346>.
69. Singh A, Singh K, Sharma A, Kaur K, Kaur K, Chadha R, et al. Recent developments in synthetic α -glucosidase inhibitors: A comprehensive review with structural and molecular insight. *J Mol Struct* 2023;1281. <https://doi.org/10.1016/j.molstruc.2023.135115>.
70. Ferreira LLG, Andricopulo AD. ADMET modeling approaches in drug discovery. *Drug Discov Today* 2019;24:1157–65. <https://doi.org/10.1016/J.DRUDIS.2019.03.015>.

71. Srivastava V, Yadav A, Sarkar P. Molecular docking and ADMET study of bioactive compounds of *Glycyrrhiza glabra* against main protease of SARS-CoV2. *Mater Today Proc* 2022;49:2999–3007. <https://doi.org/10.1016/J.MATPR.2020.10.055>.
72. Farihi A, Bouhrim M, Chigr F, Elbouzidi A, Bencheikh N, Zrouri H, et al. Exploring Medicinal Herbs' Therapeutic Potential and Molecular Docking Analysis for Compounds as Potential Inhibitors of Human Acetylcholinesterase in Alzheimer's Disease Treatment. *Medicina (Lithuania)* 2023;59. <https://doi.org/10.3390/medicina59101812>.
73. Purushothaman K, Sivasankar E, Krishnamoorthy M, Karunakaran K, Muniyan R. Computational identification of potential tau tubulin kinase 1 (TTBK1) inhibitors: a structural analog approach. *In Silico Pharmacology* 2024 12:2 2024;12:1–11. <https://doi.org/10.1007/S40203-024-00242-Z>.
74. Stillhart C, Vučićević K, Augustijns P, Basit AW, Batchelor H, Flanagan TR, et al. Impact of gastrointestinal physiology on drug absorption in special populations—An UNGAP review. *European Journal of Pharmaceutical Sciences* 2020;147:105280. <https://doi.org/10.1016/J.EJPS.2020.105280>.
75. Taşcıoğlu N, Saatçi Ç, Emekli R, Tuncel G, Eşel E, Dundar M. Investigation of cytochrome p450 CYP1A2, CYP2D6, CYP2E1 and CYP3A4 gene expressions and polymorphisms in alcohol withdrawal. *Klinik Psikiyatri Dergisi* 2021;24:298–306. <https://doi.org/10.5505/KPD.2021.60938>.
76. Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. Colorimetric Method for Determination of Sugars and Related Substances. *Anal Chem* 1956;28:350–6. <https://doi.org/10.1021/ac60111a017>.
77. Tavarini S, Degl'Innocenti E, Remorini D, Massai R, Guidi L. Antioxidant capacity, ascorbic acid, total phenols and carotenoids changes during harvest and after storage of Hayward kiwifruit. *Food Chem* 2008;107:282–8. <https://doi.org/10.1016/j.foodchem.2007.08.015>.
78. Lamaison JL, Petitjen-Freytet C, Carnat A. Teneurs en acide rosmarinique, en dérivés hydroxycinnamiques totaux et activité antioxydante chez les apiacées, les borraginacées et les laminacées médicinales. *Ann Pharm Fr*, vol. 48, 1990, p. 103–8.
79. Broadhurst RB, Jones WT. Analysis of condensed tannins using acidified vanillin. *J Sci Food Agric* 1978;29:788–94. <https://doi.org/10.1002/jsfa.2740290908>.
80. Szabo K, Zorita D, Adriana-Florinela Catoi, Vodnar DC. Screening of Ten Tomato Varieties Processing Waste for Bioactive Components and Their Related Antioxidant and Antimicrobial Activities. *Antioxidants* 2019;8(8), 292.
81. Uysal S, Zengin G, Aktumsek A, Karatas S. Chemical and biological approaches on nine fruit tree leaves collected from the Mediterranean region of Turkey. *J Funct Foods* 2016;22:518–32. <https://doi.org/10.1016/j.jff.2016.02.006>.
82. Benzie IFF, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”: The FRAP assay. *Anal Biochem* 1996;239:70–6. <https://doi.org/10.1006/abio.1996.0292>.
83. Ma T, Tian C, Luo J, Zhou R, Sun X, Ma J. Influence of technical processing units on polyphenols and antioxidant capacity of carrot (*Daucus carrot* L.) juice. *Food Chem* 2013;141:1637–44. <https://doi.org/10.1016/j.foodchem.2013.04.121>.
84. Guide for the care and use of laboratory animals. National Academies Press; 2011.
85. Elrherabi A, Bouhrim M, Abdnim R, Berraouan A, Ziyat A, Mekhfi H, et al. Antihyperglycemic potential of the *Lavandula stoechas* aqueous extract via inhibition of digestive enzymes and reduction of intestinal glucose absorption. *J Ayurveda Integr Med* 2023;14:100795. <https://doi.org/10.1016/J.JAIM.2023.100795>.
86. Srivastava V, Yadav A, Sarkar P. Molecular docking and ADMET study of bioactive compounds of *Glycyrrhiza glabra* against main protease of SARS-CoV2. *Mater Today Proc* 2022;49:2999. <https://doi.org/10.1016/J.MATPR.2020.10.055>.
87. Zrouri H, Elbouzidi A, Bouhrim M, Bencheikh N, Kharchoufa L, Ouahhoud S, et al. Phytochemical analysis, antioxidant activity, and nephroprotective effect of the *Raphanus sativus* aqueous extract. *Mediterranean Journal of Chemistry* 2021;11:84. <https://doi.org/10.13171/mjc02101211565lk>.
88. Siddique MH, Ashraf A, Hayat S, Aslam B, Fakhar-e-Alam M, Muzammil S, et al. Antidiabetic and antioxidant potentials of *Abelmoschus esculentus*: In vitro combined with molecular docking approach. *Journal of Saudi Chemical Society* 2022;26:101418. <https://doi.org/10.1016/J.JSCS.2021.101418>.
89. Mendie LE, Hemalatha S. Molecular Docking of Phytochemicals Targeting GFRs as Therapeutic Sites for Cancer: an In Silico Study. *Appl Biochem Biotechnol* 2022;194:215–31. <https://doi.org/10.1007/S12010-021-03791-7/TABLES/8>.
90. Daina A, Michielin O, Zoete V. SwissADME: A free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci Rep* 2017;7. <https://doi.org/10.1038/srep42717>.
91. van de Waterbeemd H, Gifford E. ADMET in silico modelling: towards prediction paradise? *Nat Rev Drug Discov* 2003;2:192–204. <https://doi.org/10.1038/NRD1032>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s)

disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.