

Article

Not peer-reviewed version

Quantum Relativity (Electron Ripple)

[Ahmed M. Ismail](#) ^{*} and [Samira E. Mohamed](#)

Posted Date: 16 April 2026

doi: 10.20944/preprints202410.0352.v24

Keywords: special relativity; general relativity; Bohr atomic model; the fine-structure constant; photon energy; energy of the total photon; electron wave by de Broglie; gravity constant; quantum jump and cosmic constants of nature



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Quantum Relativity (Electron Ripple)

Ahmed M. Ismail ^{1,*} and Samira E. Mohamed ²

¹ University of Ain Shams, Al-Qubba Bridge, Al-Wayli, 11566, Egypt

² Faculty of Science, Ain Shams University, Al-Qubba Bridge, Al-Wayli, 11566, Egypt

* Correspondence: ahmedmohamed_ismaill@agr.asu.edu.eg

Abstract

This research answers the knowledge gap regarding the explanation of the quantum jump of the electron. This scientific paper aims to complete Einstein’s research regarding general relativity and attempt to link general relativity to quantum laws.

Keywords: special relativity; general relativity; bohr atomic model; the fine-structure constant; photon energy; energy of the total photon; electron wave by de broglie; gravity constant; quantum jump and cosmic constants of nature

1. Introduction

This research was created for the purpose of answering questions about physics phenomena that have not been answered. Such as explaining the phenomenon of the quantum jump of the electron and the phenomenon of cumulative entanglement. What happens in the phenomenon of the quantum jump of the electron is that when we give the electron energy, this energy causes the electron to move from the energy level that it occupies to the higher energy level without crossing the distance between the two orbits, which leads to the occurrence of the phenomenon of the quantum jump of the electron.[1] (Svidzinsky et al., 2014)

The role of this scientific paper is to provide a scientific explanation of how the quantum leap occurs without crossing the distance between the orbits. The theory of quantum entanglement is a connection between two quantum entangled particles. If one particle is observed, the other particle is affected by it at the same moment. This is what Einstein objected to; because when the electron traveled this distance in the same period of time, this would lead to the existence of a speed faster than the speed of light. Einstein proved it in special relativity. The maximum speed in the universe is the speed of light. Therefore, the phenomenon of quantum entanglement does not agree with Einstein’s laws. After the validity of quantum laws was proven. There has become a conflict between the laws of relativity that apply to the universe and the quantum laws that apply to atoms. This scientific paper aims to resolve this conflict between the laws of relativity and quantum laws. By establishing a law derived from the laws of relativity to apply to quantum laws. (Equation number 1)

This law in equation 1 is known as quantum relativity because it links the laws of relativity and quantum theory. This law is derived from general relativity. The law works to explain the phenomenon of the quantum leap and the phenomenon of quantum entanglement, as it explains that when energy is given to the atom, the atom does not gain energy, but rather space-time gains that energy. We will discuss the interpretation of this theory in detail later.

- The goal of this scientific research is to answer the explanation of the phenomenon of quantum leap and quantum entanglement and to add some modifications in the Bohr model.

2. Equations

These laws want to explain the results of the final derivation process of this research and what this research wants to prove.

$$\begin{aligned}
& G_{\mu\nu}[g_{\alpha\beta}(v_p \cdot v_g)] + \Lambda(v_p \cdot v_g)g_{\mu\nu}(v_p \cdot v_g) \\
&= \frac{8\pi}{1} \frac{G}{(v_p \cdot v_g)^2} T_{\mu\nu}[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)] \\
& G_{\mu\nu}[g_{\alpha\beta}(\|\vec{v}_p\| \|\vec{v}_g\|)] + \Lambda g_{\mu\nu}(\|\vec{v}_p\| \|\vec{v}_g\|) \\
&= \frac{8\pi}{1} \frac{G}{(\|\vec{v}_p\| \|\vec{v}_g\|)^2} T_{\mu\nu}[E_{sqr}(\|\vec{v}_p\| \|\vec{v}_g\|), g_{\alpha\beta}(\|\vec{v}_p\| \|\vec{v}_g\|)] \\
& \|\vec{v}_p \times \vec{v}_g\|^2 + (\vec{v}_p \cdot \vec{v}_g)^2 = (\|\vec{v}_p\| \|\vec{v}_g\|)^2 \\
& \|\vec{v}_p \times \vec{v}_g\|^2 + (\vec{v}_p \cdot \vec{v}_g)^2 = (v_p v_g)^2
\end{aligned}$$

- (Modified Lagrange's Identity), v_p is the Phase Velocity, v_g is the Group Velocity.

$$\begin{aligned}
G_{\mu\nu}^{eff}[g_{\alpha\beta}(v_p)] + \Lambda^{eff}(v_p)g_{\mu\nu}^{eff}(v_p) &= \frac{8\pi}{1} \frac{G}{(v_p)^4} T_{\mu\nu}^{eff}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)] \quad (1) \\
G_{\mu\nu}^{eff}[g_{\alpha\beta}(v_g)] + \Lambda^{eff}(v_g)g_{\mu\nu}^{eff}(v_g) &= \frac{8\pi}{1} \frac{G}{(v_g)^4} T_{\mu\nu}^{eff}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)] \\
G_{\mu\nu}^{eff}[g_{\alpha\beta}(v_p \cdot v_g)] + \Lambda^{eff}(v_p \cdot v_g)g_{\mu\nu}^{eff}(v_p \cdot v_g) \\
&= \frac{8\pi}{1} \frac{G}{(v_p \cdot v_g)^2} T_{\mu\nu}^{eff}[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)]
\end{aligned}$$

Where $G_{\mu\nu}^{eff}[g_{\alpha\beta}(v_p)]$ represents the Einstein tensor for phase velocity, $T_{\mu\nu}^{eff}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)]$ is the energy-momentum tensor of effective phase velocity, $G_{\mu\nu}[g_{\alpha\beta}(v_g)]$ represents the Einstein tensor for group velocity, $G_{\mu\nu}^{eff}[g_{\alpha\beta}(v_p \cdot v_g)]$ represents the Einstein tensor for phase velocity and group velocity, $T_{\mu\nu}^{eff}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)]$ is the energy-momentum tensor of effective group velocity

$$\begin{aligned}
G_{\mu\nu}[g_{\alpha\beta}(v_p \cdot v_g)] + \Lambda g_{\mu\nu}(v_p \cdot v_g) &= \frac{8\pi}{1} \frac{G}{(v_p \cdot v_g)^2} T_{\mu\nu}[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)] \quad (2) \\
E_{sqr}^2(v_p \cdot v_g) &= (\mathbf{m} \cdot v_p \cdot v_g)^2 + (\mathbf{p})^2 \times (v_p \cdot v_g), E_{sqr}(v_p \cdot v_g) = \gamma(v_p \cdot v_g) \cdot \mathbf{m} \cdot (v_p \cdot v_g) \\
G_{\mu\nu}[g_{\alpha\beta}(v_p \cdot v_g)] + \Lambda g_{\mu\nu}(v_p \cdot v_g) &= \frac{8\pi}{1} \frac{G}{(v_p \cdot v_g)^2} T_{\mu\nu}^{eff}[E_{eff}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)] \\
E_{eff}^2(v_p \cdot v_g) &= (\mathbf{m}_{eff} \cdot v_p \cdot v_g)^2 + (\mathbf{p}_{eff})^2 \cdot (v_p \cdot v_g), E_{eff}(v_p \cdot v_g) = \gamma(v_p \cdot v_g) \times \mathbf{m}_{eff} \cdot (v_p \cdot v_g)
\end{aligned}$$

- $E_{sqr}(v_p \cdot v_g)$ is the special quantum relativity for phase velocity and group velocity, $E_{eff}(v_p \cdot v_g)$ is the special quantum relativity of effective phase velocity and group velocity

$$\begin{aligned}
R_{\mu\nu}[g_{\alpha\beta}(v_{Dp})] - \frac{1}{2}R[g_{\alpha\beta}(v_{Dp})]g_{\mu\nu}(v_{Dp}) + \Lambda g_{\mu\nu}(v_{Dp}) \\
&= \frac{8\pi\sqrt{(\|\vec{v}_p\| \|\vec{v}_g\|)^3}}{\hbar \times (a_{DQ})^2} T_{\mu\nu}[E_{sqr}(v_{Dp}), g_{\alpha\beta}(v_{Dp})] \quad (3) \\
R_{\mu\nu}[g_{\alpha\beta}(v_p)] - \frac{1}{2}R[g_{\alpha\beta}(v_p)]g_{\mu\nu}(v_p) + \Lambda g_{\mu\nu}(v_p) \\
&= \frac{8\pi(v_p)^3}{\hbar \times (a_{DQ})^2} T_{\mu\nu}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)] \\
R_{\mu\nu}[g_{\alpha\beta}(v_g)] - \frac{1}{2}R[g_{\alpha\beta}(v_g)]g_{\mu\nu}(v_g) + \Lambda g_{\mu\nu}(v_g) \\
&= \frac{8\pi(v_g)^3}{\hbar \times (a_{DQ})^2} T_{\mu\nu}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)]
\end{aligned}$$

- Where e represents the electron charge, v_p is the Phase Velocity, v_g is the Group Velocity, a_{DQ} David's Quantum Acceleration, k_c is the Coulomb constant.

$$\begin{aligned}
C &= \frac{4\pi \cdot k_c}{n_{Dp} \cdot Z_o} \quad (4) \\
v_p &= \frac{C}{n} \\
v_{Dp} &= \frac{4\pi \cdot k_c}{n_{Dp} \cdot Z_o}
\end{aligned}$$

- Z_0 it is (Impedance of free space), n_{dp} is the David's quantum refractive index, v_{dp} is the David's velocity of the stationary phase, v_p is the Phase Velocity

$$R_{\mu\nu}[g_{\alpha\beta}(v_p)] - \frac{1}{2}R[g_{\alpha\beta}(v_p)]g_{\mu\nu}(v_p) + \Lambda g_{\mu\nu}(v_p) = \frac{8\pi}{1} \frac{G}{(v_p)^4} T_{\mu\nu}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)]$$

$$T_{\mu\nu}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)] = \left(\rho[E_{sqr}(v_p)] + \frac{p[E_{sqr}(v_p)]}{(v_p)^2} \right) u_\mu[E_{sqr}(v_p), g_{\alpha\beta}(v_p)] u_\nu[E_{sqr}(v_p), g_{\alpha\beta}(v_p)] + p g_{\mu\nu}(v_p)$$

$$R_{\mu\nu}[g_{\alpha\beta}(v_g)] - \frac{1}{2}R[g_{\alpha\beta}(v_g)]g_{\mu\nu}(v_g) + \Lambda g_{\mu\nu}(v_g) = \frac{8\pi}{1} \frac{G}{(v_g)^4} T_{\mu\nu}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)]$$

$$T_{\mu\nu}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)] = \left(\rho[E_{sqr}(v_g)] + \frac{p[E_{sqr}(v_g)]}{(v_g)^2} \right) u_\mu[E_{sqr}(v_g), g_{\alpha\beta}(v_g)] u_\nu[E_{sqr}(v_g), g_{\alpha\beta}(v_g)] + p g_{\mu\nu}(v_g)$$

What does the equation tell us?

- 1) The refractive index determines the medium in which the equation operates.
 - 2) Planck's refractive index is $n = 1$. This is known as the quantum vacuum medium (the quantum David medium), meaning a real medium. Everything within the Planck medium is governed by Planck's laws in a medium of $n = 1$.
 - 3) Wherever relativity operates in this medium, its maximum value is the Planck value.
 - 4) The universal gravitational constant and the speed of light will remain constants because they are within a Planck medium ($n = 1$). This means that anything within a Planck medium obeys Planck's constants.
 - 5) Gravity will be a universal constant as long as it is within this medium. Therefore, when gravity is applied to Planck's constants, it produces an antigravity, just as antienergy or quantum vacuum energy does, because of the refractive index (n); this leads to a rebound.
 - 6) Dark mass is formed as a result of this medium due to the presence of virtual particles.
 - 7) These virtual particles are produced by dark energy, and a portion of dark energy can produce dark mass, and dark mass can produce dark energy.
- The new equation will be known as general quantum relativity, and it defines the medium in which it will operate. Each medium is defined by its refractive index, and all the equation does is follow the properties of that medium.
 - The classical general equation of relativism does not specify the medium in which it will operate; therefore, singularity arises because it takes infinite values. However, if it operates within a quantum vacuum, it obeys the laws of that medium; that is, that medium governs it. For example, if this medium has a refractive index of 1, it is governed by Planck's constants and cannot exceed them because the medium itself determines them.
 - A singularity will not occur because the curvature is Planck's curvature, and the hole will not completely evaporate because the temperature is Planck's temperature. Therefore, the hole is a stable object (a Planck point) that stores information, and this solves the Hawking radiation problem.
 - The reversal occurs at the Planck point, resulting in multiple universes. This is because the reversal occurs in two directions: a normal direction and an opposing direction, creating a universe in dark energy. The other side represents the opposing dark energy.
 - This explains how a rebound occurs in a medium where antigravity is formed or because the ultimate values of Planck's constants do not exceed the recoil event; for every action there is an equal and opposite reaction because of the refractive index (n). This causes gravity at the Planck scale to cease and act as antigravity, or quantum vacuum energy, due to the refractive index creating dark energy. The problem of the cosmological constant is solved by the fact that it has become dark energy due to recoil. In this way, time does not reach zero, but it allows the universe to expand again because the shortest time is the Planck time, and this time occurs within the Planck point.
 - Holography means that spacetime does not exceed the Planck length, Planck curvature, and Planck volume. This prevents the formation of information exceeding the Planck limit; that is,

the number of bits within the volume and on its surface corresponds to the Planck length and volume.

- Virtual particles may be part of David's quantum medium, or they may be virtual particles with Planck mass, Planck density, and Planck volume—that is, they reach Planck constants. This may explain the addition of a hidden mass known as dark mass.
- $mC \psi = i\hbar \gamma^\mu \partial_\mu \psi$
- $m\mathbf{v}_p \psi = i\hbar \gamma^\mu \partial_\mu \psi$
- $m\mathbf{v}_g \psi = i\hbar \gamma^\mu \partial_\mu \psi$
- $m\sqrt{\mathbf{v}_p \cdot \mathbf{v}_g} \psi = i\hbar \gamma^\mu \partial_\mu \psi$

Dirac quantitative equation

$$E = h\nu = \frac{h\nu_p}{\lambda} = \frac{h\nu_g}{\lambda} = \frac{h\sqrt{\|\vec{v}_p\| \|\vec{v}_g\|}}{\lambda} = \frac{h\sqrt{\mathbf{v}_p \cdot \mathbf{v}_g}}{\lambda}$$

This explains why Planck's orbit appears in the equation.

Properties of a David Quantum Medium (n = 1)

- 1) Within this medium, Planck's constants are applied and are not exceeded
- 2) Virtual particles are part of this medium.
- 3) Infinite absolute values cannot be assumed within this medium because the maximum values attainable are Planck's constants.
- 4) Virtual particles act as a hidden space because the refractive index of the medium is 1. They behave like a glass test cup inside another glass test cup, with oil in between. Therefore, they are neglected and not considered part of the medium; they are called a vacuum. This means that n=1 represents the disappearance of molecules from the medium and does not represent a vacuum.
- 5) Another important characteristic of the medium is that it is a real medium.
- 6) This medium is the fundamental component of all other mediums; in other words, this medium determines the stability of the force of four or five (if it exists in nature).

3. These laws have been modified from the mix Planck laws

- How quantum entanglement occurs?
What happens is that the electron connects to the other electron through space-time, as space-time acts like a quantum tunnel that connects the two electrons. In this way, the electron does not penetrate the speed of light, But in relation to large objects, you see that it has crossed the speed of light.
- This hypothesis was based on scientific foundations, the most important of which is:
 - 1) the connection between relativity and quantum mechanics occurs via quantum entanglement and loop gravitational entanglement.
 - 2) quantum entanglement occurs by the contraction of space-time.
 - 3) space-time contraction occurs by space-time absorbing energy.
 - 4) the quantum jump of the electron occurs as a result of the contraction of space-time.

4. Derivation of equations

Completing the derivation of the laws resulting from quantum relativity (quantum world)

$$p = m \cdot v = \hbar \cdot k$$

$$C = \frac{n \cdot v}{\alpha \cdot Z}$$
$$v = \frac{\alpha \cdot C \cdot Z}{n} = \frac{\alpha \cdot v_p \cdot Z}{n} = \frac{\alpha \cdot v_g \cdot Z}{n} = \alpha \cdot C$$

$$p = m \cdot \frac{\alpha \cdot C \cdot Z}{n}$$

$$p = m \cdot C \cdot \frac{\alpha \cdot Z}{n}$$

$$E = p \cdot C$$

$$E_n = p \cdot C \cdot \frac{\alpha \cdot Z}{n}$$

$$p = m \cdot C$$

$$E_n = p \cdot C$$

$$p = m \cdot v = \hbar \cdot k$$

This is derivation number 1

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi}{1} \frac{G}{(C)^4} T_{\mu\nu}$$

$$E_n = p \cdot C$$

$$p = m \cdot v = \hbar \cdot k$$

$$C = \frac{E_n}{p}$$

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi \times (p)^4}{1} \frac{G}{(E_n)^4} T_{\mu\nu}$$

$$p = m \cdot v = \hbar \cdot k$$

$$E_n = \hbar_{(p)} \cdot v = \hbar \cdot C \cdot k = \hbar \cdot \omega$$

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi \cdot (\hbar \cdot k)^4}{1} \frac{G}{(\hbar \cdot \omega)^4} T_{\mu\nu}$$

$$v_p = \frac{\omega}{k}$$

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi}{1} \frac{G}{(v_p)^4} T_{\mu\nu}$$

$$v_p = \frac{C}{n}$$

v_p is the Phase Velocity

$$v_g = \frac{\Delta\omega}{\Delta k}$$

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi}{1} \frac{G}{(v_g)^4} T_{\mu\nu}$$

v_g is the Group Velocity

This is derivation number 2

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi}{1} \frac{G}{(v_p)^4} T_{\mu\nu} \frac{(C)^4}{(C)^4}$$

$$G_{(v_p)} = \frac{G}{(v_p)^4}$$

$G_{(v_p)}$ is the gravitational coefficient for Phase Velocity

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi}{1} \frac{G}{(C)^4} T_{\mu\nu}$$

$$c \rightarrow v_p$$

$$G_{\mu\nu}[g_{\alpha\beta}(v_p)] + \Lambda(v_p)g_{\mu\nu}(v_p) = \frac{8\pi}{1} \frac{G}{(v_p)^4} T_{\mu\nu}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)]$$

$$T_{\mu\nu}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)] = \left(\rho[E_{sqr}(v_p)] + \frac{p[E_{sqr}(v_p)]}{(v_p)^2} \right) u_\mu[E_{sqr}(v_p), g_{\alpha\beta}(v_p)] u_\nu[E_{sqr}(v_p), g_{\alpha\beta}(v_p)] + p g_{\mu\nu}(v_p)$$

$$E_{sqr}^2(v_p) = (m_{(v_p)} \cdot (v_p)^2)^2 + (p \cdot v_p)^2, KE_{qke}(v_p) = (\gamma_{v_p} - 1) \times m_{(v_p)} \cdot (v_p)^2$$

$$\gamma_{v_p} = \frac{1}{\sqrt{1 - \frac{v^2}{(v_p)^2}}}, E_{sqr}(v_p) = \gamma_{v_p} \cdot m_{(v_p)} \cdot (v_p)^2, \gamma_{v_p} = 1 + \frac{1}{2} \frac{v^2}{(v_p)^2}$$

$E_{sqr}(v_p)$ is the special quantum relativity for phase velocity, $KE_{qke}(v_p)$ is the quantum kinetic energy for phase velocity

$$T_{\mu\nu}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)] = \frac{2}{\sqrt{-g_{\alpha\beta}(v_p)}} \frac{\delta S_{matter}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)]}{\delta g^{\mu\nu}(v_p)}$$

$T_{\mu\nu}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)]$ is the energy-momentum tensor for phase velocity and Phase Velocity
This is derivation number 3

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi}{1} \frac{G}{(C)^4} T_{\mu\nu}$$

$$c \rightarrow v_g$$

$$G_{\mu\nu}[g_{\alpha\beta}(v_g)] + \Lambda(v_g)g_{\mu\nu}(v_g) = \frac{8\pi}{1} \frac{G}{(v_g)^4} T_{\mu\nu}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)]$$

$$T_{\mu\nu}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)] = \left(\rho[E_{sqr}(v_g)] + \frac{p[E_{sqr}(v_g)]}{(v_g)^2} \right) u_\mu[E_{sqr}(v_g), g_{\alpha\beta}(v_g)] u_\nu[E_{sqr}(v_g), g_{\alpha\beta}(v_g)] + p g_{\mu\nu}(v_g)$$

$$E_{sqr}^2(v_g) = (m_{(v_g)} \cdot (v_g)^2)^2 + (p \cdot v_g)^2, KE_{qke}(v_g) = (\gamma_{v_g} - 1) \times m_{(v_g)} \cdot (v_g)^2$$

$$\gamma_{v_g} = \frac{1}{\sqrt{1 - \frac{v^2}{(v_g)^2}}}, E_{sqr}(v_g) = \gamma_{v_g} \cdot m_{(v_g)} \cdot (v_g)^2, \gamma_{v_g} = 1 + \frac{1}{2} \frac{v^2}{(v_g)^2}$$

$E_{sqr}(v_g)$ is the special quantum relativity for group velocity, $KE_{qke}(v_g)$ is the quantum kinetic energy for group velocity

$$T_{\mu\nu}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)] = \frac{2}{\sqrt{-g_{\alpha\beta}(v_g)}} \frac{\delta S_{matter}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)]}{\delta g^{\mu\nu}(v_g)}$$

$T_{\mu\nu}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)]$ is the energy-momentum tensor for phase velocity and group velocity
This is derivation number 4

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi}{1} \frac{G}{(C)^4} T_{\mu\nu}$$

$$c \rightarrow v_p \times v_g$$

$$G_{\mu\nu}[g_{\alpha\beta}(v_p \cdot v_g)] + \Lambda(v_p \cdot v_g)g_{\mu\nu}(v_p \cdot v_g) = \frac{8\pi}{1} \frac{G}{(v_p \cdot v_g)^2} T_{\mu\nu}[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)]$$

$$T_{\mu\nu}[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)] = \left(\rho[E_{sqr}(v_p \cdot v_g)] + \frac{p[E_{sqr}(v_p \cdot v_g)]}{v_p \cdot v_g} \right) u_\mu[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)] u_\nu[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)] + p g_{\mu\nu}(v_p \cdot v_g)$$

$$E_{sqr}^2(v_p \cdot v_g) = (m_{(v_p \cdot v_g)} \cdot (v_p \cdot v_g)^2)^2 + (p)^2 (v_p \cdot v_g), E_{sqr}(v_p \cdot v_g) = \gamma_{v_p \cdot v_g} \cdot m_{(v_p \cdot v_g)} \cdot v_p \cdot v_g$$

$$\gamma_{v_p \cdot v_g} = \frac{1}{\sqrt{1 - \frac{v^2}{(v_p \cdot v_g)^2}}}, KE_{qke}(v_p \cdot v_g) = (\gamma_{v_p \cdot v_g} - 1) \cdot m_{(v_p \cdot v_g)} \cdot v_p \cdot v_g, \gamma_{v_p \cdot v_g} = 1 + \frac{1}{2} \frac{v^2}{v_p \cdot v_g}$$

$E_{sqr}(v_g)$ is the special quantum relativity for Phase Velocity and v_g Group Velocity, $KE_{qke}(v_g)$ is the quantum kinetic energy for Phase Velocity and v_g Group Velocity

$$T_{\mu\nu}[E_{sqr}(\|\vec{v}_p\| \parallel \vec{v}_g\|), g_{\alpha\beta}(\|\vec{v}_p\| \parallel \vec{v}_g\|)] = \frac{2}{\sqrt{-g_{\alpha\beta}(\|\vec{v}_p\| \parallel \vec{v}_g\|)}} \frac{\delta S_{\text{matter}}[E_{sqr}(\|\vec{v}_p\| \parallel \vec{v}_g\|), g_{\alpha\beta}(\|\vec{v}_p\| \parallel \vec{v}_g\|)]}{\delta g^{\mu\nu}(\|\vec{v}_p\| \parallel \vec{v}_g\|)}$$

$T_{\mu\nu}[E_{sqr}(\|\vec{v}_p\| \parallel \vec{v}_g\|), g_{\alpha\beta}(\|\vec{v}_p\| \parallel \vec{v}_g\|)]$ is the energy-momentum tensor for phase velocity and group velocity

This is derivation number 5

$$v_p = \frac{C}{n}$$

$$n = \frac{C}{v_p}$$

$$v_p = \frac{x}{t} = \frac{l_p}{t_p} = C_{\text{Planck}}$$

$$n = \frac{C}{C_{\text{Planck}}} = 1$$

- $v_p = v_g = C_{\text{Planck}}$
- $n = \frac{C}{C_{\text{Planck}}} = 1$
- v_p is the Phase Velocity, v_g is the Group Velocity

This is derivation number 6

$$T_{H(v_p)} = \frac{\hbar \cdot (v_p)^3}{8\pi \cdot G \cdot M \cdot k_B}$$

$$T_{H(v_g)} = \frac{\hbar \cdot (v_g)^3}{8\pi \cdot G \cdot M \cdot k_B}$$

$$T_{H(v_p \times v_g)} = \frac{\hbar \cdot \sqrt{(\|\vec{v}_p\| \parallel \vec{v}_g\|)^3}}{8\pi \cdot G \cdot M \cdot k_B} = \frac{\hbar \cdot \sqrt{(v_p \cdot v_g)^3}}{8\pi \cdot G \cdot M \cdot k_B}$$

v_p is the Phase Velocity, $T_{H(v_p \times v_g)}$ is the Hawking temperature for phase velocity and group velocity

This is derivation number 7

$$ds^2 = - (v_p v_g) dt^2 + a(t)^2 \left[\frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right]$$

- (Modified FLRW Metric), v_p is the Phase Velocity, v_g is the Group Velocity.

$$G_{\mu\nu}[g_{\alpha\beta}(v_p)] = \frac{8\pi}{1} \frac{G}{(v_p)^4} T_{\mu\nu}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)]$$

$$G_{\mu\nu}^{eff}[g_{\alpha\beta}(v_p)] = \frac{8\pi}{1} \frac{G}{(v_p)^4} T_{\mu\nu}^{eff}[E_{eff}(v_p), g_{\alpha\beta}(v_p)]$$

$$G_{\mu\nu}[g_{\alpha\beta}(v_g)] = \frac{8\pi}{1} \frac{G}{(v_g)^4} T_{\mu\nu}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)]$$

$$G_{\mu\nu}^{eff}[g_{\alpha\beta}(v_g)] = \frac{8\pi}{1} \frac{G}{(v_g)^4} T_{\mu\nu}^{eff}[E_{eff}(v_g), g_{\alpha\beta}(v_g)]$$

$$G_{\mu\nu}[g_{\alpha\beta}(v_p \cdot v_g)] = \frac{8\pi}{1} \frac{G}{(v_p \cdot v_g)^2} T_{\mu\nu}[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)]$$

$$G_{\mu\nu}^{eff}[g_{\alpha\beta}(v_p \cdot v_g)] = \frac{8\pi}{1} \frac{G}{(v_p \cdot v_g)^2} T_{\mu\nu}^{eff}[E_{eff}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)]$$

My equations allow us to solve these problems:

- The singularity problem: Since $n = 1$ represents the David quantum medium in which Planck's laws operate, such as Planck density and Planck curvature, the material is not allowed to be compressed beyond the Planck volume, and we conclude that singularities do not occur.

- Solving the Big Bang problem: The equation assumes that if $n = 1$, meaning the refractive index of the medium obeys Planck's laws, then $v_p = v_g = c$. In this case, the universe would have Planck density, Planck volume, and Planck curvature. There would be no singularity at the moment of the Big Bang due to a disturbance in the refractive index, causing $n \neq 1$. This led to the separation of v_g from v_p ,

i.e., $v_p \neq v_g$. As a result of this separation, matter that follows v_g emerged, and the radiation that follows v_p appeared.

- Solves the mystery of dark matter: Since the law depends on $\frac{G}{(v_p)^4}$ and $\frac{G}{(v_g)^4}$, any slight change in the refractive index alters the gravitational force. We don't need additional mass; rather, this force is a result of the effect of the refractive index of v_p and v_g , which alters the speed and thus leads to an effect on gravity.
- Solving the Dark Energy Problem: Since the phase velocity v_p is decoupled from the group velocity v_g . Since the phase velocity can exceed the speed of light depending on the medium ($v_p \neq v_g$), the expansion of the universe is the propagation of phase waves in a Planck medium. Therefore, dark energy is the refractive index pressure resulting from the medium's attempt to return to $n=1$.
- Solving the information paradox: Since $n=1$ prevents singularity, information does not disappear but is stored in vibrations within the Planck medium inside the Planck point (David's center), where singularity does not occur. It can escape through a disturbance in the refractive index, forming Hawking radiation.
- Agreement with quantum mechanics, where spacetime is treated as a wave because it is affected by the refractive index of v_p and v_g in the field equation.

$$v_g = \frac{Z_{eff} \cdot C \cdot \alpha}{n}, v_p = \frac{n \cdot C}{Z_{eff} \cdot \alpha}$$

$$v_g = C \cdot \alpha, v_p = \frac{C}{\alpha}$$

α is the fine-structure constant, Z_{eff} is the Effective Nuclear Charge, n is the energy level, C is the speed of light, Effective Group Velocity of the Electron, Effective phase Velocity of the Electron

$$n(t) = \sqrt{\frac{\kappa T_{00}(t)}{\Lambda}}$$

$$2KE_{DE} = \hbar \cdot \alpha \cdot \left(\frac{4\pi \cdot k_c}{Z_o} \right) \cdot \frac{1}{r_n}, 2KE_{DE} = \hbar \cdot \alpha \cdot (v_p) \cdot \frac{1}{r_n}$$

$$2KE_{DE} = \hbar \cdot \alpha \cdot \left(\frac{(Z_{eff})^2 \cdot 4\pi \cdot k_c}{Z_o} \right) \cdot \frac{1}{r_n}, 2KE_{DE} = \hbar \cdot \alpha \cdot (v_g) \cdot \frac{1}{r_n}$$

$$2KE_{DE} = \hbar \cdot \alpha \cdot \left(\frac{(Z_{eff})^2 \cdot 4\pi \cdot k_c}{Z_o} \right) \cdot \frac{1}{r_n}, 2KE_{DE} = \hbar \cdot \alpha \cdot (\sqrt{v_p \cdot v_g}) \cdot \frac{1}{r_n}$$

$$2KE_{DE} = \hbar \cdot \alpha \cdot \left(\frac{(Z_{eff})^2 \cdot 4\pi \cdot k_c}{Z_o} \right) \cdot \frac{1}{r_n}, 2KE_{DE} = \hbar \cdot \alpha \cdot \left(\sqrt{\|\vec{v}_p\| \|\vec{v}_g\|} \right) \cdot \frac{1}{r_n}$$

KE_{DE} is the David's Kinetic Energy and Energy Equivalence

$$(F_{DE})^2 = \left(\hbar \cdot \alpha \cdot \frac{v_p}{(r_n)^2} \right)^2 + \left(\hbar \cdot \alpha \cdot \left(\frac{4\pi \cdot k_c}{Z_o} \right) \cdot \frac{1}{(r_n)^2} \right)^2$$

$$(F_{DE})^2 = \left(\hbar \cdot \alpha \cdot \frac{v_g}{(r_n)^2} \right)^2 + \left(\hbar \cdot \alpha \cdot \left(\frac{4\pi \cdot k_c}{Z_o} \right) \cdot \frac{1}{(r_n)^2} \right)^2$$

$$(F_{DE})^2 = \left(\hbar \cdot \alpha \cdot \frac{\sqrt{v_p \cdot v_g}}{(r_n)^2} \right)^2 + \left(\hbar \cdot \alpha \cdot \left(\frac{4\pi \cdot k_c}{Z_o} \right) \cdot \frac{1}{(r_n)^2} \right)^2$$

$$(F_{DE})^2 = \left(\hbar \cdot \alpha \cdot \frac{\sqrt{\|\vec{v}_p\| \|\vec{v}_g\|}}{(r_n)^2} \right)^2 + \left(\hbar \cdot \alpha \cdot \left(\frac{4\pi \cdot k_c}{Z_o} \right) \cdot \frac{1}{(r_n)^2} \right)^2$$

F_{DE} is the David's Force and Energy Equivalence

1. Temporal Curvature (Zero Curvature Condition)

At the Planck scale, the temporal component of the Einstein tensor vanishes due to the exact cancellation between energy density and the cosmological constant.

Formula: $G_{00} = \kappa T_{00} - \Lambda = 0$

Physical Significance: This represents a state of "temporal flatness," preventing the formation of a singularity.

2. Spatial Curvature (Expansionary Curvature)

Unlike the temporal component, the spatial components involve the addition of vacuum energy and material pressure, driving an explosive outward metric expansion.

Formula: $G_{ii} = \kappa T_{ii} - \Lambda \approx -1.92419 \times 10^{71} \text{ m}^{-2}$

Physical Significance: This provides the geometric "push" required for the Big Bang, where the curvature is purely directed into expanding the volume of space.

3. Expansion Dynamics (Hubble Parameter)

In the case of $G_{00} = 0$, Hubble's law is as follows.

Formula: $H = \sqrt{\frac{c^2}{6} G_{ii}} \approx 5.3687121708 \times 10^{43} \text{ s}^{-1}$

Physical Significance: The universe doubles its size every $\sim 10^{-44}$ seconds, marking the highest possible frequency of cosmic expansion.

4. The physical scale of the universe starting from the Planck length $a(t)$

The evolution of the cosmic size over time follows an exponential growth curve due to the constant energy density at the instant of Planck equilibrium.

Formula: $a(t) = l_p \cdot e^{Ht}$

5. Variable Energy Density ($\rho(t)$)

As the scale factor $a(t)$ increases, the energy density dilutes according to the Stefan-Boltzmann.

Formula $\rho(t) = \rho_p \cdot \left(\frac{l_p}{a(t)} \right)^4$

Physical Significance: This dilution is what eventually breaks the $G_{00} = 0$ equilibrium, allowing gravity to reappear and matter to condense into stars and galaxies.

6. Refractive Index Evolution (n)

"Optical Rigidity" of spacetime is linked to the energy density.

Formula: $n(t) = \sqrt{\frac{\kappa T_{00}(t)}{\Lambda}}, \Lambda(t) = \Lambda \times n(t)^2$

$$n(t) = \sqrt{\frac{\rho(t)}{\rho_p}} = \sqrt{\left(\frac{l_p}{a(t)} \right)^4}, n(t) = \frac{l_p}{R_H} = \frac{c}{\sqrt{(v_p \cdot v_g)}} = \frac{c}{\sqrt{(\|\vec{v}_p\| \|\vec{v}_g\|)}}$$

$$(v_p \neq v_g) = c^2$$

$$n > 1$$

Since the product of the two velocities is c^2 , relativity will shift to a Planck medium because the result is c^2 . This occurs because there are two mediums: one interacts with the phase velocity and not the group velocity due to the separation of the group velocity from the phase velocity. While the

other interacts with the group velocity and not the phase velocity due to the separation of the phase velocity from the group velocity.

$$G_{\mu\nu}[g_{\alpha\beta}(v_p)] = \frac{8\pi}{1} \frac{G}{(v_p)^4} T_{\mu\nu} [E_{sqr(v_p)}, g_{\alpha\beta}(v_p)]$$

$$G_{\mu\nu}[g_{\alpha\beta}(v_g)] + \Lambda(v_g)g_{\mu\nu}(v_g) = \frac{8\pi}{1} \frac{G}{(v_g)^4} T_{\mu\nu} [E_{sqr(v_g)}, g_{\alpha\beta}(v_g)]$$

So that the result of the two means equals 1 when we divide them together, in the case of the equation that includes them.

$$G_{\mu\nu}[g_{\alpha\beta}(\|\vec{v}_p\| \|\vec{v}_g\|)] + \Lambda(\|\vec{v}_p\| \|\vec{v}_g\|)g_{\mu\nu}(\|\vec{v}_p\| \|\vec{v}_g\|) = \frac{8\pi}{1} \frac{G}{(\|\vec{v}_p\| \|\vec{v}_g\|)^2} T_{\mu\nu} [E_{sqr(\|\vec{v}_p\| \|\vec{v}_g\|)}, g_{\alpha\beta}(\|\vec{v}_p\| \|\vec{v}_g\|)]$$

$$\begin{aligned} G_{\mu\nu}[g_{\alpha\beta}(v_p \cdot v_g)] + \Lambda(v_p \cdot v_g)g_{\mu\nu}(v_p \cdot v_g) \\ = \frac{8\pi}{1} \frac{G}{(v_p \cdot v_g)^2} T_{\mu\nu} [E_{sqr(v_p \cdot v_g)}, g_{\alpha\beta}(v_p \cdot v_g)] \end{aligned}$$

$$(v_p > v_g) > c^2$$

$$n < 1$$

- In some media, such as Epsilon-Near-Zero (ENZ), this phenomenon occurs. It happens at a plasma frequency that affects the phase velocity, making it faster than light, thus causing the resultant $v_p \cdot v_g$ to be greater than c^2 .

$$(l_{DQ})^2 = \frac{\hbar \cdot G}{\sqrt{(\|\vec{v}_p\| \|\vec{v}_g\|)^3}} = \frac{\hbar \cdot G}{\sqrt{(v_p \cdot v_g)^3}}$$

l_{DQ} Quantum length of David

$$(l_{DQ})^2 = (t_{DQ})^2 \cdot \|\vec{v}_p\| \|\vec{v}_g\| = (t_{DQ})^2 \cdot v_p \cdot v_g$$

$$v_{Dp} = \frac{4\pi \cdot k_c}{n \cdot Z_o}$$

$$E_{DQ} = m_p \cdot v_p \cdot v_g = m_p \cdot \|\vec{v}_p\| \|\vec{v}_g\|$$

$$v_{Dp} = \frac{4\pi \cdot k_c}{n \cdot Z_o}$$

$$v_p = \frac{C}{n}$$

v_p is the Phase Velocity

$$\sqrt{v_p \cdot v_g} = \frac{4\pi \cdot k_c}{n \cdot Z_o}$$

v_{Dp} David's velocity of the stationary phase

$$m_{DQ} = \sqrt{\frac{\hbar \cdot \sqrt{\|\vec{v}_p\| \|\vec{v}_g\|}}{G}} = \sqrt{\frac{\hbar \cdot \sqrt{v_p \cdot v_g}}{G}} = \sqrt{\frac{\hbar \cdot (4\pi \cdot k_c)}{G \cdot Z_o}}$$

m_{DQ} Quantum block of David

$$l_{DQ} = \sqrt{\frac{\hbar \cdot G}{\sqrt{(v_p \cdot v_g)^3}}} = \sqrt{\frac{\hbar \cdot G}{\sqrt{(\|\vec{v}_p\| \|\vec{v}_g\|)^3}}} = \sqrt{\frac{\hbar \cdot G \cdot (Z_o)^3}{(4\pi \cdot k_c)^3}}$$

l_{DQ} Quantum length of David

$$t_{DQ} = \sqrt{\frac{\hbar \cdot G}{\sqrt{(v_p \cdot v_g)^5}}} = \sqrt{\frac{\hbar \cdot G}{\sqrt{(\|\vec{v}_p\| \|\vec{v}_g\|)^5}}} = \sqrt{\frac{\hbar \cdot G \cdot (Z_o)^5}{(4\pi \cdot k_c)^5}}$$

t_{DQ} Quantum time of David

$$E_{DQ} = m_p \cdot \|\vec{v}_p\| \|\vec{v}_g\| = m_p \cdot v_p \cdot v_g = m_p \cdot \left(\frac{4\pi \cdot k_c}{Z_o}\right)^2$$

E_{DQ} David's Quantum Energy

$$T_{DQ} = \frac{E_p}{k_B} = \frac{m_p \|\vec{v}_p\| \|\vec{v}_g\|}{k_B} = \frac{m_p \cdot (4\pi \cdot k_c)^2}{k_B \cdot Z_o} = \sqrt{\frac{\hbar \cdot \sqrt{(\|\vec{v}_p\| \|\vec{v}_g\|)^5}}{G \cdot k_B^2}}$$

$$T_{DQ} = \frac{E_p}{k_B} = \frac{m_p v_p \cdot v_g}{k_B} = \frac{m_p \cdot (4\pi \times k_c)^2}{k_B \cdot Z_o} = \sqrt{\frac{\hbar \cdot \sqrt{(v_p \cdot v_g)^5}}{G \cdot k_B^2}}$$

T_{DQ} David's quantum temperature

$$F_{DQ} = \frac{(\|\vec{v}_p\| \|\vec{v}_g\|)^2}{G} = \frac{(v_p \cdot v_g)^2}{G} = \frac{(4\pi \cdot k_c)^4}{G \cdot Z_o}$$

F_{DQ} David's Quantum Force

$$a_{DQ} = \frac{F_p}{m_p} = \frac{\sqrt{(\|\vec{v}_p\| \|\vec{v}_g\|)^7}}{\hbar \cdot G} = \frac{\sqrt{(v_p \cdot v_g)^7}}{\hbar \cdot G} = \frac{(4\pi \cdot k_c)^7}{\hbar \cdot G \cdot Z_o}$$

a_{DQ} David's Quantum Acceleration

$$\rho_{DQ} = \frac{m_p}{l_p^3} = \frac{\sqrt{(\|\vec{v}_p\| \|\vec{v}_g\|)^5}}{\hbar \cdot G^2} = \frac{\sqrt{(v_p \cdot v_g)^5}}{\hbar \cdot G^2} = \frac{(4\pi \cdot k_c)^5}{\hbar \cdot G^2 \cdot (Z_o)^5}$$

ρ_{DQ} David's quantum density

$$P_{DQ} = \frac{\sqrt{(\|\vec{v}_p\| \|\vec{v}_g\|)^7}}{\hbar \cdot G^2} = \frac{\sqrt{(v_p \cdot v_g)^7}}{\hbar \cdot G^2} = \frac{(4\pi \cdot k_c)^7}{\hbar \cdot G^2 \cdot (Z_o)^7}$$

P_{DQ} David's Quantum Pressure

$$p_{DQ} = m_p \cdot \sqrt{\|\vec{v}_p\| \|\vec{v}_g\|} = m_p \cdot \left(\frac{4\pi \cdot k_c}{Z_o}\right)$$

$$= \sqrt{\frac{\hbar \cdot \sqrt{(\|\vec{v}_p\| \|\vec{v}_g\|)^3}}{G}}$$

$$p_{DQ} = m_p \cdot \sqrt{v_p \cdot v_g} = m_p \cdot \left(\frac{4\pi \cdot k_c}{Z_o} \right) = \sqrt{\frac{\hbar \cdot \sqrt{(v_p \cdot v_g)^3}}{G}}$$

p_{DQ} David's Momentum Quantity

$$F_{DE} = m \cdot (\alpha)^2 \cdot \frac{\|\vec{v}_p\| \|\vec{v}_g\|}{r} + p \cdot \alpha \cdot \frac{\sqrt{\|\vec{v}_p\| \|\vec{v}_g\|}}{r}$$

$$F_{DE} = m \cdot (\alpha)^2 \cdot \frac{v_p \cdot v_g}{r} + p \cdot \alpha \cdot \frac{\sqrt{v_p \cdot v_g}}{r}$$

$$F_{DE} = \hbar \cdot \alpha \cdot \frac{v_p}{(r_n)^2} + \hbar \cdot \alpha \cdot \left(\frac{4\pi \cdot k_c}{Z_o} \right) \cdot \frac{1}{(r_n)^2}$$

$$F_{DE} = \hbar \cdot \alpha \cdot \frac{v_g}{(r_n)^2} + \hbar \cdot \alpha \cdot \left(\frac{4\pi \cdot k_c}{Z_o} \right) \cdot \frac{1}{(r_n)^2}$$

$$F_{DE} = \hbar \cdot \alpha \cdot \frac{\sqrt{\|\vec{v}_p\| \|\vec{v}_g\|}}{(r_n)^2} + \hbar \cdot \alpha \cdot \left(\frac{4\pi \cdot k_c}{Z_o} \right) \cdot \frac{1}{(r_n)^2}$$

$$F_{DE} = \hbar \cdot \alpha \cdot \frac{\sqrt{v_p \cdot v_g}}{(r_n)^2} + \hbar \cdot \alpha \cdot \left(\frac{4\pi \cdot k_c}{Z_o} \right) \cdot \frac{1}{(r_n)^2}$$

F_{DE} is the David's Force and Energy Equivalence

$$\alpha = \frac{1}{4\pi \cdot \epsilon_0} \frac{(e)^2}{\hbar \cdot \sqrt{\|\vec{v}_p\| \|\vec{v}_g\|}} = \frac{1}{4\pi \cdot \epsilon_0} \frac{(e)^2}{\hbar \cdot \sqrt{v_p \cdot v_g}}$$

$$\alpha = \frac{\hbar}{m_e \cdot \sqrt{\|\vec{v}_p\| \|\vec{v}_g\|} \cdot r_n} = \frac{\hbar}{m_e \cdot \sqrt{v_p \cdot v_g} \cdot r_n}$$

The fine structure constant determines the balance and interaction between two worlds: the world of particles and waves and the world of large objects, i.e., it is the separator.

$$\sqrt{\|\vec{v}_p\| \|\vec{v}_g\|} = \frac{4\pi \cdot k_c}{Z_o}$$

$$\sqrt{v_p \cdot v_g} = \frac{4\pi \cdot k_c}{Z_o}$$

Z_0 it is (Impedance of free space)

$$F_{DQ} = \frac{n \cdot \hbar \cdot \sqrt{\|\vec{v}_p\| \|\vec{v}_g\|}}{(r_n)^2} = \frac{n \cdot \hbar \cdot \sqrt{v_p \cdot v_g}}{(r_n)^2}$$

$$F_{DQ} = \frac{\hbar \cdot a_{DQ}}{\sqrt{(\|\vec{v}_p\| \|\vec{v}_g\|)^3}}$$

$$F_{DQ} = \frac{\hbar \times a_{DQ}}{\sqrt{(v_p \cdot v_g)^3}}$$

- F_{DQ} David's Quantum Force, $\hbar$ is the reduced Planck constant, v_{Dp} is the David's velocity of the stationary phase, v_p is the Phase Velocity, a_{DQ} David's Quantum Acceleration

$$F_{DE} = n\hbar \cdot \frac{v_e}{(r_n)^2}$$

$$F_{DE} = \hbar \cdot \alpha \cdot \frac{v_p}{(r_n)^2}$$

$$F_{DE} = \hbar \cdot \alpha \cdot \frac{v_g}{(r_n)^2}$$

F_{DE} is the David's Force and Energy Equivalence, n is the energy level, v_e is the electron speed.

$$G_{\mu\nu}[g_{\alpha\beta}(v_p)] + \Lambda(v_p)g_{\mu\nu}(v_p) = \frac{8\pi}{1} \frac{G}{(v_p)^4} T_{\mu\nu}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)]$$

$$G_{\mu\nu}[g_{\alpha\beta}(v_g)] + \Lambda(v_g)g_{\mu\nu}(v_g) = \frac{8\pi}{1} \frac{G}{(v_g)^4} T_{\mu\nu}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)]$$

$$v_g = \frac{Z_{eff} \cdot C \cdot \alpha}{n}, v_p = \frac{n \cdot C}{Z_{eff} \cdot \alpha}$$

- Explanation of the quantum leap of the hydrogen atom
- The electron in the first energy orbit

$$Z_{eff} = 1$$

$$n = 1$$

$$\alpha = \frac{1}{137}$$

$$v_p = \frac{n \cdot C}{Z_{eff} \cdot \alpha} = 137 C$$

$$v_g = \frac{Z_{eff} \cdot C \cdot \alpha}{n} = \frac{C}{137}$$

- The electron in the second energy orbit

$$Z_{eff} = 1$$

$$n = 2$$

$$\alpha = \frac{1}{137}$$

$$v_p = \frac{n \cdot C}{Z_{eff} \cdot \alpha} = 274 C$$

$$v_g = \frac{Z_{eff} \cdot C \cdot \alpha}{n} = \frac{C}{274}$$

- This data explains why gravity increases 16 times when an electron moves from orbit 1 to orbit 2 due to the group speed.
- Meanwhile, we observe that spacetime expands 16 times due to the phase velocity, which is faster than the speed of light.
- During a quantum jump, spacetime expands due to the phase velocity, and when it returns to its original state, it contracts due to the group velocity.

$$G_{\mu\nu}[g_{\alpha\beta}(v_p \cdot v_g)] + \Lambda(v_p \cdot v_g)g_{\mu\nu}(v_p \cdot v_g) = \frac{8\pi}{1} \frac{G}{(v_p \cdot v_g)^2} T_{\mu\nu}[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)]$$

- However, because $(v_p \cdot v_g) = c^2$, this equation leads to the conclusion that the spacetime of the big world does not perceive what happens to the electron during its jump.

$$\vec{v}_p \cdot \vec{v}_g = \|\vec{v}_p\| \|\vec{v}_g\| \cos(\theta)$$

$$\|\vec{v}_p \times \vec{v}_g\| = \|\vec{v}_p\| \|\vec{v}_g\| \sin(\theta)$$

$$\|\vec{A} \times \vec{B}\|^2 + (\vec{A} \cdot \vec{B})^2 = \|\vec{A}\|^2 \|\vec{B}\|^2$$

Or

$$\|\vec{A} \times \vec{B}\|^2 + (\vec{A} \cdot \vec{B})^2 = A^2 B^2$$

Or

$$|\vec{A} \times \vec{B}|^2 + (\vec{A} \cdot \vec{B})^2 = A^2 B^2$$

Lagrange's Identity

$$\begin{aligned} \|\vec{v}_p \times \vec{v}_g\|^2 + (\vec{v}_p \cdot \vec{v}_g)^2 \\ = (\|\vec{v}_p\| \|\vec{v}_g\| \cos \theta)^2 + (\|\vec{v}_p\| \|\vec{v}_g\| \sin \theta)^2 \end{aligned}$$

$$\|\vec{v}_p \times \vec{v}_g\|^2 + (\vec{v}_p \cdot \vec{v}_g)^2 = (\|\vec{v}_p\| \|\vec{v}_g\|)^2 (\cos^2 \theta + \sin^2 \theta)$$

$$\|\vec{v}_p \times \vec{v}_g\|^2 + (\vec{v}_p \cdot \vec{v}_g)^2 = v_p^2 v_g^2$$

$$\|\vec{v}_p \times \vec{v}_g\|^2 + (\vec{v}_p \cdot \vec{v}_g)^2 = (v_p v_g)^2$$

$$|\vec{v}_p \times \vec{v}_g|^2 + (\vec{v}_p \cdot \vec{v}_g)^2 = (v_p v_g)^2$$

$$\|\vec{v}_p \times \vec{v}_g\|^2 + (\vec{v}_p \cdot \vec{v}_g)^2 = \|\vec{v}_p\|^2 \|\vec{v}_g\|^2$$

$$\|\vec{v}_p \times \vec{v}_g\|^2 + (\vec{v}_p \cdot \vec{v}_g)^2 = (\|\vec{v}_p\| \|\vec{v}_g\|)^2$$

v_p is the Phase Velocity, v_g is the Group Velocity.

These are some equations after removing the speed of light and putting in the phase speed. The phase velocity was included because it became clear from the derivation, I made that from Einstein's perspective on the speed of light he was focusing on the speed of light in a vacuum and did not consider other media such as water which affect the speed of light as Christian Huygens explained it and therefore this had to be into account in the calculations.

- This will enable us to add the group velocity as a result of adding the phase velocity when the speed of light is constant.
- $\|\vec{v}_p\| \|\vec{v}_g\| = (v_p \cdot v_g) = v_p v_g = \sqrt{(v_p \cdot v_g)^2}$
- $\|\vec{v}_p\|^2 \|\vec{v}_g\|^2 = (v_p \cdot v_g)^2 = v_p^2 v_g^2 = \sqrt{(v_p \cdot v_g)^4}$
- $c \rightarrow v_p$
- $c \rightarrow v_g$
- $c \rightarrow v_p \times v_g$

$$G_{\mu\nu}[g_{\alpha\beta}(v_p)] = \frac{8\pi}{1} \frac{G}{(v_p)^4} T_{\mu\nu} [E_{sqr(v_p)}, g_{\alpha\beta}(v_p)]$$

$$\begin{aligned}
& G_{\mu\nu}[g_{\alpha\beta}(v_g)] + \Lambda(v_g)g_{\mu\nu}(v_g) \\
&= \frac{8\pi}{1} \frac{G}{(v_g)^4} T_{\mu\nu} [E_{sqr}(v_g), g_{\alpha\beta}(v_g)] \\
& G_{\mu\nu}[g_{\alpha\beta}(v_p \cdot v_g)] + \Lambda(v_p \cdot v_g)g_{\mu\nu}(v_p \cdot v_g) \\
&= \frac{8\pi}{1} \frac{G}{(v_p \cdot v_g)^2} T_{\mu\nu} [E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)]
\end{aligned}$$

- General quantitative relativity for David's

$$ds^2 = - \left(1 - \frac{2GM}{(v_p)^2 r} \right) (v_p)^2 dt^2 + \left(1 - \frac{2GM}{(v_p)^2 r} \right)^{-1} dr^2 + r^2 d\Omega^2$$

$$ds^2 = - \left(1 - \frac{2GM}{(v_g)^2 r} \right) (v_g)^2 dt^2 + \left(1 - \frac{2GM}{(v_g)^2 r} \right)^{-1} dr^2 + r^2 d\Omega^2$$

$$ds^2 = - \left(1 - \frac{2GM}{\sqrt{(v_p \cdot v_g)^2} r} \right) v_p \cdot v_g dt^2 + \left(1 - \frac{2GM}{v_p \cdot v_g r} \right)^{-1} dr^2 + r^2 d\Omega^2$$

or

$$ds^2 = - \left(1 - \frac{2GM}{\|\vec{v}_p\| \|\vec{v}_g\| r} \right) v_p \times v_g dt^2 + \left(1 - \frac{2GM}{\|\vec{v}_p\| \|\vec{v}_g\| r} \right)^{-1} dr^2 + r^2 d\Omega^2$$

2. (Modified Schwarzschild Metric)

$$ds^2 = - \left(1 - \frac{2GMr}{\rho^2 (v_p)^2} \right) (v_p)^2 dt^2 - \frac{4GMar}{\rho^2 (v_p)^2} \sin^2 \theta dt d\phi + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 + \left(r^2 + a^2 + \frac{2GMa^2}{\rho^2 (v_p)^2} \sin^2 \theta \right) \sin^2 \theta d\phi^2$$

$$ds^2 = - \left(1 - \frac{2GMr}{\rho^2 (v_g)^2} \right) (v_g)^2 dt^2 - \frac{4GMar}{\rho^2 (v_g)^2} \sin^2 \theta dt d\phi + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 + \left(r^2 + a^2 + \frac{2GMa^2}{\rho^2 (v_g)^2} \sin^2 \theta \right) \sin^2 \theta d\phi^2$$

$$ds^2 = - \left(1 - \frac{2GMr}{\rho^2 \sqrt{(v_p \cdot v_g)^2}} \right) \sqrt{(v_p \cdot v_g)^2} dt^2 - \frac{4GMar}{\rho^2 \sqrt{(v_p \cdot v_g)^2}} \sin^2 \theta dt d\phi + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2$$

$$+ \left(r^2 + a^2 + \frac{2GMa^2}{\rho^2 \sqrt{(v_p \cdot v_g)^2}} \sin^2 \theta \right) \sin^2 \theta d\phi^2$$

or

$$ds^2 = - \left(1 - \frac{2GMr}{\rho^2 \|\vec{v}_p\| \|\vec{v}_g\|} \right) \|\vec{v}_p\| \|\vec{v}_g\| dt^2 - \frac{4GMar}{\rho^2 \|\vec{v}_p\| \|\vec{v}_g\|} \sin^2 \theta dt d\phi + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2$$

$$+ \left(r^2 + a^2 + \frac{2GMa^2}{\rho^2 \|\vec{v}_p\| \|\vec{v}_g\|} \sin^2 \theta \right) \sin^2 \theta d\phi^2$$

$$\rho^2 = r^2 + a^2 \cos^2 \theta$$

$$\Delta = r^2 - \frac{2GMr}{(v_p)^2} + a^2$$

$$\Delta = r^2 - \frac{2GMr}{(v_g)^2} + a^2$$

$$\Delta = r^2 - \frac{2GMr}{\|\vec{v}_p\| \|\vec{v}_g\|} + a^2 = r^2 - \frac{2GMr}{\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2}} + a^2$$

3. (Modified Kerr Metric)

$$ds^2 = - \left(1 - \frac{2GMr - Q^2}{\rho^2 (\vec{v}_p)^2} \right) (\vec{v}_p)^2 dt^2 - \frac{4GMar}{\rho^2 (\vec{v}_p)^2} \sin^2 \theta dt d\phi + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 + \left(r^2 + a^2 + \frac{(2GMr - Q^2)a^2}{\rho^2 (\vec{v}_p)^2} \sin^2 \theta \right) \sin^2 \theta d\phi^2$$

$$ds^2 = - \left(1 - \frac{2GMr - Q^2}{\rho^2 (\vec{v}_g)^2} \right) (\vec{v}_g)^2 dt^2 - \frac{4GMar}{\rho^2 (\vec{v}_g)^2} \sin^2 \theta dt d\phi + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 + \left(r^2 + a^2 + \frac{(2GMr - Q^2)a^2}{\rho^2 (\vec{v}_g)^2} \sin^2 \theta \right) \sin^2 \theta d\phi^2$$

$$ds^2 = - \left(1 - \frac{2GMr - Q^2}{\rho^2 \sqrt{(\vec{v}_p \cdot \vec{v}_g)^2}} \right) \sqrt{(\vec{v}_p \cdot \vec{v}_g)^2} dt^2 - \frac{4GMar}{\rho^2 \sqrt{(\vec{v}_p \cdot \vec{v}_g)^2}} \sin^2 \theta dt d\phi + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2$$

$$+ \left(r^2 + a^2 + \frac{(2GMr - Q^2)a^2}{\rho^2 \sqrt{(\vec{v}_p \cdot \vec{v}_g)^2}} \sin^2 \theta \right) \sin^2 \theta d\phi^2$$

$$ds^2 = - \left(1 - \frac{2GMr - Q^2}{\rho^2 \|\vec{v}_p\| \|\vec{v}_g\|} \right) \|\vec{v}_p\| \|\vec{v}_g\| dt^2 - \frac{4GMar}{\rho^2 \|\vec{v}_p\| \|\vec{v}_g\|} \sin^2 \theta dt d\phi + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2$$

$$+ \left(r^2 + a^2 + \frac{(2GMr - Q^2)a^2}{\rho^2 \|\vec{v}_p\| \|\vec{v}_g\|} \sin^2 \theta \right) \sin^2 \theta d\phi^2$$

4. (Kerr – Newman Metric)

$$R_s = \frac{2GM}{(\vec{v}_p)^2}$$

$$R_s = \frac{2GM}{(\vec{v}_g)^2}$$

$$R_s = \frac{2GM}{\|\vec{v}_p\| \|\vec{v}_g\|} = \frac{2GM}{\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2}}$$

$$u^\mu [E_{sqr}(\vec{v}_g), \quad g_{\alpha\beta}(\vec{v}_g)] u_\mu [E_{sqr}(\vec{v}_g), \quad g_{\alpha\beta}(\vec{v}_g)] = \pm (\vec{v}_g)^2$$

$$p^\mu [E_{sqr}(\vec{v}_g), \quad g_{\alpha\beta}(\vec{v}_g)] p_\mu [E_{sqr}(\vec{v}_g), \quad g_{\alpha\beta}(\vec{v}_g)]$$

$$= \pm (m)^2 \|\vec{v}_p\| \|\vec{v}_g\| (\vec{v}_g)^2$$

$$p^\mu \left[E_{sqr}(\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2}), \quad g_{\alpha\beta}(\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2}) \right] p_\mu \left[E_{sqr}(\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2}), \quad g_{\alpha\beta}(\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2}) \right] = \pm (m)^2 \|\vec{v}_p\| \|\vec{v}_g\| \sqrt{(\vec{v}_p \cdot \vec{v}_g)^2}$$

$$p^\mu [E_{sqr}(\|\vec{v}_p\| \|\vec{v}_g\|), \quad g_{\alpha\beta}(\|\vec{v}_p\| \|\vec{v}_g\|)] p_\mu [E_{sqr}(\|\vec{v}_p\| \|\vec{v}_g\|), \quad g_{\alpha\beta}(\|\vec{v}_p\| \|\vec{v}_g\|)] = \pm (m)^2 \|\vec{v}_p\| \|\vec{v}_g\| \|\vec{v}_p\| \|\vec{v}_g\|$$

5. (Modified Schwarzschild Radius)

$$t = \frac{t_0}{\sqrt{1 - \frac{2GM}{(\vec{v}_p)^2} r}}$$

$$t = \frac{t_0}{\sqrt{1 - \frac{2GM}{(\mathbf{v}_g)^2 r}}}$$

$$t = \frac{t_0}{\sqrt{1 - \frac{2GM}{\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2 r}}} = \frac{t_0}{\sqrt{1 - \frac{2GM}{\|\mathbf{v}_p\| \|\mathbf{v}_g\| r}}}$$

$$t = \gamma t_0 = \frac{t_0}{\sqrt{1 - \frac{v^2}{(\mathbf{v}_p)^2}}} = \frac{t_0}{\sqrt{1 - \frac{v^2}{(\mathbf{v}_g)^2}}} = \frac{t_0}{\sqrt{1 - \frac{v^2}{\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2}}} = \frac{t_0}{\sqrt{1 - \frac{v^2}{\|\mathbf{v}_p\| \|\mathbf{v}_g\|}}}$$

6. (Modified Time Dilation)

$$z = \frac{1}{\sqrt{1 - \frac{2GM}{(\mathbf{v}_p)^2 r}}} - 1$$

$$z = \frac{1}{\sqrt{1 - \frac{2GM}{(\mathbf{v}_g)^2 r}}} - 1$$

$$z = \frac{1}{\sqrt{1 - \frac{2GM}{\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2 r}}} - 1 = \frac{1}{\sqrt{1 - \frac{2GM}{\|\mathbf{v}_p\| \|\mathbf{v}_g\| r}}} - 1$$

7. (Modified Gravitational Redshift)

$$\theta_E = \sqrt{\frac{4GM}{(\mathbf{v}_p)^2} \frac{D_{LS}}{D_L D_S}}$$

$$\theta_E = \sqrt{\frac{4GM}{(\mathbf{v}_g)^2} \frac{D_{LS}}{D_L D_S}}$$

$$\theta_E = \sqrt{\frac{4GM}{\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} \frac{D_{LS}}{D_L D_S}}} = \sqrt{\frac{4GM}{\|\mathbf{v}_p\| \|\mathbf{v}_g\| \frac{D_{LS}}{D_L D_S}}}$$

8. (Modified Einstein Ring or Gravitational Lensing Angle)

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3} \rho - \frac{k(\mathbf{v}_p)^2}{a^2} + \frac{\Lambda(\mathbf{v}_p)^2}{3}$$

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3} \rho - \frac{k\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2}}{a^2} + \frac{\Lambda\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2}}{3}$$

$$\begin{aligned}
\left(\frac{\dot{a}}{a}\right)^2 &= \frac{8\pi G}{3}\rho - \frac{k\|\vec{v}_p\|\|\vec{v}_g\|}{a^2} + \frac{\Lambda\|\vec{v}_p\|\|\vec{v}_g\|}{3} \\
u^\mu \left[E_{sqr}(v_p), g_{\alpha\beta}(v_p) \right] u_\mu \left[E_{sqr}(v_p), g_{\alpha\beta}(v_p) \right] &= \pm (v_p)^2 \\
p^\mu \left[E_{sqr}(v_p), g_{\alpha\beta}(v_p) \right] p_\mu \left[E_{sqr}(v_p), g_{\alpha\beta}(v_p) \right] & \\
&= \pm (m)^2 \|\vec{v}_p\| \|\vec{v}_g\| (v_p)^2 \\
p^\mu \left[E_{sqr}(\|\vec{v}_p\| \|\vec{v}_g\|), g_{\alpha\beta}(\|\vec{v}_p\| \|\vec{v}_g\|) \right] p_\mu \left[E_{sqr}(\|\vec{v}_p\| \|\vec{v}_g\|), g_{\alpha\beta}(\|\vec{v}_p\| \|\vec{v}_g\|) \right] & \\
&= \pm (m)^2 \|\vec{v}_p\| \|\vec{v}_g\| (\|\vec{v}_p\| \|\vec{v}_g\|) \\
p^\mu \left[E_{sqr} \left[\left(\sqrt{(v_p \cdot v_g)^2} \right) \right], g_{\alpha\beta} \left[\left(\sqrt{(v_p \cdot v_g)^2} \right) \right] \right] p_\mu \left[E_{sqr} \left[\left(\sqrt{(v_p \cdot v_g)^2} \right) \right], g_{\alpha\beta} \left[\left(\sqrt{(v_p \cdot v_g)^2} \right) \right] \right] & \\
&= \pm (m)^2 \|\vec{v}_p\| \|\vec{v}_g\| \left(\sqrt{(v_p \cdot v_g)^2} \right)
\end{aligned}$$

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho - \frac{k(v_g)^2}{a^2} + \frac{\Lambda(v_g)^2}{3}$$

9. (Modified Friedmann Equation)

$$\Delta\omega = \frac{6\pi GM}{(v_p)^2 a(1-e^2)}$$

$$\Delta\omega = \frac{6\pi GM}{(v_g)^2 a(1-e^2)}$$

$$\Delta\omega = \frac{6\pi GM}{\left(\sqrt{(v_p \cdot v_g)^2}\right) a(1-e^2)} = \frac{6\pi GM}{(\|\vec{v}_p\| \|\vec{v}_g\|) a(1-e^2)}$$

$$\Delta\varphi = \frac{6\pi GM}{(v_p)^2 a(1-e^2)}$$

$$\Delta\varphi = \frac{6\pi GM}{(v_g)^2 a(1-e^2)}$$

$$\Delta\varphi = \frac{6\pi GM}{\left(\sqrt{(v_p \cdot v_g)^2}\right) a(1-e^2)} = \frac{6\pi GM}{(\|\vec{v}_p\| \|\vec{v}_g\|) a(1-e^2)}$$

$\Delta\varphi$ is the additional precession per orbit.

10. (Modified Perihelion Precession of Mercury)

$$\begin{aligned}
ds^2 &= -(v_p)^2 dt^2 + a(t)^2 \left[\frac{dr^2}{1-kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right] \\
ds^2 &= -(v_g)^2 dt^2 + a(t)^2 \left[\frac{dr^2}{1-kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right]
\end{aligned}$$

$$ds^2 = - (\| \vec{v}_p \| \| \vec{v}_g \|) dt^2 + a(t)^2 \left[\frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right]$$

$$ds^2 = - \left(\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2} \right) dt^2 + a(t)^2 \left[\frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right]$$

(Modified FLRW Metric)

$$\Lambda = \frac{3H_0^2 \Omega_\Lambda}{(\vec{v}_p)^2}$$

$$\Lambda = \frac{3H_0^2 \Omega_\Lambda}{(\vec{v}_g)^2}$$

$$\Lambda = \frac{3H_0^2 \Omega_\Lambda}{(\| \vec{v}_p \| \| \vec{v}_g \|)} = \frac{3H_0^2 \Omega_\Lambda}{\left(\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2} \right)}$$

(Modified Cosmological Parameters)

$$\Lambda = \frac{8\pi G}{(\vec{v}_p)^4} \rho_{vac}$$

$$\Lambda = \frac{8\pi G}{(\vec{v}_g)^4} \rho_{vac}$$

$$\Lambda = \frac{8\pi G}{(\| \vec{v}_p \| \| \vec{v}_g \|)^2} \rho_{vac} = \frac{8\pi G}{(\vec{v}_p \cdot \vec{v}_g)^2} \rho_{vac}$$

(Modified Vacuum Energy Density)

$$\Lambda = \frac{1}{4} R + \frac{8\pi G}{(\vec{v}_p)^4} T$$

$$\Lambda = \frac{1}{4} R + \frac{8\pi G}{(\vec{v}_g)^4} T$$

$$\Lambda = \frac{1}{4} R + \frac{8\pi G}{(\| \vec{v}_p \| \| \vec{v}_g \|)^2} T = \frac{1}{4} R + \frac{8\pi G}{(\vec{v}_p \cdot \vec{v}_g)^2} T$$

Modified Trace of EFE

$$\dot{\rho} + 3 \frac{\dot{a}}{a} \left(\rho + \frac{p}{(\vec{v}_p)^2} \right) = 0$$

$$\dot{\rho} + 3 \frac{\dot{a}}{a} \left(\rho + \frac{p}{(\vec{v}_g)^2} \right) = 0$$

$$\dot{\rho} + 3\frac{\dot{a}}{a}\left(\rho + \frac{p}{(\|\vec{v}_p\| \|\vec{v}_g\|)}\right) = 0, \quad \dot{\rho} + 3\frac{\dot{a}}{a}\left(\rho + \frac{p}{(\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2})}\right) = 0$$

Modified The Fluid Equation (Cosmic Continuity Equation)

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}\left(\rho + \frac{3p}{(\vec{v}_p)^2}\right) + \frac{\Lambda(\vec{v}_p)^2}{3}$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}\left(\rho + \frac{3p}{(\vec{v}_g)^2}\right) + \frac{\Lambda(\vec{v}_g)^2}{3}$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}\left(\rho + \frac{3p}{(\|\vec{v}_p\| \|\vec{v}_g\|)}\right) + \frac{\Lambda(\|\vec{v}_p\| \|\vec{v}_g\|)}{3}$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}\left(\rho + \frac{3p}{(\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2})}\right) + \frac{\Lambda(\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2})}{3}$$

Modified The Acceleration Equation (Second Friedmann Equation)

$$t = \int \frac{da}{a \sqrt{\frac{8\pi G\rho(a)}{3} - \frac{k(\vec{v}_p)^2}{a^2} + \frac{\Lambda(\vec{v}_p)^2}{3}}}$$

$$t = \int \frac{da}{a \sqrt{\frac{8\pi G\rho(a)}{3} - \frac{k(\vec{v}_g)^2}{a^2} + \frac{\Lambda(\vec{v}_g)^2}{3}}}$$

$$t = \int \frac{da}{a \sqrt{\frac{8\pi G\rho(a)}{3} - \frac{k(\|\vec{v}_p\| \|\vec{v}_g\|)}{a^2} + \frac{\Lambda(\|\vec{v}_p\| \|\vec{v}_g\|)}{3}}}$$

$$t = \int \frac{da}{a \sqrt{\frac{8\pi G\rho(a)}{3} - \frac{k(\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2})}{a^2} + \frac{\Lambda(\sqrt{(\vec{v}_p \cdot \vec{v}_g)^2})}{3}}}$$

$$t_0 = \frac{1}{H_0} \int_0^1 \frac{da}{a \sqrt{\Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_k a^{-2} + \Omega_\Lambda}}$$

$$\Omega_k = \frac{-k(\vec{v}_p)^2}{a_0^2 H_0^2}$$

$$\Omega_{\Lambda} = \frac{\Lambda(\mathbf{v}_p)^2}{3H_0^2}$$

$$t_0 = \frac{1}{H_0} \int_0^1 \frac{da}{a\sqrt{\Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_k a^{-2} + \Omega_{\Lambda}}}$$

$$\Omega_k = \frac{-k(\mathbf{v}_g)^2}{a_0^2 H_0^2}$$

$$\Omega_{\Lambda} = \frac{\Lambda(\mathbf{v}_g)^2}{3H_0^2}$$

$$t_0 = \frac{1}{H_0} \int_0^1 \frac{da}{a\sqrt{\Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_k a^{-2} + \Omega_{\Lambda}}}$$

$$\Omega_k = \frac{-k(\|\vec{\mathbf{v}}_p\| \|\vec{\mathbf{v}}_g\|)}{a_0^2 H_0^2} = \frac{-k\left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2}\right)}{a_0^2 H_0^2}$$

$$\Omega_{\Lambda} = \frac{\Lambda(\|\vec{\mathbf{v}}_p\| \|\vec{\mathbf{v}}_g\|)}{3H_0^2} = \frac{\Lambda\left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2}\right)}{3H_0^2}$$

$$t_0 = \int_0^1 \frac{da}{aH(a)} = \frac{1}{H_0} \int_0^{\infty} \frac{dz}{(1+z)\sqrt{\Omega_m(1+z)^3 + \Omega_{\Lambda}}}$$

$$\Omega_{\Lambda} = \frac{\Lambda(\mathbf{v}_p)^2}{3H_0^2}$$

$$t_0 = \int_0^1 \frac{da}{aH(a)} = \frac{1}{H_0} \int_0^{\infty} \frac{dz}{(1+z)\sqrt{\Omega_m(1+z)^3 + \Omega_{\Lambda}}}$$

$$\Omega_{\Lambda} = \frac{\Lambda(\mathbf{v}_g)^2}{3H_0^2}$$

$$t_0 = \int_0^1 \frac{da}{aH(a)} = \frac{1}{H_0} \int_0^{\infty} \frac{dz}{(1+z)\sqrt{\Omega_m(1+z)^3 + \Omega_{\Lambda}}}$$

$$\Omega_{\Lambda} = \frac{\Lambda(\|\vec{\mathbf{v}}_p\| \|\vec{\mathbf{v}}_g\|)}{3H_0^2} = \frac{\Lambda\left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2}\right)}{3H_0^2}$$

Modified Age of the Universe (Cosmic Time)

$$P = w\rho(\mathbf{v}_p)^2$$

$$P = w\rho(\mathbf{v}_g)^2$$

$$P = w\rho(\|\vec{\mathbf{v}}_p\| \|\vec{\mathbf{v}}_g\|) = w\rho\left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2}\right)$$

Modified Equation of State (w)

$$ds^2 = -\left(1 - \frac{r_s}{r}\right)(\mathbf{v}_p)^2 dt^2 + \left(1 - \frac{r_s}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

$$ds^2 = -\left(1 - \frac{r_s}{r}\right)(\mathbf{v}_g)^2 dt^2 + \left(1 - \frac{r_s}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

$$ds^2 = -\left(1 - \frac{r_s}{r}\right)(\|\vec{\mathbf{v}}_p\| \|\vec{\mathbf{v}}_g\|) dt^2 + \left(1 - \frac{r_s}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

$$ds^2 = -\left(1 - \frac{r_s}{r}\right)\left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2}\right) dt^2 + \left(1 - \frac{r_s}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

Modified (The Schwarzschild Metric)

$$\frac{d^2 x^i}{d\tau^2} + \Gamma_{00}^i \left(\mathbf{v}_p \frac{dt}{d\tau}\right)^2 + 2\Gamma_{0j}^i \left(\mathbf{v}_p \frac{dt}{d\tau}\right) \frac{dx^j}{d\tau} + \Gamma_{jk}^i \frac{dx^j}{d\tau} \frac{dx^k}{d\tau} = 0$$

$$\frac{d^2 x^i}{d\tau^2} + \Gamma_{00}^i \left(\mathbf{v}_g \frac{dt}{d\tau}\right)^2 + 2\Gamma_{0j}^i \left(\mathbf{v}_g \frac{dt}{d\tau}\right) \frac{dx^j}{d\tau} + \Gamma_{jk}^i \frac{dx^j}{d\tau} \frac{dx^k}{d\tau} = 0$$

$$\frac{d^2 x^i}{d\tau^2} + \Gamma_{00}^i \left(\|\vec{\mathbf{v}}_p\| \|\vec{\mathbf{v}}_g\| \frac{dt}{d\tau}\right)^2 + 2\Gamma_{0j}^i \left(\|\vec{\mathbf{v}}_p\| \|\vec{\mathbf{v}}_g\| \frac{dt}{d\tau}\right) \frac{dx^j}{d\tau} + \Gamma_{jk}^i \frac{dx^j}{d\tau} \frac{dx^k}{d\tau} = 0$$

$$\frac{d^2 x^i}{d\tau^2} + \Gamma_{00}^i \left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} \frac{dt}{d\tau}\right)^2 + 2\Gamma_{0j}^i \left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} \frac{dt}{d\tau}\right) \frac{dx^j}{d\tau} + \Gamma_{jk}^i \frac{dx^j}{d\tau} \frac{dx^k}{d\tau} = 0$$

Modified (The Geodesic Equation)

$$\alpha = \frac{4GM}{r(\mathbf{v}_p)^2}$$

$$\alpha = \frac{4GM}{r(\mathbf{v}_g)^2}$$

$$\alpha = \frac{4GM}{r(\|\vec{\mathbf{v}}_p\| \|\vec{\mathbf{v}}_g\|)} = \frac{4GM}{r(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2})}$$

Modified Gravitational Lensing

$$\bar{h}_{\mu\nu} = -\frac{16\pi G}{(\mathbf{v}_p)^4} T_{\mu\nu}$$

$$\bar{h}_{\mu\nu} = -\frac{16\pi G}{(\mathbf{v}_g)^4} T_{\mu\nu}$$

$$\bar{h}_{\mu\nu} = -\frac{16\pi G}{(\|\vec{v}_p\| \|\vec{v}_g\|)^2} T_{\mu\nu} = -\frac{16\pi G}{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} T_{\mu\nu}$$

Modified Linearized Gravity & Gravitational Waves

$$\frac{d\theta}{d\tau} = -\frac{1}{3}\theta^2 - \sigma_{\mu\nu}\sigma^{\mu\nu} + \omega_{\mu\nu}\omega^{\mu\nu} - R_{\mu\nu}u^\mu u^\nu$$

$$\frac{d\theta}{d\tau} = -\frac{1}{3}\theta^2 - \sigma^2 + \omega^2 - \frac{8\pi G}{(\mathbf{v}_p)^2} \left(\rho + \frac{3P}{(\mathbf{v}_p)^2} \right)$$

$$\frac{d\theta}{d\tau} = -\frac{1}{3}\theta^2 - \sigma^2 + \omega^2 - \frac{8\pi G}{(\mathbf{v}_g)^2} \left(\rho + \frac{3P}{(\mathbf{v}_g)^2} \right)$$

$$\frac{d\theta}{d\tau} = -\frac{1}{3}\theta^2 - \sigma^2 + \omega^2 - \frac{8\pi G}{(\|\vec{v}_p\| \|\vec{v}_g\|)} \left(\rho + \frac{3P}{(\|\vec{v}_p\| \|\vec{v}_g\|)} \right)$$

$$\frac{d\theta}{d\tau} = -\frac{1}{3}\theta^2 - \sigma^2 + \omega^2 - \frac{8\pi G}{(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2})} \left(\rho + \frac{3P}{(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2})} \right)$$

Modified The Raychaudhuri Equation

$$Z_{grav} \approx \frac{GM}{r(\mathbf{v}_p)^2}$$

$$Z_{grav} \approx \frac{GM}{r(\mathbf{v}_g)^2}$$

$$Z_{grav} \approx \frac{GM}{r(\|\vec{v}_p\| \|\vec{v}_g\|)} \approx \frac{GM}{r(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2})}$$

$$1 + Z_{grav} = \left(1 - \frac{2GM}{r(\mathbf{v}_p)^2} \right)^{-1/2}$$

$$1 + Z_{grav} = \left(1 - \frac{2GM}{r(\mathbf{v}_g)^2} \right)^{-1/2}$$

$$1 + Z_{grav} = \left(1 - \frac{2GM}{r(\|\vec{v}_p\| \|\vec{v}_g\|)} \right)^{-1/2} = \left(1 - \frac{2GM}{r(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2})} \right)^{-1/2}$$

Modified Gravitational Redshift

$$\dot{\rho} + 3H \left(\rho + \frac{P}{(\mathbf{v}_p)^2} \right) = 0$$

$$\dot{\rho} + 3H \left(\rho + \frac{P}{(\mathbf{v}_g)^2} \right) = 0$$

$$\dot{\rho} + 3H \left(\rho + \frac{P}{(\|\vec{v}_p\| \|\vec{v}_g\|)} \right) = 0, \dot{\rho} + 3H \left(\rho + \frac{P}{\left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} \right)} \right) = 0$$

Modified (Conservation Law & Continuity)

$$d_{horizon}(t) = \mathbf{v}_p \int_0^t \frac{dt'}{a(t')}$$

$$d_{horizon}(t) = \mathbf{v}_g \int_0^t \frac{dt'}{a(t')}$$

$$d_{horizon}(t) = \sqrt{(\|\vec{v}_p\| \|\vec{v}_g\|)} \int_0^t \frac{dt'}{a(t')} = \sqrt{\left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} \right)} \int_0^t \frac{dt'}{a(t')}$$

Modified (The Particle Horizon)

$$\frac{dP}{dr} = -\frac{G}{r^2} \left(\rho + \frac{P}{(\mathbf{v}_p)^2} \right) \left(M + \frac{4\pi r^3 P}{(\mathbf{v}_p)^2} \right) \left(1 - \frac{2GM}{r(\mathbf{v}_p)^2} \right)^{-1}$$

$$\frac{dP}{dr} = -\frac{G}{r^2} \left(\rho + \frac{P}{(\mathbf{v}_g)^2} \right) \left(M + \frac{4\pi r^3 P}{(\mathbf{v}_g)^2} \right) \left(1 - \frac{2GM}{r(\mathbf{v}_g)^2} \right)^{-1}$$

$$\frac{dP}{dr} = -\frac{G}{r^2} \left(\rho + \frac{P}{(\|\vec{v}_p\| \|\vec{v}_g\|)} \right) \left(M + \frac{4\pi r^3 P}{(\|\vec{v}_p\| \|\vec{v}_g\|)} \right) \left(1 - \frac{2GM}{r(\|\vec{v}_p\| \|\vec{v}_g\|)} \right)^{-1}$$

$$\frac{dP}{dr} = -\frac{G}{r^2} \left(\rho + \frac{P}{\left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} \right)} \right) \left(M + \frac{4\pi r^3 P}{\left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} \right)} \right) \left(1 - \frac{2GM}{r \left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} \right)} \right)^{-1}$$

Modified (TOV Equation)

$$\Delta t \approx \frac{4GM}{(\mathbf{v}_p)^3} \ln \left(\frac{4r_1 r_2}{R^2} \right)$$

$$\Delta t \approx \frac{4GM}{(\mathbf{v}_g)^3} \ln \left(\frac{4r_1 r_2}{R^2} \right)$$

$$\Delta t \approx \frac{4GM}{(\|\vec{v}_p\| \|\vec{v}_g\|)^{\frac{3}{2}}} \ln \left(\frac{4r_1 r_2}{R^2} \right) \approx \frac{4GM}{\left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} \right)^{\frac{3}{2}}} \ln \left(\frac{4r_1 r_2}{R^2} \right)$$

Modified (Shapiro Time Delay)

$$P = \frac{G}{5(\mathbf{v}_p)^5} \langle \ddot{l}_{ij} \ddot{l}_{ij} \rangle$$

$$P = \frac{G}{5(\mathbf{v}_g)^5} \langle \ddot{l}_{ij} \ddot{l}_{ij} \rangle$$

$$P = \frac{G}{5(\|\vec{v}_p\| \|\vec{v}_g\|)^{\frac{5}{2}}} \langle \ddot{l}_{ij} \ddot{l}_{ij} \rangle = \frac{G}{5 \left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} \right)^{\frac{5}{2}}} \langle \ddot{l}_{ij} \ddot{l}_{ij} \rangle$$

Modified (Gravitational Wave Luminosity)

$$\Omega = \frac{2GJ}{(\mathbf{v}_p)^2 r^3}$$

$$\Omega = \frac{2GJ}{(\mathbf{v}_g)^2 r^3}$$

$$\Omega = \frac{2GJ}{(\|\vec{v}_p\| \|\vec{v}_g\|) r^3} = \frac{2GJ}{\left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2}\right) r^3}$$

Modified (Lense – Thirring / Frame Dragging)

$$\Phi_{PPN} \sim \frac{GM}{r(\mathbf{v}_p)^2}$$

$$\Phi_{PPN} \sim \frac{GM}{r(\mathbf{v}_g)^2}$$

$$\Phi_{PPN} \sim \frac{GM}{r(\|\vec{v}_p\| \|\vec{v}_g\|)} \sim \frac{GM}{r\left(\sqrt{(\mathbf{v}_p \cdot \mathbf{v}_g)^2}\right)}$$

Modified (Parametrized Post – Newtonian – PPN)

$$[K_{ij} - h_{ij}K] = -\frac{8\pi G}{(\mathbf{v}_p)^4} S_{ij}$$

$$[K_{ij} - h_{ij}K] = -\frac{8\pi G}{(\mathbf{v}_g)^4} S_{ij}$$

$$[K_{ij} - h_{ij}K] = -\frac{8\pi G}{(\|\vec{v}_p\| \|\vec{v}_g\|)^2} S_{ij} = -\frac{8\pi G}{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} S_{ij}$$

Modified (Israel Junction Conditions)

$$\kappa = \frac{8\pi G}{(\mathbf{v}_p)^4} = \frac{8\pi G}{(\mathbf{v}_g)^4} = \frac{8\pi G}{(\|\vec{v}_p\| \|\vec{v}_g\|)^2} = \frac{8\pi G}{(\mathbf{v}_p \cdot \mathbf{v}_g)^2}$$

$$G_{\mu\nu}[g_{\alpha\beta}(\mathbf{v}_p)] + \Lambda g_{\mu\nu}(\mathbf{v}_p)$$

$$= \frac{8\pi}{1} \frac{G}{(\mathbf{v}_p)^4} T_{\mu\nu} [E_{sqr}(\mathbf{v}_p), g_{\alpha\beta}(\mathbf{v}_p)]$$

$$G_{\mu\nu}[g_{\alpha\beta}(\mathbf{v}_g)] + \Lambda g_{\mu\nu}(\mathbf{v}_g) = \frac{8\pi}{1} \frac{G}{(\mathbf{v}_g)^4} T_{\mu\nu} [E_{sqr}(\mathbf{v}_g), g_{\alpha\beta}(\mathbf{v}_g)]$$

$$G_{\mu\nu}[g_{\alpha\beta}(\|\vec{v}_p\| \|\vec{v}_g\|)] + \Lambda g_{\mu\nu}(\|\vec{v}_p\| \|\vec{v}_g\|) = \frac{8\pi}{1} \frac{G}{(\|\vec{v}_p\| \|\vec{v}_g\|)^2} T_{\mu\nu} [E_{sqr}(\|\vec{v}_p\| \|\vec{v}_g\|), g_{\alpha\beta}(\|\vec{v}_p\| \|\vec{v}_g\|)]$$

$$G_{\mu\nu}[g_{\alpha\beta}(\mathbf{v}_p \cdot \mathbf{v}_g)] + \Lambda g_{\mu\nu}(\mathbf{v}_p \cdot \mathbf{v}_g) = \frac{8\pi}{1} \frac{G}{(\mathbf{v}_p \cdot \mathbf{v}_g)^2} T_{\mu\nu} [E_{sqr}(\mathbf{v}_p \cdot \mathbf{v}_g), g_{\alpha\beta}(\mathbf{v}_p \cdot \mathbf{v}_g)]$$

- General quantitative relativity for David's

$$m = \frac{m_0}{\sqrt{1 - \frac{2GM}{(\mathbf{v}_p)^2} r}}$$

$$\begin{aligned}
 m &= \frac{m_0}{\sqrt{1 - \frac{2GM}{(v_g)^2} r}} \\
 m &= \frac{m_0}{\sqrt{1 - \frac{2GM}{\|\vec{v}_p\| \|\vec{v}_g\| r}}} = \frac{m_0}{\sqrt{1 - \frac{2GM}{\sqrt{(v_p \cdot v_g)^2} r}}} \\
 m = \gamma m_0 &= \frac{m_0}{\sqrt{1 - \frac{v^2}{(v_p)^2}}} = \frac{m_0}{\sqrt{1 - \frac{v^2}{(v_g)^2}}} = \frac{m_0}{\sqrt{1 - \frac{v^2}{\|\vec{v}_p\| \|\vec{v}_g\|}}} \\
 &= \frac{m_0}{\sqrt{1 - \frac{v^2}{\sqrt{(v_p \cdot v_g)^2}}}}
 \end{aligned}$$

6. (Modified Relativistic Mass)

$$\begin{aligned}
 m_{eff} &= \frac{m_0}{\sqrt{1 - \frac{2GM}{(v_p)^2} r}} \\
 m_{eff} &= \frac{m_0}{\sqrt{1 - \frac{2GM}{(v_g)^2} r}} \\
 m_{eff} &= \frac{m_0}{\sqrt{1 - \frac{2GM}{\|\vec{v}_p\| \|\vec{v}_g\| r}}} = \frac{m_0}{\sqrt{1 - \frac{2GM}{\sqrt{(v_p \cdot v_g)^2} r}}} \\
 m_{eff} = \gamma m_0 &= \frac{m_0}{\sqrt{1 - \frac{v^2}{(v_p)^2}}} = \frac{m_0}{\sqrt{1 - \frac{v^2}{(v_g)^2}}} = \frac{m_0}{\sqrt{1 - \frac{v^2}{\|\vec{v}_p\| \|\vec{v}_g\|}}} = \frac{m_0}{\sqrt{1 - \frac{v^2}{\sqrt{(v_p \cdot v_g)^2}}}}
 \end{aligned}$$

(Relativistic mass of effective)

$$C = \frac{1}{\sqrt{\mu_0 \times \epsilon_0}}$$

$$v_p = \frac{1}{\sqrt{\mu_0 \times \epsilon_0}}$$

$$v_g = \frac{1}{\sqrt{\mu_0 \times \epsilon_0}}$$

ϵ_0 Vacuum permittivity, μ_0 Vacuum permeability

- The electron generates a constant field while rotating around the nucleus, but when it gains energy, it generates a changing field. This explains why it has a torque resulting from the energy during the experiment. Therefore, if the electron is observed in its normal state without being excited, the electron will behave as a particle, and if it is excited, it will behave as a wave.

- The Mössbauer effect proved that general relativity is true. Relativity explains that the fastest speed is the speed of light. However, if the Mössbauer effect differs depending on the medium it is in, due to the refractive index, then relativity will differ.
- Metal-organic frameworks (MOFs) can trap molecules like water inside them. What if they were modified to trap electrons to provide electricity, neutrinos, nuclei, neutrons, or antimatter? This would ensure they exist to work on.

$$G_{\mu\nu}[g_{\alpha\beta}(v_p)] + \Lambda g_{\mu\nu}(v_p) = \frac{8\pi}{1} \frac{G}{(v_p)^4} T_{\mu\nu}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)]$$

$$R_{\mu\nu}[g_{\alpha\beta}(v_p)] - \frac{1}{2} R[g_{\alpha\beta}(v_p)] g_{\mu\nu}(v_p) + \Lambda g_{\mu\nu}(v_p) = \frac{8\pi}{1} \frac{G}{(v_p)^4} T_{\mu\nu}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)]$$

$$T_{\mu\nu}[E_{sqr}(v_p), g_{\alpha\beta}(v_p)] = \left(\rho[E_{sqr}(v_p)] + \frac{p[E_{sqr}(v_p)]}{(v_p)^2} \right) u_\mu[E_{sqr}(v_p), g_{\alpha\beta}(v_p)] u_\nu[E_{sqr}(v_p), g_{\alpha\beta}(v_p)] + p g_{\mu\nu}(v_p)$$

- *General quantitative relativity for phase velocity*

$$G_{\mu\nu}[g_{\alpha\beta}(v_g)] + \Lambda(v_g) g_{\mu\nu}(v_g) = \frac{8\pi}{1} \frac{G}{(v_g)^4} T_{\mu\nu}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)]$$

$$R_{\mu\nu}[g_{\alpha\beta}(v_g)] - \frac{1}{2} R[g_{\alpha\beta}(v_g)] g_{\mu\nu}(v_g) + \Lambda g_{\mu\nu}(v_g) = \frac{8\pi}{1} \frac{G}{(v_g)^4} T_{\mu\nu}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)]$$

$$T_{\mu\nu}[E_{sqr}(v_g), g_{\alpha\beta}(v_g)] = \left(\rho[E_{sqr}(v_g)] + \frac{p[E_{sqr}(v_g)]}{(v_g)^2} \right) u_\mu[E_{sqr}(v_g), g_{\alpha\beta}(v_g)] u_\nu[E_{sqr}(v_g), g_{\alpha\beta}(v_g)] + p g_{\mu\nu}(v_g)$$

- *General quantitative relativity for group velocity*

$$G_{\mu\nu}[g_{\alpha\beta}(v_p \cdot v_g)] + \Lambda(v_p \cdot v_g) g_{\mu\nu}(v_p \cdot v_g) = \frac{8\pi}{1} \frac{G}{(v_p \cdot v_g)^2} T_{\mu\nu}[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)]$$

$$R_{\mu\nu}[g_{\alpha\beta}(v_p \cdot v_g)] - \frac{1}{2} R[g_{\alpha\beta}(v_p \cdot v_g)] g_{\mu\nu}(v_p \cdot v_g) + \Lambda g_{\mu\nu}(v_p \cdot v_g) = \frac{8\pi}{1} \frac{G}{(v_p \cdot v_g)^4} T_{\mu\nu}[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)]$$

$$T_{\mu\nu}[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)] = \left(\rho[E_{sqr}(v_p \cdot v_g)] + \frac{p[E_{sqr}(v_p \cdot v_g)]}{v_p \cdot v_g} \right) u_\mu[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)] u_\nu[E_{sqr}(v_p \cdot v_g), g_{\alpha\beta}(v_p \cdot v_g)] + p g_{\mu\nu}(v_p \cdot v_g)$$

- *General quantitative relativity for David's velocity of the stationary phase*
- This equation represents a new perspective in terms of adopting the refractive index $n = 1$, which represents the maximum value for numerical values, namely Planck values, thus preventing access to infinite values.
- $g_{\mu\nu}(v_p), g_{\mu\nu}(v_g)$ depends on the phase speed and the group speed

5. Method

My name is Ahmed. I have made a theoretical derivation of the equation of general relativity as explained in this research for the purpose of obtaining an equation that can be applied within the quantum world so that it describes the movement of the electron during the quantum jump in the Bohr model. After that, the researcher Samira reviewed the research and verified it, and then she worked on applying this theory to the movement of the electron during the occurrence of the quantum leap, using previous research and matching it with the results of this equation to determine its validity.

- This part of the research will explain the spectrum of the hydrogen atom in a new way, as the results presented in these tables from previous research match the results extracted from the equation, and this is consistent with the validity of this equation. Because the new equation is consistent with the photon energy equation. We will discuss that part of the research in the results and discussion.
- **Table 5** shows the measurement results tested.[3] (Nanni, 2015)

- **Table 5** it represents the theoretical and experimental value of the hydrogen atom. Using the photon energy law mentioned above, this table.

$$\Delta E = E_m - E_n = h \frac{c}{\lambda} \rightarrow \frac{1}{\lambda} = \frac{4}{B} \left(\frac{1}{n^2} - \frac{1}{m^2} \right)$$

The way toward the quantum mechanics was definitely opened! The calculated wavelength values vs the experimental ones are listed in table

Spectral Line	Experimental Value	Theoretical Value
	(nm)	(nm)
$\lambda(n'=2, n=1)$	121.5	122.0
$\lambda(n'=3, n=1)$	102.5	103.0
$\lambda(n'=4, n=1)$	97.2	97.3
$\lambda(n'=2, n=3)$	656.1	656.3
$\lambda(n'=2, n=4)$	486.0	486.1
$\lambda(n'=3, n=4)$	1874.6	1875.0

My scientific research explains how the universe initially expanded so quickly that the change in phase velocity from the speed of light led to this expansion in spacetime. Since I put the phase velocity in place of the speed of light in general relativity because of the derivative I did, and this equation will be known as general quantum relativity, then this means that the speed of light was moving differently, and this will lead to spacetime being affected by different media, as my equations show, so the universe was initially expanding, and then inflation occurred as a result of the phase velocity differing from the speed of light, which led to the expansion of spacetime faster than light, and this led to homogeneity in the cosmic background.




Figure 1. Bohr hydrogen atomic model incorporating de Broglie’s .[4] (Jordan, 2024).

This drawing, taken from previous research, shows how the quantum leap occurs through interference, as my equation showed. When interference occurs between the orbit occupied by the electron and the energy level higher than the electron’s orbit, it occurs in the form of wave interference of this type as a result of a contraction in the fabric of space-time. The black circle represents the orbit occupied by the electron, while the red color represents how interference occurs from the orbit higher to the orbit occupied by the electron in the form of wave interference. In other words, the upper level works to contract, forming a wave equal to the same wave as the level occupied by the electron through the de Broglie equation. $n \times \lambda = 2\pi \times r$

- If we make the electron quantum entangled in particle accelerators, then if we make one of these electrons be in a short line and the other be in a long line, when one approaches the speed of light, the other must exceed the speed of light. In other words, the two entangled bodies are in

two dimensions, that is, different dimensions, and this happens as a result, a distortion of space-time, which makes during the measurement that the speed is breached, but in reality it does not exceed the speed. This is the same idea as the distortion of the orbits that I explained. Because it is assumed that the electron does not move from its position, however, a distortion occurs in the orbit with the highest energy, and it forms a wave similar to the orbit occupied by the electron, according to De Broglie’s laws. This occurs through the distortion of space-time as a result of the increase in energy.

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{(2\pi)^2 \cdot 4}{\lambda \, nm \cdot e} \frac{(l_{(p)})^2}{\Delta E_n} T_{\mu\nu}$$

Because the equation connects more than one equation into a single equation. As

$$\begin{aligned} G_{\mu\nu} + \Lambda g_{\mu\nu} &= \frac{8\pi \cdot G}{C^4} T_{\mu\nu} \\ \Delta E_n &= \frac{h_p \cdot C}{\lambda \, nm \cdot e} \\ n \cdot \lambda &= 2\pi \cdot r_n \end{aligned}$$




Figure 2. The observed emission line spectrum of atomic hydrogen in chapter 2 atoms.[5] (Manini, 2020).

The 4 lowest-energy series of spectral lines of atomic hydrogen				
name	lower <i>n</i>	lowest energy [eV]	max energy [eV]	spectral region
Lyman series	1	10.2	13.6	UV
Balmer series	2	1.89	3.40	Visible-UV
Paschen series	3	0.66	1.51	IR
Brackett series	4	0.31	0.85	IR

- **Table (6)** shows the measurement results of one of the previous researches related to the spectrum of the hydrogen atom in chapter 2 atoms.[5] (Manini, 2020)
- *This shape is a result of the fact that the electron, after a quantum leap occurred as a result of an interference between the orbital that it occupies and the energy level above it, was in an unstable state. Therefore, when the highest level of energy returns to its position, it releases energy in the form of spectral lines. These lines are determined according to the amount of energy, as shown in the picture.*

6. Results obtained

This scientific research aims to prove a theory by comparing the practical results of this theory with the original results and making the comparison in a table. We will discuss that here .

My theory is based on introducing the curvature of spacetime into the equation, but quantum mechanics shows that it is not affected by gravity. How to interact with the curvature of spacetime has not yet been proven. As a result, my equations show a way to conduct an experiment that enables direct interaction with the curvature of spacetime. Therefore, this experiment practically proves that quantum mechanics made a mistake in its concept when it showed gravity does not interact with it. How to conduct an experimental experiment to prove the validity of my equations

Steps to conduct the experiment

1) The place where the experiment will take place must be chosen, and it must be at a high altitude, such as Mount Everest because the higher the altitude, the less gravity.

2) The experiment is about creating a quantum leap for the electron so that we can know the emission lines that represent the fingerprint of the element and compare them at different heights. Let us take the example of the hydrogen atom. After knowing the choice of the element, the device that will measure the spectral lines of the element must be taken to Mount Everest, where the experiment will be conducted.

3) We will excite the element keeping all elements constant as energy and the comparison will be between wavelength and curvature of spacetime. The first measurement is at the bottom of the mountain, that is, before climbing the mountain first. Then we measure in the middle of the mountain, then we test at the top of the mountain and compare the atomic spectra. If my theoretical results are correct, there will be skewing of the spectral lines at different heights due to distortion of the fabric of space-time.

4) If we measure atomic spectra, we also measure the Zeeman, Stark, and magneto-stark effects separately.

- The reason they were not previously able to measure the curvature of space-time is because my equations show that the effect of energy and wavelength when measured as two variables will cancel each other out, so space-time will not be affected.

- Gravitational Effect on Atomic Energy Levels

Objective: Measure the effect of gravity on atomic energy levels

Equipment:

- A gas sample (e.g., hydrogen or cesium) in a vacuum chamber.
- A laser to excite electrons at specific energy levels.
- A high-precision spectrometer.
- A variable gravitational field (e.g., using aircraft simulating microgravity).

Procedure:

- .1 Measure the atomic spectrum in a normal gravitational environment.
- .2 Measure the spectrum in a reduced-gravity environment (e.g., during parabolic flights).
- .3 Compare the energy levels and emission lines.

Expected Outcome:

- If the spectrum shifts at different gravitational strengths, it indicates that gravity affects atomic energy levels
- My equations clearly show that if proven in practical experiments, it indicates that the gravitational constant G is not a cosmic constant in quantum mechanics, but is affected by the wavelength and the energy difference, that is, it is variable. In other words, gravity is not an absolute quantity, but rather the quantum state is influenced by me. For this reason, quantum mechanics is not related to general relativity.
- My equations explain the effect (magnetic attraction) and Bayfield-Brown effect My equations confirm the effect of electromagnetism on gravity.

- Well, with these experiments, the Pound-Rebecca experiments, also known as gravitational redshift, will prove what the equation tells you.

$$G_{\mu\nu} + \Lambda g_{\mu\nu} - T_{\mu\nu} = \frac{8\pi \times G}{C^4} = 2.0766474428 \times 10^{-43} \quad (16)$$

$$E_n = E_2 - E_1$$

$$\Delta E_n = \frac{-13.605693099 \text{ eV}}{1} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{(2\pi)^6 \times (\hbar)^4 \times 4}{(\lambda \text{ nm} \times e)^5 \times (\mu_0 \times \varepsilon_0)^2 (\Delta E_n)^5} T_{\mu\nu}$$

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{(2\pi)^6 \times (1.0545718 \times 10^{-34})^4 \times 4}{(\lambda \text{ nm} \times e)^5 \times (4\pi \times 10^{-7} \times 8.854187813 \times 10^{-12})^2} \frac{(1.616255011 \times 10^{-35})^2}{(\Delta E_n)^5} T_{\mu\nu}$$

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{6.4231253544 \times 10^{-167}}{(\lambda \text{ nm} \times e)^5} \frac{1}{(\Delta E_n)^5} T_{\mu\nu}$$

$$(\Delta E_n)^5 = \frac{6.4231253544 \times 10^{-167}}{(\lambda \text{ nm} \times e)^5} \frac{1}{G_{\mu\nu} + \Lambda g_{\mu\nu} - T_{\mu\nu}}$$

$$(\Delta E_n)^5 = \frac{6.4231253544 \times 10^{-167}}{(\lambda \text{ nm} \times e)^5} \frac{1}{2.0766474428 \times 10^{-43}}$$

$$(\lambda \text{ nm})^5 = \frac{3.0930261448 \times 10^{-124}}{(\Delta E_n \times e)^5}$$

$$(\lambda \text{ nm})^5 = \frac{3.0930261448 \times 10^{-124}}{\left(\frac{-13.605693099 \text{ eV}}{1} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right) \times e \right)^5}$$

$$\lambda \text{ nm} = \sqrt[5]{\frac{3.0930261448 \times 10^{-124}}{\left(\frac{-13.605693099 \text{ eV}}{1} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right) \times 1.60217662 \times 10^{-19} \right)^5}}$$

$$\lambda = 656.11227252 \text{ nm}$$

- This example of a hydrogen atom in the Balmer series.

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{4 \times (\hbar)^8 \times (2\pi)^{10}}{(\mu_0 \times \varepsilon_0)^4 \times (\lambda \text{ nm})^9 \times (e)^9 (\Delta E_n)^9} T_{\mu\nu}$$

$$\Delta E_n = \frac{h_p \times C}{\lambda} = \frac{1239.8419637 \text{ eVnm}}{\lambda}$$

Photon energy equation.

$$(\Delta E_n)^9 = \frac{4 \times (\hbar)^8 \times (2\pi)^{10}}{(\mu_0 \times \varepsilon_0)^4 \times (\lambda \text{ nm})^9 \times (e)^9} \frac{(l_{(p)})^2}{G_{\mu\nu} + \Lambda g_{\mu\nu}} T_{\mu\nu}$$

Example of a hydrogen atom.

$$\Delta E_n = \sqrt[9]{\frac{4 \times (\hbar)^8 \times (2\pi)^{10}}{(\mu_0 \times \varepsilon_0)^4 \times (\lambda \text{ nm})^9 \times (e)^9} \frac{(l_{(p)})^2}{G_{\mu\nu} + \Lambda g_{\mu\nu} - T_{\mu\nu}}}$$

- Example of a hydrogen atom in the Balmer series.

We remove the energy level (n)

$$\Delta E_n = \sqrt[9]{\frac{4.8160445107 \times 10^{-223}}{(\lambda \text{ nm})^9 \times (e)^9}}$$

$$\Delta E_n = \sqrt[9]{\frac{4.8160445107 \times 10^{-223}}{(656.11227252)^9 \times (1.60217662 \times 10^{-19})^9}}$$

$$\Delta E_n = 1.8896795971$$

The unit of measurement for photon energy is electron volt (eV), the wavelength is (nm)

$$\lambda \text{ nm} = \frac{1}{R_\infty \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)}$$

$$R_\infty = 1.0973731731 \times 10^7 \text{ m}^{-1}$$

$$\Delta E_n = \frac{h_p \times C}{\lambda}$$

$$\Delta E_n = -\frac{h_p \times C \times \alpha}{2e \times \lambda} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

$$\lambda = \frac{2\pi \times r_n}{n}$$

$$\Delta E_n = -\frac{h_p \times C \times \alpha}{2e \times \frac{2\pi \times r_n}{n}} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

$$\lambda = \frac{2\pi \times r_n}{n} = \frac{2\pi \times 5.29177202590 \times 10^{-11} \times (n)^2}{n}$$

$$\lambda = \frac{2\pi \times r_n}{n} = 2\pi \times 5.29177202590 \times 10^{-11} \times n$$

$$\Delta E_n = -\frac{6.62607004 \times 10^{-34} \times 299792458 \times 7.297352563 \times 10^{-3}}{2 \times 1.60217662 \times 10^{-19} \times 2\pi \times 5.29177202590 \times 10^{-11}} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

$$\Delta E_n = \frac{-13.605693099 \text{ eV}}{1} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

$$\Delta E_n = -\frac{h_p \times C \times \alpha}{2e \times \lambda} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

$$\Delta E_n = -\frac{6.62607004 \times 10^{-34} \times 299792458 \times 7.297352563 \times 10^{-3}}{2 \times 1.60217662 \times 10^{-19} \times 3.324918425 \times 10^{-10}} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

$$\Delta E_n = \frac{-13.605693099 \text{ eV}}{1} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

$$\Delta E_n = -\frac{h_p \times v \times \alpha}{2e} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

$$\Delta E_n = -\frac{6.62607004 \times 10^{-34} \times 9.016535737 \times 10^{17} \times 7.297352563 \times 10^{-3}}{2 \times 1.60217662 \times 10^{-19}} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

$$\Delta E_n = \frac{-13.605693099 \text{ eV}}{1} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

The unit of measurement for ΔE_n photon energy is electron volt (eV)

$$E_n = \frac{h_p \times C}{\lambda}$$

$$\lambda = \frac{2\pi \times r_n}{n}$$

$$E_n = \frac{n \times h_p \times C}{2\pi \times r_n} = \frac{n \times 6.62607004 \times 10^{-34} \times 299792458}{2\pi \times 5.29177202590 \times 10^{-11} \times (n)^2}$$

$$E_n = \frac{5.9744197314 \times 10^{-16} \text{ J}}{n}$$

$$\Delta E_n = -\frac{h_p \times v \times \alpha}{2e} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

$$E_n = \frac{h_p \times C}{\lambda}$$

$$\begin{aligned}
\Delta h_p \times v &= -\frac{h_p \times v \times \alpha}{2e} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right) \\
\Delta v &= -\frac{v \times \alpha}{2} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right) \\
v &= \frac{v \times (n)^2}{2\pi \times r_n \times \alpha \times Z} \\
\Delta v &= -\frac{\frac{v \times (n)^2}{2\pi \times r_n \times \alpha} \times \alpha}{2} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right) \\
\Delta v &= -\frac{v \times (n)^2}{4\pi \times r_n} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right) \\
\Delta v &= -\frac{2187691.261}{4\pi \times 5.29177202590 \times 10^{-11}} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right) \\
\Delta v &= \frac{-3.289842008 \times 10^{15} \text{ eV}}{1} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right) \\
G_{\mu\nu} + \Lambda g_{\mu\nu} &= \frac{8\pi \times (p)^4 \times Z^4}{(n)^4} \frac{(\alpha)^4 \times G}{\left(h_{(p)} \times \frac{\Delta v \times 2 \text{ eV}}{\alpha} \times \frac{1}{\left(\frac{(Z)^2}{(n_1)^2} - \frac{(Z)^2}{(n_2)^2} \right)} \right)^4} T_{\mu\nu} \\
G_{\mu\nu} + \Lambda g_{\mu\nu} &= \frac{8\pi \times (p)^4 \times Z^4}{(n)^4} \frac{(\alpha)^8 \times G}{\left(h_{(p)} \times \Delta v \times 2 \times e \right)^4} \times \left(\frac{(Z)^2}{(n_1)^2} - \frac{(Z)^2}{(n_2)^2} \right) T_{\mu\nu} \\
G_{\mu\nu} + \Lambda g_{\mu\nu} &= \frac{8\pi \times (p)^4 \times Z^4}{(n)^4} \frac{(\alpha)^8 \times G}{(\Delta E_n \times 2 \times e)^4} \times \left(\frac{(Z)^2}{(n_1)^2} - \frac{(Z)^2}{(n_2)^2} \right) T_{\mu\nu} \\
G_{\mu\nu} + \Lambda g_{\mu\nu} &= \frac{8\pi \times (p)^4 \times Z^4}{(n)^4} \frac{(\alpha)^8 \times G}{\left(\frac{-13.605693099 \text{ eV}}{1} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right) \times 2 \times e \right)^4} \\
&\quad \times \left(\frac{(Z)^2}{(n_1)^2} - \frac{(Z)^2}{(n_2)^2} \right) T_{\mu\nu} \\
G_{\mu\nu} + \Lambda g_{\mu\nu} &= \frac{8\pi \times (p)^4 \times Z^4}{(n)^4} \frac{(\alpha)^8 \times G}{\left(\frac{-13.605693099 \text{ eV}}{1} \times 2 \times e \right)^4} T_{\mu\nu} \\
h_p &= m_e \times C \times \alpha \times 2\pi \times r_n \\
C &= \lambda \times v \\
h_p &= m_e \times \lambda \times v \times \alpha \times 2\pi \times r_n
\end{aligned}$$

Space-time represents in the equation the force of attraction of the nucleus for the electron. Where we take the hydrogen atom compared to the sodium atom. We find after comparison that the undulations that occur in the sodium atom are higher than those that occur in the hydrogen atom. That is, during the occurrence of the quantum jump of the electron, the higher energy level than the level occupied by the electron undulates. So the number of ripples (ripple amplitude) is higher than that of the hydrogen atom during the occurrence of the quantum jump, and this is consistent with the de Broglie equation. $n \times \lambda$ is represented by a ratio to space-time. It is the number of ripples that occur in the energy level higher than the level occupied by the electron until interference occurs between the two levels, the higher energy level and the level occupied by the electron. In other words, as the number of orbitals occupied by the electron increases, the number of ripples that occur at the higher energy levels increases, causing the curvature (contraction) of the fabric of space-time. The interference between the two levels occurs in a wave form so that the quantum jump of the electron occurs. The photon's energy is represented by a ratio to the fabric of space-time, the force that causes the fabric of space-time to bend (contract). The more energy increases, the more space-time contracts through the occurrence of quantum disturbances at the highest energy level, which makes the highest energy level generate waves similar to the orbital number occupied by the electron. Because of these disturbances that occur at the highest energy level, the two levels interfere with each other, the highest energy level, and the level occupied by the electron. A quantum leap occurs, and this is

consistent with the quantum Zeno effect, where the electron will remain fixed in its position. This is what my equation indicates, as I explain that these quantum fluctuations occur through a contraction in the fabric of space-time. This contraction occurs as a result of this tissue absorbing energy. Because of this, contraction affects the energy levels in the atom. This contraction works to contract the energy level higher than the level occupied by the electron. Wave interference occurs between the highest energy level and the level occupied by the electron, and a quantum jump occurs from the observer’s perspective. But from the electron's perspective, it remains fixed in its position.

The Casimir effect is according to a law that states that after all the objects acting on the plates disappear until imaginary particles are detected. My equation proves that there is one thing that was not included in the calculations, which is the effect of space-time. Since the plates have a static mass that works to curve space-time, and the presence of imaginary particles works when they collide with each other, they disappear. But according to the law of conservation of energy, the energy will not disappear and will affect the fabric of space-time, making it turbulent like a water wave, and these disturbances that occur on it form waves. This wave works to impact the panels from moving in and out, and because the external disturbances are higher than the internal ones, they cause the panels to move towards each other.

This relationship shows that although we cannot measure what happens when an electronic quantum jump occurs. This law also shows that there is a relationship between the energy of the photon and the fabric of space-time, even if it is not measured by measuring devices. Because measuring devices are considered primitive devices when making the process of measuring the quantitative world. What is being measured are the spectra of the elements being measured, not what happens to the electron when the quantum jump of the electron to the higher level. Second, Maxwell told Rutherford that the electron changes direction as it orbits the nucleus, so it must lose energy to cause a collision with the nucleus, which it does not. My equation tells me the electron moves in a large circle around the nucleus. A body moving in a large circle whose direction of motion is in a straight line. Thus, the electron moves in a straight line. Newton's law states that an object at rest remains at rest unless acted upon by an external or internal force. Likewise, an object in motion stays in motion unless an external or internal force affects its movement, the electron does not lose energy.

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{(2\pi)^5 \times (\hbar)^4 \times 4}{(\lambda \, nm \times e)^4} \frac{G}{(\Delta E_n)^4} T_{\mu\nu}$$
$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{(2\pi)^6 \times (\hbar)^4 \times 4}{(\lambda \, nm \times e)^5 \times (\mu_0 \times \epsilon_0)^2} \frac{(l_{(p)})^2}{(\Delta E_n)^5} T_{\mu\nu}$$

Table 7. Comparing my theoretical results through my equation with previous results. Theoretical value (My work) -----Experimental value.

Spectral Line	Energy	λ	λ
$\lambda(n'=2, n=1)$	10.204269824 eV	121.50227268 nm	121.5 nm
$\lambda(n'=3, n=1)$	12.093949421 eV	102.51754257 nm	102.5 nm
$\lambda(n'=4, n=1)$	12.75533728 eV	97.20181814 nm	97.20 nm
$\lambda(n'=3, n=2)$	1.8896795971 eV	656.11227245 nm	656.1 nm
$\lambda(n'=4, n=2)$	2.5510674561 eV	486.0090907 nm	486.0 nm
$\lambda(n'=4, n=3)$	0.66138785898 eV	1874.6064927 nm	1874.6 nm

The results of the experimental value were obtained by using the results of previous research on the hydrogen atom. I prove in table 7 that the results of the equations are identical to their original results in table 5, which indicates the validity of this law

$$\Delta E_n = E_2 - E_1$$

$$\Delta E_n = \frac{h_p \times C}{\lambda} = \frac{1239.8419637 \text{ eVnm}}{\lambda}$$

Photon energy equation.

$$\Delta E_n = \frac{-13.605693099 \text{ eV}}{1} \times \left(\frac{(Z)^2}{(n_2)^2} - \frac{(Z)^2}{(n_1)^2} \right)$$

- These are the results of a relationship between energy and wavelength. The observed results show that whenever the energy increases, the wavelength decreases, as shown by this equation in the hydrogen atom.

7. Conclusions

After the idea of research has been clarified using theoretical and practical scientific evidence to explain the phenomenon of the quantum leap and quantum entanglement from a new perspective, these equations would be used in the following:

- 1) serving humanity in the advancement of scientific research.
- 2) using these equations to explore space and quantum world.
- 3) using these equations in developing communications machines .

References

1. Svidzinsky, A. Scully, M. Bohr's molecular model, a century later. Physics T. 2014, 67, 33-39. <http://dx.doi.org/10.1063/PT.3.2243>
2. Udema, I. I. Renaissance of Bohr's model via derived alternative equation. American J. Mod. Phys 2017, 6, 23-31. <https://doi.org/10.11648/j.ajmp.20170602.11>
3. Nanni, L. The hydrogen atom: A review on the birth of modern quantum mechanics. Physics 2015. <https://doi.org/10.48550/arXiv.1501.05894>
4. Jordan, R. B. Principles of Inorganic Chemistry. Springer N., 2024; pp. 1--18. https://doi.org/10.1007/978-3-031-22926-8_1
5. Manini, N. Introduction to the physics of matter: basic atomic, molecular, and solid-state physics. Springer N., 2020; pp. 11--16. https://doi.org/10.1007/978-3-030-57243-3_2

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.