

Article

Not peer-reviewed version

---

# Collatz Conjecture Is Analogous to an Inverse Function of Natural Number

---

[Jishe Feng](#) \*

Posted Date: 7 June 2024

doi: 10.20944/preprints202405.0310.v2

Keywords: binary string; full binary directed tree; composite function; Collatz conjecture; ultimately periodic sequence



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Article

# Collatz Conjecture is Analogous to an Inverse Function of Natural Number

Jishe Feng

School of Mathematics and Information Engineering, Longdong University, Qingyang, China; gsfs6567@126.com

**Abstract:** We propose a full binary directed tree to represent the set of natural numbers and further divide the set into three sets: pure odd, pure even, and mixed numbers. We utilize a binary string to represent a natural number and demonstrate the composite procedure of odd-number and even-number functions. We analyze the sequence of iteration (or composite) of the Collatz function and reduced Collatz function analog to the inverse function in order to test the Collatz conjecture. We do this by using the parity of a natural number. In order to prove the conjecture, we provide tabular and binary strings to the algebraic formula that states the Collatz sequence. Ultimately, we can convert discrete powers of 2 into continuous powers of 2, ultimately arrive at the smallest natural, 1. If any natural number is the beginning value, the sequence produced by the infinite iterations of the Collatz function becomes the eventually periodic sequence, proving an 87-year-old conjecture.

**Keywords:** binary string; full binary directed tree; composite function; Collatz conjecture; ultimately periodic sequence

## 1. Introduction

In the study of number theory, odd and even numbers are a fundamental pair of ideas. Natural number sets can be parted to two different sets. There are many conjectures that attempt to generalize the fact of different kinds of natural numbers discovered in a restricted range to the entire infinite set of natural numbers. This article will proof the famous Collatz conjecture, which states that for each natural number  $n$ , if it is even, divided by 2, if it is odd, multiplied by 3, and added 1, and so on, the eventual value must be 1. It is also referred to as the  $3n + 1$  conjecture and was put forth in 1937 by Lothar Collatz, also known as the  $3n + 1$  problem. The mathematician Paul Erdos once said of this conjecture: "Mathematics may not be ready for such problems" [1,2].

For the Collatz conjecture, we can describe it as a function:

$$T(n) = \begin{cases} 3n + 1, & \text{if } n \text{ is odd number,} \\ \frac{n}{2} & \text{if } n \text{ is even number.} \end{cases} \quad (1)$$

The following sequence is obtained via the composite function (iteration):  $\Lambda = \{n, T(n), T(T(n)), T(T(T(n))), \dots\} = \{n, T(n), T^2(n), T^3(n), \dots\}$ . Consequently, the Collatz conjecture can be stated as follows:

**Collatz conjecture 1:** For any natural number  $n$ , there is finite natural number  $m$ , the sequence  $\Lambda$  always leads to the integer 1, namely  $T^m(n) = 1$ .

The series  $\Lambda$  is an infinite sequence of ultimately period [3,4]. So we give another statement of the Collatz conjecture as the following.

**Collatz conjecture 2:** The series  $\Lambda$  is an infinite sequence of ultimately period, the preperiod  $\eta(n)$  varies with the initial value  $n$ , but the ultimately period is always  $\{1, 4, 2\}$ .

## 2. An Algebra and Graph Representation of Natural Numbers

### 2.1. The Composition of Odd-Number and Even-Number Functions

A natural number is considered even if it can be divided by 2; if not, it is considered odd. According to the Peano's Axiom, the smallest natural number is 1. The set  $N = \{1, 2, 3, \dots\}$  of natural

numbers can be divided into odd and even sets; in this paper, we will use the usual definition of natural numbers.  $\{\text{natural number}\} = \{\text{odd number}\} \cup \{\text{even number}\}$ .

In the set of natural numbers where 1 is the smallest odd number and 2 is the smallest even number, we can use the expression  $n = 2k - 1$  to indicate that it is an odd, and the expression  $n = 2k$  to indicate that it is an even, where  $k$  is any natural number.

We introduce two functions  $O(x) = 2x + 1$  to express odd numbers greater than 1, and  $E(x) = 2x$  to express even numbers, where  $x$  is any natural number in  $N$ .

We define a strictly increase monotonically piecewise function  $f(n)$ , from a natural number  $n$  it generates two cases: odd or even numbers:

$$f(n) = \begin{cases} 2n + 1 = O(n), & \text{the value is odd number,} \\ 2n = E(n), & \text{the value is even number.} \end{cases} \quad (2)$$

**Definition 1** A natural number  $n$  is obtained by composition of the odd-number function  $O(x) = 2x + 1$  and the even-number function  $E(x) = 2x$  several times, namely

$$n = f(f(\cdots f(1))) = f^k(1),$$

the function  $f$  is either odd-number function  $O(x)$  or even-number function  $E(x)$ .


For example,  $f(1) = O(1) = 3, f(1) = E(1) = 2$ ,  
 $7 = f^2(1) = 2 \cdot 3 + 1 = 2 \cdot (2 \cdot 1 + 1) + 1 = O(O(1))$ ,  
 $189 = f^7(1) = 2 \cdot (2 \cdot (2 \cdot (2 \cdot 4 + 1)$   
 $= 2 \cdot (2 \cdot (2 \cdot (2 \cdot (2 \cdot 1) + 1) + 1) + 1) + 1) + 1) + 1) + 1$   
 $= E(O(O(O(O(E(O(1))))))$ .

Any natural integer  $n$  is the value of the finite times composite function of the odd-number and even-number functions beginning at 1. In particular, If  $n = f(f(\cdots f(1))) = f^r(1)$ , then  $f^{-r}(n) = 1$  is the inverse function.

## 2.2. Use Binary String to Represent Natural Numbers

Using binary string to represent a natural number  $n$ , we can more clearly express the odd-number and even-number functions starting at 1 composite process of a natural number. The string indicates the order of composition of  $O(x)$  and  $E(x)$ ; 0 implicates an even-number function, and 1 implicates an odd-number function.

A natural number's binary string represents its odd-even composite function; from left to right, the  $i(i > 0)$  odd-number function  $O(x)$  is represented by the 1 in the  $i(i > 0)$ -bit, and the equivalent even-number function  $E(x)$  is represented by 0. For instance, the procedure of the composite function of 60 is displayed in Figure 1.




**Figure 1.** Natural number  $60 = (111100)_2$  is obtained starting 1 through the composition of five even-number and odd-number functions.

### 2.3. Use a Graph to Represent the Composite Procedure of the Natural Numbers

In order to give an intuitive impression, we provide a **full binary directed tree** to represent the natural number set, the root is the smallest number 1. For per vertex, its left-child is an even number which double itself, its binary string is appended by 0, right-child is an odd number which double itself and add 1, its binary string is appended by 1. For an natural number its binary string indicates the procedure of the composition of  $O(x)$  and  $E(x)$  from initial value 1 to the final accordingly binary string from the left to right. The full binary directed tree, as in Figure 2, is a very good representation of natural number  $N$ .


**Proposition 1** A binary string's length indicates its level in the full binary directed tree, and a binary string's length minus one represents the number of times the odd- and even-number composite functions occur.



**Figure 2.** The representation of natural number set is a full binary directed tree.

For given natural number  $n$ , its binary string from the second bit in left to right appending 0 or 1, indicates which comes from the root 1 of the full binary directed tree traversal according only one branch to itself. For instance, Figure 3 illustrates the procedures of composite odd-number and even-number of 21 and 29.

Given a natural number  $n$ , its binary string make of 0 or 1 from the second bit from left to right shows which originates from the root 1 of the full binary directed tree traversal, which only allows one branch to lead to itself. For example, the processes for the composite odd-number and even-number of 21 and 29 are shown in Figure 3.



**Figure 3.**  $21 = (10101)_2$  and  $29 = (11101)_2$  comes from the path from root 1 walk to 10101 and 11101 accordingly appending 1 or 0 to the vertexes in succession.

## 2.4. Another Partition of the Natural Number Set

We give the definitions of three kinds of natural numbers:

**Definition 2(i)** A natural number,  $O^m(1) = 2^m - 1 = 2^{m-1} + 2^{m-2} + \dots + 2 + 1 = (11 \dots 1)_2$ , is obtained by applying the odd-number function  $O(x)$   $m$  components. We call it as **pure odd number**. For instance, those are pure odd numbers:  $3 = (11)_2, 7 = (111)_2, 15 = (1111)_2, 31 = (11111)_2, 63 = (111111)_2, \dots$ . These are located in the full binary directed tree of Figure 2, which is the right leg of the isosceles triangle.

**(ii)** A natural number,  $E^m(1) = 2^m = (10 \dots 0)_2$ , is obtained by applying the even-number function  $E(x)$   $m$  components. We call it as **pure even number**. For instance, those are pure even numbers:  $2 = (10)_2, 4 = (100)_2, 8 = (1000)_2, 16 = (10000)_2, 32 = (100000)_2, 64 = (1000000)_2$ . Those are located in the left leg of the isosceles triangle, namely the full binary directed tree of the Figure 2.

**(iii)** The natural number obtained by the composition of odd function  $O(x)$  and even function  $E(x)$ , we call it **mixed number**. Such as,  $18 = (10010)_2, 28 = (11100)_2, 67 = (1000011)_2, 309 = (100110101)_2$ . Those are in the inside of the isosceles triangle, the full binary directed tree of the Figure 2.

In particular, the natural numbers obtained by the finite alternately composition of the odd function  $O(x)$  and the even function  $E(x)$ , namely,  $[E(O(1))]^m = (101 \dots 101)_2$ . Such as  $5 = (101)_2, 21 = (10101)_2, 85 = (1010101)_2, 341 = (101010101)_2, 1365 = (10101010101)_2, 5461 = (1010101010101)_2, \dots$ .

We call **hard numbers**.

The traversal path in the full binary directed tree from the root to down along the arcs, for each natural integer  $n$ , is its binary string  $1 \times \dots$ , where the left-child appended 0 for each vertex is an even number and the right-child appended 1 for each vertex is an odd number. For instance, in Fig. 3,  $21 = (10101)_2$  originates at the root 1 and proceeds down 2, 5, 10, ultimately reaching 21. To the vertexes,  $1 \rightarrow 10 \rightarrow 101 \rightarrow 1010 \rightarrow 10101$ , 0, 1, 0, 1 are appended. In addition, for  $29 = (11101)_2$ , the appendix 1, 1, 0, 1 is added to the vertexes,  $1 \rightarrow 11 \rightarrow 111 \rightarrow 1110 \rightarrow 11101$ , after it descends from the root 1 down 3, 7, 14, and ultimately reaches 29.

**Property 2** The set of natural numbers can be partitioned into three different sets:  $\{\text{natural number}\} = \{\text{pure even number}\} \cup \{\text{pure odd number}\} \cup \{\text{mixed number}\}$ , where  $\{\text{mixed number}\} = \{\text{mixed even number}\} \cup \{\text{mixed odd number}\}$

**Example 1** (1) 60, 97 are mixed numbers.

(2) 64, 1180591620717411303424 are pure even numbers.

(3) 63, 1180591620717411303423 are pure odd numbers.

When we convert those natural numbers from decimal to binary, the facts are obvious.

(1)  $60 = (111100)_2$  is a mixed-even number,  $97 = (1100001)_2$  is a mixed-odd number.

(2)  $64 = 2^6 = (1000000)_2$ ,  $1180591620717411303424 = 2^{70} = (10000 \dots 0)_2$  are pure even numbers:

$63 = (111111)_2$ ,  $1180591620717411303423 = 2^{70} - 1 = (11 \dots 1)_2$  are pure odd numbers.

## 3. Two Functions Compare

In order to proof the Collatz conjecture 1 in section 1, finding the beingness and finiteness of the number  $m$  in the expression  $T^m(n) = 1$  for a natural number  $n$  is the main challenge.

To visually represent the process of the composite functions of odd- and even-number functions, i.e.,  $n = f^k(1)$ , we propose a full binary directed tree. The following is the inverse functions  $f^{-1}(x)$ ,

$$f^{-1}(n) = \begin{cases} \frac{n-1}{2}, & \text{the value is odd number,} \\ \frac{n}{2}, & \text{the value is even number.} \end{cases} \quad (3)$$

The inverse function  $f^{-1}(x)$  differs from the Collatz function  $T(x)$ , but so do the results of their own finite iterations. However, for infinite or a sufficiently large number of iterations, the iteration Collatz function's value is the smallest number 1 to the cycle  $\{1, 4, 2\}$ .

If there is a formula  $f^k(1) = n$  for a given natural number  $n$ , we know that  $k$  is the length of the binary string of  $n$  minus 1. In decimal notation, we represent  $n$ , which obscures the composite process of odd- and even-number functions. When  $n$  is represented as a binary string, it can be used to understand how odd- and even-number functions work.


The iteration of the Collatz function is the key topic in discuss the proof procedure, we have the reduced Collatz function [2,5,6]  $RT(x)$

$$RT(n) = \begin{cases} \frac{3n+1}{2^m}, & \text{if } n \text{ is odd number, } \frac{3n+1}{2^m} \text{ is an odd number,} \\ \frac{n}{2^r}, & \text{if } n \text{ is even number, } \frac{n}{2^r} \text{ is an odd number.} \end{cases} \quad (4)$$


There are many different points for piecewise functions when comparing the Collatz function  $T(x)$  with the function  $f^{-1}(x)$ , the reduced Collatz function  $RT(x)$ , and the iteration function  $f^k(x) = O(E(\dots(E(x))))$ .

1) For any natural number  $x$  the function  $f^{-1}(x)$  and  $f^{-r}(x)$  are strictly monotonically decreasing.

2) The function  $T(x)$ , which is strictly monotonically rising, is merely one case of the functions where  $x$  is purely even. The function  $RT(x)$  is wavy when  $x$  is a pure or mixed odd number. It is increasing at first, then goes through one or more decreasing processes, either as "increase – decrease – increase" or "increase – decrease  $\dots$  decrease – increase." For instance, the iterated sequence of Collatz functions is plotted in Figures 4 and 5, where the beginning values are pure odd  $255 = 2^8 - 1 = (1111111)_2$  and mixed odd number  $97 = (1100001)_2$ , respectively.



**Figure 4.** Point plot of a sequence of 47 iterations of the Collatz function for pure odd 255.



**Figure 5.** Point plot of a sequence of 118 iterations of the Collatz function for mixed odd 97.



#### 4. Proof the Collatz Conjecture

We have known that mathematics formula about geometric progression with initial term 1 and common ratio  $x$ , the sum of the first  $k$  terms is

$$x^k + x^{k-1} + \cdots + x + 1 = \frac{x^{k+1} - 1}{x - 1} \quad (5)$$

when  $x = 2$ , there is a formula

$$2^k + 2^{k-1} + \cdots + 2 + 1 = 2^{k+1} - 1 \quad (6)$$

The substantive characteristics is that the powers of 2 must be continuous natural numbers, this is the key to our proof method to solve the Collatz conjecture.

**Proof.** (i) The conjecture states that if a given natural number  $n = 2^k$  is a pure even, then the smallest natural number 1 can be reached by simply repeating the  $k$  times Collatz function divided by 2.

(ii) When a natural number  $n$  is not pure even, that is, when it is either pure odd or mixed odd (a mixed even is eliminated since the end-substring 0 can become an odd number; hence, we do not include this case).

Given an odd number  $n$ , which can be expressed as follows in algebraic notation:  $n = (1 \times \cdots \times 1)_2 = 2^r + 2^m + \cdots + 1$ , then

$$\begin{aligned} 3n + 1 &= 2n + n + 1 \\ &= 2^{r+1} + 2^{m+1} + \cdots + 2 + 2^r + 2^m + \cdots + 1 + 1 \\ &= 2^{r+1} + 2^{m+1} + \cdots + 2 + 2^r + 2^m + \cdots + 2 \\ &= 2^{r+1} + 2^r + 2^{m+1} + 2^m + \cdots + \cdots + 2^h \end{aligned}$$

If the length of the end-substring of  $n$  is 1, the length of end-substring of the binary string  $3x + 1$  is either 1 or bigger than 1.

Two formulas,  $2^l + 2^l = 2^{l+1}$  and above (6), can be used to modify the structure of the binary string  $n$  to the binary string  $3n + 1$ . That is, there is an appended term  $2^{r-h}$  in two equivalent terms,  $2^{r+1-h}$  and  $2^{m+1-h}$ . When the zeros in the middle of a binary string are compared to a bubble, it means that these zeros are gradually being driven out of the rightmost end by  $3n + 1$ . It is the same as progressively removing the bubbles hidden in the sponge using a means  $3n + 1$ . Once  $3n+1$ , 0 shifts one bit to the right, i.e., the length of the associated binary substring is reduced by one bit, when the length of the end-substring is greater than 1. The end-substring length of the binary string  $3x + 1$  is either greater than or equal to 1 if the length of the end-substring of  $n$  is 1.

We shall then divide  $3n + 1$  by the last term  $2^h$ ,

$$\frac{3n + 1}{2^h} = 2^{r+1-h} + 2^{r-h} + 2^{m+1-h} + 2^{m-h} + \cdots + \cdots + 2^0$$

get another odd number, this is the value of reduced Collatz function (4). And so on, finitely steps after finally we get a pure even number  $2^t$ , this is the case in above (i), thus the Collatz conjecture hold on.

We illustrate the procedure by a mixed odd number  $n = 67$  and hard number set in the following,

$$\begin{aligned}
 67 &= (1000011)_2 = 2^6 + 2^2 + 2^0 \\
 3 \cdot 67 + 1 &= 2 \cdot 67 + 67 + 1 = 2^7 + 2^2 + 2^1 + 2^6 + 2^1 + 2^0 + 2^0 = 2^7 + 2^6 + 2^3 + 2^1 \\
 3 \cdot 101 + 1 &= 2 \cdot 101 + 101 + 1 = 2^7 + 2^6 + 2^3 + 2^1 + 2^6 + 2^5 + 2^2 + 2^0 + 2^0 = 2^8 + 2^5 + 2^4 \\
 3 \cdot 19 + 1 &= 2 \cdot 19 + 19 + 1 = 2^5 + 2^2 + 2^1 + 2^4 + 2^1 + 2^0 + 2^0 = 2^5 + 2^4 + 2^3 + 2^1 \\
 3 \cdot 29 + 1 &= 2 \cdot 29 + 29 + 1 = 2^5 + 2^4 + 2^3 + 2^1 + 2^4 + 2^3 + 2^2 + 2^0 = 2^6 + 2^4 + 2^3 \\
 3 \cdot 11 + 1 &= 2 \cdot 11 + 11 + 1 = 2^4 + 2^2 + 2^1 + 2^3 + 2^1 + 2^0 + 2^0 = 2^5 + 2^1 \\
 3 \cdot 17 + 1 &= 2 \cdot 17 + 17 + 1 = 2^5 + 2^1 + 2^4 + 2^0 + 2^0 = 2^5 + 2^4 + 2^2 \\
 3 \cdot 13 + 1 &= 2 \cdot 13 + 13 + 1 = 2^4 + 2^3 + 2^1 + 2^3 + 2^2 + 2^0 + 2^0 = 2^5 + 2^3 \\
 3 \cdot 5 + 1 &= 2 \cdot 5 + 5 + 1 = 2^3 + 2^1 + 2^2 + 2^0 + 2^0 = 2^4 \\
 1
 \end{aligned}$$

For a special class of mixed numbers, the hard number  $\frac{4^k-1}{3} = (101 \cdots 101)_2$ , then its Collatz sequent result is

$$\begin{aligned}
 a_k &= \frac{4^k - 1}{3} = \frac{4^k - 1}{4 - 1} = 4^{k-1} + 4^{k-2} + \cdots + 4 + 1 = (101 \cdots 101 \cdots 101)_2, \\
 T(a_k) &= 3a_k + 1 = 4^k = 2^{2k} = (10 \cdots 0)_2, T^{2k+1}(a_k) = 1.
 \end{aligned}$$

This means that the Collatz conjecture is valid for this case. Therefore we have proved the Collatz conjecture 1 at section 1 of this paper.  $\square$

**Proof.** For the smallest number  $1 = 2^0$

$$\begin{aligned}
 T(1) &= 3 \times 1 + 1 = 2 \times 1 + 1 + 1 = 2^1 + 2^0 + 2^0 = 2^1 + 2^1 = 2^2 = 4, \\
 T^2(1) &= 2, \\
 T^3(1) &= 1,
 \end{aligned}$$

and so on, this is a cycle  $\{1, 4, 2\}$ , This is the Collatz conjecture 2 at section 1 of this paper.  $\square$

## 5. Conclusions

From previous proof of the conjecture, it becomes a theorem.

**Theorem 1** There exists a finite natural number  $m$  for every natural number  $n$ . The Collatz sequence always arrives at the integer 1, that is,  $T^m(n) = 1$ .

**Theorem 2** For any positive integer  $n$ , the sequence of the iteration of the Collatz function is an ultimately periodic sequence, its preperiod  $\eta(n)$  is a related-to  $n$ , and the least period is  $\{1, 4, 2\}$ .

## References

1. Lagarias, Jeffrey C. The Ultimate challenge: the  $3x + 1$  problem; American Mathematical Society, Providence, RI, 2010. isbn=978-0-8218-4940-8.
2. Richard Kaufman. A reduced forward Collatz algorithm: How binary strings change their length under  $3x+1$ . arXiv 2023, arXiv:2301.07466.
3. G. Ganesan. Linear recurrences over a finite field with exactly two periods. *Advances In Applied Mathematics* 2021. 102-108.
4. D. Quijada. Periods of linearly recurring sequences. Bachelor thesis, Washingto and Lee University. 2015. isbn=978-0-8218-4940-8
5. Bulent Sukusu Proof of the Collatz Conjecture. Theoret. Math. App. 2023, 13, 1-7.
6. Jishe Feng. Proof the Collatz Conjecture by a new view of natural numbers, 2023.



7. Eldar Sultanow, Christian Koch, Sean Cox, title=Collatz Sequences in the Light of Graph Theory. 2023. doi.org/10.25932/publishup-48214.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.