

Article

Not peer-reviewed version

4+1 Gravitation in the SHP Formalism

[Martin Land](#)*

Posted Date: 2 March 2026

doi: 10.20944/preprints202603.0002.v1

Keywords: Stueckelberg-Horwitz-Piron gravitation; ADM formalism; Hamiltonian systems; KaluzaKlein theory



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a [Creative Commons CC BY 4.0 license](#), which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

4+1 Gravitation in the SHP Formalism

Martin Land 

Jerusalem Multidisciplinary College (Formerly, Hadassah College); martin@jmc.ac.il

Abstract

The Stueckelberg-Horwitz-Piron (SHP) formalism describes particles and fields traced-out as spacetime events functionally dependent on an external evolution parameter τ . This approach addresses a number of difficulties associated with the problem of time. In SHP general relativity, the state of the unconstrained phase space variables $\{x^\mu(\tau), p_\nu(\tau)\}$ specifies a 4D block spacetime $\mathcal{M}(\tau)$ that evolves to an infinitesimally close 4D block spacetime $\mathcal{M}(\tau + \delta\tau)$ under a scalar Hamiltonian. As the configuration of matter and energy evolves with τ it induces changes in the spacetime metric $\gamma_{\mu\nu}(x, \tau)$, leading to τ -dependent geodesic equations for the phase space variables. The 4+1 approach in gravitation generalizes 3+1 formalism of Arnowitt, Deser, and Misner (ADM) to construct τ -dependent Einstein field equations, a canonical Hamiltonian formalism, and an initial value problem for $\gamma_{\mu\nu}(x, \tau)$. To conform to known gravitational phenomenology, we must respect the 5D gauge symmetries associated with the free fields — the geometrical constructs relevant to $\mathcal{M}(\tau)$ as an embedded hypersurface — and the $O(3,1)$ symmetries of 4D matter. The 4+1 formalism has been discussed in a series of publications. The goal of this paper is to provide a systematic review of the subject, make a few corrections and some significant additions, and present the theory in a concise and orderly fashion.

Keywords: Stueckelberg-Horwitz-Piron gravitation; ADM formalism; Hamiltonian systems; Kaluza-Klein theory

1. Introduction

The Stueckelberg-Horwitz-Piron (SHP) formalism for classical and quantum relativistic dynamics [1–4] addresses a number of difficulties in relativity associated with the problem of time [5–7]. These difficulties are often attributed to the conceptual conflict between general relativity (GR) and quantum theory (see, for example, [8]), but the fundamental issue was already explored by Stueckelberg [9,10] in his foundational work on classical and quantum electrodynamics. In relativity, as in natural language, time refers to two distinct aspects of nature: instantaneous location and evolving chronology [11]. As one of four spacetime coordinates, time is determined by equations of motion, but it also serves as the evolution parameter by which these equations are posed and solved. The SHP framework explicitly separates these roles by introducing an external evolution parameter τ , independent of the phase space coordinates, providing the chronological “arrow of time” by which the equations of motion may be integrated.

In a series of recent papers [12–18] the author has developed a 4+1 approach to 4D gravitation within the SHP formalism. In this paper, we review this work, make a few corrections and some significant additions, and present the theory in a concise and orderly fashion.

The principal goal of the 4+1 approach is to formulate generalized field equations for a local 4D spacetime metric $\gamma_{\mu\nu}(x, \tau)$ that evolves under the influence of matter and field sources $T_{\mu\nu}(x, \tau)$, themselves evolving as spacetime events $x^\mu(\tau)$ trace out worldline trajectories under the monotonic advance of τ . As we shall see, attention to the gauge and spacetime symmetries required for such field equations leads to a 4+1 formalism analogous to the 3+1 geometrodynamics of Arnowitt, Deser, and Misner (ADM) [19]. The resulting formalism recovers standard Einstein GR in the equilibrium limit of no τ -evolution, and is compatible with standard Kaluza-Klein theory in this limit. Following a brief

review of the SHP formalism, motivating the 4+1 approach, we will discuss the geodesic equations in this framework and develop the field theory for the metric.

1.1. Hamiltonian Structure on 8D Phase Space

In 1937, Fock [20] generalized the phase space of nonrelativistic classical mechanics to 4D Minkowski spacetime as

$$\{x_n^i(t), p_i^n(t)\}, i = 1, \dots, 3 \longrightarrow \{x_n^\mu(s), p_\mu^n(s)\}, \mu = 0, \dots, 3 \quad (1)$$

where $n = 1, \dots, N$. Here, $x_n^0(s) = ct_n(s)$ represents the time registered on a laboratory clock, and Fock identified the monotonically advancing scalar parameter s with the proper time of the motion. Writing the velocity

$$\dot{x}^\mu(s) = \frac{dx^\mu}{ds} \quad (2)$$

and the Lorentz scalar action

$$S = \int ds L(x, \dot{x}, s) = \int ds \left(\frac{1}{2} M \dot{x}^\mu \dot{x}_\mu + \frac{e}{c} \dot{x}_\mu A^\mu \right) \quad (3)$$

he showed that the covariant Euler-Lagrange equations

$$\frac{d}{ds} \frac{\partial L}{\partial \dot{x}^\mu} - \frac{\partial L}{\partial x^\mu} = 0 \quad (4)$$

lead to the classical Lorentz force in Maxwell electrodynamics.

In 1941, Stueckelberg [9,10] proposed that a particle-antiparticle interaction could be described by a single evolving event interacting with an electromagnetic field in such a way that its energy becomes negative and it reverses direction in coordinate time x^0 . This description, central to quantum field theory, became known as the Feynman-Stueckelberg formalism [21,22]. But Stueckelberg observed that for classical worldlines, the evolution parameter cannot be identified with the proper time. While crossing the spacelike region that separates future-oriented trajectories from past-oriented trajectories, the sign of $\dot{x}^2$ will change twice, and so the computed proper time interval

$$\frac{1}{c} \sqrt{-\eta_{\mu\nu} dx^\mu dx^\nu} = \frac{1}{c} \sqrt{-\dot{x}^2(s)} ds \quad \eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1) \quad (5)$$

cannot provide a consistent parameterization. Therefore, Stueckelberg argued for the introduction of an external evolution parameter τ , analogous to the Newtonian time t and related to the proper time s through the dynamical relation $c^2 ds^2(\tau) = -\dot{x}^2(\tau) d\tau^2$. Implicit in the form of Fock's action (3) is that $\dot{x}^2$ is not *a priori* constrained, but it is nevertheless conserved under the electromagnetic interaction as

$$M \ddot{x}^\mu = e F^{\mu\nu} \dot{x}_\nu \longrightarrow \frac{d}{d\tau} \dot{x}^2 = \frac{e}{M} F^{\mu\nu} \dot{x}_\mu \dot{x}_\nu = 0. \quad (6)$$

To overcome this restriction, Stueckelberg proposed a modified Lorentz force that included a new vector field, but found no satisfying way to justify its presence in classical relativistic mechanics.

In 1973, Horwitz and Piron [23] advanced Stueckelberg's work, constructing a manifestly covariant relativistic mechanics with interactions on an unconstrained phase space, effectively demoting constraints associated with the presumed dynamics to the status of conservation laws. For example, one may introduce a scalar potential $V(x)$ to the action (3) producing the modified Lorentz force

$$M \ddot{x}^\mu = e F^{\mu\nu} \dot{x}_\nu - \frac{\partial V}{\partial x_\mu} \quad (7)$$

sought by Stueckelberg. More generally, by making the potentials τ -dependent and introducing a scalar gauge field $a_5(x, \tau)$

$$S_{\text{SHP}} = \int d\tau \frac{1}{2} M \dot{x}^\mu \dot{x}_\mu + \frac{e}{c} \dot{x}^\mu a_\mu(x, \tau) + \frac{e}{c} c_5 a_5(x, \tau) \quad (8)$$

the Fock action is made maximally U(1) gauge invariant [3,24] as

$$a_\mu \longrightarrow a_\mu + \partial_\mu \Lambda(x, \tau) \quad a_5 \longrightarrow a_5 + \frac{1}{c_5} \partial_\tau \Lambda(x, \tau) \quad (9)$$

where we introduce a constant fifth velocity c_5 to write $x^5 = c_5 \tau$ in analogy to the notation $x^0 = ct$. Partitioning the Greek indices as

$$\alpha, \beta, \gamma, \delta = 0, 1, 2, 3, 5 \quad \lambda, \mu, \nu, \rho \dots = 0, 1, 2, 3 \quad (10)$$

we see that the O(3,1) scalars $\dot{x}^\mu \dot{x}_\mu$, $\dot{x}^\mu a_\mu$, and a_5 leave the action

$$S_{\text{SHP}} = \int d\tau \frac{1}{2} M \dot{x}^\mu \dot{x}_\mu + \frac{e}{c} \dot{x}^\beta a_\beta(x, \tau) \quad (11)$$

4D Lorentz invariant, despite the 5D gauge invariance $a_\alpha(x, \tau) \longrightarrow a_\alpha(x, \tau) + \partial_\alpha \Lambda(x, \tau)$.

The canonical momentum is

$$p_\mu = \frac{\partial L}{\partial \dot{x}^\mu} = M \dot{x}_\mu + \frac{e}{c} a_\mu \quad (12)$$

leading to the O(3,1) scalar Hamiltonian

$$K = p_\mu \dot{x}^\mu - L = \frac{1}{2M} \left(p^\mu - \frac{e}{c} a^\mu \right) \left(p_\mu - \frac{e}{c} a_\mu \right) - \frac{e}{c} c_5 a_5 \quad (13)$$

subject to the canonical equations of motion

$$\dot{x}^\mu = \frac{dx^\mu}{d\tau} = \frac{\partial K}{\partial p_\mu} \quad \dot{p}_\mu = \frac{dp_\mu}{d\tau} = -\frac{\partial K}{\partial x^\mu} \quad (14)$$

and 4D Poisson bracket structure. The Stueckelberg-Schrodinger equation

$$i\hbar \partial_\tau \psi(x, \tau) = \left[\frac{1}{2M} \left(p^\mu - \frac{e}{c} a^\mu \right) \left(p_\mu - \frac{e}{c} a_\mu \right) - \frac{e}{c} c_5 a_5 \right] \psi(x, \tau) \quad (15)$$

provides a covariant quantum mechanics with relativistic generalizations to the standard central force problems [3,4].

1.2. Motivation for a τ -Dependent Local Metric

In the Stueckelberg-Horwitz-Piron (SHP) formalism, particle worldlines are traced out by the evolution of classical events $x^\mu(\tau)$ or quantum events $\psi(x, \tau)$ as the external evolution parameter τ advances monotonically. Even a ‘static’ classical particle in its rest frame is characterized by its uniform evolution along its t -axis as

$$x(\tau) = \left(x^0(\tau), \mathbf{x}(\tau) \right) = (c\tau, \mathbf{0}) \quad (16)$$

and event trajectories carrying electromagnetic charge induce five τ -dependent gauge fields $a_\alpha(x, \tau)$. In curved spacetime, these worldlines and fields produce a scalar event density $\rho(x, \tau)$ and an energy-momentum density $T_{\mu\nu}(x, \tau)$ that evolve with τ and act as sources for the gravitational field. Following Wheeler’s description of geometrodynamics, “spacetime tells matter how to move; matter tells spacetime how to curve” [25], the τ -evolution of the mass/energy/momentum distribution associated with these events entails the τ -evolution of spacetime curvature described by a local metric $\gamma_{\mu\nu}(x, \tau)$. In the presence of τ -dependent matter and fields, a τ -independent 4D metric would have the character of

an absolute background field, contradicting the goals of general relativity. Evolution of the metric with τ naturally leads to generalized geodesic equations, and requires generalized field equations.

General relativity (GR) begins with consideration of the squared interval

$$\delta x^2 = \gamma_{\mu\nu} \delta x^\mu \delta x^\nu = (x_2 - x_1)^2 \quad (17)$$

between two neighboring points of the spacetime manifold $\mathcal{M}$ viewed as a 4D block universe. The spacetime trajectory of a material event is a continuous sequence of events $x^\mu(s)$ of timelike separation in $\mathcal{M}$, so that s may be the proper time. Writing the invariant interval (17) as

$$\delta x^2 = \gamma_{\mu\nu} \delta x^\mu \delta x^\nu = \gamma_{\mu\nu} \frac{dx^\mu}{ds} \frac{dx^\nu}{ds} ds^2 = \gamma_{\mu\nu} \dot{x}^\mu \dot{x}^\nu ds^2 \quad (18)$$

suggests a dynamical description of the trajectory by the action

$$S = \int ds \frac{1}{2} \gamma_{\mu\nu} \dot{x}^\mu \dot{x}^\nu \quad (19)$$

to which the constraint $\dot{x}^2 = -c^2$ may be imposed *a posteriori*.

In the SHP framework, a physical event $x^\mu(\tau)$ is an irreversible occurrence at τ with spacetime coordinates x^μ . Thus, the 4D block universe $\mathcal{M}(\tau)$ similarly occurs at τ , representing the instantaneous manifold of general relativity. The scalar Hamiltonian K generates evolution of $\mathcal{M}(\tau)$ to an infinitesimally close 4D block universe $\mathcal{M}(\tau + d\tau)$ occurring at $\tau + d\tau$. We may therefore consider the interval

$$x_2(\tau + \delta\tau) - x_1(\tau) \simeq x_2(\tau) + \frac{dx(\tau)}{d\tau} \delta\tau - x_1(\tau) = \delta x + \frac{dx(\tau)}{d\tau} \delta\tau \quad (20)$$

which introduces a notion of 5D distance that combines the *geometrical distance* δx between any two arbitrary points in $\mathcal{M}(\tau)$ and the *dynamical distance* between events generated by the Hamiltonian that evolves $\mathcal{M}(\tau) \rightarrow \mathcal{M}(\tau + \delta\tau)$.

We may regard the event $X = (x, x^5) = (x^0, \dots, x^3, c_5\tau)$ as a point in a pseudo-spacetime $\mathcal{M}_5$ obtained as the image of an injective mapping

$$\Phi : \mathcal{M} \rightarrow \mathcal{M}_5 = \mathcal{M} \times R \quad X = \Phi(x, \tau) = (x, c_5\tau) . \quad (21)$$

For two neighboring points $X_1, X_2 \in \mathcal{M}_5$ we may write

$$X_1 = (x_1(\tau), c_5\tau) \quad X_2 = (x_2(\tau + \delta\tau), c_5(\tau + \delta\tau)) \quad (22)$$

so that

$$\delta X = X_2 - X_1 = \left(\delta x + \frac{dx}{d\tau} \delta\tau, c_5 \delta\tau \right) \quad (23)$$

suggests a 5D invariant interval of the form

$$\delta X^2 = \gamma_{\mu\nu} \left(\delta x^\mu + \frac{dx^\mu}{d\tau} \delta\tau \right) \left(\delta x^\nu + \frac{dx^\nu}{d\tau} \delta\tau \right) + \sigma c_5^2 \delta\tau^2 = g_{\alpha\beta}(x, \tau) \delta X^\alpha \delta X^\beta \quad (24)$$

where $\sigma = \pm 1$. This squared interval leads to the general Lagrangian

$$L = \frac{1}{2} M g_{\alpha\beta}(x^\mu, x^5) \dot{X}^\alpha \dot{X}^\beta \quad \mu = 0, 1, 2, 3 \quad \alpha, \beta = 0, 1, 2, 3, 5 \quad (25)$$

from which we may find equations of motion in the manifold determined by some local metric $g_{\alpha\beta}$. This structure permits us to characterize 4D spacetime $\mathcal{M}$ as a hypersurface embedded in $\mathcal{M}_5$. The natural foliation of $\mathcal{M}_5$ into equal- τ spacetimes $\mathcal{M}(\tau)$ is then analogous to the foliation of 4D spacetime into equal- t spacelike hypersurfaces in standard GR. As shown in Section 3, this facilitates

the generalization of 3+1 geometrodynamics to a 4+1 formalism for the field equations describing the induced 4D metric $\gamma_{\mu\nu}(x, \tau)$.

1.3. Overview of Paper

The remainder of this paper is structured as follows. In Section 2 we find the equations of motion from a Lagrangian of the form (25), noting the requirement to preserve 5D gauge symmetry but only $O(3,1)$ symmetry for particle motion. Section 3 presents the generalized Einstein field equations for the τ -dependent local metric. Once again, these equations must respect the 5D symmetries of free fields as geometric objects, while respecting the 4D symmetries of the matter terms. In Section 4 we generalize the 3+1 formalism [19] developed by Arnowitt, Deser, and Misner (ADM) to 4+1. In this approach we decompose the generalized field equations into an initial-value problem, posed as first order evolution equations for the induced 4D metric $\gamma_{\mu\nu}(x, \tau)$ along with the extrinsic curvature $K_{\mu\nu}(x, \tau)$. Section 5 presents the 4+1 formalism as a canonical Hamiltonian theory. As in ADM, strategic combinations of $\gamma_{\mu\nu}(x, \tau)$ and $K_{\mu\nu}(x, \tau)$ lead to a canonical Hamiltonian formalism for SHP GR derived from a scalar action. Section 6 discusses the perturbative theory for weak gravitation and shows that in this approximation the initial value problem can be obtained directly from the 5D Ricci tensor. In Section 7 we compare this 4+1 formalism to the Kaluza-Klein model [26,27] for 4D gravitation and electromagnetism, and establish the compatibility of the two approaches only in τ -independent equilibrium. Section 8 discusses some examples and some prospects for gravitation in the SHP formalism.

2. SHP Particle Dynamics

Varying the scalar event Lagrangian

$$L = \frac{1}{2} M g_{\alpha\beta} \dot{X}^\alpha \dot{X}^\beta \quad X^\alpha \in \mathcal{M}_5 \quad \alpha, \beta, \gamma = 0, 1, 2, 3, 5 \quad (26)$$

with respect to X^γ leads to 5D geodesic equations

$$\frac{D\dot{X}^\gamma}{D\tau} = \ddot{X}^\gamma + \Gamma_{\alpha\beta}^\gamma \dot{X}^\alpha \dot{X}^\beta = 0 \quad \Gamma_{\beta\gamma}^\alpha = \frac{1}{2} g^{\alpha\delta} \left(\frac{\partial g_{\delta\beta}}{\partial X^\gamma} + \frac{\partial g_{\delta\gamma}}{\partial X^\beta} - \frac{\partial g_{\beta\gamma}}{\partial X^\delta} \right) \quad (27)$$

with 5D Christoffel symbols in standard form. The apparent 5D symmetry of these equations is broken to $O(3,1)$ by imposing $\dot{x}^5 = c_5 \rightarrow \dot{x}^5 = 0$ as a constraint, so that

$$\frac{D\dot{x}^\mu}{D\tau} = \ddot{x}^\mu + \Gamma_{\alpha\beta}^\mu \dot{x}^\alpha \dot{x}^\beta = \ddot{x}^\mu + \Gamma_{\nu\sigma}^\mu \dot{x}^\nu \dot{x}^\sigma + 2c_5 \Gamma_{5\nu}^\mu \dot{x}^\nu + c_5^2 \Gamma_{55}^\mu = 0 \quad (28)$$

$$\frac{D\dot{x}^5}{D\tau} = \dot{x}^5 \equiv 0 \quad (29)$$

describe the dynamical degrees of freedom. In case $g_{5\alpha} = 0$ and $\partial_\tau g_{\mu\nu} = 0$, the connection components $\Gamma_{5\alpha}^\mu$ vanish, recovering the standard 4D GR geodesic equations. Writing the canonical momentum

$$p_\mu = \frac{\partial L}{\partial \dot{x}^\mu} = M(g_{\mu\nu} \dot{x}^\nu + c_5 g_{\mu 5}) \quad (30)$$

the Legendre transformation produces the Hamiltonian

$$K = \frac{1}{2M} p^2 + \frac{1}{2} c_5 g_{55} g^{5\mu} p_\mu - \frac{1}{2} c_5 g_{5\mu} g^{\mu\lambda} p_\lambda + \frac{1}{2} M c_5^2 g_{5\mu} g^{\mu\lambda} g_{\lambda 5} + \frac{1}{2} M c_5^2 g_{55} \quad (31)$$

representing the total 5D mass of the motion. It is useful here to think of mass as arising from $p^2 = M^2 \dot{x}^2$ which is not constrained to be a constant of the motion. Thus, particle energy may change with no corresponding change in 3-momentum, producing an change in the mass p^2 associated with the trajectory.

For $g^{5\mu} = 0$ the Hamiltonian takes the simple form

$$K = \frac{1}{2M} p^\mu p_\mu + \frac{1}{2} M c_5^2 g_{55} \quad (32)$$

with $g_{55}(x, \tau)$ appearing as a τ -dependent scalar potential on 4D spacetime. While such a potential would be non-geodesic in standard GR, it follows simply from the 5D geodesic equations. The Hamilton equations (14) permit us to write the Poisson bracket relations

$$\{F, G\} = \frac{\partial F}{\partial x^\alpha} \frac{\partial G}{\partial p_\alpha} - \frac{\partial F}{\partial p_\alpha} \frac{\partial G}{\partial x^\alpha} = \frac{\partial F}{\partial x^\mu} \frac{\partial G}{\partial p_\mu} - \frac{\partial F}{\partial p_\mu} \frac{\partial G}{\partial x^\mu} \quad (33)$$

where the second equality follows from $p_5 \equiv 0$. Since

$$\frac{dF}{d\tau} = \{F, K\} + \frac{\partial F}{\partial \tau} \quad (34)$$

for any scalar function F on phase space, the Hamiltonian is conserved unless it depends explicitly on τ through $g_{\alpha\beta}(x, \tau)$. Nevertheless, the mass $m = -p^\mu p_\mu / 2M$ associated with the event trajectory may vary under $g_{5\alpha}$. For a discussion of geodesic equations and quantum mechanics with a τ -independent local metric, see [28,29].

In the Newtonian limit, neglecting $\dot{x}^i/c < 1$ for $i = 1, 2, 3$ and taking $g_{\alpha\beta}$ to be t -independent, the equations of motion reduce to

$$\frac{d^2 t}{d\tau^2} = \frac{dt}{d\tau} \partial_\tau g_{00} \quad \ddot{\mathbf{x}} = \frac{1}{2} c^2 \left(\frac{dt}{d\tau} \right)^2 \nabla g_{00} + \frac{1}{2} c_5^2 \nabla g_{55} \quad (35)$$

which deviate from the standard nonrelativistic case if g_{00} is τ -dependent, producing an acceleration in the time coordinate t . A simple post-Newtonian expression is found by putting $\nabla g_{55} = 0$ and introducing an additive perturbation $\delta M(\tau)$ to the mass parameter M in the Schwarzschild metric $g_{00} = -1 + 2GM/c^2 R$. The t equation admits the solution

$$\frac{dt}{d\tau} = \exp\left(\frac{2G}{c^2 R} \delta M\right) \quad (36)$$

which recovers $\dot{t} = 0$ in the absence of the perturbation. In spherical coordinates, angular momentum $L = MR^2 \dot{\phi}$ is conserved and the radial equation is

$$\ddot{R} - \frac{L^2}{M^2 R^3} + \exp\left(\frac{4G}{c^2 R} \delta M\right) \frac{GM_0}{R^2} = 0, \quad (37)$$

which becomes Newtonian gravitation for $\delta M = 0$. We find the Hamiltonian for the test event in this coordinate system to be

$$K = \frac{1}{2} M g_{\alpha\beta} \dot{X}^\alpha \dot{X}^\beta = -\frac{1}{2} M c^2 \left(1 - \frac{2GM_0}{c^2 R}\right) \exp\left(\frac{4G}{c^2 R} \delta M\right) + \frac{1}{2} M \dot{R}^2 + \frac{1}{2} \frac{L^2}{MR^2} \quad (38)$$

with τ derivative

$$\frac{d}{d\tau} K = \exp\left(\frac{4G}{c^2 R} \delta M\right) \left(-\frac{GM}{r} + \frac{4G^2 M M_0}{c^2 R^2}\right) \frac{d}{d\tau} \delta M \quad (39)$$

so that the total mass K is not conserved in the presence of a variable mass gravitational source.

The scalar event density $\rho(x, \tau)$ is defined as the number of events per spacetime volume. The 5-component event current is

$$j^\alpha(x, \tau) = M \rho(x, \tau) \dot{x}^\alpha(\tau) \quad (40)$$

where $j^5 = Mc_5 \rho(x, \tau)$ is an $O(3,1)$ scalar and not the 5-component of a vector. Therefore, the continuity equation is

$$\nabla_\alpha j^\alpha = \frac{\partial j^\alpha}{\partial x^\alpha} + j^\gamma \Gamma_{\gamma\alpha}^\alpha = \frac{\partial \rho}{\partial \tau} + \nabla_\mu j^\mu = 0 \quad (41)$$

and by virtue of the continuity and geodesic equations, the mass-energy-momentum tensor

$$T^{\alpha\beta} = M\rho \dot{x}^\alpha \dot{x}^\beta \longrightarrow \begin{cases} T^{\mu\nu} = M\rho \dot{x}^\mu \dot{x}^\nu, \\ T^{5\beta} = \dot{x}^5 \dot{x}^\beta M\rho = c_5 j^\beta \end{cases} \quad (42)$$

is conserved as

$$\nabla_\alpha T^{\alpha\beta} = 0. \quad (43)$$

As shown in [30] the spacetime components of $T_{\mu\nu}$ have the usual interpretation, with T_{00} the energy flux and T_{0i} a 3-momentum flux for $i = 1, 2, 3$, while T_{55} is the mass flux and $T_{5\mu}$ is a 4-momentum flux for $\mu = 0, 1, 2, 3$. Varying the electromagnetic action (11) with respect to x^μ provides the Lorentz force

$$\begin{aligned} M\ddot{x}^\mu(\tau) &= \frac{e}{c} \left[f^\mu_\nu(x, \tau) \dot{x}^\nu(\tau) + c_5 f^\mu_5(x, \tau) \right] = \frac{e}{c} f^\mu_\alpha(x, \tau) \dot{x}^\alpha(\tau) \\ \frac{d}{d\tau} \left(-\frac{1}{2} M\dot{x}^2 \right) &= -M\dot{x}^\mu \ddot{x}_\mu = \frac{ec_5}{c} f_{5\mu} \dot{x}^\mu \end{aligned} \quad (44)$$

where

$$f_{\alpha\beta} = \partial_\alpha a_\beta - \partial_\beta a_\alpha \quad (45)$$

is the field strength. We see that $f_{5\mu}$ can change the mass $M\dot{x}^2$ of an evolving matter event, as anticipated by Stueckelberg. The mass-energy-momentum tensor for the electromagnetic field satisfies [24,30]

$$\partial_\alpha T^{\alpha\beta}_{field} = -\frac{e}{c} f^{\beta\alpha} \dot{x}_\alpha(\tau) \delta(x - x(\tau)) \quad T^{\alpha\beta}_{field} = \frac{1}{c} \left(f^{\alpha\gamma} f^\beta_\gamma - \frac{1}{4} \eta^{\alpha\beta} f^{\delta\epsilon} f_{\delta\epsilon} \right) \quad (46)$$

which together with the Lorentz force (44) leads to

$$\frac{d}{d\tau} \left[\int d^4x T^{55}_{field} + \eta^{55} \left(-\frac{1}{2} M\dot{x}^2 \right) \right] = 0 \quad (47)$$

expressing conservation of the combined mass of particles and fields. In these expressions, the scalar T^{55}_{field} represents the spacetime density of mass associated with the fields, in analogy to the energy density T^{00}_{field} . Similarly, $T^{5\mu}_{field}$ is a Poynting 4-vector describing the flow of electromagnetic 4-momentum into spacetime, in analogy with the standard Poynting vector T^{0i}_{field} .

Since the 5D Ricci tensor, obtained in the usual manner from the metric $g_{\alpha\beta}(x, \tau)$, is invariant under translations $x'^\alpha = x^\alpha + \Lambda^\alpha(x, \tau)$, the Bianchi identity [31]

$$\nabla_\alpha \left(R^{\alpha\beta} - \frac{1}{2} g^{\alpha\beta} R \right) = 0 \quad (48)$$

must hold. In the absence of the matter current, the first of (46) provides

$$\nabla_\alpha T^{\alpha\beta}_{field} = 0 \quad (49)$$

and so together with (43) we may write 5D Einstein equations

$$R_{\alpha\beta} - \frac{1}{2} g_{\alpha\beta} R = \frac{8\pi G}{c^4} \left(T_{\alpha\beta} + T^{\alpha\beta}_{field} \right) \quad (50)$$

with the mass-energy-momentum tensors as sources. Nevertheless, the spacetime symmetry of $T^{\alpha\beta}$ should be no larger than $O(3,1)$ as a matter of principle, and so the apparent 5D symmetry of (50) must be broken to 4+1. This symmetry breaking is most conveniently accomplished in a quintrad frame.

3. SHP Field Equations

3.1. Quintrad Frame

The tangent space $\mathcal{T}(\mathcal{M}_5)$ of the pseudo-spacetime $\mathcal{M}_5$ can be given the standard coordinate basis

$$\mathbf{g}_\alpha = \partial_\alpha = \frac{\partial}{\partial X^\alpha} \quad (51)$$

while the dual space $\mathcal{T}^*(\mathcal{M}_5)$ has the basis 1-forms

$$\mathbf{g}^\alpha = dX^\alpha \quad (52)$$

which satisfy

$$\mathbf{g}^\alpha(\mathbf{g}_\beta) = \mathbf{g}^\alpha \cdot \mathbf{g}_\beta = \delta_\beta^\alpha \quad \mathbf{g}_\alpha \cdot \mathbf{g}_\beta = g_{\alpha\beta} \quad \mathbf{g}^\alpha \cdot \mathbf{g}^\beta = g^{\alpha\beta}. \quad (53)$$

Extending the vierbein formalism [32] to 5D we define the constant quintrad frames $\{\mathbf{e}_a\}$ for $\mathcal{T}(\mathcal{M}_5)$ and $\{\mathbf{e}^a\}$ for $\mathcal{T}^*(\mathcal{M}_5)$ which satisfy

$$\mathbf{e}_a \cdot \mathbf{e}_b = \eta_{ab} \quad \mathbf{e}^a \cdot \mathbf{e}^b = \eta^{ab} \quad \partial_a \mathbf{e}_b = \partial_a \mathbf{e}^b = 0. \quad (54)$$

By convention the Latin letters

$$a, b, c, \dots = 0, 1, 2, 3, 5 \quad \eta_{ab} = \text{diag}(-1, 1, 1, 1, \sigma) \quad (55)$$

indicate reference to the quintrad, where $\sigma = \eta_{55} = \eta^{55} = \pm 1$ provides a metric pseudo-signature on $\mathcal{M}_5$. The position-dependent coordinate bases are related to the quintrad through the vielbein field

$$\mathbf{g}_\alpha = E_\alpha^a(X) \mathbf{e}_a \quad \mathbf{g}^\alpha = \bar{E}^\alpha_a(X) \mathbf{e}^a \quad (56)$$

which is invertible as

$$\mathbf{e}_a = e_a^\alpha(X) \mathbf{g}_\alpha \quad \mathbf{e}^a = \bar{e}^a_\alpha(X) \mathbf{g}^\alpha. \quad (57)$$

Consistency requires that

$$e_a^\alpha(X) E_\beta^a(X) = \delta_\beta^\alpha \quad e_a^\alpha(X) E_\alpha^b(X) = \delta_a^b \quad (58)$$

with similar relations for $\bar{E}^\alpha_a$ and $\bar{e}^a_\alpha$. Duality imposes the conditions

$$\begin{aligned} \delta_{a'}^a &= \mathbf{e}^a \cdot \mathbf{e}_{a'} = \bar{e}^a_\alpha \mathbf{g}^\alpha \cdot e_{a'}^\beta \mathbf{g}_\beta = \bar{e}^a_\alpha e_{a'}^\beta \delta_\beta^\alpha = \bar{e}^a_\alpha e_{a'}^\alpha \\ \delta_{\alpha'}^\alpha &= \mathbf{g}^\alpha \cdot \mathbf{g}_{\alpha'} = \bar{E}^\alpha_a \mathbf{e}^a \cdot E_{\alpha'}^b \mathbf{e}_b = \bar{E}^\alpha_a E_{\alpha'}^b \delta_b^a = \bar{E}^\alpha_a E_{\alpha'}^a \end{aligned} \quad (59)$$

which together with (58) put the transformation equations into the form

$$\begin{aligned} \mathbf{g}_\alpha &= E_\alpha^a \mathbf{e}_a & \mathbf{e}_a &= e_a^\alpha \mathbf{g}_\alpha \\ \mathbf{g}^\alpha &= e_a^\alpha \mathbf{e}^a & \mathbf{e}^a &= \bar{e}^a_\alpha \mathbf{g}^\alpha. \end{aligned} \quad (60)$$

The vielbein field thus induces the local metric through

$$\begin{aligned} g_{\alpha\beta} &= \mathbf{g}_\alpha \cdot \mathbf{g}_\beta = \eta_{ab} E_\alpha^a E_\beta^b & g^{\alpha\beta} &= \mathbf{g}^\alpha \cdot \mathbf{g}^\beta = \eta^{ab} e_a^\alpha e_b^\beta \\ \eta_{ab} &= \mathbf{e}_a \cdot \mathbf{e}_b = g_{\alpha\beta} e_a^\alpha e_b^\beta & \eta^{ab} &= \mathbf{e}^a \cdot \mathbf{e}^b = g^{\alpha\beta} E_\alpha^a E_\beta^b \end{aligned} \quad (61)$$

where we use (56) and (57). Since any vector in $\mathcal{T}(\mathcal{M}_5)$ can be written

$$\mathbf{V} = V^\alpha \mathbf{g}_\alpha = [V^\alpha E_\alpha^a(X)] \mathbf{e}_a = V^a \mathbf{e}_a \quad (62)$$

we may view

$$X_b^a = E_\alpha^a e_a^\beta X_\beta^\alpha \quad X_\beta^\alpha = e_a^\alpha E_\beta^b X_b^a \quad (63)$$

as transformations between the coordinate frame and the vielbein frame.

The mapping of $\mathcal{M} \times R \longrightarrow \mathcal{M}_5$ in (21) provides a natural foliation of the tangent space into spacetime hyperspaces of equal- τ . Thus, we further partition the quintrad indices, as for the coordinate indices, so that the index convention becomes

$$\begin{aligned} \alpha, \beta, \gamma, \delta &= 0, 1, 2, 3, 5 & \lambda, \mu, \nu, \rho \dots &= 0, 1, 2, 3 \\ a, b, c, d, &= 0, 1, 2, 3, 5 & k, l, m, n, \dots &= 0, 1, 2, 3 \end{aligned} \quad (64)$$

where the five index with respect to the quintrad frame will be denoted $\bar{5}$ when necessary to avoid confusion. In this notation the frame transformations can be expanded as

$$\begin{aligned} \mathbf{g}_\mu &= E_\mu^k \mathbf{e}_k + E_\mu^{\bar{5}} \mathbf{e}_5 & \mathbf{e}_k &= e_k^\mu \mathbf{g}_\mu + e_k^{\bar{5}} \mathbf{g}_5 \\ \mathbf{g}_5 &= E_5^k \mathbf{e}_k + E_5^{\bar{5}} \mathbf{e}_5 & \mathbf{e}_5 &= e_5^\mu \mathbf{g}_\mu + e_5^{\bar{5}} \mathbf{g}_5 \end{aligned} \quad (65)$$

and

$$\begin{aligned} \mathbf{g}^\mu &= e_k^\mu \mathbf{e}^k + e_5^\mu \mathbf{e}^{\bar{5}} & \mathbf{e}^k &= E_\mu^k \mathbf{g}^\mu + E_5^k \mathbf{g}^{\bar{5}} \\ \mathbf{g}^5 &= e_k^{\bar{5}} \mathbf{e}^k + e_5^{\bar{5}} \mathbf{e}^{\bar{5}} & \mathbf{e}^{\bar{5}} &= E_\mu^{\bar{5}} \mathbf{g}^\mu + E_5^{\bar{5}} \mathbf{g}^{\bar{5}} \end{aligned} \quad (66)$$

where the orthogonality relations (58) are required for consistency. In the quintrad we expect the spacetime hypersurface to be spanned by $\{\mathbf{e}_k\}$ and normal to $\mathbf{e}_5$ which points in the direction of advancing τ . We therefore denote the unit normal to spacetime and its associated 1-form as

$$n = \mathbf{e}_5 \quad \bar{n} = \sigma \mathbf{e}^{\bar{5}} \quad (67)$$

with normalizations

$$n^2 = \mathbf{e}_5 \cdot \mathbf{e}_5 = \sigma \quad \bar{n}^2 = \mathbf{e}^{\bar{5}} \cdot \mathbf{e}^{\bar{5}} = \sigma. \quad (68)$$

Introducing the ADM parameterization [19] we express the fifth basis vector for $\mathcal{T}(\mathcal{M}_5)$ as a linear combination of n and $\{\mathbf{g}_\mu\}$

$$\mathbf{g}_5 = N^\mu \mathbf{g}_\mu + Nn \quad (69)$$

where the 4-vector N^μ generalizes the shift 3-vector in 3+1 formalisms and N is the lapse function with respect to τ . Designating the spacetime part of the 5D coordinate frame metric as

$$\gamma_{\mu\nu} = g_{\mu\nu} = \mathbf{g}_\mu \cdot \mathbf{g}_\nu \quad (70)$$

we can express the 5-components of the coordinate metric as

$$\begin{aligned} g_{5\mu} &= \mathbf{g}_\mu \cdot \mathbf{g}_5 = \gamma_{\mu\mu'} N^{\mu'} = N_\mu \\ g_{55} &= (N^\mu \mathbf{g}_\mu + Nn) \cdot (N^{\mu'} \mathbf{g}_{\mu'} + Nn) = \gamma_{\mu\mu'} N^\mu N^{\mu'} + \sigma N^2 \end{aligned} \quad (71)$$

so that

$$g_{\alpha\beta} = \begin{bmatrix} \gamma_{\mu\nu} & N_\mu \\ N_\mu & \sigma N^2 + \gamma_{\mu\nu} N^\mu N^\nu \end{bmatrix} \quad (72)$$

with

$$g^{\alpha\beta} = \begin{bmatrix} \gamma^{\mu\nu} + \sigma \frac{1}{N^2} N^\mu N^\nu & -\sigma \frac{1}{N^2} N^\mu \\ -\sigma \frac{1}{N^2} N^\mu & \sigma \frac{1}{N^2} \end{bmatrix} \quad (73)$$

as its inverse.

With these choices we have

$$\mathbf{e}_5 = n = \frac{1}{N} (-N^\mu \mathbf{g}_\mu + \mathbf{g}_5) \quad \mathbf{e}^5 = \sigma \bar{n} = \sigma^2 N \mathbf{g}^5 = N \mathbf{g}^5 \quad (74)$$

so that comparison with (65) and (66) determines four vielbein components

$$e^\mu_{\bar{5}} = -\frac{1}{N} N^\mu \quad e^5_{\bar{5}} = \frac{1}{N} \quad E_\mu^{\bar{5}} = 0 \quad E_5^{\bar{5}} = N. \quad (75)$$

Orthogonality in the quintrad frame provides two additional conditions

$$\begin{aligned} 0 &= \mathbf{e}_k \cdot \mathbf{e}_5 = (e^\mu_k \mathbf{g}_\mu + e^5_k \mathbf{g}_5) \cdot \frac{1}{N} (-N^{\mu'} \mathbf{g}_{\mu'} + \mathbf{g}_5) \\ &= \frac{1}{N} (-e^\mu_k N_\mu + e^\mu_k g_{\mu 5} - e^5_k g_{\mu' 5} N^{\mu'} + e^5_k g_{55}) \end{aligned} \quad (76)$$

and

$$0 = \mathbf{e}^k \cdot \mathbf{e}^5 = (E_\mu^k \mathbf{g}^\mu + E_5^k \mathbf{g}^5) \cdot N \mathbf{g}^5 = N (E_\mu^k g^{\mu 5} + E_5^k g^{55}), \quad (77)$$

which combined with (71) provide the components

$$e^5_k = 0 \quad E_5^k = E_\mu^k N^\mu. \quad (78)$$

Inserting $e^5_k = 0$ into (65) leads to

$$\mathbf{e}_k = e^\mu_k \mathbf{g}_\mu + e^5_k \mathbf{g}_5 = e^\mu_k (E_\mu^{k'} \mathbf{e}_{k'} + E_\mu^5 \mathbf{e}_5) \quad (79)$$

from which we conclude that

$$E_\mu^{\bar{5}} = 0 \quad e^\mu_k E_\mu^{k'} = \delta_k^{k'}. \quad (80)$$

The transformations between coordinate and quintrad frames take the final form

$$\begin{aligned} \mathbf{g}_\alpha &= E_\alpha^a \mathbf{e}_a = \delta_\alpha^\mu E_\mu^k \mathbf{e}_k + \delta_\alpha^5 (E_\mu^k N^\mu \mathbf{e}_k + N \mathbf{e}_5) \\ \mathbf{e}_a &= e^\alpha_a \mathbf{g}_\alpha = \delta_a^k e^\mu_k \mathbf{g}_\mu + \delta_a^5 \frac{1}{N} (-N^\mu \mathbf{g}_\mu + \mathbf{g}_5) \end{aligned} \quad (81)$$

providing a quintrad basis for $\mathcal{T}(\mathcal{M}_5)$ with a 4+1 foliation built-in by construction. The vielbein field may be summarized as

$$\begin{aligned} E_\alpha^a &= \delta_\alpha^\mu \delta_k^a E_\mu^k + \delta_\alpha^5 (E_\mu^k N^\mu \delta_k^a + N \delta_5^a) \\ e^\alpha_a &= \delta_a^k \delta_\mu^\alpha e^\mu_k - \delta_a^5 \delta_\mu^\alpha \frac{1}{N} N^\mu + \delta_a^5 \delta_5^\alpha \frac{1}{N} \end{aligned} \quad (82)$$

and since $e^\mu_k = E^\mu_k$ by orthogonality, the vielbein is constructed from E^μ_k , N and N^μ . As shown in Sections 4 and 5, the τ -dependent lapse N and shift N^μ , play the role of Lagrange multipliers, enforcing the constraints on $g_{\alpha\beta}$ associated with the Bianchi relations, but they are not subject to second order

evolution equations. Thus, the dynamical content of the vielbein field is entirely contained in the spacetime part E_μ^k , where $\mu, k = 0, 1, 2, 3$.

3.2. Field Equations

The standard Einstein field equations

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \frac{8\pi G}{c^4}T_{\mu\nu} \quad (83)$$

enjoy $O(3,1)$ covariance, general diffeomorphism invariance, and the translation invariance of the Ricci tensor that guarantees the Bianchi identity as a 4D gauge symmetry. It follows from the Bianchi identity that of the ten components of (83) six represent dynamical degrees of freedom and four represent constraints [31]. This distinction is made explicit in 3+1 ADM formalism which decomposes the field equations into a pair of coupled t -evolution equations for the geometry of 3D space, along with 3-vector and scalar constraints among the initial conditions. As articulated by Wheeler [33], “A decade and more of work by Dirac, Bergmann, Schild, Pirani, Anderson, Higgs, Arnowitt, Deser, Misner, DeWitt, and others has taught us through many a hard knock that Einstein’s geometrodynamics deals with the dynamics of geometry: of 3-geometry, not 4-geometry.”

The 3+1 ADM decomposition is most conveniently performed using the trace-reversed form of (83) found from

$$g^{\mu\nu}R_{\mu\nu} - \frac{1}{2}g^{\mu\nu}g_{\mu\nu}R = \frac{8\pi G}{c^4}g^{\mu\nu}T_{\mu\nu} \quad (84)$$

and using $g^{\mu\nu}g_{\mu\nu} = 4$ to obtain

$$R_{\mu\nu} = \frac{8\pi G}{c^4} \left(T_{\mu\nu} - \frac{1}{2}g_{\mu\nu}T \right) \quad (85)$$

which expresses the relationship between spacetime geometry on the LHS and the distribution of matter on the RHS. But in 5D we have $g^{\alpha\beta}g_{\alpha\beta} = 5 \neq 4$ leading to the trace-reversed form

$$R_{\alpha\beta} = \frac{8\pi G}{c^4} \left(T_{\alpha\beta} + \frac{\frac{1}{2}g_{\alpha\beta}}{1 - \frac{1}{2}g^{\gamma\delta}g_{\gamma\delta}} g^{\gamma'\delta'} T_{\gamma'\delta'} \right) = \frac{8\pi G}{c^4} \left(T_{\alpha\beta} - \frac{1}{3}g_{\alpha\beta}T \right). \quad (86)$$

As we shall see in Section 6, this form cannot be made consistent with standard phenomenology in weak SHP gravitation, where the metric may be found as an additive perturbation $h_{\alpha\beta}$ to the flat space metric $\eta_{\alpha\beta}$. Correct perturbative solutions require the replacement $\eta_{\alpha\beta} \rightarrow \hat{\eta}_{\alpha\beta} = (-1, 1, 1, 1, 0)$ in the trace-reversed matter terms, which breaks the apparent 5D symmetry of the source to $O(3,1)$. For the full field equations we expect a similar substitution $g_{\alpha\beta} \rightarrow \hat{g}_{\alpha\beta}$ on the RHS of (86) where the modified metric remains to be determined. By requiring $\hat{g}^{\alpha\beta}\hat{g}_{\alpha\beta} = 4$ we recover the form

$$R_{\alpha\beta} = \frac{8\pi G}{c^4} \left(T_{\alpha\beta} - \frac{1}{2}\hat{g}_{\alpha\beta}\hat{T} \right) \quad (87)$$

where $\hat{T} = \hat{g}^{\alpha\beta}T_{\alpha\beta}$.

To find an expression for $\hat{g}_{\alpha\beta}$ we write unbroken 5D Einstein equations in the quintrad frame

$$R_{ab} - \frac{1}{2}\eta_{ab}R = \frac{8\pi G}{c^4}T_{ab} \rightarrow R_{ab} = \frac{8\pi G}{c^4} \left(T_{ab} + \frac{\frac{1}{2}\eta_{ab}}{1 - \frac{1}{2}\eta^{cd}\eta_{cd}} \eta^{c'd'} T_{c'd'} \right) \quad (88)$$

where the metric η_{ab} is flat, while R_{ab} and T_{ab} are related to the coordinate frame tensors through the vielbein field. We break the 5D symmetry in the matter terms by replacing

$$\eta_{ab} \rightarrow \hat{\eta}_{ab} = (-1, 1, 1, 1, 0) = \delta_a^k \delta_b^l \eta_{kl} \quad (89)$$

on the RHS of (88), leaving the Ricci tensor R_{ab} unchanged. The gravitational Einstein equations in the SHP formalism now take the form

$$R_{ab} = \frac{8\pi G}{c^4} \left(T_{ab} - \frac{1}{2} \hat{\eta}_{ab} \hat{T} \right) \quad (90)$$

where $\hat{T} = \hat{\eta}^{ab} T_{ab} = \eta^{kl} T_{kl}$. Using the vielbein field (82) we may transform (90) back to a coordinate frame, leading to

$$R_{\alpha\beta} = \frac{8\pi G}{c^4} \left(T_{\alpha\beta} - \frac{1}{2} \hat{g}_{\alpha\beta} \hat{T} \right) \quad (91)$$

providing the symmetry-broken local metric as

$$\hat{g}_{\alpha\beta} = E_{\alpha}^a E_{\beta}^b \hat{\eta}_{ab} = g_{\alpha\beta} - \delta_{\alpha}^5 \delta_{\beta}^5 \eta_{55} N^2 = g_{\alpha\beta} - \sigma n_{\alpha} n_{\beta} = P_{\alpha\beta}. \quad (92)$$

We notice that $P_{\alpha\beta}$ is a projection operator onto the spacetime hypersurface in $\mathcal{T}(\mathcal{M}_5)$, and satisfies

$$P_{\alpha\beta} n^{\beta} = 0 \quad P_{\alpha\beta} P^{\beta\gamma} = P_{\alpha\beta} (g^{\beta\gamma} - \sigma n^{\beta} n^{\gamma}) = g_{\alpha}^{\gamma} - \sigma n_{\alpha} n^{\gamma} = P_{\alpha}^{\gamma} \quad (93)$$

from which we may write the completeness relation

$$\delta_{\alpha}^{\beta} = g_{\alpha}^{\beta} = P_{\alpha}^{\beta} + \sigma n_{\alpha} n^{\beta}. \quad (94)$$

The breaking of 5D symmetry can thus be understood as replacing the metric in the matter terms of the field equation with the projector onto the 4D hypersurface.

Using the completeness relation (94), we may decompose the mass-energy-momentum tensor into

$$T_{\alpha\beta} = T_{\alpha'\beta'} (P_{\alpha}^{\alpha'} + \sigma n^{\alpha\alpha'} n_{\alpha}) (P_{\beta}^{\beta'} + \sigma n^{\beta\beta'} n_{\beta}) = S_{\alpha\beta} - 2\sigma n_{\alpha} p_{\beta} + n_{\alpha} n_{\beta} \kappa \quad (95)$$

where

$$S_{\alpha\beta} = P_{\alpha}^{\alpha'} P_{\beta}^{\beta'} T_{\alpha'\beta'} \quad p_{\beta} = -n^{\alpha} P_{\beta}^{\beta'} T_{\alpha\beta'} \quad \kappa = n^{\beta} n^{\alpha} T_{\alpha\beta} \quad (96)$$

representing the 4D energy momentum tensor $S_{\alpha\beta}$ found by projecting $T_{\alpha\beta}$ onto the spacetime hypersurface, a 4-momentum vector p_{β} describing the flow of mass into spacetime, and an invariant form κ of the scalar mass density T_{55} . The trace of $T_{\alpha\beta}$ is

$$T = \eta^{\alpha\beta} T_{\alpha\beta} = S + \sigma \kappa \quad (97)$$

but the symmetry broken trace is

$$\hat{T} = \hat{g}^{\alpha\beta} T_{\alpha\beta} = P^{\alpha\beta} T_{\alpha\beta} = P^{\alpha\beta} (S_{\alpha\beta} + 2\sigma n_{\alpha} p_{\beta} + n_{\alpha} n_{\beta} \kappa) = S \quad (98)$$

where we used $P^{\alpha\beta} n_{\alpha} = 0$. The SHP field equations in a coordinate frame now take the form

$$R_{\alpha\beta} = \frac{8\pi G}{c^4} \left(T_{\alpha\beta} - \frac{1}{2} P_{\alpha\beta} S \right). \quad (99)$$

4. Decomposition, Projection, and Initial Value Problem

As we will see in this section, the fifteen degrees of freedom in the symmetric 5D Einstein tensor

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2} g_{\alpha\beta} R \quad (100)$$

can be decomposed into ten components $G_{\mu\nu}$ satisfying unconstrained second order differential equations, and five components $G_{5\mu}$ that impose τ -dependent constraints satisfying algebraic equations. The differential equations require as initial conditions $\gamma_{\mu\nu}(x, \tau_0)$ and $\partial_{\tau} \gamma_{\mu\nu}(x, \tau_0)$ at some τ_0 .

A plausible argument for this decomposition can be seen [34] by writing the Bianchi relations as

$$\nabla_\alpha G^{\alpha\beta} = \partial_\alpha G^{\alpha\beta} + \text{Christoffel Symbols} \times G^{\alpha\beta} = 0 \quad (101)$$

forming a set of relations among the field entities. Expanding $\partial_\alpha G^{\alpha\beta} = \partial_\mu G^{\mu\beta} + \partial_5 G^{5\beta}$ to rewrite the Bianchi relations as

$$\frac{1}{c_5} \partial_\tau G^{5\beta} = -\partial_\mu G^{\mu\beta} - \text{Christoffel Symbols} \times G^{\alpha\beta} \quad (102)$$

we see that since the LHS cannot be more than second order in ∂_τ , we must conclude that $G^{5\beta}$ cannot be more than first order in ∂_τ . Now, if $G^{\alpha\beta}(x, \tau_0) = 0$ then $\partial_\mu G^{\alpha\beta}(x, \tau_0) = 0$ which leads to

$$\frac{1}{c_5} \partial_\tau G^{5\beta}(x, \tau_0) = -\partial_\mu G^{\mu\beta}(x, \tau_0) - \text{Christoffel Symbols} \times G^{\alpha\beta}(x, \tau_0) = 0 \quad (103)$$

and so any terms contained in $G^{5\beta}$ must be constraints on the initial conditions that propagate to future times with τ , but whose form does not evolve.

The formal decomposition of the field equations is accomplished by way of projection of the relevant geometric structures defined on $\mathcal{M}_5$ onto $\mathcal{M}(\tau)$. Among these are intrinsic curvature, which appears in the parallel transport of a vector tangent to the surface, and extrinsic curvature which appears in the parallel transport of a vector normal to the surface. As an intermediate step we define the scalar field $S(X) = \tau$ whose level surfaces provide the natural foliation of pseudo-spacetime

$$\Sigma(\tau_0) \subset \mathcal{M}_5 = \left\{ X \in \mathcal{M}_5 \mid S(X) = X^5/c_5 = \tau_0 \right\} \quad (104)$$

implied by the construction of $\mathcal{M}_5$ in (21). Since $\Sigma(\tau_0)$ is homeomorphic to $\mathcal{M}(\tau_0)$ for any τ_0 , we may drop reference to τ_0 . Thus, for any vector $V = V^\alpha g_\alpha \in \mathcal{T}(\mathcal{M}_5)$, where $g_\alpha = \partial_\alpha$ is the coordinate basis given in (51), the projection operator (92) provides the vector $V_\perp^\alpha = P_\beta^\alpha V^\beta \in \mathcal{T}(\Sigma)$. The homeomorphism is expressed by way of the vector $v \in \mathcal{T}(\mathcal{M})$ for which

$$V_\perp^\alpha = v^\mu (g_\mu)^\alpha = v^\mu \frac{\partial X^\alpha}{\partial x^\mu} = v^\mu \delta_\mu^\alpha \quad (105)$$

and so

$$V_\perp^5 = 0 \longrightarrow v^\mu = \delta_\alpha^\mu V_\perp^\alpha = \delta_\alpha^\mu P_\beta^\alpha V^\beta = P_\beta^\mu V^\beta. \quad (106)$$

To express the projection operator (92) we transform the unit normal $n = e_5$ defined in (67) for the vielbein frame to a coordinate frame as

$$n_\alpha = \left[\delta_\alpha^\mu \delta_k^a E_\mu^k + \delta_\alpha^5 \left(E_\mu^k N^\mu \delta_k^a + N \delta_5^a \right) \right] \eta_{a5} = \sigma N \delta_\alpha^5 \quad (107)$$

where we used $n_\alpha = n \cdot e_\alpha = e_5 \cdot e_\alpha = \eta_{\alpha 5}$. Similarly,

$$n^\alpha = g^{\alpha\beta} n_\beta = g^{\alpha\beta} \eta_{\beta 5} = \frac{1}{N} \left(-N^\mu g_\mu^\alpha + g_5^\alpha \right) = \frac{1}{N} \left(-N^\mu \delta_\mu^\alpha + \delta_5^\alpha \right). \quad (108)$$

The extrinsic curvature $K_{\mu\nu}$ will be defined as the projection of the covariant derivative of the unit normal onto $\mathcal{M}(\tau)$. We will see that $K_{\mu\nu}$ is closely related to $\partial_\tau \gamma_{\mu\nu}$.

In the Legendre transformation from Lagrangian to Hamiltonian mechanics, the canonical momentum is introduced as an auxiliary variable, so that the phase space $(x, \dot{x})$ becomes (x, p) containing no explicit τ -derivatives. This transforms the equations of motion from one expression involving the second order derivatives $\ddot{x}^\mu$ to a pair of expressions involving only the first order derivatives $\dot{x}^\mu$ and $\dot{p}^\mu$. By analogy, the ADM approach to geometrodynamics introduces the extrinsic curvature $K_{\mu\nu}$ as an auxiliary variable, transforming the second order field equations (99) in $g_{\alpha\beta}$ into a pair of first order differential equations for $\gamma_{\mu\nu}$ and $K_{\mu\nu}$ in spacetime. Using standard methods from the theory of embedded surfaces, the initial value problem is posed in two stages. First, we study the

evolution of the spacetime hypersurface through the Lie derivative in the direction of the unit normal to Σ , leading to an expression relating $\partial_\tau \gamma_{\mu\nu}$ to $K_{\mu\nu}$. We then decompose the Ricci tensor $R_{\alpha\beta}$ into components tangent and normal to Σ , providing an expression relating $\partial_\tau K_{\mu\nu}$ to the matter terms in the field equations and a pair of constraints on the initial conditions.

4.1. Projected Covariant Derivative and Curvature

With the compatible connection $\Gamma_{\alpha\beta}^\gamma$ the covariant derivative of $g_{\alpha\beta}$ on $\mathcal{M}_5$ vanishes, leading to the Ricci identity

$$[\nabla_\beta, \nabla_\alpha] X_\delta = X_\gamma R_{\delta\alpha\beta}^\gamma \quad (109)$$

with Riemann tensor

$$R_{\delta\alpha\beta}^\gamma = \frac{\partial}{\partial x^\alpha} \Gamma_{\delta\beta}^\gamma - \frac{\partial}{\partial x^\beta} \Gamma_{\delta\alpha}^\gamma + \Gamma_{\sigma\alpha}^\gamma \Gamma_{\delta\beta}^\sigma - \Gamma_{\sigma\beta}^\gamma \Gamma_{\delta\alpha}^\sigma \quad (110)$$

and associated Bianchi relations. The projected covariant derivative is defined as

$$\bar{\nabla}_\alpha V_\beta = P_\alpha^{\alpha'} P_\beta^{\beta'} \nabla_{\alpha'} V_{\beta'} \quad (111)$$

and since

$$\nabla_\alpha P_{\beta\gamma} = \nabla_\alpha (g_{\beta\gamma} - \sigma n_\beta n_\gamma) = -\sigma \nabla_\alpha (n_\beta n_\gamma) = -\sigma [(\nabla_\alpha n_\beta) n_\gamma + n_\beta \nabla_\alpha n_\gamma] \quad (112)$$

we have the compatibility condition

$$\bar{\nabla}_\alpha P_{\beta\gamma} = -\sigma P_\alpha^{\alpha'} P_\beta^{\beta'} P_\gamma^{\gamma'} [(\nabla_{\alpha'} n_{\beta'}) n_{\gamma'} + n_{\beta'} \nabla_{\alpha'} n_{\gamma'}] = 0 \quad (113)$$

where we used $P_\beta^\delta n_\delta = 0$. We notice that for a vector $V = V^\perp \in \mathcal{T}(\Sigma)$

$$\bar{\nabla}_\alpha V_\beta^\perp = \bar{\nabla}_\alpha (P_\beta^{\beta'} V_{\beta'}) = P_\beta^{\beta'} \bar{\nabla}_\alpha (V_{\beta'}) = P_\beta^{\beta'} (P_\alpha^{\alpha'} \nabla_{\alpha'} V_{\beta'}) = \bar{\nabla}_\alpha V_\beta \quad (114)$$

by comparison with (111). This compatibility justifies regarding $\bar{\nabla}_\alpha$ as the intrinsic covariant derivative on $\mathcal{T}(\Sigma)$, denoted as

$$D_\alpha = \bar{\nabla}_\alpha = P_\alpha^\gamma \nabla_\gamma \quad D_\mu = P_\mu^\alpha D_\alpha = P_\mu^\alpha P_\alpha^\gamma \nabla_\gamma = P_\mu^\gamma \nabla_\gamma \quad (115)$$

where we used (106), leading to $D_\mu \gamma_{\lambda\rho} = 0$. The projected curvature $\bar{R}_{\lambda\mu\nu}^\rho$ is defined through the projected covariant derivatives

$$[D_\nu, D_\mu] X_\lambda = X_\rho \bar{R}_{\lambda\mu\nu}^\rho \quad (116)$$

and will be given an explicit form below.

For vectors $V, U \in \mathcal{T}(\Sigma) \subset \mathcal{T}(\mathcal{M})$, the Weingarten map is defined as

$$\chi(V) = \nabla_V n = V \cdot (\nabla n) \quad \chi^\alpha(V) = V^\beta \nabla_\beta n^\alpha \quad (117)$$

and the extrinsic curvature on $\mathcal{T}(\Sigma)$ is defined as the projection onto a vector U of the Weingarten map along a vector V

$$K(U, V) = -U \cdot \chi(V) = -U \cdot \nabla_V n = -g_{\alpha\gamma} V^\alpha U^\beta \nabla_\beta n^\gamma \quad (118)$$

$$K_{\alpha\beta} = -g_{\alpha\gamma} \nabla_\beta n^\gamma = -\nabla_\beta n_\alpha. \quad (119)$$

This definition is extended to $\mathcal{T}(\mathcal{M})$ as

$$K(U^\perp, V^\perp) = K(PU, PV) = -g_{\gamma'\gamma} (P_{\alpha'}^{\gamma'} V^\alpha) (P_{\beta'}^\beta U^{\beta'}) \nabla_\beta n^\gamma \quad (120)$$

$$K_{\beta'\alpha} U^{\beta'} V^\alpha = V^\alpha U^{\beta'} (-g_{\gamma'\gamma} P_{\alpha'}^{\gamma'} P_{\beta'}^\beta) \nabla_\beta n^\gamma \quad (121)$$

$$K_{\alpha\beta} = -P_{\alpha'}^{\alpha'} P_{\beta'}^{\beta'} \nabla_{\beta'} n_{\alpha'} \quad (122)$$

where although n_γ is normal to $\mathcal{T}(\Sigma)$ the covariant derivative $\nabla_\delta n_\gamma$ may have both normal and tangent components. Using $n^2 = \sigma$ we have

$$0 = \nabla_\alpha n^2 = 2n^\beta (\nabla_\alpha n_\beta) \quad (123)$$

so that

$$P_{\alpha'}^{\alpha'} \nabla_\beta n_{\alpha'} = (g_{\alpha'}^{\alpha'} - \sigma n_\alpha n^{\alpha'}) \nabla_\beta n_{\alpha'} = \nabla_\beta n_\alpha \quad (124)$$

and so (122) becomes

$$K_{\alpha\beta} = -P_{\alpha'}^{\alpha'} \nabla_{\alpha'} n_\beta = -\nabla_\alpha n_\beta + \sigma n_\alpha (n^{\alpha'} \nabla_{\alpha'} n_\beta) \quad (125)$$

with the contracted form

$$K = \gamma^{\alpha\beta} K_{\alpha\beta} = \gamma^{\alpha\beta} P_{\alpha'}^{\alpha'} P_{\beta'}^{\beta'} \nabla_{\alpha'} n_{\beta'} = \gamma^{\alpha\beta} \nabla_\alpha n_\beta = \nabla_\alpha n^\alpha. \quad (126)$$

Using (107) for the unit normal n_α we expand

$$\begin{aligned} (n^\gamma \nabla_\gamma n_\beta) &= \sigma n^\gamma \nabla_\gamma (N \nabla_\beta \tau) = \sigma n^\gamma (\nabla_\gamma N) \frac{n_\beta}{\sigma N} + \sigma n^\gamma N \nabla_\beta \left(\frac{n_\gamma}{\sigma N} \right) \\ &= \frac{1}{N} [n^\gamma n_\beta \nabla_\gamma N - \sigma \delta_\beta^\gamma \nabla_\gamma N] = -\sigma \frac{1}{N} [\delta_\beta^\gamma - \sigma n^\gamma n_\beta] \nabla_\gamma N \\ &= -\sigma \frac{1}{N} P_\beta^\gamma \nabla_\gamma N = -\sigma \frac{1}{N} D_\beta N \end{aligned} \quad (127)$$

to obtain the extrinsic curvature (125) in the useful form

$$K_{\alpha\beta} = -\nabla_\alpha n_\beta - n_\alpha \frac{1}{N} D_\beta N. \quad (128)$$

4.2. Evolution of the Hypersurface Σ

With the parameterization (69) the fifth basis vector is

$$\partial_5 = N^\mu \mathbf{g}_\mu + Nn = \mathbf{N} + Nn \quad (129)$$

and we define the normal evolution vector $m = Nn$, noting that for the time function $S(X)$ which defines the foliation (104), we have

$$\mathbf{d}S(X) \cdot m = \delta_\alpha^5 (Nn)^\alpha = \delta_\alpha^5 N (g^{\alpha\beta} \sigma N \delta_\beta^5) = N^2 \sigma g^{55} = N^2 \sigma \left(\sigma \frac{1}{N^2} \right) = 1. \quad (130)$$

Under a displacement

$$X \longrightarrow X' = X + m \delta\tau(X) \quad (131)$$

the time function evolves as

$$S(X) \longrightarrow S(X') = S(X + m \delta\tau) = S(X) + \mathbf{d}S(X) \cdot m \delta\tau = S(X) + \delta\tau \quad (132)$$

showing that for $X \in \Sigma_\tau$

$$X' = X + m \delta S(X) \in \Sigma_{\tau+\delta\tau} \quad (133)$$

so that the hypersurface $\Sigma_{\tau+\delta\tau}$ is obtained from Σ_τ by the displacement of each point by $m \delta S$. The evolution of the hypersurface Σ can thus be characterized through the Lie derivative in the direction of the normal evolution vector

$$\mathcal{L}_m = \mathcal{L}_5 - \mathcal{L}_N. \quad (134)$$

The Lie derivative of a second rank tensor $A_{\alpha\beta}$ is

$$\mathcal{L}_m A_{\alpha\beta} = m^\gamma \nabla_\gamma A_{\alpha\beta} + A_{\gamma\beta} \nabla_\alpha m^\gamma + A_{\alpha\gamma} \nabla_\beta m^\gamma \quad (135)$$

where in the absence of torsion ∇_γ can be replaced by ∂_γ and we have

$$\mathcal{L}_m \delta^\alpha_\beta = -\nabla_\beta m^\alpha + \nabla_\beta m^\alpha = 0 \quad \mathcal{L}_m g_{\alpha\beta} = \nabla_\alpha m_\beta + \nabla_\beta m_\alpha \quad (136)$$

$$\mathcal{L}_5 A_{\alpha\beta} = \delta_5^\gamma \partial_\gamma A_{\alpha\beta} + A_{\gamma\beta} \partial_\alpha \delta_5^\gamma + A_{\alpha\gamma} \partial_\beta \delta_5^\gamma = \partial_5 A_{\alpha\beta} = \frac{1}{c_5} \partial_\tau A_{\alpha\beta}. \quad (137)$$

For $X \in \Sigma \subset \mathcal{M}_5$, the squared interval is

$$\delta X^2 \Big|_\Sigma = g_{\alpha\beta} \delta X^\alpha \delta X^\beta \Big|_\Sigma = g_{\alpha\beta} \frac{\partial X^\alpha}{\partial x^\mu} \frac{\partial X^\beta}{\partial x^\nu} \delta x^\mu \delta x^\nu = g_{\alpha\beta} \delta^\alpha_\mu \delta^\beta_\nu \delta x^\mu \delta x^\nu = \gamma_{\mu\nu} \delta x^\mu \delta x^\nu \quad (138)$$

and since

$$P_{\mu\nu} = g_{\mu\nu} - \sigma n_\mu n_\nu = \gamma_{\mu\nu} \quad (139)$$

we may find the Lie derivative of the metric $\gamma_{\mu\nu}$ on $\mathcal{M}$ by evaluating $\mathcal{L}_m P_{\alpha\beta}$ and pulling back the result to $\mathcal{M}$. Writing

$$\mathcal{L}_m P_{\alpha\beta} = m^\gamma \nabla_\gamma P_{\alpha\beta} + P_{\gamma\beta} \nabla_\alpha m^\gamma + P_{\alpha\gamma} \nabla_\beta m^\gamma \quad (140)$$

we find

$$m^\gamma \nabla_\gamma P_{\alpha\beta} = N n^\gamma \nabla_\gamma (g_{\alpha\beta} - \sigma n_\alpha n_\beta) = -\sigma N (n^\gamma (\nabla_\gamma n_\alpha) n_\beta + n_\alpha n^\gamma \nabla_\gamma n_\beta) \quad (141)$$

and using (128) to obtain

$$\nabla_\beta m_\alpha = N \nabla_\beta n_\alpha + n_\alpha \nabla_\beta N = -N K_{\beta\alpha} - n_\beta \nabla_\alpha N + n_\alpha \nabla_\beta N \quad (142)$$

we finally arrive at

$$\mathcal{L}_m \gamma_{\mu\nu} = \mathcal{L}_5 \gamma_{\mu\nu} - \mathcal{L}_N \gamma_{\mu\nu} \quad (143)$$

from which we can write

$$\frac{1}{c_5} \partial_\tau \gamma_{\mu\nu} = \mathcal{L}_N \gamma_{\mu\nu} - 2N K_{\mu\nu} \quad (144)$$

as the evolution equation for the metric.

4.3. Decomposition of the Riemann Tensor

The 4+1 decomposition of $R^\gamma_{\delta\alpha\beta}$ is accomplished by projecting onto Σ and n . Using the completeness relation (94) to write

$$R^\gamma_{\delta\alpha\beta} = (P^\alpha_{\alpha'} + \sigma n_\alpha n^{\alpha'}) (P^\beta_{\beta'} + \sigma n_\beta n^{\beta'}) (P^\gamma_{\gamma'} + \sigma n_\gamma n^{\gamma'}) (P^{\delta'}_{\delta} + \sigma n_\delta n^{\delta'}) R^{\gamma'}_{\delta'\alpha'\beta'} \quad (145)$$

we obtain products of the type

$$R^\gamma_{\delta\alpha\beta} = \delta^\alpha_{\alpha'} \delta^\beta_{\beta'} \delta^\gamma_{\gamma'} \delta^{\delta'}_{\delta} R^{\gamma'}_{\delta'\alpha'\beta'} \longrightarrow \begin{cases} \delta^\alpha_\mu \delta^\beta_\nu \delta^\gamma_\lambda \delta^{\delta'}_\sigma P^\alpha_{\alpha'} P^\beta_{\beta'} P^\gamma_{\gamma'} P^{\delta'}_{\delta} R^{\gamma'}_{\delta'\alpha'\beta'} = R^\lambda_{\sigma\mu\nu} \\ \delta^\alpha_\mu \delta^\beta_\nu \delta^\gamma_\lambda P^\gamma_{\gamma'} n^\delta P^\alpha_{\alpha'} P^\beta_{\beta'} R^{\gamma'}_{\delta'\alpha'\beta'} = \sigma N R^\lambda_{5\mu\nu} \\ \delta^\alpha_\mu \delta^\beta_\nu P^\alpha_{\alpha'} n^\delta P^\beta_{\beta'} n^\gamma R^{\gamma'}_{\delta'\alpha'\beta'} = N^2 R^\mu_{5\nu 5} \end{cases} \quad (146)$$

where the symmetries of the Riemann tensor lead to $R_{\delta\alpha\beta}^{\gamma} n^{\delta} n^{\alpha} n^{\beta} = 0$. To expand the projected curvature defined in (116), we write

$$D_{\alpha} D_{\beta} V^{\gamma} = D_{\alpha} (D_{\beta} V^{\gamma}) = P_{\alpha}^{\alpha'} P_{\beta}^{\beta'} P_{\gamma}^{\gamma'} \nabla_{\alpha'} (D_{\beta'} V^{\gamma'}) \quad (147)$$

and use (112) to find

$$D_{\alpha} D_{\beta} V^{\gamma} = \sigma K_{\alpha\beta} P_{\gamma}^{\gamma'} n^{\beta'} \nabla_{\beta'} V^{\gamma'} + \sigma K_{\alpha}^{\gamma} K_{\beta\delta} V^{\delta} + P_{\alpha}^{\alpha'} P_{\beta}^{\beta'} P_{\gamma}^{\gamma'} (\nabla_{\alpha'} \nabla_{\beta'} V^{\gamma'}) . \quad (148)$$

Thus,

$$[D_{\alpha}, D_{\beta}] V^{\gamma} = \bar{R}_{\delta\alpha\beta}^{\gamma} V^{\delta} = -\sigma (K_{\alpha\delta} K_{\beta}^{\gamma} - K_{\beta\delta} K_{\alpha}^{\gamma}) V^{\delta} + P_{\alpha}^{\alpha'} P_{\beta}^{\beta'} P_{\gamma}^{\gamma'} R_{\delta'\alpha'\beta'}^{\gamma'} P_{\delta}^{\delta'} V^{\delta} \quad (149)$$

which by the quotient theorem on Σ leads to

$$P_{\alpha}^{\alpha'} P_{\beta}^{\beta'} P_{\gamma}^{\gamma'} P_{\delta}^{\delta'} R_{\delta'\alpha'\beta'}^{\gamma'} = \bar{R}_{\delta\alpha\beta}^{\gamma} - \sigma (K_{\alpha}^{\gamma} K_{\beta\delta} - K_{\beta}^{\gamma} K_{\alpha\delta}) \quad (150)$$

generalizing the Gauss relation. Acting on this expression with $\delta_{\gamma}^{\mu} \delta_{\nu}^{\delta} \delta_{\lambda}^{\alpha} \delta_{\rho}^{\beta}$ we find

$$R_{\nu\lambda\rho}^{\mu} = \bar{R}_{\nu\lambda\rho}^{\mu} - \sigma (K_{\lambda}^{\mu} K_{\rho\nu} - K_{\rho}^{\mu} K_{\lambda\nu}) \quad (151)$$

providing an expression for the projected curvature $\bar{R}_{\nu\lambda\rho}^{\mu}$ defined in (116) in terms of the spacetime components of the 5D intrinsic curvature $R_{\nu\lambda\rho}^{\mu}$ and the extrinsic curvature $K_{\rho\nu}$. Contracting on α and γ in (150) leads to

$$P_{\alpha}^{\alpha'} P_{\beta}^{\beta'} R_{\alpha'\beta'} - \sigma P_{\alpha\alpha'} n^{\delta'} P_{\beta}^{\beta'} n^{\gamma'} R_{\delta'\beta'\gamma'}^{\alpha'} = \bar{R}_{\alpha\beta} - \sigma (K K_{\alpha\beta} - K_{\alpha}^{\delta} K_{\beta\delta}) \quad (152)$$

and contracting on α and β gives

$$R - 2\sigma R_{\alpha\beta} n^{\alpha} n^{\beta} = \bar{R} - \sigma (K^2 - K^{\alpha\beta} K_{\alpha\beta}) \quad (153)$$

known as the scalar Gauss relation. Some insight into (150) may be found by considering that the curvature $\bar{R}_{\nu\lambda\rho}^{\mu}$ of a hypersurface embedded in a flat space with $R_{\nu\lambda\rho}^{\mu} = 0$ is determined entirely by the extrinsic curvature.

Applying the Ricci identity (109) to the vector n as

$$(\nabla_{\beta} \nabla_{\alpha} - \nabla_{\alpha} \nabla_{\beta}) n^{\gamma} = R_{\gamma'\alpha\beta}^{\gamma} n^{\gamma'} , \quad (154)$$

projecting the LHS onto Σ as

$$P_{\alpha}^{\alpha'} P_{\beta}^{\beta'} P_{\gamma}^{\gamma'} (\nabla_{\alpha'} \nabla_{\beta'} - \nabla_{\beta'} \nabla_{\alpha'}) n^{\gamma'} , \quad (155)$$

and using the identity (125) leads us to

$$D_{\beta} K_{\alpha}^{\gamma} - D_{\alpha} K_{\beta}^{\gamma} = P_{\gamma}^{\gamma'} n^{\delta} P_{\alpha}^{\alpha'} P_{\beta}^{\beta'} R_{\delta\alpha'\beta'}^{\gamma'} \quad (156)$$

which is called the Codazzi relation. Using (107) for the unit normal n_{α} provides an interpretation of this expression as

$$n_{\delta} R_{\mu\nu\lambda}^{\delta} = \sigma N \delta_{\delta}^5 R_{\mu\nu\lambda}^{\delta} \longrightarrow R_{\mu\nu\lambda}^5 = \sigma \frac{1}{N} (D_{\lambda} K_{\nu\mu} - D_{\nu} K_{\lambda\mu}) \quad (157)$$

recalling the role of the extrinsic curvature $K_{\mu\nu}$ as the curvature of $\mathcal{M}$ mapped to the hypersurface Σ and embedded in the larger manifold $\mathcal{M}_5$. Contracting on α and γ in (156) produces

$$D_{\beta} K - D_{\alpha} K_{\beta}^{\alpha} = n^{\alpha} P_{\beta}^{\beta'} R_{\alpha\beta'} \quad (158)$$

known as the contracted Codazzi relation.

Returning to the Ricci identity for n^α , we apply (128) twice to terms $\nabla_\beta \nabla_\gamma n^\alpha$ and project onto (154) with $P_{\alpha\alpha'} n^{\gamma'} P_\beta^{\beta'}$ to obtain

$$-K_{\alpha\gamma} K_\beta^\gamma + \sigma \frac{1}{N} D_\beta D_\alpha N + P_{\alpha'}^{\alpha'} P_\beta^{\beta'} n^{\gamma'} \nabla_\gamma K_{\alpha'\beta'} = P_{\alpha\alpha'} n^{\gamma'} P_\beta^{\beta'} R_{\delta\beta'\gamma'} n^\delta. \quad (159)$$

Writing the Lie derivative of $K_{\alpha\beta}$ and using (142) for $\nabla_\beta m_\alpha$

$$\mathcal{L}_m K_{\alpha\beta} = N n^\gamma \nabla_\gamma K_{\alpha\beta} - 2N K_{\alpha\gamma} K_\beta^\gamma - K_{\alpha\gamma} D^\gamma N n_\beta - K_{\beta\gamma} D^\gamma N n_\alpha \quad (160)$$

the last two equations combine as

$$\frac{1}{N} \mathcal{L}_m K_{\alpha\beta} + \sigma \frac{1}{N} D_\alpha D_\beta N + K_{\alpha\gamma} K_\beta^\gamma = P_{\alpha\alpha'} n^\delta P_\beta^{\beta'} n^{\gamma'} R_{\delta\beta'\gamma'}^{\alpha'} \quad (161)$$

to provide an evolution equation for $K_{\alpha\beta}$. Rewriting the contracted Gauss relation (152) as

$$P_{\alpha\alpha'} n^{\delta'} P_\beta^{\beta'} n^{\gamma'} R_{\delta'\beta'\gamma'}^{\alpha'} = \sigma P_{\alpha'}^{\alpha'} P_\beta^{\beta'} R_{\alpha'\beta'} - \sigma \bar{R}_{\alpha\beta} + K K_{\alpha\beta} - K_\alpha^\delta K_{\beta\delta} \quad (162)$$

we can put (161) into the form

$$P_{\alpha'}^{\alpha'} P_\beta^{\beta'} R_{\alpha'\beta'} = \frac{1}{N} \mathcal{L}_m K_{\alpha\beta} + \sigma \frac{1}{N} D_\alpha D_\beta N + \bar{R}_{\alpha\beta} - \sigma K K_{\alpha\beta} + \sigma 2 K_\alpha^\delta K_{\beta\delta} \quad (163)$$

in which only $R_{\alpha\beta}$ on the LHS refers explicitly to the 5D geometry of $\mathcal{M}_5$.

4.4. Decomposition of the Field Equations

Moving from the mathematics of embedded surfaces to the physics of gravitation, we introduce the field equations (99) into (163) by projecting

$$P_{\alpha'}^{\alpha'} P_\beta^{\beta'} R_{\alpha'\beta'} = \frac{8\pi G}{c^4} P_{\alpha'}^{\alpha'} P_\beta^{\beta'} \left(T_{\alpha'\beta'} - \frac{1}{2} P_{\alpha'\beta'} S \right) = \frac{8\pi G}{c^4} \left(S_{\alpha\beta} - \frac{1}{2} P_{\alpha\beta} S \right) \quad (164)$$

and pulling back to $\mathcal{M}$ to obtain the evolution equation for $K_{\mu\nu}$ in the form

$$\begin{aligned} \frac{1}{c_5} \partial_\tau K_{\mu\nu} &= \mathcal{L}_N K_{\mu\nu} - \sigma D_\mu D_\nu N \\ &+ N \left[-\sigma \bar{R}_{\mu\nu} + K K_{\mu\nu} - 2K_\mu^\lambda K_{\nu\lambda} + \sigma \frac{8\pi G}{c^4} \left(S_{\mu\nu} - \frac{1}{2} \gamma_{\mu\nu} S \right) \right] \end{aligned} \quad (165)$$

where again we used $\mathcal{L}_m = \mathcal{L}_5 - \mathcal{L}_N$.

The double projection of the field equation onto the unit normal n is

$$R_{\alpha\beta} n^\alpha n^\beta = \frac{8\pi G}{c^4} \left(T_{\alpha\beta} - \frac{1}{2} P_{\alpha\beta} S \right) n^\alpha n^\beta = \frac{8\pi G}{c^4} \kappa \quad (166)$$

where we used $P_{\alpha\beta} n^\alpha = 0$ and the definition of κ in (96). Using the scalar Gauss relation (153)

$$R - 2\sigma R_{\alpha\beta} n^\alpha n^\beta = \bar{R} - \sigma \left(K^2 - K^{\alpha\beta} K_{\alpha\beta} \right) \quad (167)$$

we evaluate R by acting on the field equations (99) with the symmetry-broken metric $P^{\alpha\beta}$ leading to

$$P^{\alpha\beta} R_{\alpha\beta} = \frac{8\pi G}{c^4} P^{\alpha\beta} \left(S_{\alpha\beta} - \frac{1}{2} P_{\alpha\beta} S \right) = -\frac{8\pi G}{c^4} S. \quad (168)$$

Finally, the double projection onto n of the field is

$$R_{\alpha\beta}n^\alpha n^\beta = \frac{8\pi G}{c^4} \left(T_{\alpha\beta} - \frac{1}{2} P_{\alpha\beta} S \right) n^\alpha n^\beta = \frac{8\pi G}{c^4} T_{\alpha\beta} n^\alpha n^\beta = \frac{8\pi G}{c^4} \kappa \quad (169)$$

so that the scalar Gauss relation becomes

$$\bar{R} - \sigma \left(K^2 - K^{\mu\nu} K_{\mu\nu} \right) = -\frac{8\pi G}{c^4} (S + \sigma\kappa) \quad (170)$$

known as the Hamiltonian constraint. Since this expression has no τ -derivatives, it will be satisfied at all times if it is satisfied by the initial conditions.

The mixed projection of the field equations with $P^{\beta'}_\beta$ and n^α is

$$R_{\alpha\beta}n^\alpha P^{\beta'}_\beta = \frac{8\pi G}{c^4} \left(T_{\alpha\beta'} - \frac{1}{2} P_{\alpha\beta'} S \right) n^\alpha P^{\beta'}_\beta = \frac{8\pi G}{c^4} T_{\alpha\beta'} n^\alpha P^{\beta'}_\beta = -\frac{8\pi G}{c^4} p_\beta \quad (171)$$

where we used the definition of p_β in (96). Combining this with the contracted Codazzi relation (158) and pulling back to $\mathcal{M}$ leads to the momentum constraint

$$D_\mu K^\mu_\nu - D_\nu K = \frac{8\pi G}{c^4} p_\nu \quad (172)$$

referring to the flow of mass into spacetime. This expression again has no τ -derivatives and remains form-invariant as τ advances. We notice that the evolution equations contain only objects defined on $\mathcal{M}$ and so the connection components $\Gamma^5_{\mu\nu}$ appear only implicitly through the extrinsic curvature $K_{\mu\nu}$.

Collecting the evolution equations (144) and (165) with the constraints (170) and (172), the initial-value problem takes the final form

$$\begin{aligned} \frac{1}{c_5} \partial_\tau \gamma_{\mu\nu} &= \mathcal{L}_N \gamma_{\mu\nu} - 2N K_{\mu\nu} \\ \frac{1}{c_5} \partial_\tau K_{\mu\nu} &= \mathcal{L}_N K_{\mu\nu} - \sigma D_\mu D_\nu N \\ &\quad + N \left[-\sigma \bar{R}_{\mu\nu} + K K_{\mu\nu} - 2K^\lambda_\mu K_{\nu\lambda} + \sigma \frac{8\pi G}{c^4} \left(S_{\mu\nu} - \frac{1}{2} \gamma_{\mu\nu} S \right) \right] \\ \bar{R} - \sigma (K^2 - K^{\mu\nu} K_{\mu\nu}) &= -\frac{8\pi G}{c^4} (S + \sigma\kappa) \\ D_\mu K^\mu_\nu - D_\nu K &= \frac{8\pi G}{c^4} p_\nu \end{aligned}$$

5. The ADM Hamiltonian Formulation

The 4+1 ADM formalism can be put into Hamiltonian form with phase space variables $\gamma_{\mu\nu}$ and conjugate momentum $\pi_{\mu\nu}$. This formulation follows the general outline for the 3+1 formalism in standard GR, and so only the most important steps will be summarized here.

The ADM parameterization (69) splits the configuration space variable $g_{\alpha\beta}$ in the field equations into $g_{\alpha\beta}(\gamma_{\mu\nu}, N^\mu, N)$. Because N and N^μ are not dynamical, the phase space consists of $\gamma_{\mu\nu}$ and $\dot{\gamma}_{\mu\nu} = \partial_5 \gamma_{\mu\nu}$ where from (144) we write

$$\dot{\gamma}_{\mu\nu} = \frac{1}{c_5} \mathcal{L}_\tau \gamma_{\mu\nu} = \mathcal{L}_N \gamma_{\mu\nu} + 2N K_{\mu\nu} \quad (173)$$

with the sign in the definition of $K_{\mu\nu}$ reversed by convention. Contracting on α and β in (163) and combining with the scalar Gauss relation (153) leads to

$$R = \bar{R} - \sigma \left(K^2 - K^{\alpha\beta} K_{\alpha\beta} \right) + 2\sigma \nabla_\alpha \left(n^\beta \nabla_\beta n^\alpha - n^\alpha \nabla_\beta n^\beta \right) \quad (174)$$

so that discarding the total gradient, the Einstein-Hilbert action for GR in the absence of matter becomes

$$S_{ADM}[\gamma_{\mu\nu}, \dot{\gamma}_{\mu\nu}, N^\mu, N] = \int d\tau d^4x \sqrt{\gamma} N \left[\bar{R} - \sigma \left(K^{\mu\nu} K_{\mu\nu} - K^2 \right) \right] \quad (175)$$

where $\gamma = |\det \gamma_{\mu\nu}|$. The DeWitt metric is defined as

$$G^{\mu\nu\lambda\rho} = \frac{1}{2} \left(\gamma^{\mu\lambda} \gamma^{\nu\rho} + \gamma^{\mu\rho} \gamma^{\nu\lambda} - 2\gamma^{\mu\nu} \gamma^{\lambda\rho} \right) \quad (176)$$

with inverse in D dimensions

$$G_{\mu\nu\lambda\rho} = \frac{1}{2} \left(\gamma_{\lambda\zeta} \gamma_{\rho\kappa} + \gamma_{\lambda\kappa} \gamma_{\rho\zeta} - \frac{2}{D-1} \gamma_{\lambda\rho} \gamma_{\zeta\kappa} \right) \quad (177)$$

in terms of which

$$G^{\mu\nu\lambda\rho} K_{\mu\nu} K_{\lambda\rho} = \frac{1}{2} \left(\gamma^{\mu\lambda} \gamma^{\nu\rho} + \gamma^{\mu\rho} \gamma^{\nu\lambda} - 2\gamma^{\mu\nu} \gamma^{\lambda\rho} \right) K_{\mu\nu} K_{\lambda\rho} = K^{\mu\nu} K_{\mu\nu} - K^2 \quad (178)$$

so that

$$\mathcal{L}_{ADM}[\gamma_{\mu\nu}, \dot{\gamma}_{\mu\nu}, N^\mu, N] = \sqrt{\gamma} N \left[-\sigma G^{\mu\nu\lambda\rho} K_{\mu\nu} K_{\lambda\rho} + \bar{R} \right]. \quad (179)$$

Because $K^{\mu\nu}$ is first order in derivatives, the first term has the form of kinetic energy. The canonical conjugate momentum to $\gamma_{\mu\nu}$ is

$$\pi^{\mu\nu} = \frac{\partial \mathcal{L}_{ADM}}{\partial \dot{\gamma}_{\mu\nu}} = -2\sigma \sqrt{\gamma} N G^{\zeta\kappa\lambda\rho} K_{\lambda\rho} \frac{\partial K_{\zeta\kappa}}{\partial \dot{\gamma}_{\mu\nu}} \quad (180)$$

so that using (173) to obtain

$$\frac{\partial K_{\zeta\kappa}}{\partial \dot{\gamma}_{\mu\nu}} = \frac{1}{2N} \delta_\zeta^\mu \delta_\kappa^\nu \quad (181)$$

we find

$$\pi^{\mu\nu} = -\sigma \sqrt{\gamma} (K^{\mu\nu} - \gamma^{\mu\nu} K) \quad (182)$$

with trace

$$\pi = \gamma_{\mu\nu} \pi^{\mu\nu} = \sigma(D-1) \sqrt{\gamma} K \rightarrow K = \frac{\sigma}{(D-1) \sqrt{\gamma}} \pi. \quad (183)$$

Writing $K^{\mu\nu}$ in terms of $\pi^{\mu\nu}$

$$K^{\mu\nu} = -\frac{\sigma}{\sqrt{\gamma}} \left(\pi^{\mu\nu} - \gamma^{\mu\nu} \frac{1}{(D-1)} \pi \right) \quad (184)$$

and lowering the indices of $\pi^{\mu\nu}$

$$G_{\mu\nu\lambda\rho} \pi^{\lambda\rho} = \pi_{\mu\nu} - \frac{1}{D-1} \gamma_{\mu\nu} \pi = -\sigma \sqrt{\gamma} K_{\mu\nu} \quad (185)$$

we see that $K_{\mu\nu}$ represents the momentum conjugate to $\gamma_{\mu\nu}$. Replacing $K_{\mu\nu}$ in (173), we can write the velocity as

$$\dot{\gamma}_{\mu\nu} = -\sigma \frac{2N}{\sqrt{\gamma}} G_{\mu\nu\lambda\rho} \pi^{\lambda\rho} + L_N \gamma_{\mu\nu} \quad (186)$$

in terms of the momentum and configuration variable. Because $\bar{R}$ is independent of the lapse N and shift N^μ the Lagrangian $\mathcal{L}_{ADM}[\gamma_{\mu\nu}, \dot{\gamma}_{\mu\nu}, N^\mu, N]$ contains no derivatives of N, N^μ and these act

as Lagrange multipliers enforcing as constraints their conjugates. Thus, we find the Hamiltonian constraint from

$$0 = p_N = -\frac{\partial \mathcal{L}_{ADM}}{\partial \dot{N}} = \sqrt{\gamma} \left[-\sigma \left(K^{\mu\nu} K_{\mu\nu} - K^2 \right) - \bar{R} \right] = \mathcal{H} \quad (187)$$

and the momentum constraint is

$$0 = -\frac{\partial \mathcal{L}_{ADM}}{\partial \dot{N}_\mu} = 2\sigma \sqrt{\gamma} (D_\nu K^{\nu\mu} - D^\mu K) = \mathcal{H}^\mu. \quad (188)$$

Using (182), we can also write

$$\mathcal{H}^\nu = 2\sigma \sqrt{\gamma} D_\mu \left(G^{\mu\nu\lambda\rho} K_{\lambda\rho} \right) = \sigma 2D_\mu \pi^{\mu\nu}. \quad (189)$$

Performing the Legendre transformation to the Hamiltonian density

$$\begin{aligned} \mathcal{H}_{ADM} &= \pi^{\mu\nu} \dot{\gamma}_{\mu\nu} - \mathcal{L}_{ADM} [\gamma_{\mu\nu}, \dot{\gamma}_{\mu\nu}, N^\mu, N] \\ &= -\sigma N \sqrt{\gamma} G^{\mu\nu\lambda\rho} K_{\mu\nu} K_{\lambda\rho} + 2\pi^{\mu\nu} D_\mu N_\nu - \sqrt{\gamma} N R \end{aligned} \quad (190)$$

where the Lagrange multipliers N, N^μ do not require kinetic terms. Integrating by parts and discarding the total gradient provides

$$2\pi^{\mu\nu} D_\mu N_\nu = 2D_\mu (\pi^{\mu\nu} N_\nu) - N_\nu (2D_\mu \pi^{\mu\nu}) = N_\nu \mathcal{H}^\nu \quad (191)$$

and using (187), we arrive at

$$\mathcal{H}_{ADM} = N\mathcal{H} + N_\nu \mathcal{H}^\nu. \quad (192)$$

Writing the Hamiltonian in the form

$$\mathcal{H}_{ADM} = \pi^{\mu\nu} \dot{\gamma}_{\mu\nu} + \dot{p}_N \mathcal{H} + \dot{p}_{N^\mu} \mathcal{H}^\mu - \mathcal{L}_{ADM} \quad (193)$$

the Hamiltonian and momentum constraints $\mathcal{H} = 0$ and $\mathcal{H}^\nu = 0$ are seen to be secondary constraints arising from the requirement that the primary constraints $p_N = 0$ and $p_{N^\mu} = 0$ are preserved under time evolution,

$$\dot{p}_N = \{p_N, \mathcal{H}_{ADM}\} = 0 \quad \dot{p}_{N^\mu} = \{p_{N^\mu}, \mathcal{H}_{ADM}\} = 0. \quad (194)$$

The Einstein equations

$$R_{\mu\nu} - \frac{1}{2} \gamma_{\mu\nu} R = 0 \quad (195)$$

then follow from

$$\dot{\gamma}_{\mu\nu} = \{\gamma_{\mu\nu}, \mathcal{H}_{ADM}\} \quad \dot{\pi}^{\mu\nu} = \{\pi^{\mu\nu}, \mathcal{H}_{ADM}\} \quad (196)$$

for the canonical variables

$$\left\{ \gamma_{\mu\nu}, \pi^{\lambda\rho} \right\} = \frac{1}{2} \left(\delta_\mu^\lambda \delta_\nu^\rho + \delta_\mu^\rho \delta_\nu^\lambda \right). \quad (197)$$

The equation for $\dot{\gamma}_{\mu\nu}$ simply reproduces the definition of $\pi^{\mu\nu}$, since R does not contain $\dot{\gamma}_{\mu\nu}$ and so $\{\gamma_{\mu\nu}, R\} = 0$. The Einstein equations are thus equivalent to $\dot{\pi}^{\mu\nu}$.

6. Weak Gravitation in the SHP Formalism

6.1. Weak Field Approximation

The weak field approximation poses the metric as a small perturbation of the 5D flat metric

$$g_{\alpha\beta} = \eta_{\alpha\beta} + h_{\alpha\beta} \quad \longrightarrow \quad \partial_\gamma g_{\alpha\beta} = \partial_\gamma h_{\alpha\beta} \quad (h_{\alpha\beta})^2 \approx 0 \quad (198)$$

$$\eta_{\alpha\beta} = \text{diag}(-1, 1, 1, 1, \sigma) \quad (199)$$

with inverse

$$g^{\alpha\beta} = \eta^{\alpha\beta} - h^{\alpha\beta} \quad (200)$$

so that the 5D Ricci tensor reduces to the linear terms

$$R_{\alpha\beta} \approx \frac{1}{2} \left(\partial_\beta \partial_\gamma h_\alpha^\gamma + \partial_\alpha \partial_\gamma h_\beta^\gamma - \partial^\gamma \partial_\gamma h_{\alpha\beta} - \partial_\alpha \partial_\beta h \right). \quad (201)$$

Invariance of $R_{\alpha\beta}$ under a coordinate translation $x'^\alpha = x^\alpha + \Lambda^\alpha(x)$ allows us to impose the Lorenz gauge condition

$$\partial^\beta \left(h_{\alpha\beta} - \frac{1}{2} \eta_{\alpha\beta} h \right) = 0 \longrightarrow \partial^\beta h_{\alpha\beta} = \frac{1}{2} \partial_\alpha h \quad (202)$$

putting the 5D Ricci tensor into the form

$$R_{\alpha\beta} \approx -\frac{1}{2} \partial^\gamma \partial_\gamma h_{\alpha\beta} \quad (203)$$

and so we obtain the 5D wave equation

$$-\partial^\gamma \partial_\gamma h_{\alpha\beta} = \frac{16\pi G}{c^4} \left(T_{\alpha\beta} - \frac{1}{2} \hat{\eta}_{\alpha\beta} S \right). \quad (204)$$

A Green's function satisfying

$$\partial^\gamma \partial_\gamma G(x, \tau) = -\delta^4(x) \delta(\tau) \quad (205)$$

provides the generic solution

$$h_{\alpha\beta} = \frac{16\pi G}{c^4} \int d^4x d\tau G(x - x', \tau - \tau') \left(T_{\alpha\beta}(x', \tau') - \frac{1}{2} \hat{\eta}_{\alpha\beta} S(x', \tau') \right). \quad (206)$$

Introducing the notation

$$\xi^\alpha(\tau) = \frac{1}{c} u^\alpha(\tau) = \frac{1}{c} \frac{dx^\alpha}{d\tau} \quad (207)$$

the mass-energy-momentum tensor (42) is

$$T^{\alpha\beta} = m\rho(x, \tau) \dot{x}^\alpha \dot{x}^\beta = m\rho(x, \tau) u^\alpha u^\beta = mc^2 \rho(x, \tau) \xi^\alpha \xi^\beta \quad (208)$$

for some spacetime event density $\rho(x, \tau)$. Denoting

$$\mathcal{G}[\rho(x, \tau)] = \int d^4x d\tau G(x - x', \tau - \tau') \rho(x', \tau') \quad (209)$$

we have

$$h_{\alpha\beta} = \frac{16\pi Gm}{c^2} \mathcal{G}[\rho(x, \tau)] \left(\xi_\alpha \xi_\beta - \frac{1}{2} \hat{\eta}_{\alpha\beta} \hat{\xi}^2 \right) \quad (210)$$

where $\hat{\xi}^2 = \hat{\eta}^{\alpha\beta} \xi_\alpha \xi_\beta$. Considering a source distribution evolving along the t -axis in its rest frame, we have

$$u = (c, 0, 0, 0, c_5) \longrightarrow \xi = \left(1, 0, 0, 0, \frac{c_5}{c} \right) \longrightarrow \hat{\eta}_{\alpha\beta} \xi^\alpha \xi^\beta = -1 \quad (211)$$

from which the perturbation becomes

$$\begin{aligned} h_{00} &= \frac{8\pi Gm}{c^2} \mathcal{G}[\rho(x, \tau)] & h_{05} &= \frac{16\pi Gm}{c^2} \frac{c_5}{c} \mathcal{G}[\rho(x, \tau)] \\ h_{ij} &= \frac{8\pi Gm}{c^2} \mathcal{G}[\rho(x, \tau)] \delta_{ij} & h_{55} &= \frac{16\pi Gm}{c^2} \frac{c_5^2}{c^2} \mathcal{G}[\rho(x, \tau)] \end{aligned} \quad (212)$$

where electromagnetic phenomenology [3] indicates that $c_5/c \ll 1$. Writing the approximate Green's function used in electrodynamics [35]

$$G(x, \tau) = \frac{1}{2\pi} \delta(x^2) \delta(\tau) \quad (213)$$

and taking a uniform density $\rho(x) = \delta^3(x)$ evenly spread along the t -axis leads to the spacetime metric

$$g_{\mu\nu} = \text{diag} \left[-1 + \frac{2Gm}{c^2 r}, \left(1 + \frac{2Gm}{c^2 r} \right) \delta_{ij} \right] \quad (214)$$

which becomes the Schwarzschild metric when expressed in the isotropic spherical coordinates [36] defined through

$$R = r \left(1 + \frac{GM}{2c^2 r} \right)^2. \quad (215)$$

It is worth noting how this result would be different if we had not broken the 5D flat metric $\eta_{\alpha\beta} \rightarrow \hat{\eta}_{\alpha\beta}$ in constructing the field equations in Section 3. We would then have (86) for the field equations which produce the perturbation

$$\begin{aligned} h_{00} &= \frac{2}{3} \frac{4\pi Gm}{c^2} \mathcal{G}[\rho(x, \tau)] & h_{05} &= \frac{2}{3} \frac{4\pi Gm}{c^2} \frac{c_5}{c} \mathcal{G}[\rho(x, \tau)] \\ h_{ij} &= \frac{1}{3} \frac{4\pi Gm}{c^2} \mathcal{G}[\rho(x, \tau)] \delta_{ij} & h_{55} &= \frac{1}{3} \frac{4\pi Gm}{c^2} \frac{c_5^2}{c^2} \mathcal{G}[\rho(x, \tau)] \end{aligned} \quad (216)$$

leading to the phenomenologically unacceptable result $h_{00} = 2h_{ij}$. This shows that preserving the $O(3,1)$ covariance of the mass terms is not only physically reasonable, but actually required for a reasonable theory.

6.2. The 4+1 Decomposition for the Linearized Theory

Discarding terms of order $(h_{\alpha\beta})^2 \approx 0$, the ADM decomposition allows us to identify the terms of the perturbed metric as

$$\|g_{\alpha\beta}\| = \begin{bmatrix} \gamma_{\mu\nu} & N_\mu \\ N_\mu & \sigma N^2 + \gamma_{\mu\nu} N^\mu N^\nu \end{bmatrix} = \begin{bmatrix} \eta_{\mu\nu} + h_{\mu\nu} & h_{\mu 5} \\ h_{\mu 5} & \eta_{55} + h_{55} \end{bmatrix} \quad (217)$$

from which

$$\sigma N^2 + \gamma_{\mu\nu} N^\mu N^\nu \approx \sigma N^2 = \sigma + h_{55} \rightarrow N = \sqrt{1 + \sigma h_{55}} \approx 1 + \frac{1}{2} \sigma h_{55} \quad (218)$$

and the unit normal is

$$n_\alpha = \sigma N \delta_\alpha^5 = \sigma \sqrt{1 + \sigma h_{55}} \delta_\alpha^5 = \sigma \left(1 + \frac{1}{2} \sigma h_{55} \right) \delta_\alpha^5 \quad (219)$$

$$n^\alpha = -h_5^\mu \delta_\mu^\alpha + \left(1 - \frac{1}{2} \sigma h_{55} \right) \delta_5^\alpha. \quad (220)$$

Inserting these into (125) and discarding terms of the type $\Gamma_{\alpha\beta}^\gamma \times h_{\alpha'\beta'} \approx 0$, we may calculate the extrinsic curvature directly as

$$K_{\alpha\beta} = \frac{1}{2} \partial_\alpha h_{55} \delta_\beta^5 + \sigma \Gamma_{\alpha\beta}^5 + \delta_\alpha^5 \left(\frac{1}{2} \partial_5 h_{55} \delta_\beta^5 + \sigma \Gamma_{5\beta}^5 \right) \quad (221)$$

which using $\delta_\nu^5 = 0$ provides

$$K_{\mu\nu} = \Gamma_{5\mu\nu} \quad (222)$$

showing explicitly that the extrinsic curvature recovers the 5-components of the 5D connection not present in the 4D intrinsic curvature. By definition

$$\Gamma_{5\mu\nu} = \frac{1}{2}(\partial_\nu h_{5\mu} + \partial_\mu h_{5\nu} - \partial_5 h_{\mu\nu}) \quad (223)$$

and so (222) shows that $K_{\mu\nu} \times h_{\alpha\beta} \approx 0$. Discarding all terms of order $(h_{\alpha\beta})^2 \approx 0$, the initial value problem reduces to

$$\begin{aligned} \frac{1}{c_5} \partial_\tau \gamma_{\mu\nu} &= -2K_{\mu\nu} + \partial_\mu h_{5\nu} + \partial_\nu h_{5\mu} \\ \frac{1}{c_5} \partial_\tau K_{\mu\nu} &= \frac{1}{2} \partial_\mu \partial_\nu h_{55} - \sigma \left[\bar{R}_{\mu\nu} - \frac{8\pi G}{c^4} \left(S_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} S \right) \right] \\ \bar{R} &= -\frac{8\pi G}{c^4} (S + \sigma\kappa) \\ \partial_\mu K_\nu^\mu - \partial_\nu K &= \frac{8\pi G}{c^4} p_\nu. \end{aligned} \quad (224)$$

Writing $\partial_5 = (1/c_5) \partial_\tau$ in (223) we see the the evolution equation for $\gamma_{\mu\nu}$ simply restates (222).

6.3. Insights from the Linearized Theory

In the linearized theory, the initial value problem can be derived directly from the definition of the 5D Ricci tensor

$$R_{\alpha\beta} \approx \frac{1}{2} (\partial_\beta \partial^\gamma h_{\gamma\alpha} + \partial_\alpha \partial^\gamma h_{\gamma\beta} - \partial_\gamma \partial^\gamma h_{\alpha\beta} - \partial_\alpha \partial_\beta h). \quad (225)$$

We expand the spacetime components as

$$R_{\mu\nu} = R_{\mu\nu}^{(4)} + \frac{1}{2} (\partial_\nu \partial^5 h_{5\mu} + \partial_\mu \partial^5 h_{5\nu} - \partial_5 \partial^5 h_{\mu\nu} - \sigma \partial_\mu \partial_\nu h_{55}) \quad (226)$$

where $R_{\mu\nu}^{(4)}$ is the standard 4D intrinsic curvature. Explicit calculation of the projected Ricci tensor using the contracted Gauss relation (152) with the unit normals (219) and (220) shows that in weak gravitation $\bar{R}_{\mu\nu} = R_{\mu\nu}^{(4)}$. Using (222) and (223) in (226) we obtain

$$R_{\mu\nu} = \bar{R}_{\mu\nu} + \sigma \partial_5 K_{\mu\nu} - \frac{1}{2} \sigma \partial_\mu \partial_\nu h_{55} \quad (227)$$

and writing the field equations (99) for $R_{\mu\nu}$ this becomes

$$\frac{1}{c_5} \partial_\tau K_{\mu\nu} = \frac{1}{2} \partial_\mu \partial_\nu h_{55} - \sigma \left[\bar{R}_{\mu\nu} - \frac{8\pi G}{c^4} \left(T_{\mu\nu} - \frac{1}{2} P_{\mu\nu} S \right) \right] \quad (228)$$

which recovers the evolution equation for $K_{\mu\nu}$ in (224). Thus, as expected, the evolution equations, representing the dynamical degrees of freedom, follow from the spacetime components of the 5D field equations along with the definition of $K_{\mu\nu}$.

Imposing the Lorenz gauge condition on the 5-components of the Ricci tensor

$$R_{5\beta} = -\frac{1}{2} \partial_\gamma \partial^\gamma h_{5\beta} + \frac{1}{2} (\partial_\beta \partial^\gamma h_{\gamma 5} + \partial_5 \partial^\gamma h_{\gamma\beta} - \partial_5 \partial_\beta h) \quad (229)$$

where $h = \eta^{\alpha\beta} h_{\alpha\beta}$, we obtain from (99)

$$R_{5\beta} = -\frac{1}{2} \partial^\gamma \partial_\gamma h_{5\beta} = \frac{8\pi G}{c^4} T_{5\beta} \quad (230)$$

and neglecting $T_{\alpha\beta} \times h_{\alpha'\beta'} \approx 0$ we can write the sources as

$$\kappa = n_\alpha n_\beta T^{\alpha\beta} \simeq T^{55} \quad p_\mu = -n_\alpha P_{\mu\mu'} T^{\alpha\mu'} \simeq -\sigma T_\mu^5 \quad (231)$$

satisfying wave equations

$$\frac{1}{2} \partial^\gamma \partial_\gamma h_{5\mu} = \frac{8\pi G}{c^4} p_\mu \quad \frac{1}{2} \partial^\gamma \partial_\gamma h_{55} = -\frac{8\pi G}{c^4} \kappa. \quad (232)$$

Taking the trace of (228) with the induced metric $\gamma_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ and rearranging leads to

$$\bar{R} = -\frac{8\pi G}{c^4} S - \sigma \partial_5 K - \frac{1}{2} \partial^\mu \partial_\mu h^{55}. \quad (233)$$

We may evaluate the trace of $K_{\mu\nu}$ using (222) along with the Lorenz gauge condition in the form $\partial^\beta h_{\alpha\beta} = \frac{1}{2} \partial_\alpha h$ to find

$$K = -\frac{1}{2} \sigma \partial_5 h^{55} = -\frac{1}{2} \partial^5 h^{55} \quad (234)$$

so that (233) becomes

$$\bar{R} = -\frac{8\pi G}{c^4} S + \frac{1}{2} \sigma \partial_5 \partial^5 h^{55} + \frac{1}{2} \sigma \partial^\mu \partial_\mu h^{55} = -\frac{8\pi G}{c^4} S + \frac{1}{2} \sigma \partial^\gamma \partial_\gamma h^{55}. \quad (235)$$

Now inserting the second of (232) we obtain

$$\bar{R} = -\frac{8\pi G}{c^4} (S + \sigma \kappa) \quad (236)$$

which we recognize as the Hamiltonian constraint in the linearized form. Combining

$$\partial_\nu K = -\frac{1}{2} \partial_\nu \partial^5 h_{55} \quad (237)$$

and

$$\partial_\mu K_\nu^\mu = \frac{1}{2} \left(-\partial_5 \partial^\mu h_{\mu\nu} + \partial^\mu \partial_\mu h_{5\nu} + \frac{1}{2} \partial_\nu \partial_5 h - \partial_\nu \partial^5 h_{55} \right) \quad (238)$$

with the Lorenz gauge we obtain

$$\partial_\mu K_\nu^\mu - \partial_\nu K = \frac{1}{2} \partial^\gamma \partial_\gamma h_{5\nu} = \frac{8\pi G}{c^4} p_\nu \quad (239)$$

which we recognize as the momentum constraint. Thus, as expected, the 5-components of the field equations (99) provide the non-evolving the constraints.

The equilibrium condition for the initial value problem is $\partial_\tau \rightarrow 0$ so that

$$0 = -2K_{\mu\nu} + \partial_\mu h_{5\nu} + \partial_\nu h_{5\mu} \quad (240)$$

$$0 = \frac{1}{2} \partial_\mu \partial_\nu h_{55} - \sigma \left[\bar{R}_{\mu\nu} - \frac{8\pi G}{c^4} \left(S_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} S \right) \right] \quad (241)$$

$$\bar{R} = -\frac{8\pi G}{c^4} (S + \sigma \kappa) \quad (242)$$

$$\partial_\mu K_\nu^\mu - \partial_\nu K = \frac{8\pi G}{c^4} p_\nu. \quad (243)$$

Taking the trace of (241) we find

$$\bar{R} = \frac{1}{2} \sigma \partial^\mu \partial_\mu h_{55} - \frac{8\pi G}{c^4} S = -\frac{8\pi G}{c^4} S - \frac{1}{2} \sigma \partial^\mu \partial_\mu h^{55} = -\frac{8\pi G}{c^4} (S + \sigma \kappa) \quad (244)$$

where we used the second of (232), so that the Hamiltonian constraint is implicit in the equation relating $R_{\mu\nu}$ to the sources. At equilibrium, (234) provides $K = 0$ and the Lorenz condition reduces to

$$\partial^\mu h_{\alpha\mu} = \frac{1}{2} \partial_\alpha (h_\mu^\mu + h_5^5) \quad (245)$$

so that using (240) we find

$$\partial_\mu K_\nu^\mu - \partial_\nu K = \frac{1}{2} \partial^\mu \partial_\mu h_{5\nu} = \frac{8\pi G}{c^4} p_\nu \quad (246)$$

where we used the first of (232). Thus, the momentum constraint is also implicit in the equation for $K_{\mu\nu}$ and we see that the fields $h_{5\mu}$ and $K_{\mu\nu}$ decouple from the 4D metric $\gamma_{\mu\nu}$ so that the extrinsic curvature does not contribute to the metric at equilibrium.

We recognize the standard 4D equations in (241) along with a post-Einstein contribution from h_{55} . Writing

$$\frac{1}{2} \partial_\mu \partial_\nu h_{55} = \hat{S}_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} \hat{S} \longrightarrow \hat{S} = -\frac{1}{2} \partial^\mu \partial_\mu h_{55} = \frac{8\pi G}{c^4} \kappa \quad (247)$$

where we used the second of (232) in equilibrium, we note that wave equation for h_{55} possesses an unambiguous solution through the Green's function for 4D wave equation. In this form, the equilibrium field equations become

$$\bar{R}_{\mu\nu} = \frac{8\pi G}{c^4} \left[\left(S_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} S \right) + \sigma \left(\hat{S}_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} \hat{S} \right) \right] \quad (248)$$

in which the contribution from h_{55} now appears as an additional source term associated with a background scalar mass density that could play the role of 'dark matter' or 'dark energy'.

7. Kaluza-Klein in SHP Formalism

In 1919 and 1926 Theodor Kaluza [26] and Oscar Klein [27] proposed a unification of gravitation and electromagnetism by extending Einstein's general relativity (GR) to five dimensions. They introduced a 5D manifold with line element

$$ds^2 = g_{\alpha\beta} dx^\alpha dx^\beta \quad x^\alpha = x^0, x^1, x^2, x^3, x^5 \quad (249)$$

and decomposed the metric as

$$g_{\alpha\beta} = \begin{bmatrix} \gamma_{\mu\nu} + \phi^2 A_\mu A_\nu & \phi^2 A_\mu \\ \phi^2 A_\nu & \phi^2 \end{bmatrix} \quad g^{\alpha\beta} = \begin{bmatrix} \gamma^{\mu\nu} & -A^\mu \\ -A^\nu & \phi^{-2} + A_\rho A^\rho \end{bmatrix} \quad (250)$$

with index convention $\alpha, \beta = 0, 1, 2, 3, 5$, and $\mu, \nu = 0, 1, 2, 3$. They showed that the Einstein equations extended to 5D

$$R_{\alpha\beta} - \frac{1}{2} g_{\alpha\beta} R = 0 \quad (251)$$

decompose into standard 4D gravitation and Maxwell electromagnetism, expressed through the Einstein-Hilbert-Maxwell action

$$S = \int d^4x dx^5 \sqrt{-g} \left(\bar{R} + \frac{1}{4} F^{\mu\nu} F_{\mu\nu} \right) \quad (252)$$

where

$$\bar{R} = g^{\mu\nu} R_{\mu\nu} \quad (253)$$

is the 4D Ricci scalar and

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \quad (254)$$

is the Maxwell field strength. Kaluza and Klein found that the identification of $F_{\mu\nu}$ with the Maxwell field requires that $\phi = \text{constant}$, and that $\gamma_{\mu\nu}$ and A_μ be independent of the coordinate x^5 . Moreover, the x^5 dimension is assumed to be compactified so that the action remains finite despite the dx^5 integration over x^5 -independent fields.

Comparing with the ADM decomposition (72) of the 5D SHP metric we notice that taking $\sigma = 1$ and making the replacements $\phi \rightarrow 1/N$ and $A_\mu \rightarrow -N_\mu$, we have

$$(g^{\text{SHP}})_{\alpha\beta} = (g^{\text{Kaluza-Klein}})^{\alpha\beta} \quad (255)$$

suggesting a correspondence between SHP gravitation and Kaluza-Klein theory. But as we saw above, in SHP gravitation N and N_μ play the role of non-dynamical Lagrange multipliers, and so we expect that the electromagnetic field will not evolve with τ . In this section, we develop the Kaluza-Klein theory in the general SHP framework and show that it is consistent with SHP gravitation only in non-evolving τ -equilibrium.

In the pseudo-spacetime we defined in (21) a displacement dX leads to the interval dX^2 which Kaluza and Klein parameterized as

$$dX^2 = g_{\alpha\beta} dX^\alpha dX^\beta = \gamma_{\mu\nu} dX^\mu dX^\nu + \phi^2 (A_\mu dX^\mu + dX^5)^2. \quad (256)$$

Following the notation of (8) we rewrite this interval in the more symmetric form

$$dX^2 = g_{\alpha\beta} dX^\alpha dX^\beta = \gamma_{\mu\nu} dX^\mu dX^\nu + (a_\mu dX^\mu + a_5 dX^5)^2 \quad (257)$$

so that the metric becomes

$$g_{\alpha\beta} = \begin{bmatrix} \gamma_{\mu\nu} + a_\mu a_\nu & a_\mu a_5 \\ a_\mu a_5 & a_5^2 \end{bmatrix} \quad g^{\alpha\beta} = \begin{bmatrix} \gamma^{\mu\nu} & -\frac{1}{a_5} a^\mu \\ -\frac{1}{a_5} a^\nu & \frac{1}{a_5^2} (1 + a_\mu a^\mu) \end{bmatrix} \quad (258)$$

with the apparent correspondence

$$\sigma N^2 = \frac{1}{a_5^2} \quad N^\mu = -a_5 a_\mu \quad (259)$$

between the Kaluza-Klein fields and the Lagrange multipliers of the ADM parameterization. As an initial assumption we take $g_{\alpha\beta} = g_{\alpha\beta}(x, \tau)$ for all components. The line element (257) suggests the action

$$S = \int d\tau L(X, \dot{X}, \tau) = \int d\tau \frac{1}{2} M g_{\alpha\beta} \dot{X}^\alpha \dot{X}^\beta \quad (260)$$

where we parameterize by the external τ and unlike Kaluza and Klein, make no reference to the proper time of the motion. The Euler-Lagrange equations lead to geodesic equations

$$M \ddot{X}^\gamma + \Gamma_{\alpha\beta}^\gamma \dot{X}^\alpha \dot{X}^\beta = 0 \quad (261)$$

with the standard 5D Christoffel connection given in (27). The spacetime components of the geodesic equations expand as

$$M \ddot{X}^\mu + \Gamma_{\lambda\rho}^\mu \dot{X}^\lambda \dot{X}^\rho + 2\Gamma_{5\lambda}^\mu \dot{X}^5 \dot{X}^\lambda + \Gamma_{55}^\mu \dot{X}_5^2 = 0 \quad (262)$$

and we recall that SHP GR imposes

$$X^5 = c_5 \tau \rightarrow \dot{X}^5 \equiv 0 \quad (263)$$

as an *a priori* constraint. But here we follow Kaluza and Klein in writing the Euler-Lagrange equation for $\gamma = 5$ explicitly to obtain

$$\left(\frac{d}{d\tau} \frac{\partial}{\partial \dot{X}^5} - \frac{\partial}{\partial X^5} \right) L = \frac{d}{d\tau} (a_\mu a_5 \dot{X}^\mu + a_5^2 \dot{X}^5) - \frac{1}{2} (\partial_5 g_{\alpha\beta}) \dot{X}^\alpha \dot{X}^\beta = 0 \quad (264)$$

which we will use below. Writing the spacetime part of $g_{\alpha\beta}$ as

$$\bar{\gamma}_{\alpha\beta} = \delta_\alpha^\lambda \delta_\beta^\rho \gamma_{\lambda\rho} \quad (265)$$

we combine the metric components as

$$g_{\alpha\beta} = \bar{\gamma}_{\alpha\beta} + a_\alpha a_\beta \quad (266)$$

and readily decompose the connection into 4D metric and field terms

$$\Gamma_{\alpha\beta}^\gamma = \bar{\Gamma}_{\alpha\beta}^\gamma + \frac{1}{2} g^{\gamma\delta} [a_\delta (\partial_\beta a_\alpha + \partial_\alpha a_\beta) + a_\alpha f_{\beta\delta} + a_\beta f_{\alpha\delta}] \quad (267)$$

where

$$f_{\alpha\beta} = \partial_\alpha a_\beta - \partial_\beta a_\alpha \quad (268)$$

and

$$\bar{\Gamma}_{\gamma\alpha\beta} = \frac{1}{2} [\partial_\beta \bar{\gamma}_{\alpha\gamma} + \partial_\alpha \bar{\gamma}_{\beta\gamma} - \partial_\gamma \bar{\gamma}_{\alpha\beta}] = \frac{1}{2} [\delta_\alpha^\lambda \delta_\beta^\rho \partial_\gamma \gamma_{\lambda\rho} + \delta_\beta^\lambda \delta_\gamma^\rho \partial_\alpha \gamma_{\lambda\rho} - \delta_\alpha^\lambda \delta_\beta^\rho \partial_\gamma \gamma_{\lambda\rho}] \quad (269)$$

depends only on derivatives of the spacetime metric $\gamma_{\lambda\rho}$. The spacetime components of $\bar{\Gamma}_{\alpha\beta}^\gamma$ that enter the equations of motion (262) are

$$\bar{\Gamma}_{\alpha\beta}^\mu = \frac{1}{2} g^{\mu\gamma} [\delta_\alpha^\lambda \delta_\beta^\rho \partial_\gamma \gamma_{\lambda\rho} + \delta_\beta^\lambda \delta_\gamma^\rho \partial_\alpha \gamma_{\lambda\rho} - \delta_\alpha^\lambda \delta_\beta^\rho \partial_\gamma \gamma_{\lambda\rho}] \quad (270)$$

which we split into

$$\bar{\Gamma}_{\lambda\rho}^\mu = \Gamma_{\lambda\rho}^{(4)\mu} - \frac{1}{2} g^{\mu 5} \partial_5 \gamma_{\lambda\rho} = \Gamma_{\lambda\rho}^{(4)\mu} - \frac{1}{2} a^\mu a^5 \partial_5 \gamma_{\lambda\rho} \quad (271)$$

where $\Gamma_{\lambda\rho}^{(4)\mu}$ is the standard 4D Christoffel symbol,

$$\bar{\Gamma}_{\nu 5}^\mu = \frac{1}{2} g^{\mu\gamma} [\delta_\gamma^\lambda \partial_5 \gamma_{\nu\lambda} + \delta_5^\lambda \delta_\gamma^\rho \partial_\nu \gamma_{\lambda\rho} - \delta_\nu^\lambda \delta_5^\rho \partial_\gamma \gamma_{\lambda\rho}] = \frac{1}{2} g^{\mu\lambda} \partial_5 \gamma_{\nu\lambda} \quad (272)$$

and

$$\bar{\Gamma}_{55}^\mu = \frac{1}{2} g^{\mu\gamma} [\delta_5^\lambda \delta_\gamma^\rho \partial_5 \gamma_{\lambda\rho} + \delta_5^\lambda \delta_\gamma^\rho \partial_5 \gamma_{\lambda\rho} - \delta_5^\lambda \delta_5^\rho \partial_\gamma \gamma_{\lambda\rho}] = 0. \quad (273)$$

Combining with the field parts in (267) we obtain

$$\Gamma_{\lambda 5}^\mu = \frac{1}{2} g^{\mu\lambda} \partial_5 \gamma_{\nu\lambda} + \frac{1}{2} \left[a^\mu (\partial_5 a_\lambda + \partial_\lambda a_5) + a_\lambda f_5^\mu + a_5 f_\lambda^\mu \right] \quad (274)$$

$$\Gamma_{55}^\mu = \bar{\Gamma}_{55}^\mu + \frac{1}{2} g^{\mu\delta} [a_\delta (\partial_5 a_5 + \partial_5 a_5) + a_5 f_{5\delta} + a_5 f_{5\delta}] = a^\mu \partial_5 a_5 + a_5 f_5^\mu. \quad (275)$$

Considering the contribution of the 5-components to the geodesic equations, we have

$$\begin{aligned} 2c_5 \Gamma_{5\lambda}^\mu \dot{x}^\lambda + c_5^2 \Gamma_{55}^\mu &= (\gamma^{\mu\lambda} + a^\mu a^\lambda) \dot{x}^\lambda \partial_\tau \gamma_{\nu\lambda} \\ &\quad + 2c_5 \left[a^\mu \partial_\lambda a_5 + \frac{1}{2} \left(a^\mu f_{5\lambda} + a_\lambda f_5^\mu + a_5 f_\lambda^\mu \right) \right] \dot{x}^\lambda \\ &\quad + c_5^2 (a^\mu \partial_5 a_5 + a_5 f_5^\mu) \end{aligned} \quad (276)$$

and it is clear by inspection that the RHS will not be invariant under electromagnetic gauge transformations $a_\alpha \rightarrow a_\alpha + \partial_\alpha \Lambda$ unless we adopt the three conditions imposed by Kaluza and Klein

$$a_5 = \text{constant} \quad \partial_5 a_\alpha = 0 \rightarrow f_5^\mu = \partial_5 a^\mu - \partial^\mu a_5 = 0 \quad \partial_5 \gamma_{\nu\lambda} = 0 \quad (277)$$

which from (271) provides $\bar{\Gamma}_{\lambda\rho}^\mu \rightarrow \Gamma_{\lambda\rho}^{(4)\mu}$. Under the Kaluza-Klein conditions, the fields behave as standard Maxwell potentials $A_\mu(x)$, rather than the SHP fields $a_\alpha(x, \tau)$, and so we change notation as

$$a_\mu \rightarrow A_\mu \quad f_{\mu\nu} \rightarrow F_{\mu\nu}. \quad (278)$$

Now (276) simplifies to

$$2c_5 \Gamma_{5\lambda}^\mu \dot{x}^\lambda + c_5^2 \Gamma_{55}^\mu = c_5 a_5 F_\lambda{}^\mu \dot{x}^\lambda \quad (279)$$

where $a_5 = \text{constant}$ is no longer a gauge field, and the remaining Christoffel symbols are

$$\Gamma_{\lambda\rho}^\mu = \bar{\Gamma}_{\lambda\rho}^\mu + \frac{1}{2} \gamma^{\mu\sigma} (a_\lambda F_{\rho\sigma} + a_\rho F_{\lambda\sigma}) = \Gamma_{\lambda\rho}^{(4)\mu} + \frac{1}{2} (a_\lambda F_\rho{}^\mu + a_\rho F_\lambda{}^\mu). \quad (280)$$

Combining (279) and (280) with (262) the geodesic equations take the form

$$0 = M\ddot{x}^\mu + \left[\Gamma_{\lambda\rho}^{(4)\mu} + \frac{1}{2} (A_\lambda F_\rho{}^\mu + A_\rho F_\lambda{}^\mu) \right] \dot{x}^\lambda \dot{x}^\rho + c_5 a_5 F_\lambda{}^\mu \dot{x}^\lambda \quad (281)$$

$$= M\ddot{x}^\mu + \Gamma_{\lambda\rho}^{(4)\mu} \dot{x}^\lambda \dot{x}^\rho + (A_\lambda \dot{x}^\lambda + c_5 a_5) F_\rho{}^\mu \dot{x}^\rho \quad (282)$$

where we used the index symmetry of $\dot{x}^\lambda \dot{x}^\rho$. Since now $\partial_5 g_{\alpha\beta} = 0$, we see that the basis vector in the 5-direction is a Killing vector, so that the momentum P_5 conjugate to X^5 is a constant. This is expressed in the fifth Euler-Lagrange equation (264) as

$$\frac{d}{d\tau} \frac{\partial L}{\partial \dot{X}^5} = \frac{dP_5}{d\tau} = \frac{d}{d\tau} (a_\mu a_5 \dot{X}^\mu + a_5^2 \dot{X}^5) = 0, \quad (283)$$

imposing the condition

$$A_\mu \dot{x}^\mu + a_5 c_5 = \text{constant}. \quad (284)$$

Taking this constant to be e/c puts (282) into the form

$$M\ddot{x}^\mu + \Gamma_{\lambda\rho}^{(4)\mu} \dot{x}^\lambda \dot{x}^\rho = \frac{e}{c} F_\rho{}^\mu \dot{x}^\rho \quad (285)$$

which we recognize as the standard Lorentz force in curved 4D spacetime.

Although we began by assuming that all components of $g_{\alpha\beta}$ may be τ -dependent as in SHP theory, we found that identification of the geodesic equations derived from the Kaluza-Klein metric as the Lorentz force associated with a gauge invariant field requires that

$$a_5 = \text{constant} \quad \partial_5 g_{\alpha\beta} = \frac{1}{c_5} \partial_\tau (\gamma_{\alpha\beta} + a_\alpha a_\beta) = 0. \quad (286)$$

In light of the correspondence (259) these are the equilibrium conditions under which the metric and fields undergo no τ -evolution and SHP GR reduces to standard 4D GR. Nevertheless, we see from (284) that $A_\mu \dot{x}^\mu = \text{constant}$, which is not a condition associated with the standard Maxwell theory.

In the framework of SHP electrodynamics, the standard τ -independent Maxwell theory is recovered by integrating the field equations over τ as

$$A_\mu(x) = \int d\tau a_\mu(x, \tau) \quad F_{\mu\nu}(x) = \int d\tau f_{\mu\nu}(x, \tau) \quad J_\mu(x) = \int d\tau j_\mu(x, \tau) \quad (287)$$

where $a_\mu(x, \tau)$ and $j_\mu(x, \tau)$ are the potential and current induced by an event a spacetime point x at time τ , while $A_\mu(x)$ and $J_\mu(x)$ are the standard Maxwell potential and current. This process, known as concatenation, is understood to sum the field contributions at a given spacetime point x over all times τ , and describes an equilibrium state of the particles and fields.

Summarizing the connection under conditions (277) and (284) as

$$\Gamma_{\lambda\rho}^\mu = \frac{1}{2}\gamma^{\mu\sigma}(\partial_\rho\gamma_{\sigma\lambda} + \partial_\lambda\gamma_{\sigma\rho} - \partial_\sigma\gamma_{\lambda\rho}) + \frac{1}{2}(a_\lambda F_\rho{}^\mu + a_\rho F_\lambda{}^\mu) \quad (288)$$

$$\Gamma_{\lambda 5}^\mu = a_5 f_\lambda{}^\mu \quad (289)$$

$$\Gamma_{55}^5 = 0 \quad (290)$$

we calculate the diagonal components of the Ricci tensor

$$R_{\alpha\beta} = \partial_\gamma \Gamma_{\alpha\beta}^\gamma - \partial_\beta \Gamma_{\alpha\gamma}^\gamma + \Gamma_{\delta\gamma}^\gamma \Gamma_{\alpha\beta}^\delta - \Gamma_{\delta\beta}^\gamma \Gamma_{\alpha\gamma}^\delta \quad (291)$$

to find

$$\begin{aligned} R_{\mu\nu} &= R_{\mu\nu}^{(4)} - \frac{1}{2}(A_\mu \nabla_\lambda F_\nu^\lambda + A_\nu \nabla_\lambda F_\mu^\lambda) + \frac{1}{4}(F_\nu^\lambda F_{\lambda\mu} + F_\mu^\lambda F_{\lambda\nu}) \\ &\quad - \frac{1}{4}A_\mu A_\nu F_{\lambda\rho} F^{\lambda\rho} \end{aligned} \quad (292)$$

$$R_{55} = \frac{1}{4}F_{\lambda\rho} F^{\lambda\rho} \quad (293)$$

where $R_{\mu\nu}^{(4)}$ is the standard Ricci tensor in 4D, and the Ricci scalar is

$$R = R^{(4)} + \frac{1}{4}F_{\lambda\rho} F^{\lambda\rho}. \quad (294)$$

The Einstein-Hilbert action is then

$$S = \int d^5 X \sqrt{-g} R = \int d^5 X \sqrt{-g} \left(R^{(4)} + \frac{1}{4}F_{\lambda\rho} F^{\lambda\rho} \right) \quad (295)$$

taking the form of 4D gravitation with the electromagnetic field as its source. Since the integrand is independent of X^5 , it is here that Kaluza and Klein require that this fifth dimension must be compactified. As we saw in (276), permitting the metric $g_{\alpha\beta}$ to depend on τ introduces a variety of new non-gauge-invariant terms to $\Gamma_{\alpha\beta}^\gamma$. These terms would also appear in the Ricci tensor, disrupting identification of the action with the 4D Einstein-Hilbert action.

8. Discussion

As seen in Section 6, we can obtain a solution to the SHP field equations in weak gravitation that is recognizable as the Schwarzschild metric (214) by considering a source distribution evolving along the t -axis in its rest frame. This result, which is at equilibrium in τ and does not evolve, is induced by a static matter density $\rho(x) = \delta^3(x)$ describing matter evenly spread along the t -axis. We may understand this density as the concatenation of a time-local distribution of the type described in (287)

$$\rho(x) = \int d\tau \rho(x, \tau) = \int d\tau \delta(t - \tau) \delta^3(x) \quad (296)$$

where $\rho(x, \tau)$ describes an event in its rest frame evolving uniformly along the t -axis. In SHP electrodynamics, it is useful to model a source particle as an ensemble of events [37] located at some point x in space and narrowly distributed along the time axis according to some probability distribution

$\varphi(t - \tau)$, where $t(\tau)$ is the event's nominal time coordinate and $\varphi(s)$ has its maximum at $\varphi(0)$. Using the lowest order approximation to the Green's function for the 5D wave equation with

$$\rho(x, \tau) = \varphi(t - \tau)\delta^3(\mathbf{x}) \quad (297)$$

this model leads to a Coulomb type potential

$$a_0(x, \tau) = -\frac{\varphi(t - R/c - \tau)}{R} + o\left(\frac{1}{R^2}\right). \quad (298)$$

A test event at some spacetime point $x = (ct, R\hat{\mathbf{x}})$ will experience a potential whose support is centered around the chronological time $\tau = t - R/c$, the retarded time of the source. An observer located at this static point x will see the potential rise and fall in strength as the chronological time τ approaches and then retreats from the retarded time.

While this model has some intuitive appeal and is generally adequate in electrodynamics, it fails as a generalization of Newtonian gravitation. Writing the perturbation to the flat metric as

$$h_{\mu\nu} = \frac{2Gm}{c^2 R} \varphi(t - R/c - \tau) \delta_{\mu\nu} \quad (299)$$

the geodesic equations for a nonrelativistic test event lead to the conserved angular momentum is $L = MR^2\dot{\phi}$ and the radial equation

$$\ddot{R} = -\frac{GM}{R^2} \left(\varphi - R \partial_R \varphi \right) + \frac{L^2}{M^2 R^3}. \quad (300)$$

For a test event similarly evolving along its t -axis at a distance R and on the lightcone of the source at $t = \tau - R/c$, we have $\varphi = 1$, so that $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ recovers Newtonian gravitation. However, under the resulting gravitational force, the distance R will decrease, so that φ will deviate from its maximum value. Given a narrow distribution, close to but away from its maximum, the term $R \partial_R \varphi$ becomes large at large R and will generally change the sign of the force term in (300). We must therefore reject separable perturbations of type (299).

8.1. Ansatz Metric

As a first step in an alternative direction, we propose an ansatz for the metric possessing certain expected properties. We can then use the 5D wave equation in the weak field approximation, to derive the source that produces the ansatz metric, evaluate the Ricci tensor and extrinsic curvature, and set up the $4 + 1$ evolution equations for general mass–energy–momentum configurations. While the ansatz metric for the derived source satisfies the evolution equations exactly, the procedure leads to alternative metrics found as perturbations under general sources. The construction of general expressions characterizing perturbed metrics is discussed in standard texts on numerical relativity, such as [38].

We propose an ansatz in the form

$$g^{\alpha\beta} = H^{\alpha\beta} \Phi(t, R, \tau) \quad (301)$$

where $H^{\alpha\beta}$ is a constant kinematic term, $\Phi(t, R, \tau)$ is a spherically symmetric function containing the dependence on t , $R = |\mathbf{x}|$, and τ . We specify the component structure of $H^{\alpha\beta}$ to be

$$\begin{aligned} g^{\mu\nu} &= \text{diag}(-1 + H^{00}\Phi, (1 + H^{00}\Phi)\delta^{ij}) \\ g^{05} &= 2\zeta_5^5 H^{00}\Phi & g^{i5} &= 0 & g^{55} &= 2\zeta_5^2 H^{00}\Phi \end{aligned} \quad (302)$$

as found from (212) for a source event evolving on the t -axis.

We would like a function Φ that recovers the $1/R$ -dependence of Newtonian gravitation but whose support is restricted to a neighborhood of τ for a given test particle. We consider a source in its rest frame evolving along its time axis as $x^0(\tau) = x^5$ and a test event similarly evolving along its time axis at a spatial distance R . This suggests the choice

$$\Phi(t, R, \tau) = \frac{1}{\sqrt{R^2 + (x^0 - x^5)^2}} = \frac{1}{\sqrt{R^2 + c^2(t - \xi^5 \tau)^2}} \quad (303)$$

and we denote

$$\rho = \sqrt{R^2 + c^2(t - \xi^5 \tau)^2} \quad (304)$$

for convenience. Evaluating the connection for this metric, we study the trajectory of a test event determined by the geodesic equation (28) with the initial conditions

$$R(0) = R \quad \dot{R}(0) = 0 \quad t(0) = 0 \quad \dot{t}(0) = \xi^5 \quad (305)$$

for which $\rho \rightarrow R$, and Φ takes on its maximum value. If the test event deviates from the trajectory $(x^0, x) = (\xi^5 \tau, R\hat{x})$, then the strength of the metric will diminish. Because $\Phi(t, R, \tau)$ has a maximum at $t = \xi^5 \tau$ with respect to t but not with respect to R , this functional form does not suffer from the difficulties associated with the geodesic equation (300).

Writing the derivatives

$$\frac{\partial \Phi}{\partial t} = -\frac{c^2(t - \xi^5 \tau)}{\rho^3} \quad \frac{\partial \Phi}{\partial \tau} = \frac{\xi^5 c^2(t - \xi^5 \tau)}{\rho^3} \quad \frac{\partial \Phi}{\partial x^i} = -\frac{x_i}{\rho^3} \quad (306)$$

leads to the connection

$$\Gamma_{00}^\mu = \frac{1}{2} H^{00} \frac{1}{\rho^3} \left(\delta^{\mu 0} c(t - \xi^5 \tau) + \delta^{\mu k} x_k \right) \quad (307)$$

$$\Gamma_{i0}^\mu = H^{00} \frac{1}{2} \frac{1}{\rho^3} \left[-\delta_i^\mu c(t - \xi^5 \tau) + \delta^{\mu 0} x_i \right] \quad (308)$$

$$\Gamma_{ij}^\mu = -\frac{1}{2} H^{00} \frac{1}{\rho^3} \left[\delta_i^\mu x_j + \delta_j^\mu x_i - \delta^{\mu k} \delta_{ij} x_k + \delta^{\mu 0} \delta_{ij} c(t - \xi^5 \tau) \right] \quad (309)$$

$$\Gamma_{50}^\mu = -\frac{1}{2} H^{00} \frac{1}{\rho^3} \left[2\sigma \xi^5 \delta^{\mu k} x_k + \delta^{\mu 0} c(t - \xi^5 \tau) \right] \quad (310)$$

$$\Gamma_{5i}^\mu = -\frac{1}{2} H^{00} \frac{1}{\rho^3} \left(-\delta_i^\mu c(t - \xi^5 \tau) + \xi^5 \delta^{\mu 0} 2\sigma x_i \right) \quad (311)$$

$$\Gamma_{55}^\mu = H^{00} \frac{1}{\rho^3} \left[\delta^{\mu k} \xi_5^2 x_k + \delta^{\mu 0} c_5(t - \xi^5 \tau)(2\sigma - \xi^5) \right] \quad (312)$$

and the equations of motion in time and space components

$$0 = \ddot{t} + H^{00} \frac{1}{\rho^3} \left\{ \frac{1}{2} c^2(t - \xi^5 \tau) \dot{t}^2 + x_i \dot{x}^i \dot{t} - \frac{1}{2} (t - \xi^5 \tau) \delta_{ij} \dot{x}^i \dot{x}^j - c c_5(t - \xi^5 \tau) \dot{t} - 2\sigma \xi_5^2 x_i \dot{x}^i + c c_5 \xi_5^2 (t - \xi^5 \tau) (2\sigma - \xi_5^2) \right\} \quad (313)$$

$$0 = \ddot{x}^k + H^{00} \frac{1}{\rho^3} c^2 \left\{ \frac{1}{2} x^k \dot{t}^2 - c(t - \xi^5 \tau) \frac{\dot{x}^k}{c} \dot{t} - x_j \frac{\dot{x}^k}{c} \frac{\dot{x}^j}{c} + \frac{1}{2} x^k \frac{\dot{x}^j}{c} \frac{\dot{x}^j}{c} - 2\sigma \xi_5^2 x^k \dot{t} + c_5(t - \xi^5 \tau) \frac{\dot{x}^k}{c} + \xi_5^4 x^k \right\}. \quad (314)$$

In the neighborhood of the initial conditions, the equations of motion reduce to

$$0 = \ddot{t} + H^{00} \frac{x_i}{R^3} \frac{\dot{x}^i}{c} c \left(\dot{t} - 2\sigma\zeta_5^2 \right) \quad (315)$$

$$0 = \ddot{x}^k + H^{00} \frac{1}{R^3} c^2 \left\{ \frac{1}{2} x^k \dot{t}^2 - x_j \frac{\dot{x}^k}{c} \frac{\dot{x}^j}{c} + \frac{1}{2} x^k \frac{\dot{x}_j}{c} \frac{\dot{x}^j}{c} - 2\sigma\zeta_5^2 x^k \dot{t} + \zeta_5^4 x^k \right\} \quad (316)$$

so that neglecting $\dot{x}/c \ll 1$ in the nonrelativistic regime the time equation reduces to

$$0 = \ddot{t} \longrightarrow \dot{t}(\tau) = \zeta^5 \quad (317)$$

and the space equation becomes

$$0 = \ddot{x}^k + H^{00} \frac{1}{R^3} c^2 \zeta_5^2 \left\{ \frac{1}{2} x^k - 2\sigma\zeta^5 x^k + \zeta_5^2 x^k \right\} \simeq \ddot{x}^k + \frac{1}{2} c^2 \zeta_5^2 H^{00} \frac{1}{R^2} \hat{x}^k \quad (318)$$

which recovers the form of Newtonian gravitation if we take $H^{00} = 2GM/c^2$.

For a relativistic test event with the initial condition $t = \zeta^5 \tau$, the equations of motion become

$$0 = \ddot{t} + H^{00} \frac{1}{R^2} \hat{x}_i \dot{x}^i \dot{t} \longrightarrow c_5 \ddot{t} \simeq -\frac{2GM}{R^2} \hat{x}_i \frac{\dot{x}^i}{c} \quad (319)$$

$$0 = \ddot{x}^k + \frac{GM}{R^2} \left\{ \hat{x}^k - \left(\frac{2\hat{x}_j \dot{x}^k - \hat{x}^k \dot{x}_j}{c} \right) \frac{\dot{x}^j}{c} \frac{c^2}{c_5^2} \right\}. \quad (320)$$

Since $c/c_5 > 1$ by assumption, while for non-tachyonic particles $\dot{x}^k < c$, we see here that the event trajectory may differ from standard post-Newtonian relativistic forms. Using the wave equation (204) to calculate the source of the ansatz metric we find

$$\frac{16\pi G}{c^4} \left(T_{\mu\nu} - \frac{1}{2} \hat{\eta}_{\mu\nu} S \right) = -\frac{1}{2} H^{00} \delta_{\mu\nu} \left(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} + \sigma \frac{1}{c_5^2} \frac{\partial^2}{\partial \tau^2} \right) \Phi \quad (321)$$

for any spacetime functional Φ . Combining the derivatives of (303)

$$\nabla^2 \Phi = -\frac{3c^2(t - \zeta^5 \tau)^2}{\rho^5} \quad (322)$$

$$-\frac{1}{c^2} \frac{\partial^2}{\partial t^2} \Phi = \frac{r^2 - 2c^2(t - \zeta^5 \tau)^2}{\rho^5} \quad (323)$$

and

$$\sigma \frac{1}{c_5^2} \frac{\partial^2}{\partial \tau^2} \Phi = -\sigma \frac{r^2 - 2c^2(t - \zeta^5 \tau)^2}{\rho^5} \quad (324)$$

we are led to

$$\frac{16\pi G}{c^4} \left(T_{\mu\nu} - \frac{1}{2} \hat{\eta}_{\mu\nu} S \right) = -\frac{1}{2} \frac{H^{00}}{\rho^5} \left[(\sigma - 1)R^2 + (5 - 2\sigma)c^2(t - \zeta^5 \tau)^2 \right] \delta_{\mu\nu} \quad (325)$$

and if we again take $H^{00} = 2GM/c^2$, then

$$T_{\mu\nu} = \frac{Mc^2}{4\pi\rho^5} \left[(\sigma - 1)R^2 + (5 - 2\sigma)c^2(t - \zeta^5 \tau)^2 \right] \text{diag}(-1, 0, 0, 0) \quad (326)$$

where only the energy component T_{00} is non-vanishing. The structure of this source is easiest to see for $\sigma = 1$, in which case

$$T_{\mu\nu} = 3 \frac{Mc^2}{4\pi\rho^5} \left[c(t - \xi^5\tau) \right]^2 \text{diag}(-1, 0, 0, 0) \quad (327)$$

where $c^2(t - \xi^5\tau)^2/\rho^5$ has units of length^{-3} , as expected for a particle density in space. Although $T_{\mu\nu}$ appears to vanish at $t = \xi^5\tau$, we recall that under this condition, $\rho \rightarrow R$ so that Φ becomes independent of t and τ . In this case, the wave equation (321) from which we derive the source reduces to

$$\partial^\gamma \partial_\gamma \left(\frac{1}{\rho} \right) \rightarrow \nabla^2 \left(\frac{1}{R} \right) = -4\pi\delta^{(3)}(x) \quad (328)$$

which describes a point source evenly spread along the t -axis. In contrast, at a small spatial distance $R \ll c(t - \xi^5\tau)$, we have $c^2(t - \xi^5\tau)^2/\rho^5 \rightarrow [c^2(t - \xi^5\tau)]^{-3}$, which describes a narrow particle density along the t -axis centered at $t = \xi^5\tau$. This localization at a coordinate time t determined by the chronological time τ expresses the desired τ -evolution. As expected, the source (327) describes a matter distribution evolving with τ , leading to the metric (302), which similarly evolves with τ , and the geodesic equations (313) and (314), whose coefficients evolve with τ .

The projected Ricci tensor is found using (227)

$$\bar{R}_{\mu\nu} = H^{00} \left[-\frac{1}{2} \delta_{\mu\nu} \left(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) + \xi^5 \frac{1}{c_5} \frac{\partial}{\partial \tau} \left(\delta_{\nu 0} \frac{\partial}{\partial x^\mu} + \delta_{\mu 0} \frac{\partial}{\partial x^\nu} \right) \right] \Phi \quad (329)$$

and the extrinsic curvature is

$$K_{\mu\nu} = -H^{00} \left[\sigma \xi^5 \left(\delta_{\nu 0} \frac{\partial}{\partial x^\mu} + \delta_{\mu 0} \frac{\partial}{\partial x^\nu} \right) + \frac{1}{2} \delta_{\mu\nu} \frac{1}{c_5} \frac{\partial}{\partial \tau} \right] \Phi \quad (330)$$

from which

$$\sigma \partial_5 K_{\mu\nu} = H^{00} \left[-\xi^5 \frac{1}{c_5} \frac{\partial}{\partial \tau} \left(\delta_{\nu 0} \frac{\partial}{\partial x^\mu} + \delta_{\mu 0} \frac{\partial}{\partial x^\nu} \right) - \sigma \frac{1}{2} \delta_{\mu\nu} \frac{1}{c_5^2} \frac{\partial^2}{\partial \tau^2} \right] \Phi. \quad (331)$$

We see that the off-diagonal terms in $\bar{R}_{\mu\nu}$ and $\sigma \partial_5 K_{\mu\nu}$ mutually cancel, leaving the source diagonal as required.

Using any suitable functional $\Phi(t, R, \tau)$ to derive an ansatz metric (302), its unperturbed source (321), along with initial conditions (329) and (330), the weak field evolution equations are satisfied exactly. Writing the unperturbed source as $T_{\mu\nu}^{(0)}$, the perturbed source $T_{\mu\nu}$ can be found using standard approximation methods. For example, we may write a source parameterized by λ as

$$T_{\mu\nu} = T_{\mu\nu}^{(0)} + \lambda T_{\mu\nu}^{(1)} + \lambda^2 T_{\mu\nu}^{(2)} + \dots \quad (332)$$

where $\lambda = 0 \rightarrow T_{\mu\nu} = T_{\mu\nu}^{(0)}$, and we seek a perturbed metric

$$h_{\alpha\beta} = h_{\alpha\beta}^{(0)} + \lambda h_{\alpha\beta}^{(1)} + \lambda^2 h_{\alpha\beta}^{(2)} + \dots \quad (333)$$

up to the desired order in λ . For example, writing

$$\Phi = \frac{1}{\sqrt{r^2 + c^2(t - \xi^5\tau)^2}} \rightarrow \frac{1}{\sqrt{R^2 + c^2 A(t, \tau)}} \quad (334)$$

where

$$A(t, \tau) = (t - \xi^5\tau)^2 + \lambda \alpha(t, \tau) \quad (335)$$

leads to

$$\Phi \simeq \frac{1}{\sqrt{R^2 + c^2(t - \xi^5 \tau)^2}} - \frac{1}{2} \frac{\lambda \alpha(t, \tau)}{R^2 + c^2(t - \xi^5 \tau)^2} \quad (336)$$

to first order. In the linearized theory, we also have the first-order

$$R_{\mu\nu} = R_{\mu\nu}^{(0)} + \lambda R_{\mu\nu}^{(1)} \quad K_{\mu\nu} = K_{\mu\nu}^{(0)} + \lambda K_{\mu\nu}^{(1)} \quad (337)$$

where $R_{\mu\nu}^{(0)}$ and $K_{\mu\nu}^{(0)}$ are derived from $T_{\mu\nu}^{(0)}$. And since the evolution equations are linear and solved exactly by $h_{\mu\nu}^{(0)}$ for $T_{\mu\nu}^{(0)}$, the evolution equations for the perturbed metric reduce to the evolution equations for the perturbation itself.

$$\partial_5 h_{\mu\nu}^{(1)} = -2K_{\mu\nu}^{(1)} + \partial_\mu h_{5\nu}^{(1)} + \partial_\nu h_{5\mu}^{(1)} \quad (338)$$

$$\partial_5 K_{\mu\nu}^{(1)} = -\sigma \left[R_{\mu\nu}^{(1)} - \frac{8\pi G}{c^4} \left(S_{\mu\nu}^{(1)} - \frac{1}{2} \eta_{\mu\nu} S^{(1)} \right) \right] \quad (339)$$

where we may use (329) and (330) to express $R_{\mu\nu}^{(1)}$ and $K_{\mu\nu}^{(1)}$ in terms of the perturbation $\alpha(t, \tau)$. We note that in order to preserve the structure of the evolution equations as an initial value problem, we must preserve $\partial_5 \alpha(t, \tau)$ as an independent dynamical quantity and choose the initial values for $\alpha(t, 0)$ and $\partial_5 \alpha(t, 0)$.

Funding: This research received no external funding.

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

MDPI	Multidisciplinary Digital Publishing Institute
SHP	Stueckelberg-Horwitz-Piron formalism
ADM	Arnowitt, Deser, and Misner formalism
LHS	Left Hand Side
RHS	Right Hand Side

References

1. Horwitz, L.P. *Relativistic Quantum Mechanics*; Springer: Dordrecht, Netherlands, 2015. <https://doi.org/10.1007/978-94-017-7261-7>.
2. Horwitz, L.P.; Arshansky, R.I. *Relativistic Many-Body Theory and Statistical Mechanics*; 2053-2571, Morgan & Claypool Publishers, 2018. <https://doi.org/10.1088/978-1-6817-4948-8>.
3. Land, M.; Horwitz, L.P. *Relativistic classical mechanics and electrodynamics*; Morgan and Claypool Publishers, 2020.
4. Horwitz, L. *Concepts in Relativistic Dynamics*; World Scientific Publishing Co., 2023. Publisher Copyright: © 2023 by World Scientific Publishing Co. Pte. Ltd. All rights reserved., <https://doi.org/10.1142/11474>.
5. Isham, C. Canonical Quantum Gravity and the Problem of Time. Technical Report Imperial/TP/91-92/25, Blackett Laboratory, Imperial College, 1992. Lectures at the NATO Summer School in Salamanca.
6. Kiefer, C.; Peter, P. Time in Quantum Cosmology. *Universe* **2022**, *8*, 36. <https://doi.org/10.3390/universe8010036>.
7. Anderson, E. Problem of time in quantum gravity. *Annalen der Physik* **2012**, *524*, 757–786, [\[https://onlinelibrary.wiley.com/doi/pdf/10.1002/andp.201200147\]](https://onlinelibrary.wiley.com/doi/pdf/10.1002/andp.201200147). <https://doi.org/https://doi.org/10.1002/andp.201200147>.
8. Problem of time, Wikipedia Foundation, https://en.wikipedia.org/wiki/Problem_of_time, 2026.
9. Stueckelberg, E. La signification du temps propre en mécanique: Ondulatoire. *Helv. Phys. Acta* **1941**, *14*, 321–322. (In French).

10. Stueckelberg, E. Remarque a propos de la création de paires de particules en théorie de relativité. *Helv. Phys. Acta* **1941**, *14*, 588–594. (In French).
11. Horwitz, L.; Arshansky, R.; Elitzur, A. On the two aspects of time: The distinction and its implications. *Found. Phys.* **1988**, *18*, 1159–1193. <https://doi.org/10.1007/BF01889430>.
12. Land, M. Local metric with parameterized evolution. *Astronomische Nachrichten* **2019**, *340*, 983–988. <https://doi.org/10.1002/asna.201913719>.
13. Land, M. A 4+1 Formalism for the Evolving Stueckelberg-Horwitz-Piron Metric. *Symmetry* **2020**, *12*. <https://doi.org/10.3390/sym12101721>.
14. Land, M. A new approach to the evolving 4+1 spacetime metric. *Journal of Physics: Conference Series* **2021**, *1956*, 012010. <https://doi.org/10.1088/1742-6596/1956/1/012010>.
15. Land, M. Weak Gravitation in the 4+1 Formalism. *Universe* **2022**, *8*. <https://doi.org/10.3390/universe8030185>.
16. Land, M. A vielbein formalism for SHP general relativity. *Journal of Physics: Conference Series* **2023**, *2482*, 012006. <https://doi.org/10.1088/1742-6596/2482/1/012006>.
17. Land, M. An Evolving Spacetime Metric Induced by a ‘Static’ Source. *Symmetry* **2023**, *15*. <https://doi.org/10.3390/sym15071381>.
18. Land, M. A Spacetime Metric for the 4 + 1 Formalism. *Sci* **2025**, *7*. <https://doi.org/10.3390/sci7030086>.
19. Arnowitt, R.L.; Deser, S.; Misner, C.W. Republication of: The dynamics of general relativity. *General Relativity and Gravitation* **2004**, *40*, 1997–2027.
20. Fock, V. Proper time in classical and quantum mechanics. *Phys. Z. Sowjetunion* **1937**, *12*, 404–425.
21. Itzykson, C.; Zuber, J.B. *Quantum Field Theory*; McGraw-Hill: New York, 1980.
22. Halzen, F.; Martin, A.D. *Quarks and Leptons: An Introductory Course in Modern Particle Physics*; John Wiley and Sons: New York, 1984.
23. Horwitz, L.; Piron, C. Relativistic Dynamics. *Helv. Phys. Acta* **1973**, *48*, 316–326.
24. Saad, D.; Horwitz, L.; Arshansky, R. Off-shell electromagnetism in manifestly covariant relativistic quantum mechanics. *Found. Phys.* **1989**, *19*, 1125–1149.
25. Wheeler, J.A. *Geons, Black Holes and Quantum Foam: A Life in Physics*; W. W. Norton & Company, 2000.
26. Kaluza, T. Zum Unitatsproblem der Physik. *Sitzungsber. Preuss. Akad. Wiss* **1921**, p. 966–972.
27. Klein, O. Quantentheorie und funfdimensionale Relativitätstheorie. *Zeitschrift für Physik* **1926**, p. 895–906.
28. Horwitz, L.P. An elementary canonical classical and quantum dynamics for general relativity. *The European Physical Journal Plus* **2019**, *134*, 313. <https://doi.org/10.1140/epjp/i2019-12689-7>.
29. Horwitz, L.P. Spin and entanglement in general relativity. *The European Physical Journal Plus* **2021**, *136*. <https://doi.org/10.1140/epjp/s13360-020-00967-8>.
30. Land, M. Mass-Energy-Momentum Radiation in Stueckelberg-Horwitz-Piron (SHP) Electrodynamics. *Journal of Physics: Conference Series* **2019**, *1239*, 012005. <https://doi.org/10.1088/1742-6596/1239/1/012005>.
31. Weinberg, S. *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*; Wiley: New York, NY, 1972.
32. Yeppez, J. Einstein’s vierbein field theory of curved space, 2011, [[arXiv:gr-qc/1106.2037](https://arxiv.org/abs/gr-qc/1106.2037)].
33. Wheeler, J.A. Superspace and the nature of quantum geometrodynamics. pp 615-724 of *Topics in Nonlinear Physics. Zabusky, Norman J. (ed.). New York, Springer-Verlag New York, Inc., 1968. 1969.*
34. Blau, M. Lecture Notes on General Relativity. Technical report, Albert Einstein Center for Fundamental Physics, Universität Bern, 2020.
35. Land, M.; Horwitz, L. Green’s functions for off-shell electromagnetism and spacelike correlations. *Found. Phys.* **1991**, *21*, 299–310.
36. Misner, C.W.; Thorne, K.S.; Wheeler, J.A. *Gravitation*; San Francisco: W.H. Freeman and Co., 1973.
37. Land, M. The Particle as a Statistical Ensemble of Events in Stueckelberg–Horwitz–Piron Electrodynamics. *Entropy* **2017**, *19*, 234. <https://doi.org/10.3390/e19050234>.
- 38.ourgoulhon, E. 3+1 Formalism and Bases of Numerical Relativity. Technical report, Laboratoire Univers et Theories, C.N.R.S., 2007. Lectures given at the General Relativity Trimester held in the Institut Henri Poincare (Paris, Sept.-Dec. 2006) and at the VII Mexican School on Gravitation and Mathematical Physics (Playa del Carmen, Mexico, 26. Nov. - 2 Dec. 2006).

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.