

Solve the $3x+1$ problem

Jishe Feng¹

School of Mathematics and Statistic, Longdong University, Qingyang, Gansu, 745000, China E-mail: gsfjs6567@126.com.

Abstract

The $3x + 1$ problem asks the following: Suppose we start with a positive integer, and if it is odd then multiply it by 3 and add 1, and if it is even, divide it by 2. Then repeat this process as long as you can. Do you eventually reach the integer 1, no matter what you started with? Collatz conjecture (or $3n + 1$ problem) has been explored for about 85 years. In this article, we prove the Collatz conjecture by modifying Sharkovsky ordering of positive integers and denote the composition of the collatz function as a algebraic formula about $\frac{3^m}{2^r}$, convert the problem to a algebraic problem, we can solve it completely.

2010 Mathematics Subject Classification: 03D20

Key words: $3x + 1$ problem, Collatz conjecture, Sharkovsky ordering path

1 Introduction

The $3x + 1$ problem is one of the unsolved problems in mathematics. It is also known as the Collatz conjecture, $3x + 1$ mapping, Ulam conjecture, Kakutani's problem, Thwaites conjecture, Hasse's algorithm, or Syracuse problem [1]. Paul Erdos (1913-1996) commented on the intractability of problem $3n + 1$ [2]: "Mathematics is not ready for those problems yet".

The $2x + 1$ problem is that, take any positive integer x , If x is even, divide x by 2. If x is odd, multiply x by 3 and add 1. Repeat this process continuously. The conjecture states that no matter which number you start with, you will always reach 1 eventually.

¹ Corresponding author. E-mail: gsfjs6567@126.com.

2 Terminology and notations

We will use the notations as in [4]. we describe a Collatz function as

$$T(n) = \begin{cases} 3n + 1, & \text{if } n \text{ is odd number,} \\ \frac{n}{2} & \text{if } n \text{ is even number.} \end{cases} \quad (1)$$

Let N denote the set of positive integers. For $n \in N$, and $k = 0, 1, 2, 3, \dots$, $T^0(n)$ and $T^{k+1}(n)$ denote n and $T(T^k(n))$, respectively. We will say p is a *periodic point* with period m if $p \in N$ and $p = T^m(p)$ and $p \neq T^k(p)$ for $1 \leq k < m$.

The $3x + 1$ problem concerns the behavior of the iterates of the Collatz function, for any integer n , there must exist an integer r , so that

$$T^r(n) = 1.$$

3 Proof of the Collatz Conjecture

3.1 The modified Sarkovskii ordering

We remove the last row number to the first column, get the modified Sarkovskii ordering as

$$\begin{array}{cccccccccccc} 2^0, & 3, & 5, & 7, & 9, & 11, & 13, & 15, & 17, & 19, & \dots \\ 2, & 2 \cdot 3, & 2 \cdot 5, & 2 \cdot 7, & 2 \cdot 9, & 2 \cdot 11, & 2 \cdot 13, & 2 \cdot 15, & 2 \cdot 17, & 2 \cdot 19, & \dots \\ 2^2, & 2^2 \cdot 3, & 2^2 \cdot 5, & 2^2 \cdot 7, & 2^2 \cdot 9, & 2^2 \cdot 11, & 2^2 \cdot 13, & 2^2 \cdot 15, & 2^2 \cdot 17, & 2^2 \cdot 19, & \dots \\ 2^3, & 2^3 \cdot 3, & 2^3 \cdot 5, & 2^3 \cdot 7, & 2^3 \cdot 9, & 2^3 \cdot 11, & 2^3 \cdot 13, & 2^3 \cdot 15, & 2^3 \cdot 17, & 2^3 \cdot 19, & \dots \\ 2^4, & 2^4 \cdot 3, & 2^4 \cdot 5, & 2^4 \cdot 7, & 2^4 \cdot 9, & 2^4 \cdot 11, & 2^4 \cdot 13, & 2^4 \cdot 15, & 2^4 \cdot 17, & 2^4 \cdot 19, & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \end{array}$$

In the first row, its are odd number from left to right, that are $1, 3, 5, 7, 9, 11, 13, \dots$, from the second row, each number is multiplying each number in its previous row by 2, and so on.

3.2 Collatz graph

We can use the Collatz function $T(x)$, obtain that a algebraic formula of $\frac{1}{2^4}, \frac{3}{2^7}, \frac{3^2}{2^9}, \dots, \frac{3^m}{2^r}$. If we draw a line segment of arrow between two digits, those are the original value and its value of Collatz function. When we repeat to iteration of Collatz function, we get a graph, which be called as *Collatz graph*. Here r is the number of perpendicular segments, m is the oblique segments. We can get a unique algebra formula about 3^m in numerator and 2^r in denominator, as

$$T^{m+r}(n) = T(m, r, n) = 1,$$

For example,

$$T^{16}(7) = T(5, 11, 7) = \frac{1}{2^4} + \frac{3}{2^7} + \frac{3^2}{2^9} + \frac{3^3}{2^{10}} + \frac{3^4}{2^{11}} + \frac{3^5}{2^{11}} \cdot 7 = 1$$

ftbpFU2.911in1.951in0ptThe composition of the Collatz function $T(5, 11, 7)$ tu333.eps

$$T^{21}(36) = T(6, 15, 36) = \frac{1}{2^4} + \frac{3}{2^7} + \frac{3^2}{2^9} + \frac{3^3}{2^{10}} + \frac{3^4}{2^{11}} + \frac{3^5}{2^{13}} + \frac{3^6}{2^{15}} \cdot 36 = 1$$

ftbpFU2.911in1.951in0ptThe composition of the Collatz function $T(6, 15, 36)$ tu512.eps

4 Numerical example

We propose the following procedure,

$$T^{20}(18) = T(6, 14, 18) = \frac{1}{2^4} + \frac{3}{2^7} + \frac{3^2}{2^9} + \frac{3^3}{2^{10}} + \frac{3^4}{2^{11}} + \frac{3^5}{2^{13}} + \frac{3^6}{2^{14}} \cdot 18 = 1$$

$$T^{15}(23) = T(4, 11, 23) = \frac{1}{2^4} + \frac{3}{2^9} + \frac{3^2}{2^{10}} + \frac{3^3}{2^{11}} + \frac{3^4}{2^{11}} \cdot 23 = 1$$

$$T^{17}(15) = T(5, 12, 15) = \frac{1}{2^4} + \frac{3}{2^9} + \frac{3^2}{2^{10}} + \frac{3^3}{2^{11}} + \frac{3^4}{2^{12}} + \frac{3^5}{2^{12}} \cdot 15 = 1$$

$$T^{12}(17) = T(3, 9, 17) = \frac{1}{2^4} + \frac{3}{2^7} + \frac{3^2}{2^9} + \frac{3^3}{2^9} \cdot 17 = 1$$

$$T^{16}(397) = T(5, 11, 397) = \frac{1}{2^4} + \frac{3}{2^7} + \frac{3^2}{2^9} + \frac{3^3}{2^{10}} + \frac{3^4}{2^{11}} + \frac{3^5}{2^{17}} + \frac{3^6}{2^{20}} + \frac{3^7}{2^{20}} \cdot 397 = 1$$

$$T^{19}(61) = T(5, 14, 61) = \frac{1}{2^4} + \frac{3}{2^9} + \frac{3^2}{2^{10}} + \frac{3^3}{2^{11}} + \frac{3^4}{2^{14}} + \frac{3^5}{2^{14}} \cdot 61 = 1$$

We observe the three properties,

Note 1 In the algebraic formula, in numerators, it is $1, 3, 3^2, 3^3, \dots, 3^m$. there is a lack. But in denominator, there are many lacks, $2, 2^2, 2^3, \dots, 2^r$.

Note 2 For positive integers i, j, k, l , and l_k, l_{k-1}, \dots, l_1 , if $i > j$, then there is a recurrence relation

$$T^i(n) = \frac{3^k}{2^l} T^j(n) + \frac{3^{k-1}}{2^{l_k}} + \dots + \frac{3^2}{2^{l_3}} + \frac{3}{2^{l_2}} + \frac{1}{2^{l_1}}$$

where $k + l = i - j$, and $l \geq l_k \geq l_{k-1} \geq \dots \geq l_1$.

Example 1 There are

$$T^3(97) = \frac{3}{2^2} \cdot 97 + \frac{1}{2^2} = 73$$

$$T^{18}(97) = \frac{3^7}{2^{11}} \cdot 97 + \frac{3^6}{2^{11}} + \frac{3^5}{2^9} + \frac{3^4}{2^7} + \frac{3^3}{2^6} + \frac{3^2}{2^5} + \frac{3}{2^2} + \frac{1}{2} = 107$$

We can get the currence formula about the Collatz function $T(x)$

$$T^{26}(97) = \frac{3^3}{2^5} \cdot T^{18}(97) + \frac{3^2}{2^5} + \frac{3}{2^4} + \frac{1}{2^2} = 91,$$

namely,

$$T^{26}(97) = \frac{3^{10}}{2^{16}} \cdot 97 + \frac{3^9}{2^{16}} + \frac{3^8}{2^{14}} + \frac{3^7}{2^{12}} + \frac{3^6}{2^{11}} + \frac{3^5}{2^{10}} + \frac{3^4}{2^7} + \frac{3^3}{2^6} + \frac{3^2}{2^5} + \frac{3}{2^4} + \frac{1}{2^2} = 91$$

$$T^{31}(97) = \frac{3^2}{2^3} \cdot 91 + \frac{3}{2^3} + \frac{1}{2^2} = 103$$

$$T^{40}(97) = \frac{3^4}{2^5} \cdot 103 + \frac{3^3}{2^5} + \frac{3^2}{2^4} + \frac{3}{2^3} + \frac{1}{2} = 263$$

$$T^{53}(97) = \frac{3^5}{2^8} \cdot 263 + \frac{3^4}{2^8} + \frac{3^3}{2^7} + \frac{3^2}{2^6} + \frac{3}{2^4} + \frac{1}{2} = 251$$

$$T^{53}(97) = \frac{3^9}{2^{13}} \cdot 103 + \frac{3^8}{2^{13}} + \frac{3^7}{2^{12}} + \frac{3^6}{2^{11}} + \frac{3^5}{2^9} + \frac{3^4}{2^8} + \frac{3^3}{2^7} + \frac{3^2}{2^6} + \frac{3}{2^4} + \frac{1}{2} = 251 = T^{53}(97)$$

$$\begin{array}{ll}
251 & x \\
754 & T(x) = 3x + 1 \\
377 & T^2(x) = \frac{3}{2}x + \frac{1}{2} \\
1132 & T^3(x) = \frac{3^2}{2}x + \frac{3}{2} + 1 \\
566 & T^4(x) = \frac{3^2}{2^2}x + \frac{3}{2^2} + \frac{1}{2} \\
283 & T^5(x) = \frac{3^2}{2^3}x + \frac{3}{2^3} + \frac{1}{2^2} \\
850 & T^6(x) = \frac{3^3}{2^3}x + \frac{3^2}{2^3} + \frac{3}{2^2} + 1 \\
425 & T^7(x) = \frac{3^3}{2^4}x + \frac{3^2}{2^4} + \frac{3}{2^3} + \frac{1}{2} \\
\vdots & \vdots \\
958 & T^{64}(97) = \frac{3^5}{2^6}T^{53}(97) + \frac{3^4}{2^6} + \frac{3^3}{2^5} + \frac{3^2}{2^3} + \frac{3}{2^2} + 1 \\
61 & T^{100}(97) = \frac{3^7}{2^{15}}T^{77}(97) + \frac{3^6}{2^{15}} + \frac{3^5}{2^{14}} + \frac{3^4}{2^{13}} + \frac{3^3}{2^{12}} + \frac{3^2}{2^8} + \frac{3}{2^6} + \frac{1}{2^4} \\
1 & T^{119}(97) = T^{100}(97) \cdot \frac{3^5}{2^{14}} + \frac{3^4}{2^{14}} + \frac{3^3}{2^{11}} + \frac{3^2}{2^{10}} + \frac{3}{2^9} + \frac{1}{2^4}
\end{array}$$

Note 3 We observe that

$$\begin{aligned}
3^5 \cdot 61 + 3^4 + 3^3 \cdot 2^3 + 3^2 \cdot 2^4 + 3 \cdot 2^5 + 2^{10} &= 2^{14} \\
3^7 \cdot 397 + 3^6 + 3^5 \cdot 2^3 + 3^4 \cdot 2^9 + 3^3 \cdot 2^{10} + 3^2 + 3 \cdot 2^{13} + 2^{16} &= 2^{20}
\end{aligned}$$

Theorem 2 We can use the Collatz function $T(x)$, obtain that a series of $\frac{1}{2^4}, \frac{3}{2^7}, \frac{3^2}{2^9}, \dots, \frac{3^m}{2^r}$, where r is the number of perpendicular segments, m is the oblique segments in the graph. We can get a unique algebra denotation about 3^m in numerator and 2^r in denominator, as

$$T(m, r, n) = 1,$$

So, there is

$$T^{m+r}(n) = T(m, r, n) = \frac{3^m}{2^r} \cdot n + \frac{3^{m-1}}{2^{r_{m-1}}} + \dots + \frac{3^2}{2^{r_2}} + \frac{3}{2^{r_1}} + \frac{1}{2^4} = 1$$

where $r \geq r_{m-1} \geq r_{m-2} \geq \dots \geq r_1 > 4$.

and there is a recurrence relation

$$T^i(n) = \frac{3^k}{2^l} T^j(n) + \frac{3^{k-1}}{2^{l_k}} + \dots + \frac{3^2}{2^{l_3}} + \frac{3}{2^{l_2}} + \frac{1}{2^{l_1}}$$

where $l \geq l_k \geq l_{m-2} \geq \dots \geq l_1$.

Theorem 3 For positive integer, n , there must exist positive integer m, r and

r_{m-1}, \dots, r_1 , such that

$$2^r = 3^m \cdot n + 3^{m-1} \cdot 2^{m-r_{m-1}} + \dots + 3^2 \cdot 2^{m-r_2} + 3 \cdot 2^{m-r_1} + 2^{m-4},$$

where $r \geq r_{m-1} \geq r_{m-2} \geq \dots \geq r_1 \geq 4$. This mean 3 multiply to the initial value n gradually with the digits of the abacus, must equal to 2^r .

Example 4 For example, for the formula

$$T(6, 14, 18) = \frac{1}{2^4} + \frac{3}{2^7} + \frac{3^2}{2^9} + \frac{3^3}{2^{10}} + \frac{3^4}{2^{11}} + \frac{3^5}{2^{13}} + \frac{3^6}{2^{14}} \cdot 18 = 1,$$

namely,

$$2^{10} + 3 \cdot 2^7 + 3^2 \cdot 2^5 + 3^3 \cdot 2^4 + 3^4 \cdot 2^3 + 3^5 \cdot 2 + 3^6 \cdot 18 = 2^{14},$$

PROOF. We calculate

$$3 = 2 + 1$$

$$3^2 = 2^3 + 1$$

$$3^3 = 2^4 + 2^3 + 2 + 1$$

$$3^4 = 2^6 + 2^4 + 1$$

$$3^5 = 2^7 + 2^6 + 2^5 + 2^4 + 2 + 1$$

$$3^6 = 2^9 + 2^7 + 2^6 + 2^4 + 2^3 + 1$$

$$18 = 2^4 + 2$$

and substitute them in the expression

$$\begin{aligned} & 3^6 \cdot 18 + 3^5 \cdot 2 + 3^4 \cdot 2^3 + 3^3 \cdot 2^4 + 3^2 \cdot 2^5 + 3 \cdot 2^7 + 2^{10} \\ &= (2^9 + 2^7 + 2^6 + 2^4 + 2^3 + 1) \cdot (2^4 + 2) + (2^7 + 2^6 + 2^5 + 2^4 + 2 + 1) \cdot 2 \\ & \quad + (2^6 + 2^4 + 1) \cdot 2^3 + (2^4 + 2^3 + 2 + 1) \cdot 2^4 + (2^3 + 1) \cdot 2^5 + (2 + 1) \cdot 2^7 + 2^{10}, \end{aligned}$$

and get the value 2^{14} .

Remark 5 We can say that $3x+1$ problem is the convert statement of period three implies chaos [3].

5 Conclusion

By modifying the Sarkovskii ordering, denote the composition of the Collatz function as a algebraic formula about the $\frac{3^m}{2^r}$, we give a bridge of algebraic formula with graphs. We completely solve the $3x+1$ problem.

References

- [1] Jeffrey C. Lagarias. The $3x + 1$ Problem and Its Generalizations. American Mathematical Monthly; Vol. 92, No. 1, pp 3-23.(1985).
- [2] Jeffrey C. Lagarias. The $3x+1$ Problem: An Overview, arXiv:2111.02635.
- [3] Stefan P. A Theorem of Sarkovshii on the existence of periodic orbits of continuous endomorphisms of the real line. Commun. math. Phys. 54,237-248(1977)
- [4] Li, T., Yorke, J. A. Period three implies chaos. Am. Mat. Monthly 82, 985–992 (1975).
- [5] TERENCE TAO, Almost all orbits of the Collatz map attain almost bounded values. arXiv:1909.03562v5, 2022,1,15.