

Article

Not peer-reviewed version

Successful *Brachyspira hyodysenteriae* Eradication Through a Combined Approach of a Zinc Chelate Treatment and Adapted Management Measures

[Frédéric Vangroenweghe](#)*

Posted Date: 8 August 2025

doi: 10.20944/preprints202508.0621.v1

Keywords: eradication; *Brachyspira hyodysenteriae*; Zn-chelate; management measures; biosecurity



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Successful *Brachyspira hyodysenteriae* Eradication Through a Combined Approach of a Zinc Chelate Treatment and Adapted Management Measures

Frédéric Vangroenweghe^{1,2,*}

¹ Elanco Animal Health Belgium, BU Swine and Ruminants, Generaal Lemanstraat 55/3 (Building D, 1st floor), 2018 Antwerp, Belgium

² Ghent University, Faculty of Veterinary Medicine, Department of Department of Internal Medicine – Reproduction – Population Medicine, Unit Porcine Health Management, Salisburylaan 133, 9820 Merelbeke, Belgium

* Correspondence: vangroenweghe.frederic@telenet.be; Tel.: +32 477 558 562

Simple Summary

Brachyspira hyodysenteriae, the primary cause of swine dysentery, has recently been demonstrated less susceptible to several crucially important antimicrobials used for control and eradication of swine dysentery under field conditions. Therefore, we developed an alternative approach using a zinc chelate (IntraDysovinol® 499 mg/ml; IntraCare) treatment (14 days) combined with alternative management strategies (including partial depopulation of post-weaning facilities and improved external and internal biosecurity measures) and a thorough cleaning and disinfection protocol (including 2% NaOH) to eradicate *B. hyodysenteriae*. We evaluated this protocol in 18 *B. hyodysenteriae* infected swine farms under field conditions and concluded that the alternative approach was successful in the eradication of *B. hyodysenteriae* from these farms.

Abstract

Brachyspira hyodysenteriae is the primary cause of swine dysentery, characterized by bloody to mucoid diarrhea due to mucohaemorrhagic colitis in pigs. The disease primarily affects pigs during the growth and finishing stage. Control and prevention of *B. hyodysenteriae* consists of administration of antimicrobial drugs, besides management and adapted feeding strategies. A worldwide re-emergence of the disease has recently been reported with an increasing number of isolates demonstrating decreased susceptibility to several crucially important antimicrobials in the control of swine dysentery. This compromises the possibilities to eradicate *B. hyodysenteriae* from infected swine farms. A novel non-antibiotic zinc chelate has been reported to demonstrate positive effects on fecal quality and consistency, general clinical signs, average daily weight gain and *B. hyodysenteriae* excretion during and after a 6-day oral treatment. The objective of the present study was to evaluate the zinc chelate (IntraDysovinol® 499 mg/ml; IntraCare) within an eradication schedule with naturally occurring swine dysentery due to *B. hyodysenteriae* resistant to pleuromutilins under field conditions in Belgium. We evaluated a 14-day treatment schedule combined with alternative management measures (including partial depopulation of post-weaning facilities and improved external and internal biosecurity measures) and thorough cleaning and disinfection (including 2% NaOH) of the buildings and the sows from day 7 of treatment onwards. This alternative approach for *B. hyodysenteriae* eradication was evaluated on 18 swine farms over a 5-year period. All enrolled eradication programs were evaluated as successful at least 6-9 months after the finalization of the protocol. In conclusion, the zinc chelate product has an excellent potential for application within an eradication protocol of *B. hyodysenteriae* that were diagnosed resistant to pleuromutilins. The combined approach of zinc chelate treatment and alternative management measures demonstrated to be successful in the eradication of farrow-to-wean, farrow-to-finish and finishing swine farms under field conditions in Belgium.

Keywords: eradication; *Brachyspira hyodysenteriae*; Zn-chelate; management measures; biosecurity

1. Introduction

Brachyspira hyodysenteriae (*B. hyodysenteriae*) – a β -haemolytic Gram-negative oxygen-tolerant anaerobic spirochete – is the primary cause of swine dysentery, characterized by bloody to mucoid diarrhea due to mucohaemorrhagic colitis in pigs [1]. Swine dysentery primarily affects pigs during the growth and finishing period, while it is less frequently observed in weaner pigs or sows. Transmission of *B. hyodysenteriae* occurs through the fecal-oral route. The infection is associated with several risk factors such as introduction of colonized animals (carriers), poor external (quarantine, rodents, wild birds and other potential reservoirs) and internal biosecurity measures (adequate cleaning and disinfection protocols, mixing of age groups) [2]. The initial onset of the disease is characterized by loss of appetite and mild, yellow to grey-coloured diarrhea, further progressing to watery diarrhea with blood, mucus and pseudomembranes [3]. The disease has been reported in all major swine-rearing countries globally and contributes to significant economic losses, mainly due to growth losses, mortality, increased variation in pig weight and decreased feed conversion, in the swine industry worldwide.

Control and prevention of *B. hyodysenteriae* mainly consists of administration of antimicrobial drugs, besides management and adapted feeding strategies [4]. Currently, no commercial vaccines against *B. hyodysenteriae* are available [1], although some experimental vaccines, such as bacterins, subunit vaccines and live attenuated strains have been evaluated [5]. Recently, *B. hyodysenteriae* has been reported as a worldwide re-emerging disease with an increasing number of isolates having decreased susceptibility to several crucially important antimicrobials in the control of swine dysentery [6–19].

Overall, control and prevention of *B. hyodysenteriae* seems to become more challenging, due to the limited treatment options [2], the lack of effective preventive feeding strategies and the increased awareness on reduction of antimicrobial use in animal production [20]. A recent evaluation of different eradication strategies for *B. hyodysenteriae* under Belgian conditions revealed that only 40% of the eradication attempts were finally successful [21]. Three different eradication strategies had been enrolled, being total depopulation ($n = 2$), partial depopulation combined with antimicrobial treatment ($n = 6$), and only antimicrobial treatment without depopulation ($n = 2$). All farms implemented additional biosecurity measures during and after the eradication program. Based on these results, total depopulation was most successful (100%) followed by partial depopulation combined with an antimicrobial treatment (33%), whereas only antimicrobial treatment without depopulation did never succeed to eradicate *B. hyodysenteriae* (0%) [21]. A 3-week vacancy period for all rooms following cleaning and disinfection, combined with thorough rodent control and additional biosecurity measures appeared to be the key to a successful eradication. Unfortunately, no clear information on the *B. hyodysenteriae* resistance profile towards pleuromutilins was reported [21].

Due to the reported emergence of antimicrobial resistance, research focused on non-antibiotic alternatives to reduce *B. hyodysenteriae* infections has become more prominent. Adapted feeding strategies, including a high dietary concentration of inulin, have proven to reduce the incidence of swine dysentery due to *B. hyodysenteriae* in grower pigs [22]. A citrus extract commercialized as raw material and used as feed additive showed relevant in vitro bacteriostatic and bactericidal activity against *B. hyodysenteriae* at relatively low concentrations of 32 and 128 ppm, respectively [23]. Among others, zinc has been evaluated as a potential intervention to control *B. hyodysenteriae*. In vitro, the addition of either ZnSO₄ or CuSO₄ to the growth medium of *B. hyodysenteriae* caused complete inhibition of hemolytic activity in 3 culture cycles. Further research revealed that the inhibition of hemolysin was specifically mediated by Zn²⁺ [24]. A comparative study with ZnSO₄, Zn-methionine and ZnO only demonstrated a prophylactic effect of high concentrations of in-feed ZnO (2000 ppm or higher) against *B. hyodysenteriae* in a mouse challenge model for swine dysentery [25]. This resulted in the in-feed addition of ZnO in levels of 2500 ppm to reduce the incidence and severity of non-

specific cases of post-weaning diarrhea [26]. However, in 2017, the Committee for Medicinal Products of Veterinary Use concluded that the benefits of ZnO for the prevention of diarrhea in pigs do not outweigh the environmental risks of the product [27]. The recent withdrawal of the marketing authorization of ZnO by the European Commission limits the availability of effective alternatives to antimicrobial drugs. Consequently, there is a continuing need for new, effective, non-antibiotic innovations to further improve animal health and welfare and to help reducing economic losses due to *B. hyodysenteriae* infections in pigs.

Chelation of zinc with an anorganic molecule – instead of covalent binding of zinc to inorganic oxygen – reduces its environmental impact [28]. In addition, previous in vitro (unpublished data) and in vivo studies [28] have demonstrated that a novel organic zinc chelated complex – in relatively low concentrations – is potentially able to reduce adverse effects due to *B. hyodysenteriae* infections in pigs. The zinc chelate stimulated the in vitro growth of porcine intestinal cells and restored cell viability in the presence of *E. coli* toxins. Moreover, based on a decreased attachment of *E. coli* to cultured intestinal cells in an in vitro cell culture infection model, it is postulated that the zinc chelate might interact with the attachment of *B. hyodysenteriae* outer membrane proteins (OMPs) to the extracellular matrix proteins of the colonic epithelium, reducing the load of *B. hyodysenteriae* in the colon [28]. Another in vivo study demonstrated a positive effect of the zinc chelate to fecal quality and consistency, general clinical signs and average daily weight gain (ADWG) in *B. hyodysenteriae*-infected animals. Moreover, at the last treatment day, *B. hyodysenteriae* was not detectable by qPCR in most of the treated animals [28]. A field study using this zinc chelate (Intra Dysovinol® 499 mg/ml; IntraCare, Veghel, The Netherlands) demonstrated good clinical efficacy following a 6-day treatment with recovery of heavily *B. hyodysenteriae*-affected animals [29].

Based on these promising field experiences combined with the urgent need to develop alternative eradication strategies for swine farms infected with a highly resistant *B. hyodysenteriae* strain, we developed and evaluated a practical *B. hyodysenteriae* eradication program using a zinc chelate treatment combined with adapted management measures on 18 *B. hyodysenteriae*-infected swine farms in Belgium.

2. Materials and Methods

2.1. Selection of the Farms

Belgian pig farms with a positive diagnosis of *B. hyodysenteriae* that had been confirmed resistant to the pleuromutilin antimicrobials (tiamulin, valnemulin) were eligible for the field study. Following first contact, all relevant historic health data as well as diagnostic laboratory information, especially related to *B. hyodysenteriae* was collected. Subsequently, a farm visit was performed by the principal investigator together with the herd veterinarian to explain the eradication protocol in detail to the farmer. After the first farm visit, the principal investigator prepared a detailed, personalized farm-specific eradication protocol that was shared with the farm veterinarian and the farmer. Following internal discussion, the plan was refined and finalized during a second farm visit and specific critical control points were addressed and discussed on-farm. A total of 18 swine herds, of which 15 sow herds and 3 finishing herds, enrolled in this field study (Table 1).

2.2. Farm Description and Diagnostic Information About *B. hyodysenteriae*

A general description of the farms before the implementation of the eradication programme are shown in Table 1. The farms were representative for Belgium in terms of size, management practices, biosecurity measures, housing conditions and nutrition. Mild to severe diarrhea was present in at least on production category (sows, post-weaned piglets, fattening pigs) on all farms. Most of the farms had been suffering from swine dysentery for 2 months to several years prior to the start of the field trial. The eradication protocol was enrolled in the farms included in the field trial between May 2019 and June 2024.

Table 1. Descriptive overview of the different *Brachyspira hyodysenteriae*-infected farms with their detailed management and therapeutic approach towards the eradication of *Brachyspira hyodysenteriae*.

Farm ID	Year	Farm Type	# sows	Depopulation ¹	Zn-Chelate Treatment Group
A	2019	Farrow-to-finish	225	Partial	Sows
B	2019	Farrow-to-finish	300	Partial	Sows
C	2019	Farrow-to-finish	150	Partial	Sows
D	2019	Farrow-to-finish	500	Total finishing	Piglet/finishers
E	2019	Farrow-to-finish	800	Partial	Sows
F	2019	Farrow-to-finish	500	Partial	Sows
G	2019	Finishing	-	Partial	Piglets
H	2019	Farrow-to-finish	800	Partial	Sows
I	2019	Farrow-to-finish	40	Partial	Sows/piglets ²
J	2020	Farrow-to-finish	500	Partial	Sows
K	2020	Finishing	-	Partial	Finishers
L	2020	Farrow-to-finish	400	Partial	Sows/piglets ²
M	2020	Farrow-to-wean	500	Partial	Sows/piglets ²
N	2020	Finishing	-	Partial	Finishers
O	2020	Farrow-to-finish	250	Partial	Sows
P	2020	Farrow-to-finish	230	Partial	Sows
Q	2024	Farrow-to-finish	800	Partial	Sows/piglets ²
R	2024	Farrow-to-finish	100	Partial	Sows/piglets ²
S	2024	Farrow-to-finish	300	Partial	Sows/piglets ²

¹ Partial depopulation means the removal of 2-3 batches of post-weaned piglets from the sow farm to create a sanitary vacancy between the piglets born from treated sows and the preceding *Brachyspira hyodysenteriae* infected post-weaned piglets to be able to run a thorough cleaning and disinfection protocol and omit re-infection through internal spread. ² In case of difficulty to consistently separate sow and piglet category due to specific farm design, both sows and piglets were treated with the Zn-chelate product.

The *B. hyodysenteriae* isolates obtained from the farms were tested for their susceptibility against two antimicrobials commonly used for treatment and eradication of swine dysentery [1], namely tiamulin and valnemulin. Therefore, the agar dilution test as previously described by Vyt and Hommez [19] was used. Isolated were considered resistant if MIC-values ($\mu\text{g/ml}$) were higher than 2 for both tiamulin and valnemulin, respectively [13; 30].

2.3. Description of the Eradication Strategies

The basic eradication strategy was similar on all farms and included a partial depopulation of the post-weaning facility to obtain at least 3 weeks of sanitary vacancy necessary for thorough cleaning and disinfection, a strict cleaning and disinfection protocol including a 2% NaOH intermediate step between standard cleaning and the final disinfection step of the premises (Table 2) and a specific cleaning and disinfection protocol for the sows (Table 3), a prolonged treatment with an oral zinc chelate product (IntraDysovinol® 499 mg/ml; IntraCare, Veghel, The Netherlands) to eradicate *B. hyodysenteriae* from the colon of infected animals (Table 4), and a series of relevant additional external and internal biosecurity measures to omit re-introduction or internal spread during the eradication process at sow level (Table 5).

Table 2. Detailed cleaning and disinfection protocol to sanitize the swine farm environment during the eradication process for *Brachyspira hyodysenteriae*.

Step N°	Description of Specific Action
1	Remove all fecal material manually by shovel and brush
2	Soak the surfaces from top to bottom using a detergent solution for at least 1 h

3	Clean the surfaces using a pressure washer
4	Rinse the cleaned surfaces from top to bottom to remove remaining fecal material prior to the next step
5	Apply a 2% NaOH solution (dilute 1 part of 30% NaOH to 14 parts of water) on all non-metal surfaces (concrete, plastic, ...) and let the solution soak for 20-30 min
6	Rinse the surfaces again from top to bottom with sufficient water to remove the 2% NaOH solution after the required contact period
7	Apply an efficacious disinfectant (containing an aldehyde component) from top to bottom to all surfaces and let the solution work for the recommended contact period (see product information sheet)
8	Rinse the surface from top to bottom to remove the disinfectant solution after the required contact period
9	Dry the surfaces through sufficient ventilation of the compartment for at least 24 h after the final rinsing step

Table 3. Detailed cleaning and disinfection protocol to sanitize the sows during the eradication process for *Brachyspira hyodysenteriae*.

Step N°	Description of Specific Action
1	Soak the sows with a suitable shampoo and warm water (30-35°C)
2	Rinse the sows after the minimal required contact time
3	Specifically focus on thorough cleaning of the hoof section of the limbs
4	Disinfect the entire sow with a suitable disinfectant for topical use in animals
5	Move the cleaned and disinfected sows to a clean section of the gestation room

Table 4. Treatment protocol using a zinc chelate product to eradicate *Brachyspira hyodysenteriae* from an infected farm.

Gestating and Lactating Sow Treatment Protocol	
Determine the water intake of the sows one day prior to the start of the zinc chelate treatment	
Calculate the daily dose for the sow group with the following formula	
S = number of gestating sows	
BW = estimated/measured average body weight	
D = dose of zinc chelate product (0.023 ml/kg BW)	$V = S * BW * D$
V = volume of zinc chelate product to be administered daily	
Example for 200 sows weighing 245 kg	1127 ml
Dose the product continuously 24/24 for at least 15 days	
7 days prior to the start of cleaning & disinfection of the environment and sow washing protocol	
Extend for another 7 days after the finalization of the above cleaning & disinfection protocol	
Dose gestating and lactating sows separately due to the major difference in daily water consumption	
Post-weaned piglet treatment protocol	
Determine the water intake of the post-weaned piglets one day prior to the start of the zinc chelate treatment	
Calculate the daily dose for the sow group with the following formula	
S = number of post-weaned piglets	
BW = estimated/measured average body weight of the group	
D = dose of zinc chelate product (0.023 ml/kg BW)	$V = S * BW * D$
V = volume of zinc chelate product to be administered daily	
Example for 1000 piglets ranging from 6 – 25 kg (average at 17 kg)	391 ml

Dose the product continuously 24/24 for at least 14 days to clear all potential infection from the colon of the affected piglets

Table 5. Overview of all relevant additional management measures that should be implemented before and during the eradication process for *Brachyspira hyodysenteriae*.

Optimize External Biosecurity Measures to Prevent New Infection Entry
Create a hygienic lock with a one-way entry direction including a strict entry protocol Farm workers and external visitors ALL follow a strict entry protocol Provide clean boots in different sizes Thorough boot cleaning and disinfection protocol through installation of disinfection trays for boots using an efficient disinfectant Loading facilities for reform sow removal with strict one-way protocol Removal of dead animals with a strict one-way and end-of-day protocol Critical evaluation of manure removal process
Optimize internal biosecurity measures to prevent infection spread from 'infected' to 'clean' batches during the <i>Brachyspira hyodysenteriae</i> eradication
Provide separate sets of boots / coveralls for different animal categories <ul style="list-style-type: none"> ✓ Gestating sows ✓ Lactating sows ✓ Post-weaned piglets ✓ Fattening pigs ✓ Gilt development unit Strict walking lines among animal categories No return from older to younger age categories with the same clothing & boots Strict fly control in the premises Strict bird control in and around the premises Strict rodent control in and around the premises

2.4. Sample Size Calculations

The sample size required to achieve a 90% level of confidence of population freedom of disease was calculated using hypergeometric approximation method [31]. For this calculation, the test's sensitivity (90%) and specificity (100%), within-herd prevalence (5%; [32; 33]), prior confidence of population freedom (75%), probability of introduction (1%) and indicative population size (150-800 sows) were used. The results of this calculation showed that sample size varied between 35 samples for 150 sows and 41 samples for 800 sows to achieve a 95% confidence of freedom.

2.5. Sampling Protocol to Monitor the Infection Status Post-Eradication

Once the entire eradication programme had been completed, the farms were monitored for *B. hyodysenteriae* in the feces and for the presence of clinical signs of swine dysentery. Individual fecal samples were collected by the farm veterinarian between 6 and 9 months after the finalization of the eradication programme. Additional periodic follow-up visits were performed on a 2-monthly basis by the farm veterinarian to monitor clinical signs for a period of 1 year.

In the sow herds random samples were collected from 36 (n = 12 pools) to 42 (n = 14 pools) sows, whereas in the finishing herds a standard number of 42 (n = 14 pools) fattening pigs were sampled. Sampling was mainly performed at stressful moments (weaning, moving production group, change of diet) to increase the likelihood of *B. hyodysenteriae* detection.

The samples were transported cooled in closed plastic recipients to the laboratory within 2 h after collection.

2.6. Analysis of Fecal Samples

Individual fecal samples were pooled for analysis into pools of 3 individual samples [34]. The pooled samples were investigated using real-time PCR for the presence of *B. hyodysenteriae* (BactoReal® Kit *Brachyospira hyodysenteriae*, Ingenetix, Vienna, Austria). The PCR test detects the NADH oxidase gen (*nox*) of *B. hyodysenteriae* [35].

The samples were also used for culture of *B. hyodysenteriae* as previously described [21].

3. Results

3.1. *Brachyospira Hyodysenteriae* Sensitivity Testing

The antimicrobial sensitivity data revealed MIC-values varying between 4 µg/ml to > 16 µg/ml for both tiamulin and valnemulin. Therefore, all farms enrolled in the field trial had a moderately to highly resistant *B. hyodysenteriae* strain that did not respond to pleuromutilin treatment anymore.

3.2. Eradication Success

All farms enrolled into the field study were demonstrated *B. hyodysenteriae*-negative at diagnostic sampling on 6 to 9 months after the finalization of the eradication protocol. In 17 out of the 18 farms, a partial depopulation was implemented at the level of the post-weaning facility, creating a sanitary vacancy of at least 3 weeks to permit the farmer to perform a thorough cleaning and disinfection protocol. On one farrow-to-finish farm, due to the broad extent of the *B. hyodysenteriae* infection within the fattening unit, a total depopulation of the fattening unit was implemented to optimize the eradication success through reduction of re-infection risk from 'infected' to 'clean' fattening pigs during the eradication process.

In all 15 farrow-to-wean or farrow-to-finish herds, an extended treatment with the zinc chelate product (Intra Dysovinol® 499 mg/ml; IntraCare) was implemented for the entire sow population present on the farm. In 5 of these farms, due to the specific farm building design, post-weaned piglets were also treated with the zinc chelate product to omit potential risk of re-introduction of *B. hyodysenteriae* from the piglets back into the sows due to poor internal biosecurity measures.

In the 3 finishing units, specific batches of piglets or finisher pigs were treated with the zinc chelate product to reduce the infection pressure on the site and to omit further spread while the premises were gradually emptied and a thorough cleaning and disinfection procedure was performed.

4. Discussion

To our knowledge, this is the first report of *B. hyodysenteriae* eradication using a non-antibiotic zinc chelate product. Others have reported a variable success rate for eradication of *B. hyodysenteriae* using different strategies with or without antimicrobial treatment under Belgian conditions [21]. Critical points identified in previous *B. hyodysenteriae* eradication strategies were presence of antimicrobial resistance, too short (10-14 days) sanitary vacancy period, non-compliance to eradication protocol or new introduction from outside the farm. To tackle these critical risk factors, we designed a strict eradication protocol with the four main pillars as presented below.

Farm-adapted *B. hyodysenteriae* eradication strategies were implemented in all farms (n = 18) that were enrolled in this field study. The farm-specific strategies were the implementation of a zinc chelate treatment of the sows and/or piglets combined with adapted management measures. Based upon monitoring 6 to 9 months after the finalization of the eradication program, eradication of *B. hyodysenteriae* was successful in all of the infected farms.

The need for an alternative approach towards *B. hyodysenteriae* eradication in this field trial was mainly due to the dramatic increase in antimicrobial resistance of the isolated *B. hyodysenteriae* strains under Belgian conditions during the last few years [19]. Moreover, some of the included farms

already performed an unsuccessful eradication attempt using pleuromutilins, before we were contacted to support them with our innovative *B. hyodysenteriae* eradication strategy.

The eradication program enrolled in these *B. hyodysenteriae*-infected farms consisted of 4 essential pillars, being a zinc chelate treatment, a thorough cleaning and disinfection protocol of both premises and sows, a partial depopulation by removal of one or more production batches of piglets or total depopulation of a grow/finishing production group, and the implementation of several relevant farm-specific external and internal biosecurity measures. An important point to mention is that all programs were initiated during the spring or summer period (between end of April and end of August), as environmental survival of *B. hyodysenteriae* is much lower during the warm season [3].

The first pillar of the eradication program, a zinc chelate treatment from 7 days prior to the start of the cleaning and disinfection protocol was essential to remove the pathogen from the intestinal tract of the *B. hyodysenteriae*-infected animals. As previously demonstrated, continuous (24 h) oral treatment with a zinc chelate (IntraDysovinol® 499 mg/l) resulted in *B. hyodysenteriae*-negative fecal samples after 6 days of treatment [28]. A prolonged, off-label treatment under the conditions of the current field trial for at least 14 days could therefore result in a complete clearance of the pathogen from the infected animals. The second period of at least 7 days aimed to protect the animals from potential re-infection during washing and movement from the *B. hyodysenteriae*-contaminated environment to the clean and disinfected environment, which happened from day 7 onwards in the eradication program. In all farrow-to-wean and farrow-to-finish farms, sows were treated with the zinc chelate product to eradicate *B. hyodysenteriae*. In some farms, it was decided to include piglets into this treatment protocol, mainly due to the farm layout, which made it impossible to correctly separate both animal categories in a safe way to omit potential re-introduction from the 'infected' post-weaned piglet side to the 'clean' sow side of the farm.

The second pillar of the eradication program, thorough cleaning and disinfection was enrolled as soon as the sows were at least 7 days under treatment. Besides the standard cleaning and disinfection steps, an additional step applying 2% NaOH (Brenntag, Deerlijk, Belgium) between cleaning and disinfection was introduced. The aim of this supplementary step was to dissolve biofilm layers that might be present on concrete housing elements. These biofilm layers might protect *B. hyodysenteriae* from the subsequent disinfection step and therefore increase the risk of residual infection in the environment. Application of the 2% NaOH should be performed using correct personal protection material, such as gloves and eye protection. After a 20 to 30 minute soaking with this NaOH, the product should be removed through rinsing, prior to the implementation of disinfection. The choice of disinfection product should be based on efficacy against *B. hyodysenteriae* as stated on the label. In our experience, products containing an aldehyde component (formaldehyde, glutaraldehyde or glyoxal) were suitable for this purpose.

The third pillar of the eradication program, partial or total depopulation of one or more animal production batches (piglets or fattening pigs) was implemented to create a sanitary vacancy of at least 3 weeks between already *B. hyodysenteriae*-negative batches, born from sows that underwent the entire eradication protocol, and still *B. hyodysenteriae*-positive batches, which were considered a potential source of re-infection. Moreover, the sanitary vacancy could optimize the cleaning and disinfection protocol through a prolonged dry empty period of the different compartments. Total depopulation of the fattening unit was only implemented in one farm, mainly due to the nature of the farm layout and complexity, which made only partial depopulation and the related sanitary vacancy difficult to implement correctly.

For the fourth and last pillar of the eradication program, additional external and internal biosecurity measures were advised based on an on-farm visit. These measures were farm-tailored according to the farm layout, specific farm protocols, and motivation of the farmer. The external biosecurity aimed to reduce the risk of re-introduction of *B. hyodysenteriae* from outside the farm and consisted of a strict entry protocol, including boot cleaning and disinfection, a strict control of animal (live and dead) removal protocols, and a strict hygiene during manure removal. The internal biosecurity aimed to reduce the risk of spreading *B. hyodysenteriae* from infected compartments on

the farm to already eradicated compartments through clear separation of different animal categories, strict walking lines between animal categories, and strict rodent, bird and fly control in and around the premises.

Overall, it is important to stress that the high success rate was due to the strict follow-up of the entire eradication program from the initial farm visit, throughout the different preparation steps towards biosecurity measures, until the final execution of the treatment protocol and all the additional management measures. From this field study, even in the presence of antimicrobial resistance to pleuromutilins, eradication of *B. hyodysenteriae* may still be possible using a non-antibiotic zinc chelate for oral application. However, the eradication program should not solely be based on treatment, since all accompanying measures might be considered as crucial for the success of the current protocol. Although treatment with a zinc chelate product might clear *B. hyodysenteriae* from the colon, potential sources of re-infection on the farm remain present and therefore all presented additional measures should be implemented.

5. Conclusions

In conclusion, a zinc chelate product has excellent potential for application within an eradication protocol of *B. hyodysenteriae* diagnosed resistant to pleuromutilins. The combined approach of zinc chelate treatment and alternative management measures demonstrated to be successful in the eradication of farrow-to-wean, farrow-to-finish and finishing swine farms under field conditions in Belgium.

Author Contributions: Conceptualization, F.V.; methodology, F.V.; validation, F.V.; formal analysis, F.V.; investigation, F.V.; data curation, F.V.; writing—original draft preparation, F.V.; writing—review and editing, F.V.; visualization, F.V. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding. The registered pharmaceutical product was commercially available to all cooperating swine farms and swine veterinarians.

Institutional Review Board Statement: Not applicable. In this section, you should add the Institutional Review Board Statement and approval number, if relevant to your study. You might choose to exclude this statement if the study did not require ethical approval. Please note that the Editorial Office might ask you for further information. Please add “The study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Review Board (or Ethics Committee) of NAME OF INSTITUTE (protocol code XXX and date of approval).” for studies involving humans. OR “The animal study protocol was approved by the Institutional Review Board (or Ethics Committee) of NAME OF INSTITUTE (protocol code XXX and date of approval).” for studies involving animals. OR “Ethical review and approval were waived for this study due to REASON (please provide a detailed justification).” OR “Not applicable” for studies not involving humans or animals.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data is unavailable due to privacy restrictions of the involved farms and veterinarians.

Acknowledgments: The author would like to acknowledge all swine veterinarians and swine veterinary practices that cooperated in gathering the study results from the field study.

Conflicts of Interest: The authors declare no conflicts of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Alvarez-Ordóñez, A.; Martínez-Lobo, F.J.; Arguello, H.; Carvajal, A.; Rubio, P. Swine dysentery: aetiology, pathogenicity, determinants of transmission and the fight against the disease. *Int J Environ Res Public Health* **2013**, *10*, 1927-1947. <https://doi.org/10.3390/ijerph10051927>.
2. Hampson, D.J.; Lugsomya, K.; La, T.; Phillips, N.D.; Trott, D.J.; Abraham, S. Antimicrobial resistance in *Brachyspira* – an increasing problem for disease control. *Vet Microbiol* **2019**, *229*, 59-71. <https://doi.org/10.1016/j.vetmic.2018.12.019>.
3. Hampson, D.J. Brachyspiral colitis. In: Diseases of Swine. Zimmerman, J.J.; Karriker, L.A.; Ramirez, A.; Schwarz, K.J.; Stevenson, G.W. Eds. 10th edition. Ames, IA; Wiley-Blackwell; **2012**, 680-696.
4. Lobová, D.; Čížek, A. Bactericidal efficacy of two disinfectants against *Brachyspira hyodysenteriae* and one feed supplement against *B. hyodysenteriae* and *B. pilosicoli*. *Vet Med – Czech* **2004**, *49*, 156-160.
5. Mahu, M.; Boyen, F.; Canessa, S.; Zavala-Marchan, J.; Haesebrouck, F.; Martel, A.; Pasmans, F. An avirulent *Brachyspira hyodysenteriae* strain elicits intestinal IgA and slows down spread of swine dysentery. *Vet Res* **2017**, *48*, 59-72. <https://doi.org/10.1186/s13567-017-0465-y>.
6. De Lorenzi, G.; Gherpelli, Y.; Luppi, A.; Pupillo, G.; Bassi, P.; Dottori, M.; Di Donato, A.; Merialdi, G.; Bonilauri, P. In vitro susceptibility of *Brachyspira hyodysenteriae* strains isolated in pigs in northern Italy between 2005 and 2022. *Res Vet Sci* **2024**, *168*, 105125-105130. <https://doi.org/10.1016/j.rvsc.2024.105152>.
7. Duinhof, T.F.; Dierickx, C.M.; Koene, M.G.J.; van Bergen, M.A.P.; Mevius, D.J.; Veldman, K.T.; van Beers-Schreurs, H.M.G.; de Winne, R.T.J.A. Multiresistentie bij *Brachyspira hyodysenteriae*-isolaten op een varkensvermeerderingsbedrijf in Nederland. *Tijdschrift voor Diergeneeskunde* **2008**, *133*, 604-608.
8. Hakimi, M.; Ye, F.; Stinman, C.C.; Sahin, O.; Burrough, E.R. Antimicrobial susceptibility of U.S. porcine *Brachyspira* isolates and genetic diversity of *B. hyodysenteriae* by multilocus sequence typing. *J Vet Diagn Invest* **2024**, *36*, 62-69. <https://doi.org/10.1177/10406387231212189>.
9. Hidalgo, A.; Carvajal, A.; Vester, B.; Pringle, M.; Naharro, G.; Rubio, P. Trends towards lower antimicrobial susceptibility and characterization of acquired resistance among clinical isolates of *Brachyspira hyodysenteriae* in Spain. *Antimicrob Ag Chemother* **2011**, *55*, 3330-3337. <https://doi.org/10.1128/AAC.01749-10>.
10. Joerling, J.; Barth, S.A.; Schlez, K.; Willems, H.; Herbst, W.; Ewers, C. Phylogenetic diversity, antimicrobial susceptibility and virulence gene profiles of *Brachyspira hyodysenteriae* isolates from pigs in Germany. *PlosOne* **2018**, *13*, e0190928. <https://doi.org/10.1371/journal.pone.0190928>.
11. Karlsson, M.; Aspán, A.; Landén, A.; Franklin, A. Further characterization of porcine *Brachyspira hyodysenteriae* isolates with decreased susceptibility to tiamulin. *J Med Microbiol* **2004**, *53*, 281-285. <https://doi.org/10.1099/jmm.0.05395-0>.
12. Kirchgässner, C.; Schmitt, S.; Borgström, A.; Wittenbrink, M.M. Antimicrobial susceptibility of *Brachyspira hyodysenteriae* in Switzerland. *Schweiz Arch Tierheilk* **2016**, *158*, 405-410. <https://doi.org/10.17236/sat00066>.
13. Lobová, D.; Smola, J.; Čížek, A. Decreased susceptibility to tiamulin and valnemulin among Czech isolates of *Brachyspira hyodysenteriae*. *J Med Microbiol* **2004**, *53*, 287-291. <https://doi.org/10.1099/jmm.0.05407-0>.
14. Mahu, M.; Pasmans, F.; Vranckx, K.; De Pauw, N.; Vande Maele, L.; Vyt, P.; Vandersmissen, T.; Martel, A.; Haesebrouck, F.; Boyen, F. Presence and mechanisms of acquired antimicrobial resistance in Belgian *Brachyspira hyodysenteriae* isolates belonging to different clonal complexes. *Vet Microbiol* **2017**, *207*, 125-132. <https://doi.org/10.1016/j.vetmic.2017.05.022>.
15. Massacci, F.R.; De Luca, S.; Cucco, L.; Tentellini, M.; Perreten, V.; Pezzotti, G.; Magistrali, C.F. Multiresistant *Brachyspira hyodysenteriae* shedding by pigs during the fattening period. *Vet Rec* **2018**, *183*, 264-266. <https://doi.org/10.1136/vr.104886>.
16. Mirajkar, N.S.; Davies, P.R.; Gebhart, C.J. Antimicrobial susceptibility patterns of *Brachyspira* species isolated from swine herds in the United States. *J Clin Microbiol* **2016**, *54*, 2109-2119. <https://doi.org/10.1128/JCM.00834-16>.
17. Rugna, G.; Bonilauri, P.; Carra, E.; Bergamini, F.; Luppi, A.; Gherpelli, Y.; Magistrali, C.F.; Nigrelli, A.; Alborali, G.L.; Martelli, P.; La, T.; Hampson, D.J.; Merialdi, G. Sequence types and pleuromutilin susceptibility of *Brachyspira hyodysenteriae* isolates from Italian pigs with swine dysentery: 2003-2012. *Vet J* **2015**, *203*, 115-119. <https://doi.org/10.1016/j.tvjl.2014.10.033>.

18. Vereecke, N.; Botteldoorn, N.; Brossé, C.; Bonckaert, C.; Nauwynck, H.; Haesebrouck, F.; Boyen, F.; Maes, D.; Theuns, S. Predictive power of long-read whole-genome sequencing for rapid diagnostics of multidrug-resistant *Brachyspira hyodysenteriae* strains. *Microbiol. Spectr.* **2023**, *11*, e0412322. <https://doi.org/10.1128/spectrum.04123-22>.
19. Vyt, P.; Hommeez, J. Antimicrobial susceptibility of *Brachyspira hyodysenteriae* isolates compared with the clinical effect of treatment. *Flem. Vet. J.* **2006**, *75*, 279-285.
20. Speksnijder, D.C.; Jaarsma, A.D.C.; van der Gugten, A.C.; Verheij, T.J.M.; Wagenaar, J.A. Determinants associated with veterinary antimicrobial prescribing in farm animals in the Netherlands: a qualitative study. *Zoonosis Publ Health* **2015**, *62*, 39-51. <https://doi.org/10.1111/zph.12168>.
21. Neiryneck, W.; Boyen, F.; Chantziaras, I.; Vandersmissen, T.; Vyt, P.; Haesebrouck, F.; Dewulf, J.; Maes, D. Implementation and evaluation of different eradication strategies for *Brachyspira hyodysenteriae*. *Porc. Health Manag.* **2020**, *6*, 27-37. <https://doi.org/10.1186/s40813-020-00162-2>
22. Hansen, C.F.; Hernández, A.; Mansfield, J.; Hidalgo, A.; La, T.; Phillips, N.D.; Hampson, D.J.; Pluske, J.R. A high dietary concentration of inulin is necessary to reduce the incidence of swine dysentery in pigs experimentally challenged with *Brachyspira hyodysenteriae*. *Br J Nutr* **2011**, *106*, 1506-1513. <https://doi.org/10.1017/S000711451100208X>.
23. de Nova, P.J.G.; Carvajal, A.; Prieto, M.; Rubio, P. In vitro susceptibility of *Brachyspira hyodysenteriae* to a commercial citrus fruit extract. *Res Vet Sci* **2017**, *115*, 318-324. <https://doi.org/10.1016/j.rvsc.2017.06.010>.
24. Dupont, D.P.; Duhamel, G.E.; Carlson, M.P.; Mathiesen, M.R. Effect of divalent cations on hemolysin synthesis by *Serpulina (Treponema) hyodysenteriae*: inhibition induced by zinc and copper. *Vet Microbiol* **1994**, *41*, 63-73. [https://doi.org/10.1016/0378-1135\(94\)90136-8](https://doi.org/10.1016/0378-1135(94)90136-8).
25. Zhang, P.; Carlson, M.P.; Schneider, N.R.; Duhamel, G.E. Minimal prophylactic concentration of dietary zinc compounds in a mouse model of swine dysentery. *Anim Health Res Rev* **2001**, *2*, 67-74.
26. Poulson, H.D. Zinc oxide for weanling piglets. *Acta Agr Scand – Anim Sci* **1995**, *45*, 159-167.
27. European Medicinal Agency. Questions and answers on veterinary medicinal products containing zinc oxide to be administered orally to food-producing species. Outcome of a referral procedure under Article 35 of Directive 2001/82/EC (EMEA/V/A/118). 2017; EMA/394961/2017.
28. Lammers, G.; van Berkel, R.; Roijackers, D.; Brouwer-Middeltesch, H.; van Hout, J. Treatment of clinical *Brachyspira hyodysenteriae* with zinc chelate in pigs: a blinded, randomised controlled trial. *Vet Rec* **2019**, *185*, 659-665. <https://doi.org/10.1136/vetrec-2019-105523>.
29. Vangroenweghe, F.; Allais, L.; Van Driessche, E.; van Berkel, R.; Lammers, G.; Thas, O. Evaluation of a zinc chelate on clinical swine dysentery under field conditions. *Porc. Health Manag.* **2020**, *6*, 1-10. <https://doi.org/10.1186/s40813-019-0140-y>.
30. Burch, D.G. Pharmacokinetic, pharmacodynamic and clinical correlations related to the therapy of colonic infections in the pig and breakpoint determinations. *Pig J.* **2005**, *56*, 8-24.
31. Sergeant E.S.G. Epitools epidemiological calculators: Ausvet. **2018**. <http://epitools.ausvet.com.au>.
32. Duff, J.W.; Pittman, J.S.; Hammer, J.; Kinyon, J.M. Prevalence of *Brachyspira hyodysenteriae* in sows and suckling piglets. *J. Swine Health Prod.* **2014**, *22*, 71-77.
33. Lobert, S.; Zimmermann, W.; Burki, S.; Frey, J.; Nathues, H.; Scheer, P.; Doherr, M.; Stalder, U.; Zeeh, F. Occurrence of *Brachyspira hyodysenteriae* in multiplier pig herds in Switzerland. *Tierarztl. Prax. Ausg. G. Grosstiere Nutztiere* **2016**, *44*, 13-18. <https://doi.org/10.15653/TPG-150674>.
34. Fellström, C.; Zimmermann, U.; Aspan, A.; Gunnarsson, A. The use of culture, pooled samples and PCR for identification of herds infected with *Brachyspira hyodysenteriae*. *Anim. Health Res. Rev.* **2001**, *2*, 37-43.
35. Atyeo, R.F.; Stanton, T.B.; Jensen, N.S.; Suriyaarachichi, D.S.; Hampson, D.J. Differentiation of *Serpulina* species by NADH-oxidase gene (*nox*) sequence comparisons and *nox*-based polymerase chain reaction tests. *Vet. Microbiol.* **1999**, *67*, 47-60. [https://doi.org/10.1016/s0378-1135\(99\)00030-9](https://doi.org/10.1016/s0378-1135(99)00030-9).

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.