

Article

Not peer-reviewed version

# Epidemiology of Suicide Mortality in Paraguay from 2005 to 2019: A Descriptive Study

Ethel Santacruz , Derlis Duarte-Zoilan , Gilda Benitez Rolandi , Felicia Canete , Dins Smits , [Noel C. Barengo](#) , [Guillermo Seguera](#) \*

Posted Date: 30 October 2023

doi: 10.20944/preprints202310.1849.v1

Keywords: Suicide; trends; mental health; Paraguay; potential years of life lost



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Article

# Epidemiology of Suicide Mortality in Paraguay from 2005 to 2019: A Descriptive Study

Ethel Santacruz <sup>1</sup>, Derlis Duarte-Zoilan <sup>1</sup>, Gilda Benitez <sup>1,2</sup>, Felicia Cañete <sup>1,2</sup>, Dins Smits <sup>3</sup>,  
Noël C Barengo <sup>4</sup> and Guillermo Sequera <sup>1,2,\*</sup>

<sup>1</sup> Dirección de Vigilancia de Enfermedades No Transmisibles (DVENT), Ministerio de Salud y Bienestar Social, Paraguay

<sup>2</sup> Catedra de Salud Pública, Facultad de Ciencias Médicas, Universidad Nacional de Asunción, Paraguay

<sup>3</sup> Faculty of Medicine, Riga Stradins University, Riga, Latvia; Dins.Smits@rsu.lv

<sup>4</sup> Department of Medical Education, Herbert Wertheim College of Medicine, Florida International University, Miami, Florida, 33199, USA; Escuela Superior de Medicina, Universidad Nacional de Mar del Plata, Mar del Plata, Provincia de Buenos Aires, 7600, Argentina; nbarengo@fiu.edu

\* Correspondence: gsequera@med.una.py

**Abstract:** Suicide is an important public health problem, fundamentally affecting the younger population and responding to multiple biological, psychological, and social causes. The objective of this study was to characterize changes suicide mortality, suicide methods and years of potential life lost during 2005 and 2019 in Paraguay. This observational, descriptive study used data from the Vital Statistics Information Subsystem of the Ministry of Public Health and Social Welfare. The average mortality rate from suicide was 4.9 per 100,000 inhabitants, with an increase from 4.2 between 200-2009 to 5.8 during 2014 and 2019. Suicide was more common in men (75%) than in women. In men, the highest mortality rate was observed among the 20-24 years-old whereas in women the age most affected were the 15–19-year-old age group. The most used method for suicide was strangulation. The most frequent places of suicide occurrence were at home (73%). The seasonality of suicide occurrence showed a slight increase in the spring-summer months compared with autumn winter (53% vs 47%). The rate of potential years of life lost statistically significantly increased from 2005 to 2019. Public health measures need to be implemented to investigate the underlying reasons and to implement interventions in the population to decrease suicide mortality in Paraguay.

**Keywords:** Suicide; trends; mental health; Paraguay; potential years of life lost

## 1. Introduction

At a global level, suicide is a complex public health problem, fundamentally affecting the younger population and responding to multiple biological, psychological, and social causes [1,2]. It presents a significant global burden of disease for all societies [3,4]. It has been estimated that suicides were responsible for over 700,000 deaths worldwide in 2019 [5]. This represents approximately 1.3% of all deaths globally, making it the 17th leading cause of death in 2019 [5]. Furthermore, in 2019 the global age-standardized suicide mortality rate was estimated to be 9.0/100,000 population with a higher rate in men than women (12.6 in males vs 5.4 in females) [4]. Decreasing trends in suicide mortality were observed in most countries across the world. Unfortunately, the mortality of suicide shows an increasing trend in many countries and populations [4]. Specifically, the North, Central and South America experienced a significant increase in suicide mortality over the past 20 years [4]. Illic et al reported that between 2000 and 2019 suicide mortality increased from 3.6/100,000 to 6.2/100,000 in Paraguay.

In Paraguay, although cardiovascular diseases and cancer continue to be the leading causes of death in adults, external causes such as traffic accidents and suicide are becoming more frequent every day and represent a significant burden of premature mortality [6,7]. In recent years there has

been a linear increase in deaths by suicide, especially among young people in Paraguay, but the characteristics of these events are still unknown [8]. The methods used for suicide vary according to country, culture, time, and demographic characteristics, such as race, sex, age and rural or urban environment [9]. The most common suicide methods used in Paraguay was intentional self-inflicted injury by hanging, strangulation, or suffocation (all 67.6%) [8]. Suicide also presents large geographical and temporal variations according to the countries [10–12]. Previous studies have suggested a relationship between suicide rates and other factors such as economic, environmental, and temporal factors such as seasonality [10,12,13]. Unfortunately, epidemiological data in years of potential life lost due to suicide, changes in suicide methods and seasonal patterns over time as well as trends of suicide mortality in different states of Paraguay is scant.

The objective of this study was to describe changes suicide mortality, suicide methods and years of potential life lost during 2005 and 2019 in Paraguay. An understanding of changes over time is important to guide public health professionals in developing, implementing, and monitoring public policies to decrease suicide attempts and suicide mortality.

## 2. Material and methods

### 2.1. Study design and data

A descriptive, observational study was conducted using record from the General Directorate of Strategic Health Information of the Paraguayan Ministry of Public Health and Social Welfare, which is the official institution responsible for the national health and disease statistics[14,15]. Validated records were from 2005 to 2019. The International Classification of Diseases 10 (ICD-10) codes X60 - X84 were used to define all death by suicide occurred in Paraguay during 2005 and 2019.

### 2.2. Measures of suicide mortality and characteristics

The variables available from the death certificates and used in this study were age, sex, date of death, level of education, main cause of death, state of residence of the deceased, place of suicide occurrence, place of suicide, area of residence and suicide method used by the victim. Sex at birth was recorded as male versus female. Age was categorized into the following age groups: (i) <10 years, (ii) 10-14 years-old, (iii) 15-24 years-old, (iv), 25-44 years-old, (v)45-64 years-old) and (vi) ≥ 65 years-old. Educational level was divided into none, primary, high school, College/University and unknown. Suicide method was defined according to ICD-10 codes and were as follows: poisoning (X60-X69), ICD-10 strangulation (X70), ICD-10 drowning (X71), ICD-10 firearm firing (X72-X74) and other ICD-10 (X76-X84). Information on the event site was defined as at home, at health-care institutions, public location/area, and others. The last recorded residence of the suicide victim was dichotomized into rural versus urban. The season of the suicide occurrence was recorded as spring/summer and autumn/winter. Finally, the time between 2005 and 2019 was divided into three intervals (i) 2005-2009, (ii) 2010-2014 and (iii) 2015-2019.

### 2.3. Data analysis

The data was analyzed using Microsoft Excel® and Epi-Info7 version 7.2.1.1. The data was analyzed according to time interval, age-groups, sex, and state. For the analysis of general trends in suicides, specific rates calculated by the number of deaths in each age group and sex at the national and departmental levels were used, divided by the annual population of the corresponding five-year period for each age group. Direct standardization was used to calculate age-specific mortality rates for the different time intervals. Population estimates were obtained from the Paraguayan General Office of Statistics, Survey and Censuses. National projection of population by sex and disaggregated by age group (2000-2050) based on the 2002 Census were used for the time interval 2006-2014. Population estimates starting from 2015 used data from the 2012 population census and projections for 2000-2025 [16]. The world population was used as the WHO standard population for rate adjustments (2000-2025). The years potential of life lost (YPLL) were calculated considering 0 years as a lower limit for the cut-off point and 70 years as the upper age limit for both sex per thousand

inhabitants. Suicide mortality rates were calculated per hundred thousand inhabitants as denominator. The trend of suicide mortality at the national level during 2005 to 2019 was analyzed with the Joinpoint Trend Analysis Software [17]. The software took suicide trend data at the national level (increases or decreases) and the rate or rate of annual average change and fit the simplest joinpoint model that the data allow. The minimum and maximum number of joinpoints were supplied by the investigators. The program started with the minimum number of joinpoints and tests whether more joinpoints were statistically significant and must be added to the model up to that maximum number. This enabled us to test whether an apparent change in trend was statistically significant. The tests of significance used the Monte Carlo Permutation method.

3. Results

3.1. Sociodemographic characteristics of deaths due to suicide

Table 1 shows the sociodemographic characteristics of deaths due to suicide in Paraguay during 2005 and 2019. In total, 4,776 deaths by suicide were recorded corresponding to 1.4% of all deaths recorded. A 62.1% increase between the first-time interval (2005-2009) and the last one (2015-2019) was observed. The percentage of men among all suicides increased from 71.2% (2005-2009) to 75% in 2015-2019. No statistically significant changes were observed in the distribution of age groups according to the three intervals. Most victims of suicide were recorded in the 15- to 24-year-old population followed by the 25 to 44 years-old age group. There was an increase in the percentage of victims with high school or University degrees during 2005 and 2019. Whereas 13.1% of all suicide victims during 2005 and 2010 had high school and 3.9% a university degree, the corresponding percentages of suicide victims with high school degree (16.6%) and university degree (5.3%) was higher in 2015-2019. There seemed to be a change in the site where suicide occurred during 2005 and 2019. The percentage of suicides that occurred in health care service decreased from 23% (2005-2009) to 14% (2010-2014) and 10% in 2015-2019. Suicides at home increased from 63% (2005-2009) to 73% (2015-2019). Whereas during 2005-2009 every second recorded suicide was by strangulation (50%), strangulation already accounted for three out of four suicides in 2015-2019. During the same time interval, the share of suicide by poisoning decreased from 19% to 3.5% and the sue of firearms from 24% to 15%. An increase in the prevalence of suicide in the urban region (from 72% to 77%) was observed between 2005 and 2019. Finally, 56-58% of the suicides occurred the during spring/summer period during 2005-2019.

Table 1. Sociodemographic characteristics of deaths due to suicide in Paraguay during 2005 and 2019.


Five-year periods	2005 - 2009			2010 - 2014			2015-2019		
	No.	%	95%CI	No.	%	95%CI	No.	%	95%CI
Total	1241			1513			2022		
Sex									
Man	884	71,2	(68.6-73.6)	1096	72,4	(70.1-74.6)	1517	75,0	(73.0-76.8)
Woman	357	28,8	(26.3-31.3)	417	27,6	(25.3-29.8)	505	25,0	(23.1-26.9)
Age Group									
< 10 years	24	1,9	(1.3-2.8)	10	0,7	(0.3-1.2)	2	0,1	(0.0-0.3)
10 - 14 years	63	5,1	(3.9-6.4)	74	4,9	(3.9-6.1)	83	4,1	(3.3-5.0)
15 - 24 years	447	36,0	(33.4-38.7)	567	37,8	(35.340.2)	714	35,5	(33.4-37.6)
25 - 44 years	406	32,7	(30.1-35.3)	508	33,8	(31.4-36.2)	661	32,9	(30.8-34.9)

45 - 64 years	202	16,3	(14,3-18,4)	236	15,7	(13,9-17,6)	380	18,9	(17,2-20,6)
> 65 years old	99	8,0	(6,6-9,6)	107	7,1	(6,9-8,5)	172	8,6	(7,4-9,8)
<b>Level of Education</b>									
None	46	3,7	(2,7-4,9)	68	4,5	(3,5-5,6)	94	4,7	(3,8-5,1)
Primary	615	49,6	(46,7-52,3)	682	45,1	(42,5-47,5)	886	43,8	(41,6-45,9)
High school	163	13,1	(11,3-15,1)	217	14,3	(12,6-16,2)	336	16,6	(15,0-18,3)
University	48	3,9	(2,9-5,0)	79	5,2	(4,2-6,4)	108	5,3	(4,4-6,4)
Doesn't know	369	29,7	(27,2-32,3)	467	30,9	(28,5-33,2)	598	29,6	(27,6-31,6)
<b>Types of Methods</b>									
Poisoning	230	18,5	(16,4-20,7)	95	6,3	(5,1-7,6)	71	3,5	(2,7-4,4)
Strangulation	622	50,1	(47,3-52,9)	1021	67,8	(65,0-69,8)	1508	74,6	(72,6-76,4)
Drowning	30	2,4	(1,7-3,4)	29	1,9	(1,3-2,7)	42	2,1	(1,5-2,8)
Firearm	298	24,0	(21,7-26,4)	301	19,9	(17,9-21,9)	303	15,0	(13,5-16,6)
Other causes*	61	4,9	(3,8-6,2)	67	4,4	(3,5-5,5)	98	4,9	(3,9-5,8)
*Sharps and fall trauma									
<b>Event site</b>									
Health Services	283	22,8	(20,5-25,2)	207	13,7	(12,0-15,5)	199	9,8	(8,6-11,2)
Home	779	62,8	(60,0-65,4)	1069	70,7	(68,3-72,8)	1471	72,8	(70,7-74,6)
Public area	125	10,1	(8,5-11,8)	77	5,1	(4,0-6,3)	113	5,6	(4,6-6,6)
Other*	54	4,4	(3,3-5,6)	160	10,6	(9,1-12,2)	239	11,8	(10,4-13,3)
*on the bridge, unspecified									
<b>Area of residence</b>									
Urban	889	71,6	(69,0-74,0)	1021	67,5	(65,0-69,8)	1559	77,1	(75,1-78,8)
Rural	352	28,4	(25,9-30,9)	492	32,5	(30,2-34,9)	463	22,1	(21,1-24,7)
<b>Season of the year</b>									
Spring/Summer	729	58,7	(54,0-61,2)	812	53,7	(50,1-57,4)	1112	55,0	(52,1-58,7)
Autumn/Winter	512	41,3	(39,5-42,5)	701	46,3	(43,8-49,7)	910	45,0	(40,1-48,7)

### 3.2. Standardized mortality rate for suicides

Of the 18 health regions, 12 experienced increases in suicide rates during the 2005-2019 period (Figure 1). The regions that had increases of suicide rates above the national average were Presidente Hayes, Central, Boquerón, Caazapá, Capital, Amambay, Canindeyú, and Concepción.






**Figure 1.** Crude suicide mortality rate per 100,000 inhabitants according to health regions and five-year period in Paraguay during 2005 and 2019.

The average suicide mortality rate nationwide during 2005 and 2019 was 4.9/100,000 with a gradual increase per five-year period, from 4.3/100,000 (2005-2009) to 5.7/100,000 between 2015 and 2019. Table 2 presents the standardized mortality rate for suicides according to 5-year time intervals in Paraguay during 2005 and 2019. The age standardized mortality rate for suicides increased in men from 6.1/100,000 (2005-2009) to 8.5/100,00 during 2015-2019. The corresponding increase in women was much lower (from 2.4/100,000 in 2005-2009 to 2.9/100,000 in 2015-2019). Also, the gap in mortality rates for suicide increased between men and women during 2005 and 2019.

**Table 2.** Standardized mortality rate for suicides according to 5-year time intervals in Paraguay during 2005 and 2019.

Sex	Year 2005 – 2009				Year 2010 – 2014			Year 2015 - 2019		
	Numbers	Age rate per 100 thousand	Standard age rate per 100 thousand		Numbers	Age rate per 100 thousand	Standard age rate per 100 thousand	Numbers	Age rate per 100 thousand	Standard age rate per 100 thousand
Men	828	5,6	6,1		1031	6,5	6,7	1419	8,3	8,5
Women	347	2,4	2,4		403	2,7	2,6	487	2,9	2,9
TOTAL	1175	4,7	4,3		1434	4,6	4,7	1906	5,6	5,7

Joinpoint analysis revealed that suicide rates linearly increased by 3.82% per year in Paraguay during 2005 and 2019 (Figure 2; p-value<0.001).



**Figure 2.** Average annual percentage change trend of suicide mortality 2005 -2019. Indicates that Annual Percent of Change (APC) is significantly different from zero at the  $\alpha = 0.05$  level. This figure is select model with 0 Joinpoints.

### 3.3. Rate of years of potential life lost by suicide

During the 15 years of observation, 172,612 years of potential life were lost due to suicide (Table 3). The YPLL rate due to suicide increased in the Paraguayan population between the study period. Whereas the YPLL was 1.5/1,000, the corresponding rates for 2010-2014 and 2015-2019 were 1.8/1,000 and 2.1/1,000. Whereas the YPLL rate increased in men from 2.1/1,000 (2005-2009) to 3.02/1,000 during 2015-2019, it remained unchanged in women (1-1.1/1,000) during the three time intervals.

**Table 3.** Rate of years of potential life lost by suicide according to five years by age group under 70 years, Paraguay during 2005 and 2019.

Five-year periods	2005 - 2009			2010 - 2014			2015 - 2019		
Sex	Number of deaths	YPLL <sup>1</sup> Number	YPLL Rate <sup>2</sup>	Number of deaths	YPLL Number	YPLL Rate	Number of deaths	YPLL Number	YPLL Rate
Men	828	30905	2,1	1031	39352	2,5	1419	51562	3,02
Women	347	14802	1,0	403	16627	1,1	487	19362	1,1
<b>Total</b>	<b>1175</b>	<b>45707</b>	<b>1,5</b>	<b>1434</b>	<b>55980</b>	<b>1,8</b>	<b>1906</b>	<b>70925</b>	<b>2,1</b>

<sup>1</sup>Years of Potential Life Lost; <sup>2</sup>per 1,000.

## 4. Discussion

Our data revealed that there was a linear increase in suicide mortality in Paraguay between 2005 and 2019. The age standardized mortality rate for suicides increased more in men than in women widening the gap in suicide mortality rates between men and women during 2005 and 2019. The YPLL due to suicide increased in men by 50% but remained stable in women. Three out of four

suicides occurred by strangulation whereas suicide by poisoning and firearms showed a decreasing trend.

Even though the suicide mortality rate has increased, it is still lower than the ones reported from Argentina, Brazil, Chile, and Uruguay [4]. However, a recent study that used preliminary data predicted a further increase of suicide mortality in Paraguay and that the expected number of suicides in the following years from 2023 to 2027 will range between 462 and 530 per year [8]. There are several reasons that may explain the increase in suicide mortality rate in Paraguay. First, the increase in suicide mortality may be due to an increase in mood and alcohol use disorders. These disorders are the most important modifiable predictors of suicide and [18,19] and both have shown to increase in the region [20–22]. Moreover, recent scientific evidence has suggested that acute alcohol use increases the risk of suicide attempts and that the risk of suicidal ideation, suicidal attempts, and death by suicide are each 2–3 times higher in people with alcohol use disorders compared with the general population [23,24]. A small part of the higher mortality rates observed may be explained by improved methods to differentiate suicide mortality from injury and accident-related mortality as some suicide deaths tend to be reported as accidental or underdetermined due to shame, stigma, or lack of evidence [25].

The data that men had a higher suicide mortality rate than women is consistent with the scientific literature [26,27]. The gender difference in suicide intention or mortality rates are known as the “Gender Paradox” [27]. A systematic review and meta-analysis of longitudinal studies revealed that women had a two-fold higher risk of suicide attempt than men, whereas the mortality rate of suicide was 2.5 times higher in men compared with women [28,29]. Another study revealed that women had higher rates of suicidal thinking, non-fatal suicidal behavior, and suicide attempts compared with men [30]. Thus, women seem to exhibit less serious intent to die than men [31]. Probably, the most important reasons for the observed difference between suicide attempts and suicide mortality between men and women is the method of suicide used. According to scientific evidence, men tend to choose more lethal suicide methods (firearms, hanging, and asphyxiation for instance) while women were more prone to overdose on medications or drugs than men [32]. Our data regarding higher suicide mortality rates in those with lower levels of education is consistent with previous studies. Several studies in different populations have revealed an inverse association between educational levels and suicide mortality rates in different populations [33–37]. It has been suggested that this may be due to better opportunities of being employed for people with higher levels of education resulting in higher level of social integration and lower suicide rates [33,37,38]. While having lower levels of education may increase the chance of unemployment leading to less responsibilities, more narrow networks, and less social ties [37].

Naturally, our study has some limitations. Under-reporting of suicide may introduce bias in the assessment of suicide mortality. In addition, the quality of mortality statistics (coverage, accuracy, and completeness of data) may have introduced some bias in estimating suicide mortality. For instance, data on suicide mortality in the indigenous population was not available. The department of vital statistics reported that there is an underreporting of deaths that accounts up to 18% in the last available year (2019). Also, the validity of death certification for suicide is a major problem as some of the suicides may be classified as undetermined intent or accident or violent deaths. The vital statistics department revealed that approximately 7% of deaths were incorrectly classified [39].

## 5. Conclusions

In conclusion, the increase in suicide mortality in Paraguay is of concern and an important cause of burden of premature death. Future analytical, observational studies should be conducted to identify what the most important predictors of suicide are. Also, more data is needed on the associations between social determinants of health and increasing suicide mortality trends. Moreover, a surveillance system monitoring the characteristics and trends of suicide in Paraguay needs to be implemented to assist political decision makers in developing policies and population-wide interventions to identify people at risk early to offer counseling and support. Finally, in order



to decrease the observed suicide mortality, interdisciplinary collaboration is needed including health professionals, political decision makers, educators and others.

**Author Contributions:** Conceptualization, ES, DDZ, GB, GS, DS, NB, FC; methodology, ES, DDZ, GB, GS, FC; software, ES, DDZ, formal analysis, ES, DDZ, BG, GS; investigation, ES, DDZ; resources, DDZ; data curation, ES, GS; writing—original draft preparation, ES, NB; writing—review and editing, ES, DDZ, GB, GS, DS, NB and FC; supervision, GS, FC; project administration, GS. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** The three basic principles of ethics in research involving human beings were respected: the principle of respect for persons, the principle of beneficence and the principle of justice. The data gathered for this study was consistently collected by the national surveillance system from the Ministry of Health. The database used ensures that the information is anonymized.

**Informed Consent Statement:** The information used for the study was derived from data that is regularly collected by the Paraguayan Ministry of Health's mortality surveillance system (SSIEV-<https://ssiev.msps.gov.py/>). The records analyzed for this study have been anonymized.

**Data Availability Statement:** The authors do not have permission to share their study data.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. WHO Global Tuberculosis Report 2014. Geneva, Switzerland: World Health Organization, 2014.
2. Fazel, S.; Runeson, B. Suicide. *New England Journal of Medicine* **2020**, *382*, 266–274, doi:10.1056/NEJMra1902944.
3. Naghavi, M. Global, Regional, and National Burden of Suicide Mortality 1990 to 2016: Systematic Analysis for the Global Burden of Disease Study 2016. *BMJ* **2019**, *l94*, doi:10.1136/bmj.l94.
4. Ilic, M.; Ilic, I. Worldwide Suicide Mortality Trends (2000–2019): A Joinpoint Regression Analysis. *World J Psychiatry* **2022**, *12*, 1044–1060, doi:10.5498/wjp.v12.i8.1044.
5. World Health Organization *Mental Health and Substance Use*; Geneva, 2021;
6. Directorate for Surveillance of Noncommunicable Diseases *Surveillance Bulletin Number 2 Noncommunicable Diseases and Risk Factors (2017)*; Asuncion, 2017;
7. Mansilla Izquierdo F *Suicide and Prevention*; 1st ed.; Palma de Mallorca, 2010;
8. Torales, J.; Barrios, I.; Melgarejo, O.; Tullo-Gómez, J.E.; Díaz, N.R.; O'Higgins, M.; Maggi, C.; Adorno, V.; Medina, A.; Villalba-Arias, J.; et al. Suicides among Adults in Paraguay: An 18-Year National Exploratory Study (2004–2022). *International Journal of Social Psychiatry* **2023**, 002076402311696, doi:10.1177/00207640231169650.
9. World Health Organization *Suicide Worldwide in 2019: Global Health Estimates*; Geneva, 2021;
10. Phillips, J.A.; Nugent, C.N. Suicide and the Great Recession of 2007–2009: The Role of Economic Factors in the 50 U.S. States. *Soc Sci Med* **2014**, *116*, 22–31, doi:10.1016/j.socscimed.2014.06.015.
11. Kim, N.; Mickelson, J.B.; Brenner, B.E.; Haws, C.A.; Yurgelun-Todd, D.A.; Renshaw, P.F. Altitude, Gun Ownership, Rural Areas, and Suicide. *American Journal of Psychiatry* **2011**, *168*, 49–54, doi:10.1176/appi.ajp.2010.10020289.
12. Yip, P.S.F.; Yang, K.C.T.; Qin, P. Seasonality of Suicides with and without Psychiatric Illness in Denmark. *J Affect Disord* **2006**, *96*, 117–121, doi:10.1016/j.jad.2006.05.012.
13. Laanani, M.; Ghosn, W.; Jougl, E.; Rey, G. Impact of Unemployment Variations on Suicide Mortality in Western European Countries (2000–2010). *J Epidemiol Community Health (1978)* **2015**, *69*, 103–109, doi:10.1136/jech-2013-203624.
14. DIGIES Sub-Sistema de Estadísticas Vitales Available online: <https://ssiev.msps.gov.py/> (accessed on 6 October 2023).
15. DIGIES Indicadores Básicos de Salud Available online: <http://portal.msps.gov.py/digies/publicaciones/indicadores/> (accessed on 6 October 2023).
16. Instituto Nacional de Estadística Proyección de La Población Por Sexo y Edad, Según Distrito, 2000–2025.
17. Surveillance Research Program, N.C.Institute. *Joinpoint Regression Program Version 4.8.0.1 - April 2020; Statistical Methodology and Applications Branch*; 2020;
18. Nordentoft, M. Absolute Risk of Suicide After First Hospital Contact in Mental Disorder. *Arch Gen Psychiatry* **2011**, *68*, 1058, doi:10.1001/archgenpsychiatry.2011.113.
19. Nock, M.K. Future Directions for the Study of Suicide and Self-Injury. *Journal of Clinical Child & Adolescent Psychology* **2012**, *41*, 255–259, doi:10.1080/15374416.2012.652001.

20. Mental Health and Substance Use *Depression and Other Common Mental Disorders: Global Health Estimates.*; 2017;
21. Manthey, J.; Shield, K.D.; Rylett, M.; Hasan, O.S.M.; Probst, C.; Rehm, J. Global Alcohol Exposure between 1990 and 2017 and Forecasts until 2030: A Modelling Study. *The Lancet* **2019**, *393*, 2493–2502, doi:10.1016/S0140-6736(18)32744-2.
22. Martinez-Ales, G.; Hernandez-Calle, D.; Khauli, N.; Keyes, K.M. Why Are Suicide Rates Increasing in the United States? Towards a Multilevel Reimagination of Suicide Prevention. In; 2020; pp. 1–23.
23. Borges, G.; Bagge, C.L.; Cherpitel, C.J.; Conner, K.R.; Orozco, R.; Rossow, I. A Meta-Analysis of Acute Use of Alcohol and the Risk of Suicide Attempt. *Psychol Med* **2017**, *47*, 949–957, doi:10.1017/S0033291716002841.
24. Darvishi, N.; Farhadi, M.; Haghtalab, T.; Poorolajal, J. Alcohol-Related Risk of Suicidal Ideation, Suicide Attempt, and Completed Suicide: A Meta-Analysis. *PLoS One* **2015**, *10*, e0126870, doi:10.1371/journal.pone.0126870.
25. BEAUTRAIS, A.L.; JOYCE, P.R.; MULDER, R.T. Risk Factors for Serious Suicide Attempts among Youths Aged 13 through 24 Years. *J Am Acad Child Adolesc Psychiatry* **1996**, *35*, 1174–1182, doi:10.1097/00004583-199609000-00015.
26. Cervantes, C.A.D.; Montaña, A.M.P. Estudio de La Carga de La Mortalidad Por Suicidio En México 1990-2017. *Revista Brasileira de Epidemiologia* **2020**, *23*, doi:10.1590/1980-5497202000069.
27. Canetto, S.S.; Sakinofsky, I. The Gender Paradox in Suicide. *Suicide Life Threat Behav* **1998**, *28*, 1–23, doi:10.1111/j.1943-278X.1998.tb00622.x.
28. Miranda-Mendizabal, A.; Castellví, P.; Parés-Badell, O.; Alayo, I.; Almenara, J.; Alonso, I.; Blasco, M.J.; Cebrià, A.; Gabilondo, A.; Gili, M.; et al. Gender Differences in Suicidal Behavior in Adolescents and Young Adults: Systematic Review and Meta-Analysis of Longitudinal Studies. *Int J Public Health* **2019**, *64*, 265–283, doi:10.1007/s00038-018-1196-1.
29. Vijayakumar, L. Suicide in Women. *Indian J Psychiatry* **2015**, *57*, 233, doi:10.4103/0019-5545.161484.
30. Crosby, A.E.; Han, B.; Ortega, L.A.G.; Parks, S.E.; Gfroerer, J.; Centers for Disease Control and Prevention (CDC) Suicidal Thoughts and Behaviors among Adults Aged ≥18 Years--United States, 2008-2009. *MMWR Surveill Summ* **2011**, *60*, 1–22.
31. Freeman, A.; Mergl, R.; Kohls, E.; Székely, A.; Gusmao, R.; Arensman, E.; Koburger, N.; Hegerl, U.; Rummel-Kluge, C. A Cross-National Study on Gender Differences in Suicide Intent. *BMC Psychiatry* **2017**, *17*, 234, doi:10.1186/s12888-017-1398-8.
32. Callanan, V.J.; Davis, M.S. Gender Differences in Suicide Methods. *Soc Psychiatry Psychiatr Epidemiol* **2012**, *47*, 857–869, doi:10.1007/s00127-011-0393-5.
33. Øien-Ødegaard, C.; Hauge, L.J.; Reneflot, A. Marital Status, Educational Attainment, and Suicide Risk: A Norwegian Register-Based Population Study. *Popul Health Metr* **2021**, *19*, 33, doi:10.1186/s12963-021-00263-2.
34. Bálint, L.; Osváth, P.; Rihmer, Z.; Döme, P. Associations between Marital and Educational Status and Risk of Completed Suicide in Hungary. *J Affect Disord* **2016**, *190*, 777–783, doi:10.1016/j.jad.2015.11.011.
35. Dare, A.J.; Irving, H.; Guerrero-López, C.M.; Watson, L.K.; Kolpak, P.; Reynales Shigematsu, L.M.; Sanches, M.; Gomez, D.; Gelband, H.; Jha, P. Geospatial, Racial, and Educational Variation in Firearm Mortality in the USA, Mexico, Brazil, and Colombia, 1990–2015: A Comparative Analysis of Vital Statistics Data. *Lancet Public Health* **2019**, *4*, e281–e290, doi:10.1016/S2468-2667(19)30018-0.
36. Næss, E.O.; Mehlum, L.; Qin, P. Marital Status and Suicide Risk: Temporal Effect of Marital Breakdown and Contextual Difference by Socioeconomic Status. *SSM Popul Health* **2021**, *15*, 100853, doi:10.1016/j.ssmph.2021.100853.
37. Choi, M.; Sempungu, J.K.; Lee, E.H.; Chang, S.-S.; Lee, Y.H. Single and Combined Effects of Marital Status, Education Attainment, and Employment Status on Suicide among Working-Age Population: A Case-Control Study in South Korea. *SSM Popul Health* **2022**, *19*, 101246, doi:10.1016/j.ssmph.2022.101246.
38. Denney, J.T.; Rogers, R.G.; Krueger, P.M.; Wadsworth, T. Adult Suicide Mortality in the United States: Marital Status, Family Size, Socioeconomic Status, and Differences by Sex \*. *Soc Sci Q* **2009**, *90*, 1167–1185, doi:10.1111/j.1540-6237.2009.00652.x.
39. Ministerio de Salud Pública y Bienestar Social *Indicadores Básicos de Salud, Paraguay*; Asunción, 2019;

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.