

1 *Review*

## 2 **Nerve, Muscle and Synaptogenesis**

3 **Lauren Eric Swenarchuk**

4 580 Christie Street, Suite 914, Toronto, Ontario M6G 3E3

5 \* Correspondence: l.swenarchuk@utoronto.ca, Tel.: 1-416-658-7747

6 **Abstract:** The vertebrate skeletal neuromuscular junction (NMJ) has long served as a model system  
7 for studying synapse structure, function and development. Over the last several decades a neuron-  
8 specific isoform of agrin, a heparan sulfate proteoglycan, has been identified as playing a central  
9 role in synapse formation at all vertebrate skeletal neuromuscular synapses. While agrin was  
10 initially postulated to be the inductive molecule that initiates synaptogenesis, this model has been  
11 modified in response to work showing that postsynaptic differentiation can develop in the absence  
12 of innervation, and that synapses can form in transgenic mice in which the agrin gene is ablated. In  
13 place of a unitary mechanism for neuromuscular synapse formation, studies in both mice and  
14 zebrafish have led to the proposal that two mechanisms mediate synaptogenesis, with some  
15 synapses being induced by nerve contact while others involve the incorporation of prepatterned  
16 postsynaptic structures. Moreover, the current model also proposes that agrin can serve two  
17 functions, to induce synaptogenesis and to stabilize new synapses, once these are formed. This  
18 review examines the evidence for these propositions, and concludes that it remains possible that a  
19 single molecular mechanism mediates synaptogenesis at all NMJs, and that agrin acts as a stabilizer,  
20 while its role as inducer is open to question. Moreover, if agrin does not act to initiate  
21 synaptogenesis, it follows that as yet uncharacterized molecular interactions are required to play  
22 this essential inductive role. Several alternatives to agrin for this function are suggested, including  
23 focal pericellular proteolysis and integrin signaling, but all require experimental validation.

24 **Keywords:** synapse; agrin; MuSK; Lrp4; neuromuscular junction

25

---

### 26 **1. Introduction**

27 How is the vertebrate skeletal neuromuscular junction (NMJ) made? Specifically, what  
28 molecular interaction initiates synaptogenesis when nerve and muscle meet? Despite almost half a  
29 century of work, a detailed picture has yet to emerge, and it is unclear as to whether a single process  
30 or multiple mechanisms are involved. This review does not aspire to resolve matters, but aims rather  
31 to outline the basic arguments at issue.

32 The NMJ acts as a simple relay, and unlike many central synapses, has no capacity for modifying  
33 its performance. However, this very simplicity, coupled with its large size and relative accessibility,  
34 has made it a favored object of study for elucidating basic principles of synapse structure and  
35 physiology that are relevant not only to its own functioning, but to that of central synapses as well.  
36 This expectation has certainly proved to be true historically, as studies of the NMJ gave us the first  
37 ever concept of a membrane receptor (“receptive substance”) [1], convincing evidence for the  
38 chemical nature of synaptic transmission [2] and, ultimately, the key physiological and  
39 ultrastructural evidence for the quantal mechanism of transmitter release [3].

40 Throughout millions of years of vertebrate evolution, the NMJ has remained relatively constant,  
41 with no evidence that selective pressures have led to alterations in molecular components of the  
42 synaptic machinery or to any evolutionary improvement in function. For example, few if any  
43 mammals can match frogs in their ability to leap from a standing start [4]. In studies of NMJ  
44 development, this structural and functional equivalence has been reflected in an implicit assumption,

45 invoking Occam's razor, that the molecular mechanisms underlying its formation have also been  
46 conserved throughout the vertebrate realm.

## 47 2. The agrin hypothesis

### 48 2.1. Historical development of the model

49 Early investigations in adult mammals and frogs demonstrated that transplanted nerves could  
50 form synapses on any part of the muscle surface. These studies usually involved inactivation of  
51 existing synapses, either by denervation [5] or treatment with botulinum toxin [6], although this  
52 inactivation was not always required [7]. In the late 1970s, these findings were extended through  
53 the integration of two lines of work. Using the *Xenopus in vitro* system, Anderson and Cohen [8]  
54 found that synapses formed anywhere on the surface of embryonic muscle, leading to the dispersal  
55 of existing clusters of acetylcholine receptors (AChRs) and their lateral migration to re-assemble  
56 along the path of nerve-muscle contact; this conclusion was supported by other *in vitro* studies in  
57 chick [9] and mammals [10], and was consistent with the earlier studies in adult muscle. Meanwhile,  
58 in U. J. McMahan's laboratory, work involving ablation of frog nerve and muscle demonstrated that  
59 regenerating nerves grew back to pre-existing sites of synaptic differentiation in the muscle basal  
60 lamina [11], and that AChR clusters developed under these same sites in the membrane of  
61 regenerating muscle [12]. Thus the basal lamina was implicated in two functions, providing  
62 termination signals for the nerve and AChR aggregation signals for the muscle. In subsequent work,  
63 the larger size of the muscle, permitting a relatively straightforward examination of the distribution  
64 of key synaptic markers, led to a principal focus on the latter capability.

65 These findings led to the formulation of the "agrin hypothesis" to account for postsynaptic  
66 differentiation [13]. According to this account, a protein, named agrin for its ability to induce AChR  
67 aggregation, is deposited by developing nerve in the muscle basal lamina, and not only induces the  
68 aggregation of AChRs under the nerve during embryogenesis, but also provides inductive signals to  
69 regenerating muscle. In the literature this function is typically characterized as "organizing" the  
70 postsynaptic membrane, with agrin designated as "organizer", whose action results in the  
71 localization of existing components to the point of nerve-muscle contact. It should be noted, however,  
72 that this inductive process may include the new synthesis of synaptic components as well, such as  
73 basal lamina constituents [14]. In the present discussion the term "inducer" is preferred to include  
74 this aspect, while explicitly referring to the initial interaction between nerve and muscle membranes.

75 Over a decade of studies led to agrin's isolation and cloning, a body of work which is especially  
76 notable for the underlying assumption of a unitary molecular mechanism, one that was meant to  
77 apply across the span of vertebrate classes and skeletal fiber types. While this point was rarely if ever  
78 emphasized explicitly, it was implicit in the potpourri of vertebrate species involved. Thus the  
79 biochemical isolation of agrin employed tissue from a ray, assays for activity were carried out in avian  
80 muscle cultures [15], and immunocytochemical localization was done in frog, ray and rat [16,17]. In  
81 addition, at a time when cloning a gene was a significant undertaking, agrin was cloned no fewer  
82 than three times, employing libraries from rat [18], chick [19] and ray [20]. In parallel with these  
83 efforts, the molecule which ultimately proved to be the key postsynaptic transducer of agrin-  
84 mediated signaling, a receptor tyrosine kinase, was cloned first from *Torpedo* [21], and subsequently  
85 from human, rat and mouse libraries, and named MuSK (muscle specific kinase) [22].

86 Ultimately this led to gene ablation studies with transgenic mice. Not only were mice lacking  
87 agrin unable to make synapses [23], but the cause of this defect was localized to the absence of one  
88 neuron-specific isoform possessing an exon coding for only eight amino acids [24]. In parallel, the  
89 role of MuSK was also delineated [25], and evidence was also adduced supporting the involvement  
90 of another postsynaptic protein as an essential cofactor [26]; this was later identified as low density  
91 lipoprotein receptor-related protein 4 [Lrp4; 27-29]. The agrin/MuSK findings were treated as a  
92 signal event, meriting comment in publications as diverse as *Cell* [30] and the *New York Times* [31].  
93 As we entered the new millennium, the basic problem thus seemed solved, and the agrin hypothesis  
94 reigned supreme [32].

## 95 2.2. Challenge to the model

96 Acceptance of the standard version of the agrin hypothesis proved short-lived, however, as a  
97 number of studies, whether *in vivo*, focused on the rodent diaphragm [33-35], or *in vitro*, using  
98 mammalian myotubes [36], showed that significant postsynaptic differentiation could proceed in the  
99 absence of nerve. Moreover, these studies provided evidence that the role of agrin might be to  
100 stabilize existing AChR aggregates in the face of a neurally derived dispersive agent [34,35]. This  
101 work culminated with evidence provided by two different groups, one of which, including J.R. Sanes  
102 and colleagues, had carried through the key agrin knockouts discussed above [37], and another,  
103 working independently [38]. Using double knockouts of agrin and choline acetyl transferase, the  
104 enzyme responsible for ACh synthesis, these authors found that synapses now formed. Furthermore,  
105 both groups adduced evidence in support of a mechanism whereby agrin is required to maintain  
106 synaptic AChR clusters, which in its absence would be dispersed, owing to an inhibitory mechanism  
107 mediated by the neuronal release of ACh in the region of the developing synapse. This dispersal was  
108 attributed in part to AChR endocytosis, as well as to lateral diffusion through the membrane [37]. In  
109 these transgenics, given the ablation of choline acetyl transferase expression and consequent lack of  
110 ACh, agrin was no longer essential, as it was not required to antagonize any ACh-mediated  
111 dispersion.

112 Research into the mechanism mediating this dispersal by ACh has included differing research  
113 emphases, as some studies have concentrated on AChRs at the synapse [37], while others focus on  
114 receptors that were not contacted by nerve [38]. It remains possible, however, that the molecular  
115 mechanisms underlying the dispersion of both synaptic and nonsynaptic receptors may be related,  
116 in that distant AChRs, like those at the synapse, are associated with specialized extracellular matrices  
117 [39,40], and these links must be severed to permit lateral migration of AChRs toward the synapse [8,  
118 14]. In any event, a full account of these phenomena remains to be elaborated; to date the activities of  
119 several postsynaptic enzymes have been implicated in AChR dispersion, including Cdk5 [38], calpain  
120 [41], and caspase-3 [42].

## 121 2.3. Revised model for agrin

122 The formation of synapses in mice lacking the agrin gene clearly called for a revision of its  
123 proposed mode of action. Rather than being the key synaptic organizer, agrin was assigned a role as  
124 stabilizing agent, similar to that already proposed in the interpretation of the experiments where  
125 postsynaptic differentiation was shown to occur in the absence of nerve [34,35,43]. Specifically,  
126 Misgeld et al. [37] summarized their results as follows: "In terms of agrin, we believe that in addition  
127 to its role in clustering AChRs, which has been demonstrated *in vitro*, it also acts to antagonize the  
128 effect of ACh. Thus, it is an 'antideclustering' agent, instead of or in addition to being a clustering  
129 factor."

130 Here, Misgeld et al., having just shown that synaptogenesis could proceed in agrin's absence *in*  
131 *vivo*, still remained unwilling to abandon an inductive role completely. Now, however, they based  
132 their conclusions for this capacity purely on *in vitro* data, rather than on the earlier knockouts that  
133 had seemed so compelling. With their phraseology, "instead of or in addition to being a clustering  
134 factor", Misgeld et al., drawing on two decades of *in vitro* and *in vivo* studies, posed a fundamental  
135 issue to be examined in the present review: does agrin act only as a stabilizer ("instead of") or does  
136 it both induce new synapses as well as stabilize them ("in addition to"). Given the history of the field,  
137 it is of particular note that these authors, whose transgenic studies were crucial in establishing the  
138 idea that agrin is an inducer, now accepted that this concept might have to be jettisoned altogether.

139 The overall state of matters was soon addressed in a major review by Sanes and colleagues [44];  
140 the essence of this account is summarized in their title, evoking both Milton [45] and T.S. Kuhn [46]:  
141 "*Assembly of the postsynaptic membrane at the neuromuscular junction: paradigm lost.*" There were three  
142 principal aspects of the accepted account which required alteration: (i) the evidence for muscle's  
143 ability to generate postsynaptic differentiation in the absence of nerve; (ii) the conclusion that  
144 neuregulin-1 was not involved in the induction of synaptogenesis; and (iii) the abandonment of  
145 agrin's straightforward role as the inducer of synaptogenesis. Of these three issues, the most

146 important feature of the “lost paradigm” was the last, the need to reconsider the role of agrin, since  
147 this meant that we no longer had a unifying concept, formulated at the molecular level, that could  
148 account for the initiation of synapse formation at all vertebrate NMJs.

### 149 3. Synapse formation: two mechanisms – agrin and pre patterning

150 The verdict rendered in “paradigm lost” is striking, both in terms of its incisiveness and the fact  
151 that it originated from one of the leading groups investigating the NMJ. The logic was clear enough,  
152 however, and the two reports that synapses could form in the absence of agrin [37, 38], were soon  
153 joined by a third study in which agrin’s ablation was accompanied by a second genetic  
154 transformation, leading to elevated postsynaptic concentrations of MuSK. In this last study, the mice  
155 survived, albeit runted, for several weeks [47]. It might have been expected that the modified view  
156 of agrin’s action would foster a search for an alternative that could account for the induction of  
157 postsynaptic differentiation in agrin’s absence. Surprisingly, this has not happened, and in this  
158 respect the contrast with another candidate inducer, neuregulin-1, is notable.

159 As reviewed by Kummer et al. [44], a number of studies had implicated neuregulin-1 as having  
160 an important role, but two transgenic studies, in which synapses continued to form in the absence of  
161 either neuregulin-1’s receptors [48], or neuregulin-1 itself [49], essentially refuted this concept,  
162 although neuregulin-1 does play key roles in peripheral nervous system function, especially through  
163 actions on glia [50]. With respect to agrin, however, many reports have simply ignored the findings  
164 that synapses can form in its absence. It is not uncommon for later papers, be they introductions of  
165 studies [51,52] or reviews [53-55], to characterize agrin’s role in terms of its traditional function as an  
166 organizer that promotes AChR clustering, with no reference to an alternative mode of action, and  
167 without discussing the concerns of Misgeld et al. [37] and Kummer et al. [44].

168 Instead, in the years following, focus shifted to the role of pre-existing postsynaptic  
169 differentiation, termed “pre patterning”. The findings of pre-existing postsynaptic differentiation in  
170 rodent muscles were extended by work in zebrafish, where it was reported that some synapses were  
171 induced *de novo*, while others incorporated pre-existing AChR clusters [56,57]. Together with the  
172 earlier studies of pre patterning, this led to the conclusion that two mechanisms mediate  
173 synaptogenesis at NMJs, one with pre patterning and one without. This assumption is made explicit  
174 in a Cell SnapShot, generally considered to provide an authoritative perspective, which asserts that  
175 “the final cohort of synapses on the muscle fiber includes those recognized by motor axons and those  
176 induced by motor axons” [58]. It should be noted, however, that there is an implicit bias toward the  
177 set of synapses that are recognized, as opposed to induced, in the general introductory statement that  
178 “Muscles are pre patterned ... prior to and independent of innervation”; overall, the revised view of  
179 agrin as stabilizer is given prominence in the SnapShot. Regarding those synapses that are induced,  
180 there is only the statement that “agrin can also induce new postsynaptic specializations” [58]. No  
181 other inductive mechanism is proposed for the cases where pre patterning is missing, such as the *in*  
182 *vivo* transplantation experiments [5-7], and the various *in vitro* studies where synapses formed  
183 anywhere on the muscle surface [8-10].

184 In addition, this model has been accorded adaptive significance: “it is not unexpected that  
185 control of something as important as the ability to move is overdetermined” [59]. It is fair to note,  
186 however, that this interpretation had in fact been unexpected for over a quarter century - hence  
187 “paradigm lost”. In any event, instead of a straightforward single mechanism for synapse formation  
188 and for the role of agrin, the model posits synaptogenesis as proceeding by two mechanisms and  
189 agrin having two functions.

190 While it is not unusual for cellular processes to reflect the operation of redundant mechanisms  
191 [60], the question remains as to whether this approach provides a satisfactory account for the  
192 induction of postsynaptic differentiation, the subject of “paradigm lost”. The following sections will  
193 analyze this model with respect to two questions: (i) whether it is reasonable to conclude that these  
194 two types of synapse formation proceed through a common underlying molecular mechanism, and  
195 (ii) whether the available evidence supports two roles for agrin. The conclusions, while provisional  
196 and dependent on further investigations for confirmation, are affirmative for the first question, and

197 negative for the second, as it is argued that the available evidence favors agrin's action as a stabilizer,  
198 while its inductive role is open to serious question. A final section suggests alternative inductive  
199 mechanisms, which, taken together with a role for agrin as a stabilizing factor, may restore the  
200 concept of a unitary mechanism for the initiation of postsynaptic differentiation at all vertebrate  
201 NMJs.

202 Presynaptic differentiation will not be treated, as current conclusions regarding presynaptic  
203 differentiation appear more definitive. Evidence has been adduced to implicate all three members of  
204 the agrin/MuSK/Lrp4 complex as contributing to induction of presynaptic differentiation [61-64] with  
205 particularly strong evidence in favor of Lrp4 [63,64].

### 206 3.1. Postsynaptic prepatterning and synapse formation

207 Postsynaptic prepatterning, the aggregation of AChRs in the region where synapses form prior  
208 to innervation, provides the substrate for one of the two proposed mechanisms of synapse formation,  
209 neuronal incorporation of pre-existing postsynaptic differentiation. Prepatterning was first reported  
210 by Braithwaite and Harris [65], who found that ablation of innervation by injection of  $\beta$ -bungarotoxin  
211 still left AChRs assembled in the center of the rat diaphragm. Very similar findings were reported  
212 by Yang et al. [33] in a mutant lacking the enzyme topoisomerase- $\beta$ , and in transgenics lacking either  
213 the transcription factor HB9 or neuronal agrin; prepatterning was, however found to be dependent  
214 on MuSK [34,35]. Moreover, in cultured mammalian muscle, elaborate postsynaptic structures form  
215 in the absence of nerve [36]. Follow-up studies have been especially informative in detailing the  
216 molecular interactions involving agrin, MuSK, and Lrp4, and, in the clinical sphere, the relevance of  
217 autoimmune responses to these proteins to the pathophysiology of myasthenia gravis [51,66,67].

218 This work has yielded two surprises. First, in mammals it has been established that Lrp4 rather  
219 than agrin plays the central role in the activation of MuSK, although neuronally supplied agrin does  
220 play an essential supportive function, acting as an allosteric regulator of the Lrp4-MuSK interaction,  
221 by binding to the extracellular N-terminal region of Lrp4 [29,52,68-71]. In addition, in zebrafish the  
222 activation of MuSK in forming the prepatter is mediated through its binding to Wnt, not Lrp4,  
223 although Lrp4 is still required for synapse formation [72,73].

224 In rodent muscles such as the diaphragm, prepattered AChR clusters are restricted to a narrow  
225 band in the center of the fiber, toward which the nerve terminals grow [34,35,47]. Furthermore,  
226 studies in zebrafish have provided examples where the nerve incorporates these pre-existing  
227 structures into new synapses [56,57]. At present, however, the number of muscles which have been  
228 examined for prepatterning is relatively small. It remains to be seen whether prepatterning plays  
229 any role in such large fast muscles as the frog sartorius that lack a defined central band of synapses,  
230 or in tonic fibers, that are multiply innervated [74,75].

231 As to whether prepatterning is essential for synaptogenesis, early reviews were especially  
232 cautious, pointing out the studies, outlined in Section 2.1, that synapse formation proceeds in its  
233 absence, while suggesting that the prepatter establishes a preferred region for the termination of the  
234 final nerve branches [76]. This concept was supported in a later study involving MuSK over-  
235 expression, which resulted in the destruction of the prepatter, and axons wandering randomly over  
236 the muscle surface [47]. Overall, the evidence appears strong that prepatterning, where it occurs, can  
237 direct the final terminating branches on the muscle surface [63,64,72].

### 238 3.1.2. Refinement of the prepatter: implications for a unitary molecular mechanism

239 The question remains as to whether the appropriation of pre-existing structures by the incoming  
240 nerve involves different molecular mechanisms from the case where these structures are absent. In  
241 short, how are the synapses, "those recognized by motor axons" [58], actually formed? From first  
242 principles we might reasonably assume that if the incorporation of existing structures is a different  
243 process from the induction of a novel postsynaptic apparatus, then different macromolecules on the  
244 nerve and muscle surfaces would be expected to mediate the respective interactions between the two  
245 cells. It should be noted, however, that the existing prepatter of AChRs is known to be refined by  
246 nerve contact, since the final geometry of the mammalian synapse is modified in response to

247 innervation [34,43,44,77]. To date the mechanisms mediating this rearrangement have yet to be  
248 examined, raising the question: is this process any different, at the molecular level, from the induced  
249 migration of distant AChRs [8], with the exception that the receptors are now, thanks to  
250 prepatterning, close at hand?

251 Several studies suggest that it is not. If different mechanisms were involved in synaptogenesis  
252 mediated by incorporation of a prepattern, as opposed to *de novo* generation of the postsynaptic  
253 apparatus, we might anticipate that disrupting the prepattern would have direct impacts on synapse  
254 induction. However, such interference does not prevent synapse formation, even in the very  
255 preparations where prepatterning has been most intensively studied. For example, in the mouse  
256 diaphragm, prepatterning was disrupted in muscle fibers lacking the dihydropyridine receptor  
257 essential for excitation-contraction coupling; nevertheless, functional NMJs were formed [78]. Similar  
258 results were seen in mice whose muscles expressed elevated levels of MuSK; destruction of the  
259 prepattern was accompanied by motor neurons wandering over the entire muscle surface, but  
260 functional synapses still developed [47]. Moreover, detailed experimental manipulation of Wnt  
261 signaling in zebrafish was found to eliminate prepatterning, but synapse formation still occurred;  
262 these authors concluded that prepatterning of AChR is dispensable for the induction of  
263 synaptogenesis in this system, even though it plays a central role in restricting the pathfinding of  
264 growth cones [72]; see Section 4.2.

265 The most parsimonious interpretation of these investigations is that the two modes of synapse  
266 formation proposed by Burden [58] and by Sanes and Yamagata [59] both reflect an underlying unity  
267 of molecular mechanism, expressed in the three general cases examined: (i) in the simplest case,  
268 where there is no prepattern, as in the early experimental studies *in vitro* [8-10], and *in vivo* [5-7],  
269 postsynaptic structures are induced at the point of cell-cell contact; (ii) if prepatterning is present, the  
270 same inductive interactions involved in *de novo* synaptogenesis mediate the final remodeling of  
271 postsynaptic components so as to align prepatterned AChRs with the terminal nerve branches; and  
272 (iii) when normal prepatterning is abolished experimentally, so that there are no longer any pre-  
273 existing AChR clusters to be re-aligned, the muscle simply uses the same molecular mechanisms to  
274 aggregate AChRs *de novo*.

275 Taken together, these studies lead to two general conclusions. First, whatever assistance  
276 prepatterning may provide toward direction of nerve terminal growth, it is not essential for the  
277 development of postsynaptic differentiation, the focus of the “lost paradigm”. Second, the  
278 fundamental assumption underlying the agrin hypothesis, that a single molecular mechanism  
279 mediates postsynaptic differentiation at all vertebrate skeletal NMJs, whether or not prepatterning is  
280 present, continues to be a viable proposal. It now remains to analyze whether the original  
281 formulation of this concept can be sustained, namely that the secretion of neuronal agrin is the key  
282 inductive step.

### 283 3.2. Agrin – stabilizer, inducer, or both?

284 In principle, both of the two actions proposed for agrin, induction of new synapses and  
285 stabilization of existing ones, are vital, and the loss of either one could account for the lethality of the  
286 first agrin knockouts [23,24]. In their discussion of these actions, Misgeld et al. [37] were unwilling  
287 to abandon the concept of an inductive role for agrin, even as they provided evidence for a stabilizing  
288 function. These possibilities, stabilizer and inducer, are here assessed in turn, drawing not only on  
289 the transgenic approaches which have predominated in recent studies, but also on earlier findings  
290 from the amphibian culture system.

#### 291 3.2.1. Agrin as stabilizer

292 Acting as an “antideclustering” stabilizing agent, agrin could in principle either be deposited  
293 together with an inductive stimulus, or at some time later, after AChR clustering has begun. Since  
294 Misgeld et al. [37] proposed this view of agrin’s role, there have been two data sets that might be  
295 interpreted as a challenge to the concept. The first was the double transgenic where agrin was  
296 ablated and MuSK expression in muscle was simultaneously elevated. In this study motor nerve

297 growth over the muscle surface was aberrantly profuse, but the agrin-less mice, although runted,  
298 survived for several weeks, implying a general level of functionality of the NMJs [47]. Thus, even in  
299 the face of continued ACh release, which should have led to rapid dispersal of synaptic AChR  
300 aggregates with agrin absent [37,38], synapses had formed and were clearly stable for an extended  
301 time. Since MuSK over-expression was known to promote MuSK dimerization and consequent  
302 MuSK kinase activity, allowing for activation in the absence of agrin [79], persistence of synapses was  
303 attributed to MuSK self-activation, owing to its elevated concentration [47]. Although this mechanism  
304 has not been explicitly confirmed, it can in principle compensate for the stabilizing action that would  
305 normally be provided by the neuronal agrin isoform, and the concept of agrin as stabilizer is  
306 accordingly preserved.

307 The interpretation of MuSK self-activation has been supported by the case of a second double  
308 transgenic where agrin was ablated and the mice survived; in this study, the accompanying genetic  
309 modification was the elevation of another postsynaptic protein, Downstream of kinases-7 (Dok-7), an  
310 activator of MuSK [80]. Once again functional NMJs formed, with the mice surviving for up to two  
311 months; moreover, biochemical analyses confirmed that postsynaptic MuSK activity was in fact  
312 significantly increased, reflecting the increased synthesis of Dok-7. Ultimately, however, the synapses  
313 disintegrated. Thus it appears that elevation of MuSK activity can obviate the need for neuronal agrin  
314 over the time period required for induction and consolidation of postsynaptic differentiation, even  
315 though over the longer term the system cannot be maintained. Remaining to be explained, however,  
316 is the mechanism by which the interaction of nerve and muscle was initiated in the absence of agrin.

### 317 3.2.2. Agrin as inducer

318 The classic agrin hypothesis was especially attractive in that a single factor was assumed both to  
319 initiate a complex process and also to differentiate motor neurons from other nerve types. Thus  
320 specificity [81] was in principle accounted for, as it was assumed that motor neurons are specialized  
321 for providing neuronal agrin [16,17], even though other studies indicated that sensory neurons do  
322 apparently transcribe the gene [82]. Close examination of the evidentiary basis for an inductive  
323 action of agrin, however, suggests that it is less strongly supported than is often averred.

#### 324 3.2.2.1. Early studies: issues of interpretation

325 Even at the time of its formulation, the agrin hypothesis in practice held a special position, being  
326 the only proposal with any general currency, effectively constituting a default hypothesis. In this  
327 context a number of nonphysiological approaches, including bath application [83] and postsynaptic  
328 expression from injected transgenes [84,85], gave results that were consistent with the agrin  
329 hypothesis, and accordingly were taken as being as supportive of it. Ultimately these conclusions  
330 appeared to be validated with the agrin knockouts [23,24] and studies with nerve-muscle cultures  
331 [86]. It is safe to say, however, that such interpretations would have been subject to more scrutiny  
332 if it had been known that synapse formation could proceed in animals lacking the agrin gene.  
333 Simple logic would have called for discriminating between inductive and stabilizing roles for agrin,  
334 or at least, as in the case of Misgeld et al. [37], making a case for both.

335 The key element in confirming agrin as an inducing agent is the question of time, meaning that  
336 to be the key inducer it must be deposited at an early stage of nerve-muscle contact. To establish this  
337 rigorously requires examination of synapse induction as it occurs during nerve-muscle interaction.  
338 In this context we should note one fact that may lead to over-interpretation of many experimental  
339 approaches that do not observe this condition, namely the apparent promiscuity of the muscle surface  
340 in responding to exogenous stimuli. For example, two nonphysiological examples that can lead to  
341 AChR clustering include local electric fields [87,88], and uncoated polystyrene beads [89]. Thus such  
342 approaches as local fixation onto culture surfaces [90] are open to the possibility that the muscle  
343 membrane is responding to a signal that is not the same as occurs during synaptogenesis. In fact,  
344 this caveat may even apply to the original finding that AChR clustering occurs at specialized basal  
345 lamina sites of regenerating muscle [12] despite the ultimate productivity of this approach in  
346 identifying agrin as a key actor in synapse formation.

347 In any event, the several reports [37,38,47,80] that mice lacking the agrin gene could form  
348 synapses present an obvious challenge that must be answered. In two cases [37,38] one could argue  
349 that the completeness of the inductive process was suspect, since the lack of ACh, given the  
350 concomitant knockout of choline acetyl transferase, precluded functional analysis of the outcome.  
351 The other cases, however, in which either MuSK or Dok-7 was elevated while agrin was ablated  
352 [47,80], are more definitive, in that the mice survived for many weeks, so that functional synapses  
353 clearly formed. In addition, in one of these reports postsynaptic differentiation could not have  
354 proceeded through the incorporation of existing prepatterned structures, since pre patterning was  
355 also obliterated [47]. Given that induction by neuronal agrin was obviously precluded, some other  
356 process mediated the initiation of synaptogenesis, and what this might be remains a mystery.

357 There is one major caveat, however, to applying any inferences to NMJ formation more  
358 generally, given the nature of the double transgenics in question. Specifically, the elevation in activity  
359 of the key postsynaptic transducing molecule can reasonably be taken as such an aberrant  
360 circumstance as to preclude using such transgenics to draw definitive conclusions pertaining to wild-  
361 type mice. To answer this objection properly requires a context in which agrin is missing at the key  
362 inductive juncture, even though there are no major distortions in gene expression of key participating  
363 molecules. Fortunately, this condition is satisfied by studies in the amphibian *in vitro* system,  
364 especially when viewed in the light of transgenic data showing that agrin's action is critically  
365 dependent on its binding to laminin in the basal lamina.

#### 366 3.2.2.2. Agrin in the amphibian *in vitro* system

367 To establish biological relevance of a potential inductive stimulus, the challenge is not just to  
368 demonstrate that it is capable of inducing synaptic differentiation, but that it actually does so in a  
369 physiological context, and at the required time. In particular, a candidate inductive process, such as  
370 the secretion of neuronal agrin, clearly must occur early in the sequence of events that ultimately  
371 generates a synapse. To date the preparation which remains the most closely examined step-by-step,  
372 during the process of synapse induction, is the original *in vitro* system with embryonic cells from  
373 *Xenopus laevis* [8]. In addition to facilitating the assessment of the timing of events during  
374 synaptogenesis, this culture system is of additional interest in that, like the *in vivo* nerve  
375 transplantation experiments in adult muscle [5-7], synaptogenesis proceeds in the absence of  
376 pre patterning. In the scheme of Sanes and Yamagata [59] and Burden [58], this is clearly a case of  
377 induction of a new synapse, rather than incorporation of existing postsynaptic structures.  
378 Accordingly, if agrin acts as an inducer in such cases, as argued in the Cell Snapshot [58], this role  
379 should be evident here.

380 In this preparation it is noteworthy that synaptic differentiation does not occur at an interface  
381 with a culture substrate, but rather between nerve and muscle membranes, in essence replicating the  
382 *in vivo* situation. The only difference from the *in vivo* case is that synapses form without nerve  
383 termination; nonetheless they show all the expected features of a differentiated contact by  
384 physiological, immunocytochemical, and ultrastructural criteria [91,92]. Using a pulse-chase  
385 protocol to stain AChRs first with labeled  $\alpha$ -bungaroxin, then following with unlabeled toxin after  
386 nerve addition, AChR aggregation under the nerve was shown to include receptors that were stained  
387 before nerve addition. These were interpreted as having migrated laterally through the muscle  
388 membrane [8], although endocytosis and re-insertion of labeled AChRs may also have contributed to  
389 their ultimate localization at the synapse [93]. This aggregation of AChRs was subsequently shown  
390 to proceed in parallel with a series of defined stages in the development of a specialized synaptic  
391 basal lamina [14,92].

392 These studies form the backdrop for assessing the role of agrin during synapse formation in this  
393 system. The work is almost entirely observational, but taken together with detailed functional  
394 analyses of the agrin molecule, provides three related lines of evidence that all point towards agrin  
395 as acting to stabilize, rather than induce. These include (i) the role of agrin as a basal lamina  
396 constituent, viewed in the context of basal lamina remodeling; (ii) the time course of agrin deposition



397 during synaptogenesis in living cultures; and (iii) the relationship of agrin to stabilization of AChR  
398 aggregates at the synapse following denervation.


### 399 3.2.2.3. Neuronal agrin as basal lamina constituent: implications for function

400 Agrin is a large heparan sulfate proteoglycan, approximately 400 Kd, of which half is composed  
401 of heparan sulfate side chains [94]; the final protein is subject to differential splicing of the mRNA  
402 transcript, but for the purposes of defining its functional role as stabilizer or inducer we are only  
403 concerned with the N- and C-terminal regions. First, as discussed in Section 3.1, Lrp4 and MuSK  
404 interact directly, with agrin acting to potentiate this interaction by binding to Lrp4 [29,52,68-70]. This  
405 binding to Lrp4 is effected through the C-terminal region of agrin containing the key 8-amino acid  
406 insert that is essential for synapse formation [24]. Elucidating the nature of this interaction has been  
407 the major focus of functional studies [54,69], inasmuch as it ultimately leads to downstream  
408 postsynaptic interactions.

409 In contrast, the function of the N-terminal region has been much less studied in recent years, but  
410 work in the previous millennium demonstrated that there are two principle variants, isoforms with  
411 an extended 150 amino acid sequence, termed LN, and those without, termed SN [95]. While SN-  
412 agrin mediates integration into cell membranes throughout the nervous system [95], LN-agrin has a  
413 more restricted distribution, with a particular concentration in motor neurons. Moreover, agrin's  
414 incorporation into the muscle basal lamina is a property of the LN-isoforms [95-97]. Based on *in vitro*  
415 studies with chick nerve and muscle, as well as transfected mammalian (Cos-7) cells, this longer N-  
416 terminal binds to the  $\gamma$ 1 chain found in most laminin heterotrimers [98], including those characteristic  
417 of the specialized synaptic basal lamina [99-101].

418 Of central importance for the present discussion, this binding is critical for synaptogenesis, as  
419 transgenic mice lacking only the N-terminal extension were as deficient in NMJ formation as mice  
420 lacking all forms of agrin [95]. This deficiency is unlikely to reflect interference with interactions with  
421 Lrp4, since tissue extracts from mice lacking the N-terminal extension retained AChR clustering  
422 activity on cultured myotubes [95]. It is most likely therefore, that during synaptogenesis neuronal  
423 agrin is not simply secreted onto the muscle surface in a freely diffusible form; instead its C-terminal  
424 is presented to Lrp4 while its N-terminal is bound to laminin in the extracellular matrix.

425 The relevance of this finding for agrin function becomes clear when we examine the sequence of  
426 events during synaptogenesis in the amphibian culture system. Here the first indication that a given  
427 nerve-muscle contact will form a synapse is the removal of existing basal lamina constituents along  
428 the path of the nerve. These constituents include the basal lamina heparan sulfate proteoglycan  
429 (HSPG) and, importantly, laminin [102; see Figure 1]. Both molecules, derived from the muscle [103,  
430 104], are then re-deposited at high concentration in the synaptic basal lamina [102; see Figure 2], a  
431 local specialization that frog NMJs share with those of mammals [94,105]. Meanwhile AChR  
432 aggregates accumulate in the adjacent muscle membrane [14,102; see Figure 2].



433

434

**Figure 1.** Nerve-induced removal of basal lamina constituents prior to synapse formation.

435 Living nerve-muscle cultures of *Xenopus laevis* were examined approximately one day after the addition of nerve.  
 436 Two constituent proteins of the muscle basal lamina, a heparan sulfate proteoglycan (HSPG) and laminin, (A  
 437 and B, respectively), were stained with monoclonal antibodies labeled with contrasting fluorochromes. Both  
 438 show a dramatic loss of fluorescence intensity along the path of the nerve, shown in the phase contrast image  
 439 (C). Note that the width of reduced fluorescence in A and B is several-fold greater than the width of the nerve  
 440 itself.

441 This reduced fluorescence is seen only at those nerve-muscle contacts that go on to make synapses, a 1:1  
 442 correlation suggesting that localized pericellular proteolysis of basal lamina constituents may contribute to the  
 443 transmission of inductive signals between nerve and muscle.


444

Scale: 20  $\mu$ m

445

Reproduced with permission from Anderson et al., Mol. Cell. Biol. 1996;16:4972-84.

446 Both molecules, derived from the muscle [103, 104], are then re-deposited at high concentration  
 447 in the synaptic basal lamina [102; see Figure 2], a local specialization that frog NMJs share with those  
 448 of mammals [94,105]. Meanwhile AChR aggregates accumulate in the adjacent muscle membrane  
 449 [14,102; see Figure 2].



450

451 **Figure 2.** Deposition of a specialized basal lamina parallels synaptic aggregation of acetylcholine  
 452 receptors in the muscle membrane.

453 A. In this living *Xenopus* culture, examined after several days of nerve-muscle co-culture, laminin, stained with  
 454 a monoclonal antibody, is beginning to fill in the region where it had formerly been removed along the path of  
 455 nerve-muscle contact, shown in the phase reference image (C). This newly deposited laminin is at a higher site-  
 456 density than is present elsewhere on the muscle cell. In favorable preparations the entire synaptic zone will be  
 457 filled in.

458 B. Acetylcholine receptors (AChRs), stained with fluorescent  $\alpha$ -bungaroxin, conjugated to a contrasting  
 459 fluorochrome, have aggregated in the muscle membrane along the path of cell-cell contact, co-localizing with  
 460 the synaptic laminin deposits seen in A. AChR staining is not evident elsewhere on the muscle surface.

461 Scale: 20  $\mu$ m

462 Reproduced with permission from Anderson et al., Mol. Cell. Biol. 1996;16:4972-84.

463 Taken together with the findings that agrin is presented in a laminin-bound form, this sequence  
 464 of removal and re-deposition implies that the incorporation of neuronal agrin into the synaptic cleft  
 465 follows an earlier inductive process, one that involves basal lamina remodeling. At the earliest stages  
 466 of nerve-muscle interaction, the removal of laminin from the synaptic cleft precludes the  
 467 incorporation of agrin into the matrix. Only after laminin is re-deposited can this incorporation  
 468 proceed, thereby allowing the interaction of agrin with Lrp4. Whether the development of a  
 469 specialized synaptic basal lamina follows the same time course in other vertebrate species remains to  
 470 be established, but here in the amphibian *in vitro* case, where the result is a clearly defined synapse,


471 the sequence provides evidence that agrin deposition occurs relatively late, consistent with agrin's  
472 playing a stabilizing role, rather than acting as an inducer.

473 It might be thought that drawing inferences from such disparate data, involving as they do *in*  
474 *vivo* transgenic studies [95], electron microscopic studies of the interacting macromolecules [97], *in*  
475 *vitro* binding experiments with expression plasmids [98], and *in vitro* formation of living synapses  
476 [102], is forcing the issue. In reality, however, the conclusions depend on only one basic assumption,  
477 namely that the molecular interactions between laminin, agrin and Lrp4 are conserved from *Xenopus*  
478 to mammals and birds. While ultimately requiring confirmation using *Xenopus*-derived  
479 macromolecules, this assumption is reasonable, akin to that which underlay the various studies  
480 leading to the identification of agrin: given similarity of organelle structure and function, there is no  
481 reason to posit differences at the molecular level.

482 Finally, the amphibian culture system comprises a second instance in which NMJs are induced  
483 without the involvement of either pre patterning or agrin. The first, described in mice by Kim and  
484 Burden [47] (Section 3.2.2.1), eliminated both as a result of the transgenic modifications. In the  
485 amphibian cultures, which lack pre patterning, agrin's absence at the early inductive phase is implicit,  
486 given the very early removal of its basal lamina binding partner, laminin [102]. Both cases, therefore,  
487 preclude invoking a fallback position for initiating postsynaptic differentiation, whereby either agrin  
488 or postsynaptic pre patterning can be absent, since the other's presence will suffice. In these examples  
489 both are absent, and postsynaptic differentiation still proceeds; the straightforward implication is  
490 that some other mechanism mediates its induction.

#### 491 3.2.2.4. Time course of agrin deposition during synaptogenesis in living cultures

492 The second line of evidence in support of a stabilizing role for agrin comes from direct  
493 observation of its deposition during synapse formation. This question of timing was examined in  
494 two studies in the amphibian culture preparation, both carried out in the early 1990's, prior to the  
495 agrin knockouts. Cohen and Godfrey [106], examining synapses on the upper muscle surface, found  
496 an excellent correlation between the appearance of agrin and concomitant postsynaptic  
497 differentiation. However, these results were challenged on technical grounds by Anderson et al. [107],  
498 who found that the protocols employed in the earlier study could not discriminate between agrin  
499 deposits derived from nerve-muscle contact and those secreted into the culture medium by  
500 surrounding epithelial cells. When Anderson et al. examined synapses occurring on the underside of  
501 the muscle, they did not find a consistent correlation between synaptogenesis and agrin deposition,  
502 even though a strict correlation was observed for deposition of other basal lamina constituents.  
503 Moreover, they found that agrin deposition, when it occurred, usually lagged behind other synaptic  
504 markers, as shown in the examples of Figure 3.



505

506

507

**Figure 3.** Variation in agrin deposition at different neuromuscular junctions, despite their accessibility to staining with antibodies to the basal lamina HSPG.

508

509

510

511

512

**A and B.** After several days of nerve-muscle co-culture, staining with an antibody to agrin is marked in upper nerve-muscle contact of panel A, corresponding to the nerve running from 11 o'clock to 4 o'clock in the phase reference image (arrows in E), while staining is much fainter in the second contact, corresponding to the nerve running from 10 o'clock to 6 o'clock in E. In panel B, however, neither of the two nerve-muscle contacts corresponding to the arrows in the phase reference image (F) shows any sign of agrin deposition.

513

514

**C and D.** The basal lamina HSPG, stained with an antibody conjugated to a contrasting fluorochrome, is brightly stained at all four nerve-muscle contacts, in contrast to the variability in staining of agrin.

515

Scale: 20  $\mu$ m.

516

Reproduced with permission from Anderson et al., *Dev. Biol.* 1995;170:1-20.

517

518

519

520

521

522

523

524

525

526

527


Here four different nerve-muscle contacts are visible in the phase contrast reference images (E and F). In panels A and B, all stages of agrin deposition are evident, ranging from absence to high concentration at the synapse. Note that in panel A there is significant agrin deposition corresponding to the upper nerve of panel E, while for the contact in the lower left, there is very little. Meanwhile, in panel B, there is virtually no sign of agrin whatever for either of the nerves visible in F. In contrast, the basal lamina marker, in this case HSPG (panels C and D), is present in high concentration for all four contacts. Had it been possible to employ a third fluorochrome, it would have been evident that AChR aggregation under the nerve, the classic marker for synaptic differentiation, had already occurred [92,102]; see Figure 2 above.

The most straightforward interpretation of these images is that they are essentially snapshots taken of a dynamic process during which the various nerves have contacted the myocytes at different

528 times, with the intensity of agrin staining dependent on the length of time that the two cells have  
529 been interacting. In each of the nerve-muscle contacts, basal lamina remodeling has already resulted  
530 in heavy deposition of HSPG. The contrasting variability in agrin staining implies that agrin  
531 deposition comes relatively late in the sequence of events, after HSPG deposition and AChR  
532 aggregation, rather than at the early time point expected for an inducer/organizer. Accordingly,  
533 Anderson et al. concluded that the developmental sequence is most consistent with agrin's acting to  
534 stabilize new synapses rather than to induce their formation [107], in effect presaging proposals made  
535 in the following decade [43,44].

#### 536 3.2.2.5. Behavior of AChR clusters after denervation

537 The third line of evidence for agrin as a stabilizing factor in these cultures derives from studies  
538 of the response to denervation. Anderson et al. [92] reported that when denervation was carried out  
539 in the first day after the synapse formed, synaptic AChR clusters invariably disappeared. For  
540 technical reasons all the experiments were carried out with curare in the medium; it is required in  
541 order to block the effects of large spontaneous miniature end-plate potentials that otherwise induce  
542 contractures of the myocyte, leading to its death through tearing off the culture substratum [91].  
543 Unlike the studies of the mammalian situation, where the dispersive effect of neurally-released ACh,  
544 reacting with AChRs, had to be counteracted by agrin [37,38], here in the amphibian cultures the  
545 AChRs were already blocked, precluding any action of ACh. Even so, the synaptic AChR  
546 accumulations were inherently unstable at the earliest stages of synaptogenesis, and the presence of  
547 an intact nerve was required to prevent their loss. Similar results were reported by Kuromi and  
548 Kidokoro [108], who also examined older nerve-muscle contacts, and found that after three days of  
549 co-culture, approximately half the synaptic AChR clusters remained intact after denervation,  
550 suggesting that a process of stabilization was taking hold at later times.



551

552 **Figure 4. Stability of junctional HSPG deposits after denervation.** The *Xenopus* nerve-muscle  
 553 culture was exposed to continued presence of a fluorescently-labeled monoclonal antibody, and  
 554 photographed at daily intervals. The nerve (arrowheads in A) disappeared between the first (A and  
 555 B) and second day (C and D) in culture. HSPG organization increased along the path of nerve-muscle  
 556 contact between the first and second observation (C and D) and remained static thereafter (F, H, and  
 557 J) despite loss of the nerve (C, E, and G). The intensity of HSPG staining decreased in panels F, H, and  
 558 J owing to fluorochrome bleaching in response to the repeated excitation.

559

Scale: 40  $\mu$ m.

560

Reproduced with permission from Anderson et al., *J. Cell Biol.* 1984;99:1769-84.

561


562

563

564

The denervation-induced dispersal of synaptic AChR clusters was not paralleled by  
 corresponding changes in the basal lamina however; as shown in Figure 4, synaptic HSPG deposits  
 became more pronounced between day 1 and day 2, and after denervation remained essentially  
 unchanged for several days thereafter [92]. As the remodeled basal lamina was clearly not able on

565 its own to prevent AChR dispersal, the question arises as to what agent or process led to the apparent  
566 increase in stability of synaptic AChRs in older contacts. Given the evidence that agrin makes a  
567 delayed appearance (Figure 3), it is an obvious candidate. This suggestion is further supported by  
568 evidence that once deposited, agrin can stabilize AChR clusters, as shown in Figure 5. In this older  
569 culture the nerve has died, but the typical fluorescence profile of a late-stage culture is evident, in  
570 that there are advanced synaptic AChR aggregates, together with co-localized deposition of agrin  
571 [107]. It is tempting to conclude that the appearance of agrin, relatively late in the overall process  
572 (Figure 3), has stabilized the synaptic AChR clusters so that they no longer depend on the presence  
573 of an intact nerve.



574

575 **Figure 5.** Survival of synaptic agrin and AChR accumulations after nerve degeneration in an older  
576 *Xenopus* nerve-muscle culture.

577 Agrin was stained with a monoclonal antibody (A) and AChRs were labeled with  $\alpha$ -bungarotoxin (B), using  
578 contrasting fluorochromes. Note the intense AChR staining across the center of the two myocytes (B), and the  
579 co-localized agrin deposits (A). The nerve is no longer visible in the phase contrast reference image (C).



580 In this figure, where the nerve has persisted long enough for agrin deposits to be laid down, synaptic AChR  
581 aggregates have remained. In contrast, synaptic AChR aggregates disperse if denervation occurs within a day  
582 after synaptic differentiation becomes evident.

583 Scale: 20  $\mu$ m

584 Adapted with permission from Anderson et al., *Dev. Biol.* 1995;170:1-20.

585 At first glance, these findings suggest that different mechanisms for the maintenance of synaptic  
586 AChR clusters may be involved in the amphibian cultures, as compared to those operative in the  
587 mammalian system [37,38]. The amphibian synaptic AChR aggregates do not require any neuronal  
588 stimulus such as ACh to mediate their dispersal; the nerve, rather than acting to disperse them,  
589 clearly acts to maintain them. This difference between the two experimental systems may be more  
590 apparent than real, however, if, as suggested in Section 3.1.2, synaptic AChRs at mammalian  
591 synapses are actually re-aligned receptors, newly aggregated under the nerve, having moved there  
592 from their positions in the prepattern. If so, they too may be unstable at the earliest stages of their re-  
593 alignment.

594 Such a mechanism would be supported if future studies determine that mammalian  
595 postsynaptic differentiation also involves a proteolysis-driven remodeling of the basal lamina, similar  
596 to that of the amphibian culture system. Since, as discussed above, laminin is first removed and  
597 then redeposited during this process, we would expect a similarly late appearance of agrin, given its  
598 dependence on laminin for anchoring in the basal lamina. Conversely, if no such remodeling occurs,  
599 it would strengthen the case that the AChRs in the prepattern are simply co-opted directly, in  
600 accordance with the model of Burden [58] and Sanes and Yamagata [59].

#### 601 3.2.2.6. The amphibian culture system: conclusions, caveats and recommendations

602 Taken together, several lines of evidence all incline toward the conclusion that neuronal agrin  
603 appears too late to act as an inducer of postsynaptic differentiation in the amphibian cultures, but  
604 that once deposited it acts to stabilize synapses that are already relatively advanced in their  
605 formation. The loss of this stabilizing function would be more than sufficient to account for the  
606 devastating impacts of the early agrin knockouts [23, 24]. As with the synapses of the transgenic  
607 described by Kim and Burden [47; – see Section 3.2.2.4 above], these data are most consistent with an  
608 inductive process that does not involve either agrin or pre patterning.

609 At present these studies in the amphibian system stand as the only attempts to assess the  
610 sequence of events such as basal lamina remodeling and agrin deposition during the development of  
611 the NMJ. Several caveats must be acknowledged respecting these observations, however, in  
612 particular regarding the key finding that agrin appears to come in too late to be an inducer. First,  
613 while Anderson et al. [107] went to considerable lengths to control for penetration artifacts that might  
614 lead to inconsistent staining of agrin deposits, technical difficulties, whether reflecting issues of  
615 penetration or local proteolytic action on specific epitopes, can never be excluded definitively. For  
616 example, focal proteolysis of matrix constituents is a constant feature attending synaptogenesis in the  
617 amphibian culture system [14,109] and agrin is known to be subject to hydrolysis by several  
618 metalloproteinases [110,111], as well as by neurotrypsin, a serine proteinase [112,113]. More  
619 generally, in a context where staining is, in the words of Anderson et al., “erratic” [107], definitive  
620 conclusions will require a complete description that can account for the presence of agrin at some  
621 synapses and its absence in others.

622 A second issue derives from the detailed molecular analyses of agrin’s interactions, specifically  
623 the need for laminin to be present in order for the LN-agrin isoforms to bind to the extracellular  
624 matrix. Although these findings of Burgess et al. [95] reinforce the longstanding view that agrin acts  
625 physiologically as a constituent of the remodeled basal lamina, implying a late appearance of agrin  
626 in the amphibian culture system, they do not in themselves definitively rule out an inductive role in  
627 postsynaptic differentiation. In principle there remains an early temporal window in which agrin  
628 could still mediate the initial inductive interaction, one that precedes its incorporation into the basal  
629 lamina, and initiate the entirety of the succeeding developmental sequence; ultimately this would

630 even include further accumulation of the LN-agrin isoform through its binding to laminin. While this  
631 scenario seems highly unlikely, especially in view of the *in vivo* knockouts where synapse formation  
632 persists in agrin's absence [37,38,47,80], it remains a formal possibility.

633 These considerations require a re-examination of the exact sequence of events that follow contact  
634 between nerve and muscle, in particular whether the deposition of neuronal agrin precedes or follows  
635 basal lamina remodeling. Given the technical constraints operative a generation ago, the necessary  
636 repeated observations of a given synaptic contact were difficult to perform. Modern approaches,  
637 however, include the use GFP-modified constructs that obviate concerns regarding the use of staining  
638 reagents, as well as advances in the sensitivity of fluorescence microscopy that permit examining  
639 multiple channels while minimizing fluorochrome bleaching [114]. This should facilitate repeated  
640 assessments of several key molecular actors at all stages of synapse formation, not only in studies  
641 with the amphibian culture system, but ultimately with *in vivo* preparations, the gold standard for  
642 any proposed mechanism.

#### 643 4. Synapse induction: alternatives to agrin

644 If deposition of neuronal agrin by motor neurons is not the mechanism for conferring specificity  
645 and inducing postsynaptic differentiation, the question arises as to what an alternative process might  
646 be. At the level of the nervous system as a whole, numerous classes of mechanisms have been  
647 defined for ensuring appropriate connections between neurons and their targets, including such  
648 processes as inhibitory interactions between surface molecules, placeholder and guidepost cells to  
649 direct axon growth, and actions to eliminate inappropriate synapses [59,115]. Many of these are not  
650 relevant for the present discussion, however, which will focus on the simplest of cases, that of nerve  
651 on undifferentiated muscle, as described both *in vivo* and *in vitro* (see Section 2.1). The logic in so  
652 doing is not that the answers will necessarily provide a definitive conclusion for all nerve-muscle  
653 synapses; rather, it is that the simplest case does require explanation for itself, and that it is likely that  
654 mechanisms so revealed will provide insights into the wider spectrum of NMJs. This review will  
655 examine four potential mechanisms, namely ligand receptor interactions, Wnt signaling, focal  
656 pericellular proteolysis, and integrin signaling. All four have either been examined in other contexts  
657 in the nervous system, with potential relevance to NMJ formation, or have been studied directly at  
658 the NMJ.

##### 659 4.1. Ligand receptor interactions

660 Based on work done elsewhere in the nervous system, there is in principle a plethora of potential  
661 candidates for mediating nerve-muscle interaction through homophilic or heterophilic binding,  
662 including such families as the cadherins, protocadherins, neuroligins and neuroligins, ephrins and eph  
663 kinases, semaphorins and neuropilins or plexins, and members of the immunoglobulin family (for  
664 reviews see [59,115,116]). One example that has been examined is neural cell adhesion molecule  
665 (NCAM), for which different isoforms were found to mediate stabilized contacts between growth  
666 cones and cultured myotubes [117]; however, complete ablation of NCAM does not prevent  
667 formation of NMJs, although they are somewhat reduced in size and manifest some physiological  
668 deficits [118,119]. Other families have not been subject to close examination as candidates for  
669 initiating postsynaptic differentiation, however, for the simple reason that the concept of a ligand-  
670 receptor interaction was dominated for so long by the focus on agrin. As discussed above, however,  
671 there are reasons to believe that this focus is misplaced, as agrin's binding to Lrp4 is more likely to  
672 serve as a stabilizing rather than an inducing function, thus leaving the situation open for  
673 consideration of other possibilities.

674 At present, therefore, there are no positive data to discuss regarding alternative ligand-receptor  
675 pairings, but there are at least two candidates of potential interest that can be ruled out, namely the  
676 amyloid precursor protein, together with its two related amyloid precursor-like proteins, and  
677 presynaptic Lrp4. In both cases gene knockout data argue against their involvement in the initial  
678 inductive process. Regarding the amyloid precursor and the two precursor-like proteins, several  
679 studies have implicated them as being essential for proper synapse formation [120,121], reflecting

680 molecular interactions with Lrp4 and agrin [122]. Nonetheless, simultaneous ablation of all three  
681 proteins, while leading to perinatal lethality, still leaves the mice able to breathe for many hours or  
682 even days [123], so that, as argued by Caldwell et al. [124], the initial steps involved in NMJ formation  
683 must already have taken place without them.

684 With respect to neuronally-derived Lrp4, an elegant study, involving the ablation of Lrp4 in  
685 nerve, muscle or both, demonstrated that soluble and active fragments of presynaptic Lrp4 are  
686 generated through local proteolysis [63]. In the absence of postsynaptic Lrp4, these fragments  
687 mediate the formation of AChR clusters, although the latter are significantly reduced in size relative  
688 to the normal case. Interestingly, in the context of overall postsynaptic abnormality, they are still  
689 sufficient to prevent neonatal mortality, with the mice surviving for as long as a year. However, in  
690 this paper the authors also demonstrated that synapse formation proceeds normally when only  
691 neuronal Lrp4 is ablated, implying that the presence of presynaptic Lrp4 fragments is irrelevant in  
692 the presence of an intact postsynaptic complement of Lrp4, one that swamps out any neuronal  
693 contribution. In all other studies this condition is met, including those involving synapse induction  
694 in the absence of agrin (see Section 2.2), meaning that it is unlikely that neuronally-derived Lrp4  
695 provides an inductive signal.

#### 696 4.2. *Wnt signaling*

697 The relevance of Wnt signaling to NMJ formation has been the subject of numerous studies,  
698 beginning with the finding that Dishevelled, a protein implicated in Wnt-mediated pathways,  
699 interacts with MuSK to regulate AChR clustering [125]; for reviews see [71,126]. Detailed analysis of  
700 the underlying processes has proved difficult, however, owing to the sheer number of Wnt proteins,  
701 numbering some 19 in mammals, with yet more in other vertebrate and invertebrate species [126].  
702 Moreover, Wnt signaling is especially complex, involving as it does the canonical pathway leading  
703 to regulation of gene transcription, as well as the two noncanonical pathways affecting the  
704 cytoskeleton (planar cell polarity pathway) and intracellular calcium metabolism [127]. This  
705 complexity has been reflected in work on the NMJ, as studies using *in vivo* and *in vitro* protocols have  
706 variously reported that some Wnts can promote formation of AChR clusters, while other Wnts are  
707 inhibitory; see [71] for a recent review.

708 With respect to the particular focus of the present review, namely the nature of the inductive  
709 interaction when nerve and muscle meet, studies in both mice and zebrafish have suggested that Wnt  
710 signaling does not play an essential role. As noted in Section 3.1.2, experiments in zebrafish, involving  
711 transgenic manipulation of *unplugged* (the zebrafish version of MuSK), as well as morpholino-  
712 mediated knockdown of *Wnt11r*, which binds to *unplugged*, demonstrated that interruption of Wnt  
713 signaling leads to abolition of AChR prepatterning and impaired axonal targeting. However, these  
714 perturbations did not prevent the formation of functioning NMJs, demonstrating not only that  
715 synapse induction had occurred, but also that prepatterning is not essential for synapse formation in  
716 this system [72]. Furthermore, in a recent study employing mutation of the Wnt ligand secretion  
717 mediator (*Wls*) gene, a strategy aimed at abolition of signaling by a multitude of mammalian Wnts,  
718 Shen et al. [128] found that ablation of *Wls* function in muscle or Schwann cells did not result in any  
719 significant phenotype. In contrast, mutation in motoneurons led to both presynaptic and postsynaptic  
720 abnormalities, but once again NMJs did form, and the mutant mice, while exhibiting muscle  
721 weakness and reduced growth after birth, still survived. Given that some Wnts may not be  
722 dependent on *Wls*, these results cannot definitively rule out Wnt-mediated signaling as a key  
723 inductive process, but they strongly suggest that, as with zebrafish, some other mechanism is  
724 involved.

725 Very similar results were reported by Remédios et al. [73] who found that conditional deletion of  
726 *Wls* did not prevent synapses from forming in mice, nor did ablation of the cysteine-rich-domain  
727 (CRD) of MuSK that is required for Wnt binding. Thus, Wnt signaling was dispensable for synapse  
728 formation, and was not even required for generation of the AChR prepatterning. Although another  
729 group, using mice with a different genetic background, reported that deletion of the MuSK CRD did  
730 lead to profound deficits in prepatterning and synapse morphology, functional synapses were again

731 induced, inasmuch as these mice also survived [129]. Taken together, the studies from both mice and  
732 zebrafish reinforce the conclusion that Wnt signaling, like the amyloid precursor family, may be  
733 important for synapse maturation and maintenance, but is unlikely to be essential for the induction  
734 of postsynaptic differentiation. Accordingly, in a recent review of the relationship of Wnt signaling  
735 to synapse formation at the NMJ, Li et al. [71] argued that its role is essentially modulatory, in contrast  
736 to the central role played by agrin. On present evidence, this conclusion seems justified.

#### 737 4.3. Focal pericellular proteolysis

738 As discussed in Section 3.2.2.3, synapse formation in the amphibian *in vitro* system is invariably  
739 associated with a removal of pre-existing basal lamina constituents and their ultimate re-deposition,  
740 at highly elevated concentrations, along the path of nerve-muscle contact (see Figures 1 and 2). The  
741 initial removal satisfies two key requisites for an inductive process, time of occurrence and specificity  
742 with respect to cell type. In cultures with a mixed neuronal population, it occurs only at nerve-  
743 muscle contacts that go on to make synapses, and synapses only form at contacts where this removal  
744 is observed [14]. Moreover, the removal of HSPG and laminin occurs early, and it is the first indication  
745 that a synapse will form.

746 The most reasonable interpretation of these observations is that the removal of laminin and the  
747 basal lamina HSPG reflects a local proteolysis that is specific to contacts between muscle and motor  
748 neurons; this process is clearly regulated, inasmuch as the same constituents that were removed are  
749 subsequently laid down at high concentration. As is evident from Figure 1, the width of the zone of  
750 depleted basal lamina constituents is approximately five times wider than the diameter of the motor  
751 neuron, a typical finding in this system [14,102,109]. In contrast, the zone of removal of fluorescent  
752 substrates adsorbed to the cover glass is only as wide as the nerve itself [109], suggesting that the  
753 wider zone of proteolysis of laminin and HSPG in the synaptic cleft reflects not just the action of  
754 neuronal proteinases, but activation of muscle proteinases as well.

755 The essential issue, however, is to establish that the apparent proteolysis serves a functional role.  
756 It is possible that motor neurons in these cultures are inherently more proteolytically active than other  
757 nerve types, and that basal lamina removal, while correlated with subsequent synapse formation,  
758 reflects no causal link whatever. In this case, the apparent regulation of this activity, as evidenced by  
759 the ultimate re-deposition of laminin and HSPG at high concentration in the synaptic cleft, would  
760 have nothing to do with regulation of cell-cell signaling; instead it would simply be a necessary  
761 shutting down of local extracellular proteinase activity in order that a specialized synaptic basal  
762 lamina may form. In principle this question can be approached through the use of proteinase  
763 inhibitors of varying specificity, both naturally occurring and synthetic [130–132], to assess not only  
764 whether synapse formation is affected, but also, in the event of blockage, to determine the time point  
765 at which the inhibition is functionally relevant.

#### 766 4.3.1. Proteinases as modulators of nervous system processes

767 If focal proteolysis is a key mechanism in mediating cell-cell interaction at the NMJ, this would  
768 hardly be a novel phenomenon for the nervous system. The involvement of proteinases has been  
769 has been extensively reviewed, owing to their importance for a multitude of physiological and  
770 pathological processes [133,134]; these include long term potentiation [135–137], growth cone  
771 interactions [138], hippocampal cell differentiation [139], amyloid clearance [140,141], and axonal  
772 regeneration [142]. The two principle classes of proteinases involved in these processes are the  
773 metalloproteinases, including the matrix metalloproteinases (MMPs) and the ADAM (a disintegrin  
774 and metalloproteinase) family [134], and serine proteinases such as neurotrypsin and the tissue  
775 plasminogen activator/plasmin system [130,133].

776 Several factors contribute to the capability of proteinases to mediate the extraordinary variety of  
777 interactions that are required for establishing specific synaptic connections. First, there is a plethora  
778 of potential mechanisms by which pericellular proteolysis may contribute to synapse induction. Local  
779 proteolysis can modulate the activity of kinase signaling networks, cell surface receptors, and  
780 agonists such as chemokines, cytokines and growth factors; depending on the proteinase and target

781 protein, these can take the form of activating or degrading cleavages [143–146]. There is even an entire  
782 family, the proteinase-activated receptors, which explicitly respond to proteolytic removal of an N-  
783 terminal blocking sequence [147].

784 This complexity extends to the range of action of individual enzymes. A recent tabulation of  
785 reported MMP substrates ranged as high as 94 for MMP2, with 10 members of the MMP family  
786 having 39 substrates or more [148], while a proteomic analysis of the neuronal ablation of another  
787 key metalloproteinase, ADAM10, identified 91 candidate substrates [149], in addition to several  
788 dozen already tabulated in earlier reviews [150,151]. Secondly, many proteinases are secreted as  
789 inactive zymogens, and are activated only upon cleavage by another proteinase, either of the same  
790 or a different class [152,153], with reactions further modulated through the action of endogenous  
791 inhibitors [130,131]. This allows for precise specificity in generating a final product, as exemplified  
792 by the clotting system [154]. Moreover, proteinases derived from different tissues can cooperate to  
793 effect tissue remodeling [155], to the point where proteinases from different tissues can participate in  
794 a single cascade [156]. Thus an interaction between nerve and muscle surface proteinases to generate  
795 basal lamina remodeling, as proposed by Champaneria et al. [109], is well within established  
796 phenomenologies.

#### 797 4.3.2. Proteolysis as mediator of synapse maintenance

798 Finally, it should be noted that even if focal proteolysis ultimately proves not to be important  
799 for the induction of a postsynaptic apparatus, it may still play another key role, as it is well established  
800 that the activity of MMPs can mediate signaling through the release of growth factors [157]. At the  
801 NMJ it is known that a number of growth factors are bound to the heparan sulfate side chains of the  
802 basal lamina HSPG, and it is likely that their local release is a significant contributor to long-term  
803 motor neuron maintenance. The first evidence for this mechanism was provided by Fischbach and  
804 colleagues, supporting a model for localized proteolytic release of neuregulin-1 from the basal lamina  
805 [158]; since then a number of proteins have been shown to promote motor neuron survival in the  
806 context of diseases such as amyotrophic lateral sclerosis, and, like neuregulin-1 [159], to possess  
807 heparin-binding domains. These supporting proteins could be produced by nerve, glia, or muscle,  
808 and their lack could be felt either directly by the nerve or indirectly by modulating the activity of the  
809 enveloping Schwann cell. They include vascular endothelial growth factors A [160] and B [161,162],  
810 hepatocyte growth factor [163,164], pleiotrophin [165,166], glial cell-line-derived neurotrophic factor  
811 [167, 168], bone morphogenic protein 4 [169,170], and, through the intermediation of their dedicated  
812 binding proteins, insulin growth factors 1 and 2 [171–175]. In addition, several members of the FGF  
813 family, another group known to react with glycosaminoglycan side chains [164], have been shown to  
814 promote presynaptic differentiation [176]. Thus it is likely that the findings regarding the local release  
815 of neuregulin-1 apply to these other growth factors as well.

816 Local proteolytic release of growth factors could even contribute to the initial inductive process,  
817 but at present there is no evidence that any of these factors acts at this level. In contrast, they have all  
818 been shown to enhance motor neuron survival or differentiation. In principle, therefore, aberrant  
819 regulation of their release could be important in pathological conditions where a connection to  
820 proteolysis has been identified; for example, it would be of interest to examine whether growth factor  
821 availability is linked to the involvement of MMP9 in the etiology of motor neuron disease [177].


822 In summary, depending on the outcome of further researches into synapse induction, we are  
823 thus left with the distinct possibility that focal proteolysis, advanced herein as an alternative to agrin  
824 as an inductive process, may instead, like agrin, turn out to be a contributor to the stability of synaptic  
825 connections, while the key inductive interaction remains to be found elsewhere.

#### 826 4.4. Integrin signaling

827 Integrins are membrane spanning heterodimers, composed of non-covalently linked  $\alpha$  and  $\beta$   
828 subunits, that act as adhesion receptors to mediate signaling through interactions with extracellular  
829 matrix ligands such as fibronectin and laminin [178–180]. This signaling is essential for proper  
830 synaptic function in several loci of the nervous system [181]. In the hippocampus, dendritic spine

831 enlargement and remodeling is mediated through  $\beta$ 1 integrin signaling that in turn is dependent on  
832 the activity of the matrix metalloproteinase MMP9 [135,182], while at the NMJ blockage of  $\beta$ 1 integrin  
833 function in the early postnatal period or following injury leads to motor neuron death [183]. In  
834 addition, studies with cultured myotubes demonstrated that  $\beta$ 1 integrins co-localize with AChR  
835 clusters, and are essential for their formation in response to bath application of agrin and laminin  
836 [184–186].

837 Using the amphibian culture system, Anderson et al. [102,187] found that the early removal and  
838 subsequent re-deposition of basal lamina constituents is accompanied by re-organization of  
839 postsynaptic  $\beta$ 1 integrin. First, it is removed (Figure 6) and ultimately it is concentrated at the synapse,  
840 concomitant with the accumulation of laminin in the basal lamina (Figure 7). This synaptic  
841 accumulation of integrin resembles that of AChRs, with the exception that, unlike the situation with  
842 AChRs, integrin is also found over the myocyte surface (compare Figures 2 and 7). The synaptic  
843 accumulation is not surprising, since laminin is a key basal lamina ligand for integrin [180]. Thus,  
844 correlations between the two molecules may simply represent the attachment of the  $\beta$ 1 integrin to its  
845 basal lamina binding partner, without there being any direct relation between integrin signaling and  
846 subsequent synaptogenesis; this linkage may also lead to the aberrant basal lamina deposition that  
847 has been reported in mice lacking the  $\alpha$ 3 integrin subunit [188].



848

849

850

**Figure 6.** Elimination of  $\beta 1$  integrin and laminin accumulations before the onset of postsynaptic differentiation.

851

852

853

854


Integrin (A) and laminin (B) were stained with antibodies conjugated with contrasting fluorochromes, prior to nerve addition. The cultures were examined before the onset of synaptic differentiation. Note that integrin (A) and laminin (B) accumulations have both been eliminated along the entire path of nerve-muscle contact, shown by the arrows in the phase contrast reference image (C).

855

Scale: 20  $\mu$ m

856

Reproduced with permission from Anderson et al., *Mech. Dev.* 1997;67:125-39.



857

858  
859

**Figure 7.** Aggregation of  $\beta 1$  integrin in the muscle membrane matches that of laminin in the basal lamina.

860  
861  
862  
863

In this older *Xenopus* nerve-muscle co-culture, laminin (A) and  $\beta 1$  integrin (B), are stained with monoclonal antibodies conjugated to contrasting fluorochromes. The distribution of the two antigens is essentially congruent, both along the path of this well-developed synaptic contact, shown in C, and over the remainder of the muscle cell as well.

864

Scale: 20  $\mu$ m

865

Reproduced with permission from Anderson et al., Mol. Cell. Biol. 1996;16:4972-84.

866  
867  
868  
869  
870  
871  
872  
873

However, the possibility remains that the impacts of innervation on integrin distribution are causative of later events in synaptogenesis, specifically AChR clustering. If so, the use of integrin inhibitors [189,190] may shed further light on the underlying molecular interactions with key actors such as agrin, Lrp4, and MuSK. In this connection, it is of particular interest that while genetic ablation of  $\beta 1$  integrin in motor neurons has little effect, loss of the same gene in muscle abolishes nerve termination and prevents formation of the NMJ in mice, with consequent lethality [191]. Of particular note, the lethality is embryonic, in contrast to the perinatal death that results from ablating all three amyloid precursor and precursor-like proteins (see Section 4.1), meaning that, at very least,



874 integrin signaling remains a candidate for involvement in the key inductive interactions occurring  
875 between nerve and muscle at their first contact.

876 Finally, a recent report from a completely unrelated field of cell biology, the etiology of  
877 hepatocellular carcinoma, presents an intriguing possibility for a unified mechanism for  
878 synaptogenesis involving integrin, agrin and Lrp4-MuSK signaling. Agrin is overexpressed in  
879 hepatocellular carcinoma and promotes cellular proliferation [192]; further detailed analysis of this  
880 interaction has revealed that agrin acts as a bridge between both the integrin-linked kinase and Lrp4-  
881 MuSK pathways, transducing changes in extracellular matrix rigidity and ultimately resulting in the  
882 activation of the nuclear transcription factor YAP [193]; see commentary of Xiong and Mei [194]. YAP  
883 in turn has been implicated as a downstream regulator of NMJ formation [195]. While much  
884 remains to be elucidated as to the molecular details, the overall picture is consistent with the  
885 conclusions of Section 3, that agrin acts as a stabilizing factor. A possible model would thus involve  
886 an initial proteolysis-mediated remodeling of the extracellular matrix and concomitant integrin  
887 distribution, leading to signaling through integrin-mediated pathways. Agrin would act not only to  
888 potentiate these but, similar to events in liver cancer, also serve as a bridge to activate Lrp4-MuSK  
889 signaling that further stabilizes the nascent synapse.

## 890 5. Conclusions

891 In their “paradigm lost” analysis, Kummer et al. refer to the vertebrate skeletal neuromuscular  
892 junction, a reference both specific, ruling out heart and smooth muscle but including twitch and tonic,  
893 as well as general, meant to apply throughout the Vertebrata [44]. This is the traditional view, one  
894 that emphasized evolutionary parsimony at the molecular level, and that underlay the elaboration of  
895 the agrin hypothesis, recently modified to include both inductive and stabilizing functions for agrin.  
896 The principal conclusion of the present review is that there is reason to question the relevance of the  
897 inductive action of agrin. This conclusion is derived primarily from *in vivo* studies, showing that  
898 NMJs form in mice lacking the agrin gene [37,38,47,80], as well as from *in vitro* data suggesting that  
899 agrin’s deposition occurs relatively late, rather than early, during synapse formation [102,107]. The  
900 clear implication is that another molecular mechanism, as yet undefined, begins the process of  
901 postsynaptic differentiation. Several candidates for this role have been advanced herein. Only  
902 additional studies can confirm or refute the involvement of any of these proposed mechanisms, and  
903 establish whether or not synapse induction involves a fundamental molecular process that operates  
904 at every vertebrate skeletal nerve-muscle interface. Irrespective of the answer, it is safe to assume  
905 that these findings will be relevant not only to the NMJ, but also to cell-cell interaction elsewhere in  
906 the nervous system.

907 **Funding:** This research received no external funding

908 **Acknowledgments:** I thank J.W. Bloom for his critical review and essential comments on the manuscript.

909 **Conflicts of Interest:** The author declares no conflict of interest.

## 910 References

- 911 1. Langley, J.N. On the reaction of cells and of nerve-endings to certain poisons, chiefly as regards the reaction  
912 of striated muscle to nicotine and to curari. *J. Physiol.* **1905**, *33*, 374-413.
- 913 2. Dale, H.H.; Feldberg, W.; Vogt, M. Release of acetylcholine at voluntary motor nerve endings. *J. Physiol.*  
914 **1936**, *86*, 353-380.
- 915 3. Katz, B. *The release of neural transmitter substances*; Charles C. Thomas: Springfield, Illinois, USA, 1969.
- 916 4. Twain M. *The Celebrated Jumping Frog of Calaveras County. 1865*. Centennial ed.; Filter Press: Palmer Lake,  
917 Colorado, USA, 1965.
- 918 5. Gwyn, D.G.; Aitken, J.T. The formation of new motor endplates in mammalian skeletal muscle. *J. Anat.*  
919 **1996**, *100 (Pt 1)*, 111-126.

- 920 6. Fex, S.; Sonesson, B.; Thesleff, S.; Zelená, J. Nerve implants in botulinum poisoned mammalian muscle. *J.*  
921 *Physiol.* **1966**, *184*, 872-882.
- 922 7. Miledi, R. Formation of extra nerve-muscle junctions in innervated muscle. *Nature* **1963**, *199*, 1191-1192.
- 923 8. Anderson, M.J.; Cohen, M.W. Nerve-induced and spontaneous redistribution of acetylcholine receptors on  
924 cultured muscle cells. *J. Physiol.* **1977**, *268*, 757-773.
- 925 9. Frank, E.; Fischbach, G.D. Early events in neuromuscular junction formation in vitro: induction of  
926 acetylcholine receptor clusters in the postsynaptic membrane and morphology of newly formed synapses.  
927 *J. Cell Biol.* **1979**, *83*, 143-158.
- 928 10. Ziskind-Conhaim, L.; Geffen, I.; Hall, Z.W. Redistribution of acetylcholine receptors on developing rat  
929 myotubes. *J. Neurosci.* **1984**, *4*, 2346-2349.
- 930 11. Sanes, J.R.; Marshall, L.M.; McMahan, U.J. Reinnervation of muscle fiber basal lamina after removal of  
931 myofibers. Differentiation of regenerating axons at original synaptic sites. *J. Cell Biol.* **1978**, *78*, 176-198.
- 932 12. Burden, S.J.; Sargent, P.B.; McMahan, U. Acetylcholine receptors in regenerating muscle accumulate at  
933 original synaptic sites in the absence of the nerve. *J. Cell Biol.* **1979**, *182*, 412-425.
- 934 13. McMahan, U.J. The agrin hypothesis. *Cold Spring Harb. Symp. Quant. Biol.* **1990**, *55*, 407-418.
- 935 14. Anderson, M.J. Nerve-induced remodeling of muscle basal lamina during synaptogenesis. *J. Cell Biol.* **1986**,  
936 *102*, 863-877.
- 937 15. Nitkin, R.M.; Smith, M.A.; Magill, C.; Fallon, J.R.; Yao, Y.M.; Wallace, B.G.; McMahan, U.J. Identification of  
938 agrin, a synaptic organizing protein from Torpedo electric organ. *J. Cell Biol.* **1987**, *105*(6 Pt 1), 2471-2478.
- 939 16. Magill-Solc, C.; McMahan, U.J. Motor neurons contain agrin-like molecules. *J. Cell Biol.* **1988**, *107*, 1825-  
940 1833.
- 941 17. Magill-Solc, C.; McMahan, U.J. Synthesis and transport of agrin-like molecules in motor neurons. *J. Exp.*  
942 *Biol.* **1990**, *153*, 1-10.
- 943 18. Rupp, F.; Payan, D.G.; Magill-Solc, C.; Cowan, D.M.; Scheller, R.H. Structure and expression of a rat agrin.  
944 *Neuron* **1991**, *6*, 811-823.
- 945 19. Tsim, K.W.; Ruegg, M.A.; Escher, G.; Kröger, S.; McMahan, U.J. cDNA that encodes active agrin. *Neuron*  
946 **1992**, *8*, 677-689.
- 947 20. Smith, M.A.; Magill-Solc, C.; Rupp, F.; Yao, Y.M.; Schilling, J.W.; Snow, P.; McMahan, U.J. Isolation and  
948 characterization of a cDNA that encodes an agrin homolog in the marine ray. *Mol. Cell. Neurosci.* **1992**, *3*,  
949 406-417.
- 950 21. Jennings, C.G.; Dyer, S.M.; Burden, S.J. Muscle-specific trk-related receptor with a kringle domain defines  
951 a distinct class of receptor tyrosine kinases. *Proc. Natl. Acad. Sci. USA* **1993**, *90*, 2895-2899.
- 952 22. Valenzuela, D.M.; Stitt, T.N.; DiStefano, P.S.; Rojas, E.; Mattsson, K.; Compton, D.L.; Nuñez, L.; Park, J.S.;  
953 Stark, J.L.; Gies, D.R.; et al. Receptor tyrosine kinase specific for the skeletal muscle lineage: expression in  
954 embryonic muscle, at the neuromuscular junction, and after injury. *Neuron* **1995**, *15*, 573-584.
- 955 23. Gautam, M.; Noakes, P.G.; Moscoso, L.; Rupp, F.; Scheller, R.H.; Merlie, J.P.; Sanes, J.R. Defective  
956 neuromuscular synaptogenesis in agrin-deficient mutant mice. *Cell* **1996**, *85*, 525-535.
- 957 24. Burgess, R.W.; Nguyen, Q.T.; Son, Y.J.; Lichtman, J.W.; Sanes, J.R. Alternatively spliced isoforms of nerve-  
958 and muscle-derived agrin: their roles at the neuromuscular junction. *Neuron* **1999**, *23*, 33-44.
- 959 25. DeChiara, T.M.; Bowen, D.C.; Valenzuela, D.M.; Simmons, M.V.; Poueymirou, W.T.; Thomas, S.; Kinetz, E.;  
960 Compton, D.L.; Rojas, E.; Park, J.S.; et al. The receptor tyrosine kinase MuSK is required for neuromuscular  
961 junction formation in vivo. *Cell* **1996**, *85*, 501-512.
- 962 26. Glass, D.J.; Bowen, D.C.; Stitt, T.N.; Radziejewski, C.; Bruno, J.; Ryan, T.E.; Gies, D.R.; Shah, S.; Mattsson,

- 963 K.; Burden, S.J.; et al. Agrin acts via a MuSK receptor complex. *Cell* **1996**, *85*, 513-523.
- 964 27. Weatherbee, S.D.; Anderson, K.V.; Niswander, L.A. LDL-receptor-related protein 4 is  
965 crucial for formation of the neuromuscular junction. *Development* **2006**, *133*, 4993-5000. doi: 10.1242/dev.02696
- 966 28. Kim, N.; Stiegler, A.L.; Cameron, T.O.; Hallock, P.T.; Gomez, A.M.; Huang, J.H.; Hubbard, S.R.; Dustin,  
967 M.L.; Burden, S.J. Lrp4 is a receptor for Agrin and forms a complex with MuSK. *Cell* **2008**, *135*, 334-342. doi:  
968 10.1016/j.cell.2008.10.002
- 969 29. Zhang, B.; Luo, S.; Wang, Q.; Suzuki, T.; Xiong, W.C.; Mei, L. LRP4 serves as a coreceptor of agrin. *Neuron*  
970 **2008**, *60*, 285-297. doi: 10.1016/j.neuron.2008.10.006
- 971 30. Kleiman, R.J.; Reichardt, L.F. Testing the agrin hypothesis. *Cell* **1996**, *85*, 461-464.
- 972 31. Blakeslee S. How nerve meets muscle and begins to talk. Available online:  
973 <http://www.nytimes.com/1996/05/21/science/how-nerve-meets-muscle-and-begins-to-talk.html> (accessed on  
974 23/06/2019).
- 975 32. Sanes, J.R.; Lichtman, J.W. Induction, assembly, maturation and maintenance of a postsynaptic apparatus.  
976 *Nat. Rev. Neurosci.* **2001**, *2*, 791-805. doi: 10.1038/35097557
- 977 33. Yang, X.; Li, W.; Prescott, E.D.; Burden, S.J.; Wang, J.C. DNA topoisomerase IIbeta and neural development.  
978 *Science* **2000**, *287*, 131-134.
- 979 34. Yang, X.; Arber, S.; William, C.; Li, L.; Tanabe, Y.; Jessell, T.M.; Birchmeier, C.; Burden, S.J. Patterning of  
980 muscle acetylcholine receptor gene expression in the absence of motor innervation. *Neuron* **2001**, *30*, 399-  
981 410.
- 982 35. Lin, W.; Burgess, R.W.; Dominguez, B.; Pfaff, S.L.; Sanes, J.R.; Lee, K.F. Distinct roles of nerve and muscle  
983 in postsynaptic differentiation of the neuromuscular synapse. *Nature* **2001**, *410*, 1057-1064.  
984 doi:10.1038/35074025
- 985 36. Kummer, T.T.; Misgeld, T.; Lichtman, J.W.; Sanes, J.R. Nerve-independent formation of a topologically  
986 complex postsynaptic apparatus. *J. Cell Biol.* **2004**, *164*, 1077-1087. doi: 10.1083/jcb.200401115
- 987 37. Misgeld, T.; Kummer, T.T.; Lichtman, J.W.; Sanes, J.R. Agrin promotes synaptic differentiation by  
988 counteracting an inhibitory effect of neurotransmitter. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 11088-11093.  
989 doi: 10.1016/j.conb.2005.12.003
- 990 38. Lin, W.; Dominguez, B.; Yang, J.; Aryal, P.; Brandon, E.P.; Gage, F.H.; Lee, K.F. Neurotransmitter  
991 acetylcholine negatively regulates neuromuscular synapse formation by a Cdk5-dependent mechanism.  
992 *Neuron* **2005**, *46*, 569-579. doi: 10.1016/j.neuron.2005.04.002
- 993 39. Anderson, M.J.; Fambrough, D.M. Aggregates of acetylcholine receptors are associated with plaques of a  
994 basal lamina heparan sulfate proteoglycan on the surface of skeletal muscle fibers. *J. Cell Biol.* **1983**, *97*, 1396-  
995 1411.
- 996 40. Bayne, E.K.; Anderson, M.J.; Fambrough, D.M. Extracellular matrix organization in developing muscle:  
997 correlation with acetylcholine receptor aggregates. *J. Cell Biol.* **1984**, *99*, 1486-1501.
- 998 41. Chen, F.; Qian, L.; Yang, Z.H.; Huang, Y.; Ngo, S.T.; Ruan, N.J.; Wang, J.; Schneider, C.; Noakes, P.G.;  
999 Ding, Y.Q.; et al. Rapsyn interaction with calpain stabilizes AChR clusters at the neuromuscular junction.  
1000 *Neuron* **2007**, *55*, 247-260. doi: 1187 10.1016/j.neuron.2007.06.031
- 1001 42. Wang, J.Y.; Chen, F.; Fu, X.Q.; Ding, C.S.; Zhou, L.; Zhang, X.H.; Luo, Z.G. Caspase-3 cleavage of  
1002 dishevelled induces elimination of postsynaptic structures. *Dev. Cell.* **2014**, *28*, 670-684. doi: 1648  
1003 10.1016/j.devcel.2014.02.009
- 1004 43. Burden, S.J. Building the vertebrate neuromuscular synapse. *J. Neurobiol.* **2002**, *53*, 501-11. doi:  
1005 10.1002/neu.10137

- 1006 44. Kummer, T.T.; Misgeld, T.; Sanes, J.R. Assembly of the postsynaptic membrane at the neuromuscular  
1007 junction: paradigm lost. *Curr. Opin. Neurobiol.* **2006**, *16*, 74-82. doi: 10.1016/j.conb.2005.12.003
- 1008 45. Milton, J. *Paradise Lost; A Poem in Twelve Books*. II ed.; S. Simmons: London, UK, 1674.
- 1009 46. Kuhn, T.S. *The structure of scientific revolutions*; University of Chicago Press: Chicago, Illinois, 1962.
- 1010 47. Kim, N.; Burden, S.J. MuSK controls where motor axons grow and form synapses. *Nat. Neurosci.* **2008**, *11*,  
1011 19-27. doi: 10.1038/nn2026
- 1012 48. Escher, P.; Lacazette, E.; Courtet, M.; Blindenbacher, A.; Landmann, L.; Bezakova, G.; Lloyd, K.C.; Mueller,  
1013 U.; Brenner, H.R. Synapses form in skeletal muscles lacking neuregulin receptors. *Science* **2005**, *308*, 1920-  
1014 1923. doi: 10.1126/science.1108258
- 1015 49. Jaworski, A.; Burden, S.J. Neuromuscular synapse formation in mice lacking motor neuron- and skeletal  
1016 muscle-derived neuregulin-1. *J. Neurosci.* **2006**, *26*, 655-661.  
1017 doi: 10.1523/JNEUROSCI.4506-05.2006
- 1018 50. Newbern, J.; Birchmeier, C. Nrg1/ErbB signaling networks in Schwann cell development and myelination.  
1019 *Semin. Cell Dev. Biol.* **2010**, *21*, 922-928. doi: 10.1016/j.semcdb.2010.08.008
- 1020 51. Ohkawara, B.; Cabrera-Serrano, M.; Nakata, T.; Milone, M.; Asai, N.; Ito, K.; Ito, M.; Masuda, A.; Ito, Y.;  
1021 Engel, A.G.; Ohno, K. LRP4 third  $\beta$ -propeller domain mutations cause novel congenital myasthenia by  
1022 compromising agrin-mediated MuSK signaling in a position-specific manner. *Hum. Mol. Genet.* **2014**, *23*,  
1023 1856-1868. doi: 10.1093/hmg/ddt578
- 1024 52. Gomez, A.M.; Burden, S.J. The extracellular region of Lrp4 is sufficient to mediate neuromuscular synapse  
1025 formation. *Dev. Dyn.* **2011**, *240*, 2626-2633. doi: 10.1002/dvdy.22772
- 1026 53. Wu, H.; Xiong, W.C.; Mei, L. To build a synapse: signaling pathways in neuromuscular junction assembly.  
1027 *Development* **2010**, *137*, 1017-1033. doi: 10.1242/dev.038711
- 1028 54. Zong, Y.; Jin R. Structural mechanisms of the agrin-LRP4-MuSK signaling pathway in neuromuscular  
1029 junction differentiation. *Cell. Mol. Life Sci.* **2013**, *270*, 3077-3088. doi: 10.1007/s00018-012-1209-9
- 1030 55. Burden, S.J.; Huijbers, M.G.; Remedio, L. Fundamental molecules and mechanisms for forming and  
1031 maintaining neuromuscular synapses. *Int. J. Mol. Sci.* **2018**, *19*, 490. doi: 10.3390/ijms19020490
- 1032 56. Flanagan-Steet H.; Fox M.A.; Meyer, D.; Sanes, J.R. Neuromuscular synapses can form in vivo by  
1033 incorporation of initially aneural postsynaptic specializations. *Development* **2005**, *132*, 4471-4481. doi:  
1034 10.1242/dev.02044
- 1035 57. Panzer, J.A.; Song, Y.; Balice-Gordon, R.J. In vivo imaging of preferential motor axon outgrowth to and  
1036 synaptogenesis at prepatterned acetylcholine receptor clusters in embryonic zebrafish skeletal muscle. *J.*  
1037 *Neurosci.* **2006**, *26*, 934-947. doi: 10.1523/JNEUROSCI.3656-05.2006
- 1038 58. Burden, S.J. SnapShot: Neuromuscular Junction. *Cell* **2011**, *144*, 826-826.e1. doi: 10.1016/j.cell.2011.02.037
- 1039 59. Sanes, J.R.; Yamagata, M. Many paths to synaptic specificity. *Annu. Rev. Cell Dev. Biol.* **2009**, *25*, 161-195.  
1040 doi: 10.1146/annurev.cellbio.24.110707.175402
- 1041 60. Sainath, R.; Gallo, G. Cytoskeletal and signaling mechanisms of neurite formation. *Cell Tissue Res.* **2015**,  
1042 *359*, 267-278. doi: 10.1007/s00441-014-1955-0
- 1043 61. Campagna, J.A.; Ruegg, M.A.; Bixby, J.L. Evidence that agrin directly influences presynaptic differentiation  
1044 at neuromuscular junctions in vitro. *Eur. J. Neurosci.* **1997**, *9*, 2269-2283.
- 1045 62. Dimitropoulou, A.; Bixby, J.L. Motor neurite outgrowth is selectively inhibited by cell surface MuSK and  
1046 agrin. *Mol. Cell. Neurosci.* **2005**, *28*, 292-302. doi: 10.1016/j.mcn.2004.09.013
- 1047 63. Wu, H.; Lu, Y.; Shen, C.; Patel, N.; Gan, L.; Xiong, W.C.; Mei, L. Distinct roles of muscle and motoneuron  
1048 LRP4 in neuromuscular junction formation. *Neuron* **2012**, *75*, 94-107. doi: 10.1016/j.neuron.2012.04.033

- 1049 64. Yumoto, N.; Kim, N.; Burden, S.J. Lrp4 is a retrograde signal for presynaptic differentiation at  
1050 neuromuscular synapses. *Nature* **2012**, *489*, 438-442. doi: 10.1038/nature11348
- 1051 65. Braithwaite, A.W.; Harris, A.J. Neural influence on acetylcholine receptor clusters in embryonic  
1052 development of skeletal muscles. *Nature* **1979**, *279*, 549-551.
- 1053 66. Verschuuren, J.J.; Huijbers, M.G.; Plomp, J.J.; Niks, E.H.; Molenaar, P.C.; Martinez-Martinez, P.; Gomez,  
1054 A.M.; De Baets, M.H.; Losen, M. Pathophysiology of myasthenia gravis with antibodies to the acetylcholine  
1055 receptor, muscle-specific kinase and low-density lipoprotein receptor-related protein 4. *Autoimmun. Rev.*  
1056 **2013**, *12*, 918-923. doi: 10.1016/j.autrev.2013.03.001
- 1057 67. Gasperi, C.; Melms, A.; Schoser, B.; Zhang, Y.; Meltoranta, J.; Risson, V.; Schaeffer, L.; Schalke, B.; Kröger,  
1058 S. Anti-agrin autoantibodies in myasthenia gravis. *Neurology* **2014**, *82*, 1976-1983. doi:  
1059 10.1212/WNL.0000000000000478
- 1060 68. Zhang, W.; Coldefy, A.S.; Hubbard, S.R.; Burden, S.J. Agrin binds to the N-terminal region of Lrp4 protein  
1061 and stimulates association between Lrp4 and the first immunoglobulin-like domain in muscle-specific  
1062 kinase (MuSK). *J. Biol. Chem.* **2011**, *286*, 40624-40630. doi: 10.1074/jbc.M111.279307
- 1063 69. Zong, Y.; Zhang, B.; Gu, S.; Lee, K.; Zhou, J.; Yao, G.; Figueiredo, D.; Perry, K.; Mei, L.; Jin, R. Structural  
1064 basis of agrin-LRP4-MuSK signaling. *Genes Dev.* **2012**, *26*, 247-258. doi: 10.1101/gad.180885.111
- 1065 70. Burden, S.J.; Yumoto, N.; Zhang, W. The role of MuSK in synapse formation and neuromuscular disease.  
1066 *Cold Spring Harb. Perspect. Biol.* **2013**, *5*, a009167. doi: 10.1101/cshperspect.a009167
- 1067 71. Li, L.; Xiong, W.C.; Mei, L. Neuromuscular junction formation, aging, and disorders. *Annu. Rev. Physiol.*  
1068 **2018**, *80*, 159-188. doi: 10.1146/annurev-physiol-022516-034255
- 1069 72. Jing, L.; Lefebvre, J.L.; Gordon, L.R.; Granato, M. Wnt signals organize synaptic prepattern and axon  
1070 guidance through the zebrafish unplugged/MuSK receptor. *Neuron* **2009**, *61*, 721-733. doi:  
1071 10.1016/j.neuron.2008.12.025
- 1072 73. Remédio, L.; Gribble, K.D.; Lee, J.K.; Kim, N.; Hallock, P.T.; Delestrée, N.; Mentis, G.Z.; Froemke, R.C.;  
1073 Granato, M.; Burden, S.J. Diverging roles for Lrp4 and Wnt signaling in neuromuscular synapse  
1074 development during evolution. *Genes Dev.* **2016**, *30*, 1058-1069. doi: 10.1101/gad.279745.116
- 1075 74. Hunt, C.C.; Kuffler, S.W. Motor innervation of skeletal muscle: multiple innervation of individual muscle  
1076 fibres and motor unit function. *J. Physiol.* **1954**, *126*, 293-303.
- 1077 75. Ogata, T. Structure of motor endplates in the different fiber types of vertebrate skeletal muscles. *Arch.*  
1078 *Histol. Cytol.* **1988**, *51*, 385-424.
- 1079 76. Arber, S.; Burden, S.J.; Harris, A.J. Patterning of skeletal muscle. *Curr. Opin. Neurobiol.* **2001**, *12*, 100-103.
- 1080 77. Sanes, J.R.; Lichtman, J.W. Development of the vertebrate neuromuscular junction. *Annu. Rev. Neurosci.*  
1081 **1999**, *22*, 389-442. doi: 10.1146/annurev.neuro.22.1.389
- 1082 78. Chen, F.; Liu, Y.; Sugiura, Y.; Allen, P.D.; Gregg, R.G.; Lin, W. Neuromuscular synaptic patterning requires  
1083 the function of skeletal muscle dihydropyridine receptors. *Nat. Neurosci.* **2011**, *14*, 570-577. doi:  
1084 10.1038/nn.2792
- 1085 79. Sander, A.; Hesser, B.A.; Witzemann, V. MuSK induces in vivo acetylcholine receptor clusters in a ligand-  
1086 independent manner. *J. Cell Biol.* **2001**, *155*, 1287-1296. doi: 10.1083/jcb.200105034
- 1087 80. Tezuka, T.; Inoue, A.; Hoshi, T.; Weatherbee, S.D.; Burgess, R.W.; Ueta, R.; Yamanashi, Y. The MuSK  
1088 activator agrin has a separate role essential for postnatal maintenance of neuromuscular synapses. *Proc.*  
1089 *Natl. Acad. Sci. USA* **2014**, *111*, 16556-16561. doi: 10.1073/pnas.1408409111
- 1090 81. Cohen, M.W.; Weldon, P.R. Localization of acetylcholine receptors and synaptic ultrastructure at nerve-  
1091 muscle contacts in culture: dependence on nerve type. *J. Cell Biol.* **1980**, *86*, 388-401.

- 1092 82. Ma, E.; Morgan, R.; Godfrey, E.W. Agrin mRNA variants are differentially regulated in developing chick  
1093 embryo spinal cord and sensory ganglia. *J. Neurobiol.* **1995**, *26*, 585-597. doi: 10.1002/neu.480260411
- 1094 83. Bezakova, G.; Helm, J.P.; Francolini, M.; Lømo, T. Effects of purified recombinant neural and muscle agrin  
1095 on skeletal muscle fibers in vivo. *J. Cell Biol.* **2001**, *153*, 1441-1452.
- 1096 84. Jones, G.; Meier, T.; Lichtsteiner, M.; Witzemann, V.; Sakmann, B.; Brenner, H.R. Induction by agrin of  
1097 ectopic and functional postsynaptic-like membrane in innervated muscle. *Proc. Natl. Acad. Sci. USA* **1997**,  
1098 *94*, 2654-2659. doi: 10.1073/pnas.94.6.2654
- 1099 85. Bezakova, G.; Rabben, I.; Sefland, I.; Fumagalli, G.; Lømo, T. Neural agrin controls acetylcholine receptor  
1100 stability in skeletal muscle fibers. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 9924-9929. doi: 10.1073/pnas.171539698
- 1101 86. Ma, J.; Lugo, B.; Shah, S.; Godfrey, E.W.; Daniels, M.P. Synaptic localization and axonal targeting of agrin  
1102 secreted by ventral spinal cord neurons in culture. *J. Neurobiol.* **2000**, *43*, 338-351.
- 1103 87. Peng, H.B.; Baker, L.P.; Dai, Z. A role of tyrosine phosphorylation in the formation of acetylcholine receptor  
1104 clusters induced by electric fields in cultured *Xenopus* muscle cells. *J. Cell Biol.* **1993**, *120*, 197-204.
- 1105 88. Zhang, H.L.; Peng, H.B. Mechanism of acetylcholine receptor cluster formation induced by DC electric  
1106 field. *PLoS ONE* **2011**, *6*, e26805. doi: 10.1371/journal.pone.0026805
- 1107 89. Anderson, M.J.; Champaneria, S.; Swenarchuk, L.E. Synaptic differentiation can be evoked by polymer  
1108 microbeads that mimic localized pericellular proteolysis by removing proteins from adjacent surfaces. *Dev.*  
1109 *Biol.* **1991**, *147*, 464-479.
- 1110 90. Jones, G.; Herczeg, A.; Ruegg, M.A.; Lichtsteiner, M.; Kröger, S.; Brenner, H.R. Substrate-bound agrin  
1111 induces expression of acetylcholine receptor epsilon-subunit gene in cultured mammalian muscle cells.  
1112 *Proc. Natl. Acad. Sci. USA* **1996**, *93*, 5985-5990.
- 1113 91. Kidokoro, Y.; Anderson, M.J.; Gruener, R. Changes in synaptic potential properties during acetylcholine  
1114 receptor accumulation and neurospecific interactions in *Xenopus* nerve-muscle cell culture. *Dev. Biol.* **1980**,  
1115 *78*, 464-483.
- 1116 92. Anderson, M.J.; Klier, F.G.; Tanguay, K.E. Acetylcholine receptor aggregation parallels the deposition of a  
1117 basal lamina proteoglycan during development of the neuromuscular junction. *J. Cell Biol.* **1984**, *99*, 1769-  
1118 1784.
- 1119 93. Bruneau, E.; Sutter, D.; Hume, R.I.; Akaaboune, M. Identification of nicotinic acetylcholine receptor  
1120 recycling and its role in maintaining receptor density at the neuromuscular junction in vivo. *J. Neurosci.*  
1121 **2005**, *25*, 9949-9959. doi: 10.1523/JNEUROSCI.3169-05.2005
- 1122 94. Singhal, N.; Martin, P.T. Role of extracellular matrix proteins and their receptors in the development of the  
1123 vertebrate neuromuscular junction. *Dev. Neurobiol.* **2011**, *71*, 982-1005. doi: 10.1002/dneu.20953
- 1124 95. Burgess, R.W.; Skarnes, W.C.; Sanes, J.R. Agrin isoforms with distinct amino termini: differential  
1125 expression, localization, and function. *J. Cell Biol.* **2000**, *151*, 41-52.
- 1126 96. Denzer, A.J.; Brandenberger, R.; Gesemann, M.; Chiquet, M.; Ruegg, M.A. Agrin binds to the nerve-muscle  
1127 basal lamina via laminin. *J. Cell Biol.* **1997**, *137*, 671-683.
- 1128 97. Denzer, A.J.; Schulthess, T.; Fauser, C.; Schumacher, B.; Kammerer, R.A.; Engel, J.; Ruegg, M.A. Electron  
1129 microscopic structure of agrin and mapping of its binding site in laminin-1. *EMBO J.* **1998**, *17*, 335-343. doi:  
1130 10.1093/emboj/17.2.335
- 1131 98. Kammerer, R.A.; Schulthess, T.; Landwehr, R.; Schumacher, B.; Lustig, A.; Yurchenco, P.D.; Ruegg, M.A.;  
1132 Engel, J.; Denzer, A.J. Interaction of agrin with laminin requires a coiled-coil conformation of the agrin-  
1133 binding site within the laminin gamma1 chain. *EMBO J.* **1999**, *18*, 6762-6770. doi: 10.1093/emboj/18.23.6762

- 1134 99. Patton, B.L, Miner, J.H.; Chiu, A.Y.; Sanes, J.R. Distribution and function of laminins in the neuromuscular  
1135 system of developing, adult, and mutant mice. *J. Cell Biol.*, **1997**, *139*, 1507-1521.
- 1136 100. Patton, B.L. Laminins of the neuromuscular system. *Microsc. Res. Tech.* **2000**, *51*, 247-61. doi: 10.1002/1097-  
1137 0029
- 1138 101. Rogers, R.S.; Nishimune, H. The role of laminins in the organization and function of neuromuscular  
1139 junctions. *Matrix Biol.* **2017**, *57-58*, 86-105. doi: 10.1016/j.matbio.2016.08.008
- 1140 102. Anderson, M.J.; Shi, Z.Q.; Zackson, S.L. Proteolytic disruption of laminin-integrin complexes on muscle  
1141 cells during synapse formation. *Mol. Cell. Biol.* **1996**, *16*, 4972-4984.
- 1142 103. Swenarchuk, L.E.; Champaneria, S.; Anderson, M.J. Induction of a specialized muscle basal lamina at  
1143 chimaeric synapses in culture. *Development* **1990**, *110*, 51-61.
- 1144 104. Nishimune, H.; Valdez, G.; Jarad, G.; Moulson, C.L.; Müller, U.; Miner, J.H.; Sanes, J.R. Laminins promote  
1145 postsynaptic maturation by an autocrine mechanism at the neuromuscular junction. *J. Cell Biol.* **2008**, *182*,  
1146 1201-1215. doi: 10.1083/jcb.200805095
- 1147 105. Patton, B.L. Basal lamina and the organization of neuromuscular synapses. *J. Neurocytol.* **2003**, *32*, 883-903.  
1148 doi: 10.1023/B:NEUR.0000020630.74955.19
- 1149 106. Cohen, M.W.; Godfrey, E.W. Early appearance of and neuronal contribution to agrin-like molecules at  
1150 embryonic frog nerve-muscle synapses formed in culture. *J. Neurosci.* **1992**, *12*, 2982-2992.
- 1151 107. Anderson, M.J.; Shi, Z.Q.; Grawel, R.; Zackson, S.L. Erratic deposition of agrin during the formation of  
1152 *Xenopus* neuromuscular junctions in culture. *Dev. Biol.* **1995**, *170*, 1-20. doi: 10.1006/dbio.1995.1191
- 1153 108. Kuromi, H.; Kidokoro, Y. Denervation disperses acetylcholine receptor clusters at the neuromuscular  
1154 junction in *Xenopus* cultures. *Dev. Biol.* **1984**, *104*, 421-427.
- 1155 109. Champaneria, S.; Swenarchuk, L.E.; Anderson, M.J. Increases in pericellular proteolysis at developing  
1156 neuromuscular junctions in culture. *Dev. Biol.* **1992**, *149*, 261-277.
- 1157 110. Werle, M.J. Cell-to-cell signaling at the neuromuscular junction: the dynamic role of the extracellular matrix.  
1158 *Ann. N. Y. Acad. Sci.* **2008**, *1132*, 13-18. doi: 10.1196/annals.1405.035
- 1159 111. Patel, T.R.; Butler, G.; McFarlane, A.; Xie, L.; Overall, C.M.; Stetefeld, J. Site specific cleavage mediated by  
1160 MMPs regulates function of agrin. *PLoS ONE* **2012**, *7*, e43669. doi: 10.1371/journal.pone.0043669
- 1161 112. Reif, R.; Sales, S.; Hettwer, S.; Dreier, B.; Gisler, C.; Wolfel, J.; Luscher, D.; Zurlinden, A.; Stephan, A.;  
1162 Ahmed, S.; et al. Specific cleavage of agrin by neurotrypsin, a synaptic protease linked to mental  
1163 retardation. *FASEB J.* **2007**, *21*, 3468-3478. doi: 10.1096/fj.07-8800com
- 1164 113. Almenar-Queralt, A.; Kim, S.N.; Benner, C.; Herrera, C.M.; Kang, D.E.; Garcia-Bassets, I.; Goldstein, L.S.  
1165 Presenilins regulate neurotrypsin gene expression and neurotrypsin-dependent agrin cleavage via cyclic  
1166 AMP response element-binding protein (CREB) modulation. *J. Biol. Chem.* **2013**, *288*, 35222-35236. doi:  
1167 10.1074/jbc.M113.513705
- 1168 114. Valm, A.M.; Cohen, S.; Legant, W.R.; Melunis, J.; Hershberg, U.; Wait, E.; Cohen, A.R.; Davidson, M.W.;  
1169 Betzig, E.; Lippincott-Schwartz, J. Applying systems-level spectral imaging and analysis to reveal the  
1170 organelle interactome. *Nature* **2017**, *546*, 162-167. doi: 10.1038/nature22369
- 1171 115. Yogevev, S.; Shen, K. Cellular and molecular mechanisms of synaptic specificity. *Annu. Rev. Cell Dev. Biol.*  
1172 **2014**, *30*, 417-437. doi: 10.1146/annurev-cellbio-100913-012953
- 1173 116. de Wit, J.; Ghosh, A. Specification of synaptic connectivity by cell surface interactions. *Nat. Rev. Neurosci.*  
1174 **2016**, *17*, 22-35. doi: 10.1038/nrn.2015.3

- 1175 117. Hata, K.; Maeno-Hikichi, Y.; Yumoto, N.; Burden, S.J.; Landmesser, L.T. Distinct roles of different  
1176 presynaptic and postsynaptic NCAM isoforms in early motoneuron-myotube interactions required for  
1177 functional synapse formation. *J. Neurosci.* **2018**, *38*, 498-510. doi: 10.1523/JNEUROSCI.1014-17.2017
- 1178 118. Moscoso, L.M.; Cremer, H.; Sanes, J.R. Organization and reorganization of neuromuscular junctions in mice  
1179 lacking neural cell adhesion molecule, tenascin-C, or fibroblast growth factor-5. *J. Neurosci.* **1998**, *18*, 1465-  
1180 1477.
- 1181 119. Rafuse, V.F.; Polo-Parada, L.; Landmesser, L.T. Structural and functional alterations of neuromuscular  
1182 junctions in NCAM-deficient mice. *J. Neurosci.* **2000**, *20*, 6529-6539.
- 1183 120. Wang, P.; Yang, G.; Mosier, D.R.; Chang, P.; Zaidi, T.; Gong, Y.D.; Zhao, N.M.; Dominguez, B.; Lee, K.F.;  
1184 Gan, W.B.; Zheng, H. Defective neuromuscular synapses in mice lacking amyloid precursor protein (APP)  
1185 and APP-Like protein 2. *J. Neurosci.* **2005**, *25*, 1219-1225. doi: 10.1523/JNEUROSCI.4660-04.2005
- 1186 121. Klevanski, M.; Saar, M.; Baumkötter, F.; Weyer, S.W.; Kins, S.; Müller, U.C. Differential role of APP and  
1187 APLPs for neuromuscular synaptic morphology and function. *Mol. Cell. Neurosci.* **2014**, *61*, 201-210. doi:  
1188 10.1016/j.mcn.2014.06.004
- 1189 122. Choi, H.Y.; Liu, Y.; Tennert, C.; Sugiura, Y.; Karakatsani, A.; Kröger, S.; Johnson, E.B.; Hammer, R.E.; Lin,  
1190 W.; Herz, J. APP interacts with LRP4 and agrin to coordinate the development of the neuromuscular  
1191 junction in mice. *Elife* **2013**, *2*, e00220. doi: 10.7554/eLife.00220
- 1192 123. Herms, J.; Anliker, B.; Heber, S.; Ring, S.; Fuhrmann, M.; Kretschmar, H.; Sisodia, S.; Müller, U. Cortical  
1193 dysplasia resembling human type 2 lissencephaly in mice lacking all three APP family members. *EMBO J.*  
1194 **2004**, *23*, 4106-4115. doi: 10.1038/sj.emboj.7600390
- 1195 124. Caldwell, J.H.; Klevanski, M.; Saar, M.; Müller, U.C. Roles of the amyloid precursor protein family in the  
1196 peripheral nervous system. *Mech. Dev.* **2013**, *130*, 433-446. doi: 10.1016/j.mod.2012.11.001
- 1197 125. Luo, Z.G.; Wang, Q.; Zhou, J.Z.; Wang, J.; Luo, Z.; Liu, M.; He, X.; Wynshaw-Boris, A.; Xiong, W.C.; Lu, B.;  
1198 Mei, L. Regulation of AChR clustering by Dishevelled interacting with MuSK and PAK1. *Neuron* **2002**, *35*,  
1199 489-505.
- 1200 126. Barik, A.; Zhang, B.; Sohal, G.S.; Xiong, W.C.; Mei, L. Crosstalk between Agrin and Wnt signaling pathways  
1201 in development of vertebrate neuromuscular junction. *Dev. Neurobiol.* **2014**, *74*, 828-838. doi:  
1202 10.1002/dneu.22190
- 1203 127. He, C-W.; Liao, C-P.; Pan, C-L. Wnt signalling in the development of axon, dendrites and synapses. *Open*  
1204 *Biol.* **2018**, *8*, 180116. doi: 10.1098/rsob.180116
- 1205 128. Shen, C.; Li, L.; Zhao, K.; Bai, L.; Wang, A.; Shu, X.; Xiao, Y.; Zhang, J.; Zhang, K.; Hui, T.; et al. Motoneuron  
1206 Wnts regulate neuromuscular junction development. *Elife* **2018**, *7*, e34625. doi: 10.7554/eLife.34625
- 1207 129. Messéant, J.; Dobbertin, A.; Girard, E.; Delers, P.; Manuel, M.; Mangione, F.; Schmitt, A.; Le Denmat, D.;  
1208 Molgó, J.; Zytnicki, D.; et al. MuSK frizzled-like domain is critical for mammalian neuromuscular junction  
1209 formation and maintenance. *J. Neurosci.* **2015**, *35*, 4926-4941. doi: 10.1523/JNEUROSCI.3381-14.2015
- 1210 130. Almonte, A.G.; Sweatt, J.D. Serine proteases, serine protease inhibitors, and protease-activated receptors:  
1211 roles in synaptic function and behavior. *Brain Res.* **2011**, *1407*, 107-122. doi: 10.1016/j.brainres.2011.06.042
- 1212 131. Baranger, K.; Rivera, S.; Liechti, F.D.; Grandgirard, D.; Bigas, J.; Seco, J.; Tarrago, T.; Leib, S.L.;  
1213 Khrestchatsky, M. Endogenous and synthetic MMP inhibitors in CNS physiopathology. *Prog. Brain.*  
1214 *Res.* **2014**, *214*, 313-351. doi:10.1016/B978-0-444-63486-3.00014-1
- 1215 132. Vandembroucke, R.; Libert, C. Is there new hope for therapeutic matrix metalloproteinase inhibition? *Nat.*  
1216 *Rev. Drug Discov.* **2014**, *13*, 904-927. doi: 10.1038/nrd4390
- 1217 133. Tsilibary, E.; Tzinia, A.; Radenovic, L.; Stamenkovic, V.; Lebitko, T.; Mucha, M.; Pawlak, R.; Frischknecht,



- 1218 R.; Kaczmarek, L. Neural ECM proteases in learning and synaptic plasticity. *Prog. Brain Res.* **2014**, *214*, 135-  
1219 157. doi: 10.1016/B978-0-444-63486-3.00006-2
- 1220 134. De Stefano, M.E.; Herrero, M.T. The multifaceted role of metalloproteinases in physiological and  
1221 pathological conditions in embryonic and adult brains. *Prog. Neurobiol.* **2017**, *155*, 36-56. doi:  
1222 10.1016/j.pneurobio.2016.08.002
- 1223 135. Wang, X.B.; Bozdagi, O.; Nikitczuk, J.S.; Zhai, Z.W.; Zhou, Q.; Huntley, G.W. Extracellular proteolysis by  
1224 matrix metalloproteinase-9 drives dendritic spine enlargement and long-term potentiation coordinately.  
1225 *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 19520-19525. doi: 10.1073/pnas.0807248105
- 1226 136. Attwood, B.; Bourgognon, J.M.; Patel, S.; Mucha, M.; Schiavon, E.; Skrzypiec, A.E.; Young, K.W.; Shiosaka,  
1227 S.; Korostynski, M.; Piechota, M.; et al. Neuropsin cleaves EphB2 in the amygdala to control anxiety. *Nature*  
1228 **2011**, *473*, 372-375. doi: 10.1038/nature09938
- 1229 137. Huntley, G.W. Synaptic circuit remodelling by matrix metalloproteinases in health and disease. *Nat. Rev.*  
1230 *Neurosci.* **2012**, *13*, 743-757. doi: 10.1038/nrn3320
- 1231 138. Lin, K.T.; Sloniowski, S.; Ethell, D.W.; Ethell, I.M. Ephrin-B2-induced cleavage of EphB2 receptor is  
1232 mediated by matrix metalloproteinases to trigger cell repulsion. *J. Biol. Chem.* **2008**, *283*, 28969-28979. doi:  
1233 10.1074/jbc.M804401200
- 1234 139. Valente, M.M.; Allen, M.; Bortolotto, V.; Lim, S.T.; Conant, K.; Grilli, M. The MMP-1/PAR-1 axis enhances  
1235 proliferation and neuronal differentiation of adult hippocampal neural progenitor cells. *Neural Plast.* **2015**,  
1236 *2015*, 646595. doi: 10.1155/2015/646595
- 1237 140. Yin, K.J.; Cirrito, J.R.; Yan, P.; Hu, X.; Xiao, Q.; Pan, X.; Bateman, R.; Song, H.; Hsu, F.F.; Turk, J.; et al. Matrix  
1238 metalloproteinases expressed by astrocytes mediate extracellular amyloid-beta peptide catabolism. *J.*  
1239 *Neurosci.* **2006**, *26*, 10939-10948. doi: 10.1523/JNEUROSCI.2085-06.2006
- 1240 141. Ridnour, L.A.; Dhanapal, S.; Hoos, M.; Wilson, J.; Lee, J.; Cheng, R.Y.; Brueggemann, E.E.; Hines, H.B.;  
1241 Wilcock, D.M.; Vitek, M.P.; et al. Nitric oxide-mediated regulation of  $\beta$ -amyloid clearance via alterations of  
1242 MMP-9/TIMP-1. *J. Neurochem.* **2012**, *123*, 736-749. doi: 10.1111/jnc.12028
- 1243 142. Andries, L.; Van Hove, I.; Moons, L.; De Groef, L. Matrix metalloproteinases during axonal regeneration, a  
1244 multifactorial role from start to finish. *Mol. Neurobiol.* **2017**, *54*, 2114-2125. doi: 10.1007/s12035-016-9801-x
- 1245 143. Rodríguez, D.; Morrison, C.J.; Overall, C.M. Matrix metalloproteinases: what do they not do? New  
1246 substrates and biological roles identified by murine models and proteomics. *Biochim. Biophys. Acta* **2010**,  
1247 *1803*, 39-54. doi: 10.1016/j.bbamcr.2009.09.015
- 1248 144. Mason, S.D.; Joyce J.A. Proteolytic networks in cancer. *Trends Cell Biol.* **2011**, *21*, 228-237. doi:  
1249 10.1016/j.tcb.2010.12.002
- 1250 145. Atapattu, L.; Lackmann, M.; Janes, P.W. The role of proteases in regulating Eph/ephrin signaling. *Cell Adh.*  
1251 *Migr.* **2014**, *8*, 294-307. doi: 10.4161/19336918.2014.970026
- 1252 146. Austin, K.M.; Covic, L.; Kuliopulos, A. Matrix metalloproteases and PAR1 activation. *Blood* **2013**, *121*, 431-  
1253 439. doi: 10.1182/blood-2012-09-355958
- 1254 147. Ramachandran, R.; Altier, C.; Oikonomopoulou, K.; Hollenberg, M.D. Proteinases, their extracellular  
1255 targets, and inflammatory signaling. *Pharmacol. Rev.* **2016**, *68*, 1110-11142. doi: 10.1124/pr.115.010991
- 1256 148. Dufour, A.; Overall, C.M. Subtracting matrix out of the equation: new key roles of matrix  
1257 metalloproteinases in innate immunity and disease. In *Matrix metalloproteinase biology*; Sagi, I., Gaffney, J.P.,  
1258 Eds.; Wiley-Blackwell: Hoboken, NJ, USA, 2015. pp. 131-152.
- 1259 149. Kuhn, P.H.; Colombo, A.V.; Schusser, B.; Drey Mueller, D.; Wetzel, S.; Schepers, U.; Herber, J.; Ludwig, A.;  
1260 Kremmer, E.; Montag, D.; et al. Systematic substrate identification indicates a central role for the

- 1261 metalloprotease ADAM10 in axon targeting and synapse function. *Elife* **2016**, *5*, e12748. doi:  
1262 10.7554/eLife.12748
- 1263 150. Weber, S.; Saftig, P. Ectodomain shedding and ADAMs in development. *Development* **2012**, *139*, 3693-3709.  
1264 doi: 10.1242/dev.076398
- 1265 151. Saftig, P.; Lichtenthaler, S.F. The alpha secretase ADAM10: A metalloprotease with multiple functions in  
1266 the brain. *Prog. Neurobiol.* **2015**, *135*, 1-20. doi: 10.1016/j.pneurobio.2015.10.003
- 1267 152. Wilkins-Port C.E.; Higgins, S.P.; Higgins, C.E.; Kobori-Hotchkiss, I.; Higgins, P.J. Complex regulation of  
1268 the pericellular proteolytic microenvironment during tumor progression and wound repair: functional  
1269 interactions between the serine protease and matrix metalloproteinase cascades. *Biochem. Res. Int.* **2012**,  
1270 *2012*, 454368. doi: 10.1155/2012/454368
- 1271 153. Gaffney, J.; Solomonov, I.; Zehorai, E.; Sagi I. Multilevel regulation of matrix metalloproteinases in tissue  
1272 homeostasis indicates their molecular specificity in vivo. *Matrix Biol.* **2015**, *44-46*, 191-199. doi:  
1273 10.1016/j.matbio.2015.01.012
- 1274 154. Ben Shimon, M.; Lenz, M.; Ikenberg, B.; Becker, D.; Shavit Stein, E.; Chapman, J.; Tanne, D.; Pick, C.G.;  
1275 Blatt, I.; et al. Thrombin regulation of synaptic transmission and plasticity: implications for health and  
1276 disease. *Front. Cell. Neurosci.* *2015*, *9*, 151. doi: 10.3389/fncel.2015.00151
- 1277 155. Danø, K.; Rømer, J.; Nielsen, B.S.; Bjørn, S.; Pyke, C.; Rygaard, J.; Lund, L.R. Cancer invasion and tissue  
1278 remodeling--cooperation of protease systems and cell types. *APMIS* **1999**, *107*, 120-127.
- 1279 156. He, C.S.; Wilhelm, S.M.; Pentland, A.P.; Marmor, B.L.; Grant, G.A.; Eisen, A.Z.; Goldberg, G.I. Tissue  
1280 cooperation in a proteolytic cascade activating human interstitial collagenase. *Proc Natl. Acad. Sci. USA*  
1281 **1989**, *86*, 2632-2636.
- 1282 157. Crawford, H.C.; Stack, M.S. Matrix metalloproteinase modification of extracellular matrix-mediated  
1283 signaling. In *Matrix metalloproteinase biology*; Sagi, I., Gaffney, J.P., Eds.; Wiley-Blackwell: Hoboken, NJ,  
1284 USA, 2015. pp. 103-113.
- 1285 158. Loeb, J.A.; Fischbach, G.D. ARIA can be released from extracellular matrix through cleavage of a heparin-  
1286 binding domain. *J. Cell Biol.* **1995**, *130*, 127-135.
- 1287 159. Li, Q.; Loeb, J.A. Neuregulin-heparan-sulfate proteoglycan interactions produce sustained erbB receptor  
1288 activation required for the induction of acetylcholine receptors in muscle. *J. Biol. Chem.* **2001**, *276*, 38068-  
1289 38075. doi: 10.1074/jbc.M104485200
- 1290 160. Ruiz de Almodovar, C.; Lambrechts, D.; Mazzone, M.; Carmeliet, P. Role and therapeutic potential of  
1291 VEGF in the nervous system. *Physiol. Rev.* **2009**, *289*, 607-648.
- 1292 161. Poesen, K.; Lambrechts, D.; Van Damme, P.; Dhondt, J.; Bender, F.; Frank, N.; Bogaert, E.; Claes, B.; Heylen,  
1293 L.; Verheyen, A.; et al. Novel role for vascular endothelial growth factor (VEGF) receptor-1 and its ligand  
1294 VEGF-B in motor neuron degeneration. *J. Neurosci.* **2008**, *28*, 10451-10459. doi: 10.1523/JNEUROSCI.1092-  
1295 08.2008
- 1296 162. Bry, M.; Kivelä, R.; Leppänen, V.M.; Alitalo, K. Vascular endothelial growth factor-B in physiology and  
1297 disease. *Physiol. Rev.* **2014**, *94*, 779-794. doi: 10.1152/physrev.00028.2013
- 1298 163. Ishigaki, A.; Aoki, M.; Nagai, M.; Warita, H.; Kato, S.; Kato, M.; Nakamura, T.; Funakoshi, H.; Itoyama, Y.  
1299 Intrathecal delivery of hepatocyte growth factor from amyotrophic lateral sclerosis onset suppresses  
1300 disease progression in rat amyotrophic lateral sclerosis model. *J. Neuropathol. Exp. Neurol.* **2007**, *66*, 1037-  
1301 1044. doi: 10.1097/nen.0b013e318159886b
- 1302 164. Shute, J. Glycosaminoglycan and chemokine/growth factor interactions. *Handb. Exp. Pharmacol.* **2012**, *207*,  
1303 307-324. doi: 10.1007/978-3-642-23056-1\_13

- 1304 165. Deuel, T.F.; Zhang, N.; Yeh, H.J.; Silos-Santiago, I.; Wang, Z.Y. Pleiotrophin: a cytokine with diverse  
1305 functions and a novel signaling pathway. *Arch. Biochem. Biophys.* **2002**, *397*, 162-171. doi:  
1306 10.1006/abbi.2001.2705
- 1307 166. Mi, R.; Chen, W.; Höke, A. Pleiotrophin is a neurotrophic factor for spinal motor neurons. *Proc. Natl. Acad.*  
1308 *Sci. USA* **2007**, *104*, 4664-4669. doi: 10.1073/pnas.0603243104
- 1309 167. Henderson, C.E.; Phillips, H.S.; Pollock, R.A.; Davies, A.M.; Lemeulle, C.; Armanini, M.; Simmons, L.;  
1310 Moffet, B.; Vandlen, R.A.; Koliatsos, V.E.; Rosenthal, A. GDNF: a potent survival factor for motoneurons  
1311 present in peripheral nerve and muscle. *Science* **1994**, *266*, 1062-1064.
- 1312 168. Alfano, I.; Vora, P.; Mummery, R.S.; Mulloy, B.; Rider, C.C. The major determinant of the heparin binding  
1313 of glial cell-line-derived neurotrophic factor is near the N-terminus and is dispensable for receptor binding.  
1314 *Biochem. J.* **2007**, *404*, 131-140. doi: 10.1042/BJ20061747
- 1315 169. Chou, H.J.; Lai, D.M.; Huang, C.W.; McLennan, I.S.; Wang, H.D.; Wang, P.Y. BMP4 is a peripherally-  
1316 derived factor for motor neurons and attenuates glutamate-induced excitotoxicity in vitro. *PLoS ONE* **2013**,  
1317 *8*, e58441. doi: 10.1371/journal.pone.0058441
- 1318 170. Mercier, F.; Douet, V. Bone morphogenetic protein-4 inhibits adult neurogenesis and is regulated by  
1319 fractone-associated heparan sulfates in the subventricular zone. *J. Chem. Neuroanat.* **2014**, *57-58*, 54-61. doi:  
1320 10.1016/j.jchemneu.2014.03.005
- 1321 171. Russo, V.C.; Schütt, B.S.; Andaloro, E.; Ymer, S.I.; Hoeflich, A.; Ranke, M.B.; Bach, L.A.; Werther, G.A.  
1322 Insulin-like growth factor binding protein-2 binding to extracellular matrix plays a critical role in  
1323 neuroblastoma cell proliferation, migration, and invasion. *Endocrinology* **2005**, *146*, 4445-4455. doi:  
1324 10.1210/en.2005-0467
- 1325 172. Miyamoto, S.; Nakamura, M.; Yano, K.; Ishii, G.; Hasebe, T.; Endoh, Y.; Sangai, T.; Maeda, H.; Shi-Chuang,  
1326 Z.; Chiba, T.; Ochiai, A. Matrix metalloproteinase-7 triggers the matricrine action of insulin-like growth  
1327 factor-II via proteinase activity on insulin-like growth factor binding protein 2 in the extracellular matrix.  
1328 *Cancer Sci.* **2007**, *98*, 685-691. doi: 10.1111/j.1349-7006.2007.00448.x
- 1329 173. Sullivan, K.A.; Kim, B.; Feldman, E.L. Insulin-like growth factors in the peripheral nervous system.  
1330 *Endocrinology* **2008**, *149*, 5963-5971. doi: 10.1210/en.2008-1020
- 1331 174. Allodi, I.; Comley, L.; Nichterwitz, S.; Nizzardo, M.; Simone, C.; Benitez, J.A.; Cao, M.; Corti, S.; Hedlund,  
1332 E. Differential neuronal vulnerability identifies IGF-2 as a protective factor in ALS. *Sci. Rep.* **2016**, *6*, 25960.  
1333 doi: 10.1038/srep25960
- 1334 175. Eleftheriadou, I.; Manolaras, I.; Irvine, E.E.; Dieringer, M.; Trabalza, A.; Mazarakis, N.D.  $\alpha$ CAR IGF-1 vector  
1335 targeting of motor neurons ameliorates disease progression in ALS mice. *Ann. Clin. Transl. Neurol.* **2016**, *3*,  
1336 752-768. doi: 10.1002/acn3.335
- 1337 176. Fox, M.A.; Sanes, J.R.; Borza, D.B.; Eswarakumar, V.P.; Fässler, R.; Hudson, B.G.; John, S.W.; Ninomiya, Y.;  
1338 Pedchenko, V.; Pfaff, S.L.; et al. Distinct target-derived signals organize formation, maturation, and  
1339 maintenance of motor nerve terminals. *Cell* **2007**, *129*, 179-193. doi: 10.1016/j.cell.2007.02.035
- 1340 177. Kaplan, A.; Spiller, K.J.; Towne, C.; Kanning, K.C.; Choe, G.T.; Geber, A.; Akay, T.; Aebischer, P.;  
1341 Henderson, C.E. Neuronal matrix metalloproteinase-9 is a determinant of selective neurodegeneration.  
1342 *Neuron* **2014**, *81*, 333-348. doi: 10.1016/j.neuron.2013.12.009
- 1343 178. Campbell, I.D.; Humphries, M.J. Integrin structure, activation, and interactions. *Cold Spring Harb. Perspect.*  
1344 *Biol.* **2011**, *3*, a004994. doi:10.1101/cshperspect.a004994
- 1345 179. Wolfenson, H.; Lavelin, I.; Geiger, B.; Dynamic regulation of the structure and functions of integrin  
1346 adhesions. *Dev. Cell* **2013**, *24*, 447-458. doi: 10.1016/j.devcel.2013.02.012

- 1347 180. Yamada, M.; Sekiguchi, K. Molecular Basis of laminin-integrin interactions. *Curr. Top. Membr.* **2015**, *76*,  
1348 197-229. doi: 10.1016/bs.ctm.2015.07.002
- 1349 181. Park, Y.K.; Goda, Y. Integrins in synapse regulation. *Nat. Rev. Neurosci.* **2016**, *17*, 745-756. doi:  
1350 10.1038/nrn.2016.138
- 1351 182. Michaluk, P.; Wawrzyniak, M.; Alot, P.; Szczot, M.; Wyrembek, P.; Mercik, K.; Medvedev, N.; Wilczek, E.;  
1352 De Roo, M.; Zuschratter, W.; et al. Influence of matrix metalloproteinase MMP-9 on dendritic spine  
1353 morphology. *J. Cell Sci.* **2011**, *124* (Pt 19), 3369-3380. doi: 10.1242/jcs.090852
- 1354 183. Wong, K.C.; Meyer, T.; Harding, D.I.; Dick, J.R.; Vrbová, G.; Greensmith, L. Integrins at the neuromuscular  
1355 junction are important for motoneuron survival. *Eur. J. Neurosci.* **1999**, *11*, 3287-3292.
- 1356 184. Martin, P.T.; Sanes, J.R. Integrins mediate adhesion to agrin and modulate agrin signaling. *Development*  
1357 **1997**, *124*, 3909-3917.
- 1358 185. Burkin, D.J.; Gu, M.; Hodges, B.L.; Campanelli, J.T.; Kaufman, S.J. A functional role for specific spliced  
1359 variants of the alpha7beta1 integrin in acetylcholine receptor clustering. *J. Cell Biol.* **1998**, *143*, 1067-1075.
- 1360 186. Burkin, D.J.; Kim, J.E.; Gu, M.; Kaufman, S.J. Laminin and alpha7beta1 integrin regulate agrin-induced  
1361 clustering of acetylcholine receptors. *J. Cell Sci.* **2000**, *113*, 2877-2886.
- 1362 187. Anderson, M.J.; Shi, Z.Q.; Zackson, S.L. Nerve-induced disruption and reformation of beta1-integrin  
1363 aggregates during development of the neuromuscular junction. *Mech. Dev.* **1997**, *67*, 125-139.
- 1364 188. Ross, J.A.; Webster, R.G.; Lechertier, T.; Reynolds, L.E.; Turmaine, M.; Bencze, M.; Jamshidi, Y.; Cetin, H.;  
1365 Muntoni, F.; Beeson, D.; et al. Multiple roles of integrin- $\alpha 3$  at the neuromuscular junction. *J. Cell Sci.* **2017**,  
1366 *130*, 1772-1784. doi: 10.1242/jcs.201103
- 1367 189. Sun, C.C.; Qu, X.J.; Gao, Z.H. Arginine-Glycine-Aspartate-Binding Integrins as Therapeutic and Diagnostic  
1368 Targets. *Am. J. Ther.* **2016**, *23*, e198-207. doi: 10.1097/MJT.0000000000000053
- 1369 190. Miller, L.M.; Pritchard, J.M.; Macdonald, S.J.F.; Jamieson, C.; Watson, A.J.B. Emergence of small-molecule  
1370 non-RGD-mimetic inhibitors for RGD integrins. *J. Med. Chem.* **2017**, *60*, 3241-3251. doi:  
1371 10.1021/acs.jmedchem.6b01711
- 1372 191. Schwander, M.; Shirasaki, R.; Pfaff, S.L.; Müller, U. Beta1 integrins in muscle, but not in motor neurons, are  
1373 required for skeletal muscle innervation. *J. Neurosci.* **2004**, *24*, 8181-8191. doi: 10.1523/JNEUROSCI.1345-  
1374 04.2004
- 1375 192. Chakraborty, S.; Lakshmanan, M.; Swa, H.L.; Chen, J.; Zhang, X.; Ong, Y.S.; Loo, L.S.; Akinclar, S.C.;  
1376 Gunaratne, J.; Tergaonkar, V.; et al. An oncogenic role of agrin in regulating focal adhesion integrity in  
1377 hepatocellular carcinoma. *Nat. Commun.* **2015**, *6*, 6184. doi: 10.1038/ncomms7184
- 1378 193. Chakraborty, S.; Njah, K.; Pobbati, A.V.; Lim, Y.B.; Raju, A.; Lakshmanan, M.; Tergaonkar, V.; Lim, C.T.;  
1379 Hong, W. Agrin as a mechanotransduction signal regulating YAP through the Hippo pathway. *Cell Rep.*  
1380 **2017**, *18*, 2464-2479. doi: 10.1016/j.celrep.2017.02.041
- 1381 194. Xiong, W.C.; Mei, L. Agrin to YAP in cancer and neuromuscular junctions. *Trends Cancer* **2017**, *3*, 247-248.  
1382 doi: 10.1016/j.trecan.2017.03.005
- 1383 195. Zhao, K.; Shen, C.; Lu, Y.; Huang, Z.; Li, L.; Rand, C.D.; Pan, J.; Sun, X.D.; Tan, Z.; Wang, H.; et al. Muscle  
1384 Yap Is a regulator of neuromuscular junction formation and regeneration. *J. Neurosci.* **2017**, *37*, 3465-3477.  
1385 doi: 10.1074/jbc.M111.279307