

Review

Not peer-reviewed version

---

# The Genetic Landscape of Sleep Disorders in Parkinson's Disease

---

[Kallirhoe Kalinderi](#) <sup>\*</sup>, [Vasileios Papaliagkas](#), [Liana Fidani](#)

Posted Date: 1 December 2023

doi: 10.20944/preprints202312.0057.v1

Keywords: Parkinson's disease; sleep disorders; GBA; Parkin



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Review

# The Genetic Landscape of Sleep Disorders in Parkinson's Disease

Kallirhoe Kalinderi <sup>1,\*</sup>, Vasileios Papaliagkas <sup>2</sup> and Liana Fidani <sup>1</sup>

<sup>1</sup> Laboratory of Medical Biology-Genetics, School of Medicine, Faculty of Health Sciences, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece. Email: kkalinde@auth.gr

<sup>2</sup> Department of Biomedical Sciences, School of Health Sciences, International Hellenic University, 57400 Thessaloniki, Greece email: vpapaliagkas@gmail.com

\* Correspondence: kkalinde@auth.gr

**Abstract:** Parkinson's disease (PD) is the second most common neurodegenerative disease, affecting 1% of people aged over 60. PD is characterized by a wide range of motor symptoms, however the clinical spectrum of PD covers a wide range of nonmotor symptoms, as well. Sleep disorders are among the most common non-motor symptoms of PD, can occur at any stage of the disease and significantly affect quality of life. These include, rapid eye movement sleep behavior disorder (RBD), restless legs syndrome (RLS), excessive daytime sleepiness (EDS), insomnia, obstructive sleep apnea (OSA) and circadian rhythm disturbances. One of the main challenges in PD research is identifying individuals during the prodromal phase of the disease. Combining genetic and prodromal data may aid the early identification of individuals susceptible to PD. This review highlights current data regarding the genetic component of sleep disorders in PD patients, focusing on genes that have currently been associated with this PD co-morbidity.

**Keywords:** Parkinson's disease; sleep disorders; GBA; Parkin

## 1. Introduction

Parkinson's disease (PD) is the second most common neurodegenerative disease, after Alzheimer's disease. It affects 1% of people aged over 60 and 3% of those older than 80 years [1]. The pathological hallmarks of the disease are the loss of the dopaminergic neurons in the substantia nigra, causing dopamine depletion in the striatum, and the presence of Lewy bodies (LBs) in the remaining neurons. PD is characterized by a wide range of motor symptoms such as resting tremor, rigidity, bradykinesia, postural instability and freezing episodes. However, the clinical spectrum of PD is more extensive, covering a wide range of nonmotor symptoms, including cognitive and behavioral symptoms, sleep disorders, autonomic symptoms, sensory symptoms and fatigue [2]. Destruction of dopaminergic neurons precedes the onset of motor symptoms by many years and usually non-motor symptoms of PD occur earlier compared to motor symptoms. Current research is focused on the recognition of biological markers that could identify the disease in early stages, allowing early diagnosis and treatment of PD.

Sleep disorders, such as REM sleep behavior disorder (RBD), restless legs syndrome (RLS), excessive daytime sleepiness (EDS), insomnia, obstructive sleep apnea and circadian rhythm disturbances are common in idiopathic PD (iPD). Particularly, individuals suffering from RBD carry a risk of >85% to manifest PD after 15-20 years, and the associated neurodegenerative process is a-synucleinopathy in 95% of cases [3]. The contribution of common genetic variants to the clinical variability in sporadic PD is a hot topic of scientific research. This review highlights current data regarding the genetic component of sleep disorders in PD patients, focusing on genes that have currently been associated with this PD co-morbidity.

## 2. Genes involved in PD patients' sleep disorders

### *GBA*

*GBA* is located on chromosome 1 (1q21) and encodes for the lysosomal enzyme glucocerebrosidase (GCase). GCase catalyzes the hydrolysis of glucocerebroside into glucose and ceramide. Mutations of *GBA* are associated with Gaucher's disease (GD), a recessive lysosomal storage disorder that is characterized from reduced GCase activity and accumulation of glucocerebroside in the liver, spleen and bone marrow. Interestingly, it has been observed that both homozygote and heterozygote carriers of *GBA* mutations are at increased risk for developing PD. More specifically, the risk is significantly higher among those who are carriers of severe mutations compare to mild mutation carriers, with up to 10-fold and 2-3 fold increased risk for developing PD, respectively [4]. *GBA* variants are classified into severe variants (such as L444P, W291X, H225Q and IVS2 + 1G > A) and mild variants (such as N370S). In addition, several nonpathogenic variants of *GBA* in GD, such as E326K and T369M, have been found to increase the risk of PD. PD patients with severe variants of *GBA* have a younger age of onset, faster progression and more severe cognitive impairment than those with mild variants. *GBA* mutations are present in about 2–30% of PD patients and carrier frequency can be very different across different populations [5].

RBD is a sleep disorder characterized by abnormal motor behavior associated with dream mentation and loss of atonia during REM sleep [3]. Scientific attention over RBD has increased due to its association with  $\alpha$ -synucleinopathies. More specifically, there is accumulating evidence that the idiopathic/isolated form of RBD (IRBD) constitutes the prodromal stage of the  $\alpha$ -synucleinopathies, as most of these cases are eventually diagnosed with PD or dementia with Lewy bodies (DLB), with an estimated rate of conversion of 34% after five years from the IRBD diagnosis, 74% after ten and 91% after fourteen years. Additionally, postmortem neuropathological studies in subjects initially diagnosed with IRBD showed widespread LBs and neurites containing  $\alpha$ -synuclein as their main component [6]. RBD has been associated with *GBA* mutations both in GD patients and in PD patients. Gan-Or et al. found that *GBA* mutations carriers had an OR of 6.24 for RBD, and among PD patients, the OR for mutation carriers to have probable RBD (pRBD) was 3.13 [7]. Jesus et al. also found a higher prevalence of RBD in both deleterious *GBA*-carriers and benign *GBA*-carriers [8]. A "dose effect" of *GBA* mutations on PD phenotype was also described by Thaler et al., with a more severe PD phenotype in patients with GD and PD (GD-PD) as compared to *GBA*-related PD patients (*GBA*-PD) and iPD. These patients except for an earlier age of onset, more severe motor impairment, poorer cognition and lower olfactory scores had also a higher prevalence of RBD and higher frequencies of hallucinations compared to both *GBA*-PD and iPD [9]. Additionally, a recent study, confirmed that non-motor symptoms and a more severe and rapidly progressive disease course are seen more frequently among *GBAPD* patients in comparison to iPD. A higher occurrence of diurnal sleepiness was also observed in the carriers' group [10]. The fact that the severity of the *GBA* mutation would correspond to the PD phenotype was also addressed in the study of Thaler et al. in 2018. A total of 355 PD patients were included in this study; 152 iPD patients, 139 mild *GBA* (m*GBA*) mutation carriers, 48 severe *GBA* (s*GBA*) mutation carriers and 16 patients with GD and PD (GD-PD). Both s*GBA* and GD-PD had higher frequencies of RBD and hallucinations compared to the other groups of patients. Motor, cognitive, olfactory and psychiatric symptoms were also more severe in s*GBA* and GD-PD compared to m*GBA* and iPD, thus the severity of the PD phenotype is associated with the severity of the mutation in the *GBA* gene [11]. A recent meta-analysis study showed that PD patients with heterozygous *GBA* variants are at high risk of developing RBD. PD patients with *GBA* variants, like N370S and L444P, are at an even higher RBD risk than PD patients without these variants [3,12]. Moreover, in a recent study Perez-Lloret et al. found that *GBA*\_N370S\_rs76763715 was associated with more frequent pRBD among PD patients. In addition, some *GBA* variants, like rs2230288/E326K, rs75548401/T369M, and rs369068553/V460L, were also associated with more frequent non-motor symptoms, including RBD, as well as with a more aggressive motor disease [13].

## SNCA

The *a*-synuclein (*SNCA*) gene has been mapped to human chromosome 4q21.3-q22 and encodes for *a*-synuclein, a 140 amino acid soluble protein expressed primarily in neural tissue and especially in presynaptic terminals. *a*-Synuclein was the first gene implicated in the genetic pathogenesis of PD, especially with autosomal dominant PD. The A53T missense mutation was first identified in a large Italian–American family and in a number of Greek families [2]. Interestingly, increased expression of *a*-synuclein has been found to reduce the age of PD onset and increase the disease severity. *a*-Synuclein in its native state is a soluble and unfolded protein. Due to its central hydrophobic region *a*-synuclein has an inclination to aggregate. It initially forms an intermediate structure called an oligomer or protofibril and in succession insoluble polymers or fibrils. These insoluble fibrils are the major component of LBs [14]. The regulation of the levels of monomeric and/or oligomeric *a*-synuclein in neurons appears to be critical; this regulation might be altered in the common PD forms due to mutations in the *a*-syn gene that can lead to increased *a*-synuclein expression or decreased clearance or both [15,16].

Toffoli et al., examined 113 consecutive patients with a diagnosis of iRBD (56 patients) or PD (with or without RBD, 57 patients). The *SNCA* 3'UTR regulatory region, which is involved in gene expression, was sequenced and the *SNCA* rs356165, rs3857053, rs1045722 polymorphisms were found to be more frequent in PD patients compared to iRBD patients. Different regulating sequences are present in the 3'UTR, such as those bound by microRNAs (miRNAs); thus, genetic variants in this region can affect miRNA activity, alter the process of transcription and in the end protein production [17]. In another study, *SNCA* variants conferring risk of PD have also been found to predispose to an RBD phenotype and RBD has been proposed as a relevant marker for distinct PD subtypes. More specifically, pRBD was significantly associated with a known GWAS hit variant in the 5' region of *SNCA*, highlighting how distinctive clinical features in PD may depend on the genetic basis of disease [18]. In a recent study, RBD and other sleep abnormalities were examined in carriers of the p.A53T *SNCA* mutation, using both subjective and objective measures, such as questionnaires and polysomnography (PSG) recordings. Sleep disorders were very common in A53T-PD carriers. More specifically, RBD was highly represented and was associated with hyposmia and cognitive decline. The increased presence of RBD may correlate with the high incidence of cognitive dysfunction in this group. Thus, impaired olfaction and RBD seem to evolve in parallel with the motor symptoms of the disease [19]. In a previous study, hyposmia was associated with RBD in PD patients with the minor G allele of rs894278, thus PD patients harboring this allele were more likely to have hyposmia and RBD simultaneously and worse disease progression, highlighting that some genotypic characteristics could participate in the co-occurrence of hyposmia and RBD in PD patients [20]. Clinical-pathological studies have also suggested that both hyposmia and RBD may be related to *a*-synuclein deposition [21]. Moreover, Lahut et al. examined a large *PARK4* PD pedigree and proposed that the risk for a future manifestation of PD is reflected in the global transcriptome of blood. More specifically, subtle complexin 1 (*CPLX1*) downregulations were shown to distinguish prodromal PD cohorts (presymptomatic *PARK4* heterozygotes and RBD individuals) from controls, suggesting that *CPLX1* loss of function may act as biomarker and modifier gene of PD risk [22].

## LRRK2

The leucine-rich repeat kinase 2 (*LRRK2*) gene is located on chromosome 12q12. *LRRK2* variants have been recognized as the most common cause of both sporadic and familial PD to date. The available data suggest that the prevalence of the *LRRK2* mutation varies markedly across populations. The most prevalent *LRRK2* amino acid substitution, G2019S, is responsible for ~ 40% of familial and sporadic PD in Arab samples from North Africa, ~30% of familial PD in Ashkenazi Jewish populations, up to 6% of familial cases in Europe and up to 3% of apparently sporadic PD in Europe and North America [2,23]. The polymorphic variant G2385R has been strongly correlated to PD in the Chinese population. This variant is observed in 11.9% of PD patient, but is absent in Caucasian or Jewish patients [24]. *LRRK2* mutations have been associated with autosomal dominant PD and the clinical presentation is compatible with typical sporadic PD. However, the penetrance of



*LRRK2* mutations is incomplete and age dependent, thus a proportion of asymptomatic individuals carrying mutations can be at the premotor stage of *LRRK2*-PD. The pathological findings in patients with *LRRK2* mutations are diverse comprising LB pathology and tauopathy with neurofibrillary tangles [24]. This diversity in pathology strongly suggests that *LRRK2* is involved in multiple cellular processes and may be a central component of multiple signalling pathways that are crucial for proper functioning of neurons.

Sleep complaints have been found to be frequent in *LRRK2*-PD patients; in the study of Pont-Sunyer et al., 78% of them reported poor sleep quality, 33% sleep onset insomnia, 56% sleep fragmentation and 39% early awakening. Compared to iPD more frequent sleep onset insomnia, similar EDS and less prominent RBD were observed in *LRRK2* PD patients. RBD and EDS seem not to occur in the premotor stage of *LRRK2*-PD. Compared to iPD, sleep onset insomnia was more frequent in *LRRK2*-PD and it was associated with depressive symptoms and poor sleep quality. Moreover, the fact that RBD in *LRRK2*-PD was less frequent could suggest that damage of the brainstem structures that regulate REM sleep atonia is less prominent in *LRRK2*-PD compared to iPD patients. This might be attributed to the heterogeneous neuropathological substrate found in *LRRK2*-PD [25]. In an observational cross sectional multi-center study, 253 subjects were enrolled and standard questionnaires assessing anxiety, depression, cognition, smell, non-motor symptoms and RBD were administered. On the RBD questionnaire of G2019S *LRRK2* mutation carriers, slightly more sleep problems were observed [26]. In the Chinese population, G2385R variant carriers had higher RBD screening questionnaire score and more RBD symptoms compared with non-carriers. As RBD had been associated with a non-tremor predominant phenotype, and a higher rate of motor complications in PD patients, a relationship between RBD and motor phenotypes in PD patients with *LRRK2* variants may exist [27]. A case of a G2019S mutation carrier, affected by RLS with a positive family history of PD has also been reported. RLS is a stressful sensorimotor disorder characterized by an unpleasant urge to move the legs, that is worsened in the evening or the night, by resting or inactivity eg lying or sitting and partially or totally relieved by movement. The prevalence rate of RLS is 5%–15% in the general population. Most studies have shown a higher rate of RLS among PD patients compared to a control population [3]. The exact pathogenesis of RLS is not well understood, but the beneficial effects of dopaminergic drugs and the results of several functional imaging studies support the hypothesis of a dopaminergic nigrostriatal pathway involvement. RLS may be part of the spectrum of the clinical phenotypes related to the dopaminergic dysfunction caused by G2019S *LRRK2* mutation [28].

#### “CLOCK” GENES

The biological rhythm of the human body is regulated by the circadian system. Sleep disorders are related to alterations of the circadian rhythm that is a physiological and behavioral cycle regulated by the suprachiasmatic nucleus (SCN) [29]. The circadian rhythm is executed by the interaction network of various circadian rhythm genes in almost all organs and tissues. The, so called, “clock” genes that regulate the circadian rhythm have recently attracted a lot of scientific attention [30]. Among these genes is, the circadian locomotor output cycle Kaput (*CLOCK*) gene that regulates the circadian rhythm in the brain and periphery and can modulate dopamine synthesis [31]. Importantly, *CLOCK* can regulate the transcription of tyrosine hydroxylase [32], which is an enzyme implicated in dopamine synthesis. In the Chinese population, the *CLOCK* 3111T/C variant carriers were found to be more susceptible to the development of sleep disorders, after adjusting for related factors to sleep, including depression, severity of motor symptoms, age, sex, and disease duration. Thus, the *CLOCK* 3111T/C polymorphism could be an independent risk factor for sleep disorders in PD. Moreover, the proportion of patients with early onset PD in the carrier group was significantly higher than that in the noncarrier group, suggesting that the *CLOCK* 3111T/C variant may lower the PD age of onset [33].

The levels of expression of two other “clock” genes that regulate the circadian system, brain and muscle Arnt-like protein 1 (BMAL1) and period 1 (PER1), have been also examined through administration of melatonin (MEL). MEL is a hormone produced by the pineal gland that regulates the sleep-wake cycle and at pharmacological doses it is used to reduce sleep disorders. In rats, the

administration of rotenone shows a decreased expression of the clock genes that are restored by administering MEL [34]. Interestingly, in the study of Delgado-Lara et al. MEL was administered to PD patients and was found to increase the BMAL1 levels, without affecting PER1 levels [35]. Moreover, several studies have found abnormal expression of circadian rhythm genes in peripheral blood lymphocytes (PBLs) from PD patients. Bmal1 and brain and muscle Arnt-like protein 2 (Bmal2) have been found significantly decreased in PD, while Bmal1 levels have been positively correlated with PD severity and sleep quality [36,37]. In another study, the expression levels of “clock” genes *BMAL1*, *CLOCK*, cryptochrome 1 (*CRY1*), *PER1* and period 2 (*PER2*) were significantly decreased in the peripheral blood mononuclear cells (PBMCs) of PD as compared to controls. Plasma MEL levels were also decreased in PD patients compared to controls. Based on these results, *BMAL1*, *CLOCK*, *CRY1*, *PER1*, *PER2* and MEL were suggested to be potential biomarkers for evaluating the sleep-wake rhythm disturbances in PD patients [38,39]. Ghrelin, an orexigenic hormone, has also been found to have a strong endogenous circadian effect. Ghrelin concentration varies, with a peak in the day and a nadir at night. Its level also depends on food intake [40]. Circulating ghrelin may act on the circadian system as a potential feedback signal for the SCN. Importantly, the genetic deletion *Bmal1* abolishes the circadian rhythmicity of ghrelin signaling and affects ghrelin mRNA production. Thus, ghrelin may be linked with “clock” genes expression [41]. Interestingly, the therapeutic potential of mesenchymal stem cell-derived exosomes (MSC-EXOs) on sleep disorder in an 6-hydroxydopamine model of PD rats was recently investigated and MSC-EXOs were found to improve circadian rhythm in PD. More specifically, BMSC-derived exosomes rich in Wnt5 were effective to treat PD-related circadian rhythm dysfunction via enhanced peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) activity [42]. PPAR $\gamma$  is a member of the steroid hormone receptor superfamily, is distributed throughout the basal ganglia and is co-expressed with dopamine receptors at local brain sites. Interestingly, treatment of PD rats with BMSC quiescent-EXOs and BMSC induced-EXOs significantly reversed the effects of 6-OHDA on SCN expression of *CLOCK*, *Bmal* and *Per2* [42].

### *TEF*

Thyrotroph embryonic factor (*TEF*) gene is located on chromosome 22q13. *TEF*, is a transcription factor of the proline and acidic amino acid-rich basic leucine zipper (PAR bZip) family, which plays a major role in circadian rhythm regulation [43]. Circadian dysfunction may be a key cause of non-motor symptoms, especially for pre-motor symptoms, like sleep disturbances. The TT genotype of rs738499 *TEF* polymorphism has been associated with rapid deterioration in sleep quality in PD patients. Importantly, PD patients with more severe sleep disturbance at baseline presented a faster deterioration rate of sleep quality [43,44]. *TEF* is one of the downstream genes in the circadian gene network is expressed in various cells and tissues and could be used as a predictive marker of the progression of sleep disturbances in PD patients. Notably, *TEF* may regulate other neurotransmitters, like dopamine and serotonin [45].

### *Preprohypocretin*

A possible link between the dopaminergic and hypocretinergic system with respect to sleep-wake regulation has been suggested. Hypocretins play a major role in the pathophysiology of narcolepsy [46]. Narcoleptic patients suffer from symptoms such as EDS and shortened latency of sleep. In patients with PD, hypocretin levels were negatively correlated with disease severity. Dopamine is believed to play a crucial role in sleep regulation. Animal and postmortem studies in humans have shown that the dopaminergic system is disturbed in narcolepsy [47]. The D2-receptor binding has been also found elevated in narcolepsy and has been associated with the frequency of sleep attacks [48]. Moreover, COMT activity one of the major enzymes in dopamine metabolism has been disease implicated in narcolepsy severity [49]. Hypocretin-1 and hypocretin-2 are neuropeptides processed from a common precursor, preprohypocretin [46]. Interestingly, in the study of Rissling et al., the T allele of the -909T/C preprohypocretin polymorphism was associated with a significantly increased risk of developing sudden onset of sleep in PD patients [50].

### *Parkin*

*Parkin* is one of the largest genes in the genome mapped to human chromosome 6q25.2-27. *Parkin* mutations have been found in PD patients of different ethnicity and have been considered as the major mutant factor for familial autosomal recessive juvenile parkinsonism (ARJP), with 50% of ARJP beyond the age of 25 and 3–7% of PD patients at ages of 30–45 years, carrying a mutation in this gene [2,51]. Until now, more than 100 different mutations have been identified, including deletions, insertions, duplications, triplications and point mutations [52,53]. Interestingly, in most PD cases with *parkin* mutations LB pathology is absent, however exceptions have been observed [54]. Interestingly, in 10 patients with *parkin* mutation, RBD was frequently observed, suggesting that mechanisms other than synuclein deposition can cause RBD in PD patients [55]. Similarly, in another study in 11 consecutive patients with two *parkin* mutations, their sleep phenotype was similar to that in iPD, except that RLS was more prevalent and secondary narcolepsy was absent [56]. Thus, the  $\alpha$ -synuclein deposit appears not to be mandatory for developing RBD in neurodegenerative diseases, such as PD. In a systematic review and meta-analysis, no correlation between *parkin* variants and the risk and severity of sleep disturbances in PD patients was found [12]. Additional studies examining sleep disorders in patients with *parkin* mutations are needed.

### *ZNF184*

Zinc finger protein 184 (*ZNF184*) is located on chromosome 6 and encodes a Kruppel C2H2-type zinc-finger protein family member, which participates in gene expression regulation [57]. The genetic basis of pRBD in PD was recently examined and the *ZNF184*\_rs9468199 polymorphism was found to increase the risk of pRBD [13]. This variant was previously associated with more frequent RBD in the Chinese population [58]. Further studies are needed to elucidate the exact role of *ZNF184* in RBD in PD.

### *ANK2.CAMK2D*

Ankyrin 2 (*ANK2*) is located on chromosome 4 and encodes a member of the ankyrin family of proteins that link the integral membrane proteins to the underlying spectrin-actin cytoskeleton. *Ca<sup>2+</sup>/calmodulin-dependent protein kinase II (CAMK2)* is located on chromosome 5 and the product of this gene belongs to the serine/threonine protein kinases family, and to the *Ca(2+)/calmodulin-dependent protein kinases subfamily* [59]. *CAMK2* is a key player in synaptic plasticity and memory formation. Interestingly, *Camk2b* knockout mice show reduced sleep, whereas the phosphorylation states of *CaMK2* appear to control sleep induction and maintenance processes differently [60]. The *ANK2.CAMK2D*\_rs78738012 polymorphism was recently found to increase the risk of pRBD [13].

### *SYT17*

Synaptotagmin 17 (*SYT17*) gene is located on chromosome 16. *SYT17* has calcium ion binding, phospholipid binding and syntaxin binding activity [61]. *SYT17* is believed to be implicated in regulating circadian rhythms. The suprachiasmatic nucleus has been found to express high levels of *SYT17* [62]. Interestingly, in the study of Perez-Lloret, the *COQ7.SYT17*\_rs11343 polymorphism was found to reduce pRBD risk [13]. Further studies are required to confirm these findings.

### *USP25*

Ubiquitin Specific Peptidase 25 (*USP25*) gene is located on chromosome 21. *USP25* has been found to have a crucial role in innate immunity [63] and to regulate microglial homeostasis and neuroinflammation during neurodegeneration [64]. In a small population study, Gan-Or et al. indicated that the *USP25* rs2823357 polymorphism was associated with faster progression to synucleinopathy from RBD. More specifically, homozygous carriers of the *USP25* rs2823357 polymorphism progressed to synucleinopathies faster than others [65]. Notably, PD is a well-known synucleinopathy; *USP25* was previously implicated in a GWAS PD study [66], however this finding was not further replicated [67].

### miRNAs

MicroRNAs (miRNAs) are coded in genes, expressed, and finally processed through a multi-step pathway. miRNAs are short double-stranded ribonucleotides that down-regulate mRNAs by RNA interference (RNAi). Dysregulation of miRNAs has been reported in PD [68]. Interestingly, dysregulation of the miR-19b has been previously observed in the prodromal stage of synucleinopathies. More specifically miRNAs were examined in patients with iRBD who converted to either PD or *Dementia with Lewy bodies* (DLB) compared to those who remained disease-free. miR-19b was downregulated almost five years before the diagnosis of overt synucleinopathy, suggesting its possible relevance as a predictive biomarker for PD or DLB [69]. Interestingly, in a recent study that investigated possible biomarkers in the blood of patients at risk of developing PD, at the prodromal stage the expressions of 19a, 19b and 29a miRNAs decreased in plasma in patients at risk of developing PD [70]. Additional studies are awaited to elucidate if miRNAs can be considered as useful prodromal disease progression markers for PD/DLB.

### Prions

Prion protein (PrP) is a membrane-binding glycoprotein which is involved in molecular pathways that are very crucial for neuronal differentiation, synaptic development, cellular oxidative stress response and to cell–cell adhesion. Due to changes in its structure, PrP forms an infectious pattern called PrP<sup>Sc</sup> and has been associated with various diseases including bovine spongiform encephalopathy, scrapie and Creutzfeldt-Jakob disease [71]. Importantly, *cerebrospinal fluid* (CSF) PrP levels have been found altered in various neurodegenerative diseases [72]. Zhang et al., examined the correlation between the CSF PrP level and sleeping behavior in PD. Patients with PD complicated with RBD had significantly elevated CSF PrP mRNA and protein levels compared to either PD patients without sleeping disorder or healthy individuals [73]. The validation of PrP CSF biomarker to predict the development of PD in patients with iRBD is still in its infancy. Additional studies are needed be carried out in patients with iRBD in order to define the role of CSF biomarkers as predictors of this conversion.

## 3. Conclusions

Over the last few years, there has been a major shift in investigating the genetic background of specific PD subtypes, focusing especially on non-motor symptoms that most often precede the diagnosis of PD [45]. Sleep disorders, such as RBD, RLS, EDS and insomnia, are the most common non-motor feature of PD and often antedate PD, suggesting that sleep disorders are closely related to PD pathophysiology. One of the main challenges in PD research is identifying individuals during the prodromal phase of the disease. Combining genetic and prodromal data may aid the identification of individuals susceptible to PD. Currently, a number of genes have been associated with sleep disorders in PD, however additional studies on the genetics of this PD co-morbidity are of great importance, as genetic stratification of patients may substantially contribute to the early recognition of individuals at risk for PD. Mapping the genetic landscape of PD is awaited to lead to major advances in our understanding of the pathophysiology of this movement disorder and is anticipated to have important implications for future PD therapeutic strategies.

**Author Contributions:** writing—original draft preparation, K.K., V.P.; writing—review and editing, K.K., L.F.; visualization, L.F.; supervision, L.F.;. All authors have read and agreed to the published version of the manuscript.

**Institutional Review Board Statement:** Not applicable

**Informed Consent Statement:** Not applicable

**Conflicts of Interest:** The authors declare no conflict of interest.



## References

- de Lau, L.; Breteler, M. Epidemiology of Parkinson's disease. *Lancet Neurol.* 2006, 5, 525-535.
- Kalinderi, K.; Bostantjopoulou, S.; Fidani, L. The genetic background of Parkinson's disease: current progress and future prospects. *Acta. Neurol. Scand.* 2016, 134, 314-326.
- Thangaleela, S.; Sivamaruthi, B.S.; Kesika, P.; Mariappan, S.; Rashmi, S.; Choeisoongnarn, T.; Sittiprapaporn, P.; Chaiyasut, C. Neurological Insights into Sleep Disorders in Parkinson's Disease. *Brain Sci.* 2023, 13, 1202.
- Riboldi GM, Di Fonzo AB. GBA, Gaucher Disease, and Parkinson's Disease: From Genetic to Clinic to New Therapeutic Approaches. *Cells* 2019, 8, 364.
- Gan-Or, Z.; Amshalom, I.; Kilarski, L.L.; Bar-Shira, A.; Gana-Weisz, M.; Mirelman, A.; Marder, K.; Bressman, S.; Giladi, N.; Orr-Urtreger, A. Differential effects of severe vs mild GBA mutations on Parkinson disease. *Neurology.* 2015, 84, 880-887.
- Iranzo, A.; Tolosa, E.; Gelpi, E.; Molinuevo, J.L.; Valldeoriola, F.; Serradell, M.; Sanchez-Valle, R.; Vilaseca, I.; Lomeña, F.; Vilas, D.; Lladó, A.; Gaig, C. Santamaria J. Neurodegenerative disease status and post-mortem pathology in idiopathic rapid-eye-movement sleep behaviour disorder: an observational cohort study. *Lancet Neurol.* 2013,12:443-453.
- Gan-Or, Z.; Mirelman, A.; Postuma, R.B.; Arnulf, I.; Bar-Shira, A.; Dauvilliers, Y.; Desautels, A.; Gagnon, J.F.; Leblond, C.S.; Frauscher, B.; Alcalay, R.N.; Saunders-Pullman, R.; Bressman, S.B.; Marder, K.; Monaca, C.; Högl, B.; Orr-Urtreger, A.; Dion, P.A.; Montplaisir, J.Y.; Giladi, N.; Rouleau, G.A. GBA mutations are associated with Rapid Eye Movement Sleep Behavior Disorder. *Ann Clin Transl Neurol.* 2015, 2, 941-945.
- Jesús, S.; Huertas, I.; Bernal-Bernal, I.; Bonilla-Toribio, M.; Cáceres-Redondo, M.T.; Vargas-González, L.; Gómez-Llamas, M.; Carrillo, F.; Calderón, E.; Carballo, M.; Gómez-Garre, P.; Mir, P. GBA Variants Influence Motor and Non-Motor Features of Parkinson's Disease. *PLoS One.* 2016, 11, e0167749.
- Thaler, A.; Gurevich, T.; Bar Shira, A.; Gana Weisz, M.; Ash, E.; Shiner, T.; Orr-Urtreger, A.; Giladi, N.; Mirelman, A. A "dose" effect of mutations in the GBA gene on Parkinson's disease phenotype. *Parkinsonism Relat Disord.* 2017, 36, 47-51.
- De Michele, G.; Palmieri, G.R.; Pane, C.; Valente, E.M.; Palmieri, I.; Dello Iacovo, C.D.P.; Cuomo, N.; Giglio, A.; De Lucia, N.; Fico, T.; Perillo, S.; De Michele, G.; De Rosa, A. Motor and non-motor features in Parkinson's Disease patients carrying GBA gene mutations. *Acta Neurol Belg.* 2023, 123, 221-226.
- Thaler, A.; Bregman, N.; Gurevich, T.; Shiner, T.; Dror, Y.; Zmira, O.; Gan-Or, Z.; Bar-Shira, A.; Gana-Weisz, M.; Orr-Urtreger, A.; Giladi, N.; Mirelman, A. Parkinson's disease phenotype is influenced by the severity of the mutations in the GBA gene. *Parkinsonism Relat Disord.* 2018, 55, 45-49.
- Huang, J.; Cheng, Y.; Li, C.; Shang, H. Genetic heterogeneity on sleep disorders in Parkinson's disease: a systematic review and meta-analysis. *Transl Neurodegener.* 2022, 11, 21.
- Perez-Lloret, S.; Chevalier, G.; Bordet, S.; Barbar, H.; Capani, F.; Udovin, L.; Otero-Losada, M. The Genetic Basis of Probable REM Sleep Behavior Disorder in Parkinson's Disease. *Brain Sci.* 2023, 13, 1146.
- Ye, H.; Robak, L.A.; Yu, M.; Cykowski, M.; Shulman, J.M. Genetics and Pathogenesis of Parkinson's Syndrome. *Annu Rev Pathol.* 2023, 18, 95-121.
- Lundvig, D.; Lindersson, E.; Jensen, P. Pathogenic effects of  $\alpha$ -synuclein aggregation. *Mol Brain Res.* 2005, 134, 3-17.
- Bennett, C. The role of  $\alpha$ -synuclein in neurodegenerative diseases. *Pharmacol Ther.* 2005, 105, 311-331.
- Toffoli, M.; Dreussi, E.; Cecchin, E.; Valente, M.; Sanvilli, N.; Montico, M.; Gagno, S.; Garziera, M.; Polano, M.; Savarese, M.; Calandra-Buonaura, G.; Placidi, F.; Terzaghi, M.; Toffoli, G.; Gigli, G.L. SNCA 3'UTR genetic variants in patients with Parkinson's disease and REM sleep behavior disorder. *Neurol Sci.* 2017, 38, 1233-1240.
- Bjørnarå, K.A.; Pihlstrøm, L.; Dietrichs, E.; Toft, M. Risk variants of the  $\alpha$ -synuclein locus and REM sleep behavior disorder in Parkinson's disease: a genetic association study. *BMC Neurol.* 2018, 18, 20.
- Simitsi, A.M.; Koros, C.; Stamelou, M.; Papadimitriou, D.; Leonardos, A.; Bougea, A.; Papagiannakis, N.; Pachi, I.; Angelopoulou, E.; Lourentzos, K.; Bonakis, A.; Stefanis, L. REM sleep behavior disorder and other sleep abnormalities in p. A53T SNCA mutation carriers. *Sleep* 2021, 44, zsa248.
- Li, Y.; Kang, W.; Zhang, L.; Zhou, L.; Niu, M.; Liu, J. Hyposmia Is Associated with RBD for PD Patients with Variants of SNCA. *Front Aging Neurosci.* 2017, 9, 303.
- Ubeda-Bañon, I.; Saiz-Sanchez, D.; de la Rosa-Prieto, C.; Argandoña-Palacios, L.; Garcia-Muñozguren, S.; Martinez-Marcos, A.  $\alpha$ -Synucleinopathy in the human olfactory system in Parkinson's disease:

- involvement of calcium-binding protein- and substance P-positive cells. *Acta Neuropathol.* 2010, 119, 723-735.
22. Lahut, S.; Gispert, S.; Ömür, Ö.; Depboylu, C.; Seidel, K.; Domínguez-Bautista, J.A.; Brehm, N.; Tireli, H.; Hackmann, K.; Pirkevi, C.; Leube, B.; Ries, V.; Reim, K.; Brose, N.; den Dunnen, W.F.; Johnson, M.; Wolf, Z.; Schindewolf, M.; Schrempf, W.; Reetz, K.; Young, P.; Vadasz, D.; Frangakis, A.S.; Schröck, E.; Steinmetz, H.; Jendrach, M.; Rüb, U.; Başak, A.N.; Oertel, W.; Auburger, G. Blood RNA biomarkers in prodromal PARK4 and rapid eye movement sleep behavior disorder show role of complexin 1 loss for risk of Parkinson's disease. *Dis Model Mech.* 2017, 10, 619-631.
  23. Rocha, E.M.; Keeney, M.T.; Di Maio, R.; De Miranda, B.R.; Greenamyre, J.T. LRRK2 and idiopathic Parkinson's disease. *Trends Neurosci.* 2022, 45, 224-236.
  24. Monfrini, E.; Di Fonzo, A. Leucine-Rich Repeat Kinase (LRRK2) Genetics and Parkinson's Disease. *Adv Neurobiol.* 2017, 14, 3-30.
  25. Pont-Sunyer, C.; Iranzo, A.; Gaig, C.; Fernández-Arcos, A.; Vilas, D.; Valldeoriola, F.; Compta, Y.; Fernández-Santiago, R.; Fernández, M.; Bayés, A.; Calopa, M.; Casquero, P.; de Fàbregues, O.; Jaumà, S.; Puente, V.; Salameiro, M.; José Martí, M.; Santamaría, J.; Tolosa, E. Sleep Disorders in Parkinsonian and Nonparkinsonian LRRK2 Mutation Carriers. *PLoS One.* 2015, 10, e0132368.
  26. Mirelman, A.; Alcalay, R.N.; Saunders-Pullman, R.; Yasinovsky, K.; Thaler, A.; Gurevich, T.; Mejia-Santana, H.; Raymond, D.; Gana-Weisz, M.; Bar-Shira, A.; Ozelius, L.; Clark, L.; Orr-Urtreger, A.; Bressman, S.; Marder, K.; Giladi, N.; LRRK2 AJ consortium. Nonmotor symptoms in healthy Ashkenazi Jewish carriers of the G2019S mutation in the LRRK2 gene. *Mov Disord.* 2015, 30, 981-986.
  27. Sun, Q.; Wang, T.; Jiang, T.F.; Huang, P.; Li, D.H.; Wang, Y.; Xiao, Q.; Liu, J.; Chen, S.D. Effect of a Leucine-rich Repeat Kinase 2 Variant on Motor and Non-motor Symptoms in Chinese Parkinson's Disease Patients. *Aging Dis.* 2016, 7, 230-236.
  28. De Rosa, A.; Guacci, A.; Peluso, S.; Del Gaudio, L.; Massarelli, M.; Barbato, S.; Criscuolo, C.; De Michele, G. A case of restless leg syndrome in a family with LRRK2 gene mutation. *Int J Neurosci.* 2013, 123, 283-285.
  29. Dibner, C.; Schibler, U.; Albrecht, U. The mammalian circadian timing system: organization and coordination of central and peripheral clocks. *Annu Rev Physiol.* 2010, 72, 517-549.
  30. Leng, Y.; Musiek, E.S.; Hu, K.; Cappuccio, F.P.; Yaffe, K. Association between circadian rhythms and neurodegenerative diseases. *Lancet Neurol.* 2019, 18, 307-318.
  31. Parekh, P.K.; Ozburn, A.R.; McClung, C.A. Circadian clock genes: effects on dopamine, reward and addiction. *Alcohol* 2015, 49, 341-349.
  32. Kawarai, T.; Kawakami, H.; Yamamura, Y.; Nakamura, S. Structure and organization of the gene encoding human dopamine transporter Gene 1997, 195; 11-18.
  33. Lou, F.; Li, M.; Luo, X.; Ren, Y. CLOCK 3111T/C Variant Correlates with Motor Fluctuation and Sleep Disorders in Chinese Patients with Parkinson's Disease. *Parkinsons Dis.* 2018, 2018, 4670380.
  34. Mattam, U.; Jagota, A. Daily rhythms of serotonin metabolism and the expression of clock genes in suprachiasmatic nucleus of rotenone-induced Parkinson's disease male Wistar rat model and effect of melatonin administration, *Biogerontology* 2014, 16, 109-123.
  35. Delgado-Lara, D.L.; González-Enríquez, G.V.; Torres-Mendoza, B.M.; González-Usigli, H.; Cárdenas-Bedoya, J.; Macías-Islas, M.A.; de la Rosa, A.C.; Jiménez-Delgado, A.; Pacheco-Moisés, F.; Cruz-Serrano, J.A.; Ortiz, G.G. Effect of melatonin administration on the PER1 and BMAL1 clock genes in patients with Parkinson's disease. *Biomed Pharmacother.* 2020, 129, 110485.
  36. Cai, Y.; Liu, S.; Sothorn, R.B.; Xu, S.; Chan, P. Expression of clock genes Per1 and Bmal1 in total leukocytes in health and Parkinson's disease. *Eur J Neurol.* 2010, 17, 550-554.
  37. Ding, H.; Liu, S.; Yuan, Y.; Lin, Q.; Chan, P.; Cai, Y. Decreased expression of Bmal2 in patients with Parkinson's disease. *Neurosci Lett.* 2011, 499, 186-188.
  38. Li, T.; Cheng, C.; Jia, C.; Leng, Y.; Qian, J.; Yu, H.; Liu, Y.; Wang, N.; Yang, Y.; Al-Nusaif, M.; Le, W. Peripheral Clock System Abnormalities in Patients With Parkinson's Disease. *Front Aging Neurosci.* 2021, 13, 736026.
  39. Ding, H.; Liu, S.; Yuan, Y.; Lin, Q.; Chan, P.; Cai, Y. Decreased expression of Bmal2 in patients with Parkinson's disease. *Neurosci Lett.* 2011, 499, 186-188.
  40. Pradhan, G.; Samson, S.L.; Sun, Y. Ghrelin: much more than a hunger hormone. *Curr Opin Clin Nutr Metab Care* 2013, 16, 619-624.

41. Tarianyk, K.A.; Lytvynenko, N.V.; Shkodina, A.D.; Kaidashev, I.P. THE ROLE OF CIRCADIAN REGULATION OF GHRELIN LEVELS IN PARKINSON'S DISEASE (LITERATURE REVIEW). *Wiad Lek.* 2021, 74, 1750-1753.
42. Li, Z.; Li, Y.; Xu, X.; Gu, J.; Chen, H.; Gui, Y. Exosomes rich in Wnt5 improved circadian rhythm dysfunction via enhanced PPAR $\gamma$  activity in the 6-hydroxydopamine model of Parkinson's disease. *Neurosci Lett.* 2023, 802, 137139.
43. Hua, P.; Liu, W.; Zhao, Y.; Ding, H.; Wang, L.; Xiao, H. Tef polymorphism is associated with sleep disturbances in patients with Parkinson's disease. *Sleep Med.* 2012, 13, 297-300.
44. Hua, P.; Cui, C.; Chen, Y.; Yao, Y.; Yu, C.Y.; Xu, L.G.; Liu, W.G. Thyrotroph embryonic factor polymorphism predicts faster progression of Parkinson's disease in a longitudinal study. *J Integr Neurosci.* 2021, 20, 95-101.
45. Kalinderi, K.; Papaliagkas, V.; Fidani, L. Current genetic data on depression and anxiety in Parkinson's disease patients. *Parkinsonism Relat Disord.* 2023, 105922.
46. Mignot, E. Genetic and familial aspects of narcolepsy. *Neurology* 1998, 50, S16-22.
47. Korotkova, T.M.; Sergeeva, O.A.; Eriksson, K.S.; Haas, H.L.; Brown, R.E. Excitation of ventral tegmental area dopaminergic and nondopaminergic neurons by orexins/hypocretins. *J Neurosci.* 2003, 23, 7-11.
48. Eiseensehr, I.; Linke, R.; Tatsch, K.; von Lindeiner, H.; Kharraz, B.; Gildehaus, F.J.; Eberle, R.; Pollmacher, T.; Schuld, A.; Noachtar, S. Alteration of the striatal dopaminergic system in human narcolepsy. *Neurology* 2003, 60, 1817-1819.
49. Dauvilliers, Y.; Neidhart, E.; Lecendreux, M.; Billiard, M.; Tafti, M. MAO-A and COMT polymorphisms and gene effects in narcolepsy. *Mol Psychiatry* 2001, 6, 367-372.
50. Rissling, I.; Körner, Y.; Geller, F.; Stiasny-Kolster, K.; Oertel, W.H.; Möller, J.C. Preprohypocretin polymorphisms in Parkinson disease patients reporting "sleep attacks". *Sleep* 2005, 28, 871-875.
51. Gasser T. Genetic basis of Parkinson's dis-ease: inheritance, penetrance, and expression. *Appl Clin Genet.* 2011, 4, 67-80.
52. Abbas, N.; Lücking, C.B.; Ricard, S.; Dürr, A.; Bonifati, V.; De Michele, G.; Bouley, S.; Vaughan, J.R.; Gasser, T.; Marconi, R.; Broussolle, E.; Brefel-Courbon, C.; Harhangi, B.S.; Oostra, B.A.; Fabrizio, E.; Böhme, G.A.; Pradier, L.; Wood, N.W.; Filla, A.; Meco, G.; Deneffe, P.; Agid, Y.; Brice, A. A wide variety of mutations in the parkin gene are responsible for autosomal recessive parkinsonism in Europe. *Hum Mol Genet.* 1999, 8, 567-574.
53. Hedrich, K.; Eskelson, C.; Wilmot, B.; Marder, K.; Harris, J.; Garrels, J.; Meija-Santana, H.; Vieregge, P.; Jacobs, H.; Bressman, S.B.; Lang, A.E.; Kann, M.; Abbruzzese, G.; Martinelli, P.; Schwinger, E.; Ozelius, L.J.; Pramstaller, P.P.; Klein, C.; Kramer, P. Distribution, type, and origin of Parkin mutations: review and case studies. *Mov Disord.* 2004, 19, 1146-1157.
54. Farrer, M.; Chan, P.; Chen, R.; Tan, L.; Lincoln, S.; Hernandez, D.; Forno, L.; Gwinn-Hardy, K.; Petrucelli, L.; Hussey, J.; Singleton, A.; Tanner, C.; Hardy, J.; Langston, J.W. Lewy bodies and parkinsonism in families with parkin mutations. *Ann Neurol.* 2001, 50, 293-300.
55. Kumru, H.; Santamaria, J.; Tolosa, E.; Valldeoriola, F.; Muñoz, E.; Marti, M.J.; Iranzo, A. Rapid eye movement sleep behavior disorder in parkinsonism with parkin mutations. *Ann Neurol.* 2004, 56, 599-603.
56. Limousin, N.; Konofal, E.; Karroum, E.; Lohmann, E.; Theodorou, I.; Dürr, A.; Arnulf, I. Restless legs syndrome, rapid eye movement sleep behavior disorder, and hypersomnia in patients with two parkin mutations. *Mov Disord.* 2009, 24, 1970-1976.
57. Goldwurm, S.; Menzies, M.L.; Banyer, J.L.; Powell, L.W.; Jazwinska, E.C. Identification of a novel Krueppel-related zinc finger gene (ZNF184) mapping to 6p21.3. *Genomics* 1997, 40, 486-489.
58. Zhang, B.; Cui, C.; Yu, H.; Li, G. Association between ZNF184 and symptoms of Parkinson's disease in southern Chinese. *Neurol. Sci.* 2020, 41, 2121-2126.
59. Stevens, S.R.; Rasband, M.N. Ankyrins and neurological disease. *Curr Opin Neurobiol.* 2021, 69, 51-57.
60. Tone, D.; Ode, K.L.; Zhang, Q.; Fujishima, H.; Yamada, R.G.; Nagashima, Y.; Matsumoto, K.; Wen, Z.; Yoshida, S.Y.; Mitani, T.T.; et al. Distinct phosphorylation states of mammalian CaMKII $\alpha$  control the induction and maintenance of sleep. *PLoS Biol.* 2022, 20, e3001813.
61. Wolfes, A.C.; Dean, C. The diversity of synaptotagmin isoforms. *Curr Opin Neurobiol.* 2020, 63, 198-209.
62. Fujioka, A.; Nagano, M.; Ikegami, K.; Masumoto, K.H.; Yoshikawa, T.; Koinuma, S.; Nakahama, K.I.; Shigeyoshi, Y. Circadian expression and specific localization of synaptotagmin17 in the suprachiasmatic nucleus, the master circadian oscillator in mammals. *Brain Res.* 2023, 1798, 148129.

63. Zhong, B.; Liu, X.; Wang, X.; Chang, S.H.; Liu, X.; Wang, A.; Reynolds, J.M.; Dong, C. Negative regulation of IL-17-mediated signaling and inflammation by the ubiquitin-specific protease USP25. *Nat Immunol.* 2012, 13, 1110–1117.
64. Zheng, Q.; Li, G.; Wang, S.; Zhou, Y.; Liu, K.; Gao, Y.; Zhou, Y.; Zheng, L.; Zhu, L.; Deng, Q.; Wu, M.; Di, A.; Zhang, L.; Zhao, Y.; Zhang, H.; Sun, H.; Dong, C.; Xu, H.; Wang, X. Trisomy 21-induced dysregulation of microglial homeostasis in Alzheimer's brains is mediated by USP25. *Sci Adv.* 2021, 7, eabe1340.
65. Gan-Or, Z.; Girard, S.L.; Noreau, A.; Leblond, C.S.; Gagnon, J.F.; Arnulf, I.; Mirarchi, C.; Dauvilliers, Y.; Desautels, A.; Mitterling, T.; Cochen De Cock, V.; Frauscher, B.; Monaca, C.; Hogg, B.; Dion, P.A.; Postuma, R.B.; Montplaisir, J.Y.; Rouleau, G.A. Parkinson's Disease Genetic Loci in Rapid Eye Movement Sleep Behavior Disorder. *J Mol Neurosci.* 2015, 56, 617–262.
66. Do, C.B.; Tung, J.Y.; Dorfman, E.; Kiefer, A.K.; Drabant, E.M.; Francke, U.; Mountain, J.L.; Goldman, S.M.; Tanner, C.M.; Langston, J.W.; Wojcicki, A.; Eriksson, N. Web-based genome-wide association study identifies two novel loci and a substantial genetic component for Parkinson's disease. *PLoS Genet.* 2011, 7, e1002141.
67. Nalls, M.A.; Pankratz, N.; Lill, C.M. Do, C.B.; Hernandez, D.G.; Saad, M.; et al. Large-scale meta-analysis of genome-wide association data identifies six new risk loci for Parkinson's disease. *Nat Genet.* 2014, 46, 989–993.
68. Titze-de-Almeida, R.; Titze-de-Almeida, S.S.; Ferreira, G.G.; Brito Silva, A.P.; de Paula Brandão, P.R.; Oertel, W.H.; Schenck, C.H.; Delgado Rodrigues, R.N. microRNA signatures in prodromal REM sleep behavior disorder and early Parkinson's disease as noninvasive biomarkers. *Sleep Med.* 2021, 78, 160–168.
69. Fernández-Santiago, R.; Iranzo, A.; Gaig, C.; Serradell, M.; Fernández, M.; Tolosa, E.; Santamaría, J.; Ezquerra, M. MicroRNA association with synucleinopathy conversion in rapid eye movement behavior disorder. *Ann Neurol.* 2015, 77, 895–901. Katunina, E.A.; Blokhin, V.; Nodel, M.R.; Pavlova, E.N.; Kalinkin, A.L.; Kucheryanu, V.G.; Alekperova, L.; Selikhova, M.V.; Martynov, M.Y.; Ugrumov, M.V. Searching for Biomarkers in the Blood of Patients at Risk of Developing Parkinson's Disease at the Prodromal Stage. *Int J Mol Sci.* 2023, 24, 1842.
70. Mays, C.E.; Soto, C. The stress of prion disease. *Brain Res.* 2016, 1648, 553–560.
71. Satoh, K. CSF biomarkers for prion diseases. *Neurochem Int.* 2022, 155, 105306.
72. Zhang, W.J.; Shang, X.L.; Peng, J.; Zhou, M.H.; Sun, W.J. Expression of prion protein in the cerebrospinal fluid of patients with Parkinson's disease complicated with rapid eye movement sleep behavior disorder. *Genet Mol Res.* 2017, 16.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.