

## Article

# The radio/gamma-ray connection from 120 MHz to 230 GHz

Marcello Giroletti<sup>1,\*</sup>, Filippo D’Ammando<sup>1,2</sup> and Monica Orienti<sup>1</sup> on behalf of the *Fermi*-LAT collaboration, and Rocco Lico<sup>1,2</sup>

<sup>1</sup> INAF Istituto di Radioastronomia, via Gobetti 101, I-40129, Bologna, Italy

<sup>2</sup> Dipartimento di Fisica e Astronomia, Università di Bologna, via Ranzani 1, 40127, Bologna, Italy

\* Correspondence: giroletti@ira.inaf.it

**Abstract:** Radio loud active galactic nuclei are composed of different spatial features, each one characterized by different spectral properties in the radio band. Among them, blazars are the most common class of sources detected at gamma-rays by *Fermi*, and their radio emission is dominated by the flat spectrum compact core. In this contribution, we explore the connection between emission at high energy revealed by *Fermi* and at radio frequency. Taking as a reference the strong and very highly significant correlation found between gamma rays and cm- $\lambda$  radio emission, we explore different behaviours found as we change the energy range in gamma rays and in radio, therefore changing the physical parameters of the zones involved in the radiation emission. We find that the correlation weakens when we consider (1) gamma rays of energy above 10 GeV (except for high synchrotron peaked blazars) or (2) low frequency radio data taken by the Murchison Widefield Array; on the other hand, the correlation strengthens when we consider mm- $\lambda$  data taken by ALMA.

**Keywords:** BL Lacertae objects: general, catalogues, gamma rays: galaxies, quasars: general, radiation mechanisms: non-thermal, radio continuum: galaxies

## 1. Introduction

The existence, significance, and interpretation of a correlation between radio and gamma-ray emission in extragalactic sources has attracted interest from several authors over the last few decades [5,8,11,12,20,26,32,34].

The basic arguments favouring the existence of such correlation go from a purely theoretical reasoning, i.e. that emission in both bands naturally arises through processes involving a population of relativistic particles, to observational matters, such as the fact that radio loud (RL) active galactic nuclei (AGN) dominate the census of gamma-ray catalogues; they made up about one half of the historical EGRET third catalogue [17], and nearly 40% of the third *Fermi* Large Area Telescope (LAT) catalogue [3FGL, 3], the largest and most recent compilation of gamma-ray sources. In more detail, 98% of the RL AGNs in the third catalogue of AGN detected by *Fermi*-LAT [3LAC, 7] belong to the subclass of blazars, which are RL AGNs whose jet axis is closely aligned with our line of sight; this indicates that relativistic beaming is a key element in both the radio and high energy emission, suggesting a strong connection between the two domains. Finally, simple one-zone synchrotron self-Compton (SSC) models successfully reproduce the basic features of the broadband spectral energy distribution (SED) of several blazars (particularly of BL Lac flares, [27]), and radio luminosity itself was shown to govern the physical properties and radiation mechanisms in relativistic jets up to gamma rays in the early formulation of the blazar sequence framework [10,13].

On the other hand, there are several issues that suggest a much more complex picture. First, the gamma-ray variability time scales, in particular during flares, often reach short values, down to  $\sim 1$  hour [4,9,18,38]. This indicates that the high energy emission region must be extremely compact and thus optically thick at radio frequency, because of the synchrotron self-absorption (SSA) mechanism. Indeed, dedicated systematic radio monitoring of gamma-ray blazars shows that there are very few

sources in which there is a significant correlation between the light curves in the two regimes [31]; so-called “orphan” flares are also quite common [2,39,40]. Moreover, while it is true that most of the gamma-ray sources are RL AGNs, and in particular blazars, the opposite is definitely not true: radio galaxies are very seldom detected in gamma rays, and even among blazars the detection rate at high energy is far from unity, both for luminous and not-so-luminous objects [24,25]. Therefore, while radio emission and beaming are necessary conditions for the presence of high energy radiation, they are not sufficient.

In this paper, we review some recent results and prospects about the existence and implications of such correlation, including the study of unidentified gamma-ray sources (UGS). In Sect. 2, we review the results based on *Fermi* and GHz-frequency radio data [5], which constitute the most comprehensive dataset ever used to address this topic. In the following sections, we consider variations of the data analysis as gamma-ray or radio data of different wavelengths are considered: in Sect. 3, we focus on sources detected at energies  $E > 10$  GeV; in Sect. 4, on sources detected at radio frequencies as low as 120 MHz [16] by the Murchison Widefield Array (MWA); in Sect. 5, on sources detected as high as 230 GHz with the Atacama Large Millimeter Array (ALMA).


## 2. Radio-gamma ray connection between $\sim$ GHz and $E > 100$ MeV data

The most comprehensive approach ever used for the assessment of the connection between radio and gamma-ray data in extragalactic radio sources was presented in [5]. In that work, we considered all 599 “clean” AGNs (i.e. gamma-ray sources with a single high confidence association and no analysis flag) contained in the first catalogue of AGN detected by *Fermi*-LAT [1LAC, 1]; this was not only the largest sample ever considered but also the one in which both blazar types (BL Lacs and flat spectrum radio quasars, FSRQ) were both adequately represented; most other works were limited to bright sources, thus excluding most of the BL Lacs. Archival interferometric data with sub-arcsecond resolution in the GHz domain (typically, at 8 GHz) were available, as well as simultaneous 15 GHz data from the Owens Valley Radio Observatory (OVRO) monitoring project, at least for a subsample [37, see also Pearson, this conference]. Finally, a dedicated statistical analysis was developed in order to assess the significance of the correlation independent of the various biases that can occur in such kind of studies [35]. The main results can be summarised as follows:

- The entire 599-source sample shows a correlation characterised by a Pearson’s coefficient  $r = 0.47$ ;
- The chance probability of obtaining this value from two intrinsically uncorrelated quantities with the same dynamic range in flux density is smaller than  $10^{-7}$ ;
- Both BL Lacs and FSRQ considered separately display a strong and highly significant correlation; the correlation is stronger for BL Lacs ( $r_{\text{BL}} = 0.62$  vs  $r_{\text{FSRQ}} = 0.42$ );
- If we classify blazars according to the synchrotron component peak frequency as low-, intermediate-, high-synchrotron peaked blazars (LSP, ISP, HSP, respectively, for peak frequencies  $\nu_{\text{peak}}$  in Hz such that  $\log \nu_{\text{peak}} < 14$ ,  $14 < \log \nu_{\text{peak}} \leq 15$ ,  $\log \nu_{\text{peak}} > 15$ ), HSP blazars are the type that shows the strongest correlation;
- If we consider gamma-ray data in sub-energy bands, we find that the energy band showing the strongest correlation with radio data increases from LSP (which have the strongest correlation when gamma rays of energy between 100 and 300 MeV are considered), to ISP, to HSP (which have the peak in the band between 1 and 3 GeV);
- Considering the subset of OVRO-monitored sources, both the correlation strength  $r$  and significance  $P$  improve when considering simultaneous vs archival data ( $1.9 \times 10^{-6}$  to  $9 \times 10^{-8}$ ); we note however that this sample contains “only” 161 sources.

## 3. Radio-gamma ray connection between VLBI and $E > 10$ GeV data

While the work of [5] demonstrated the existence of a strong and highly significant correlation between radio data and high energy gamma rays, the situation is far more open when we consider




**Figure 1.** Gamma-ray energy flux at  $E > 100$  MeV vs 8 GHz radio flux density for the 1LAC sample. Adapted from [5]. Magenta symbols indicate sources without a measured redshift.

Very High Energy (VHE,  $E > 100$  GeV) gamma rays. In this domain, observations are carried out in targeted mode with imaging atmospheric Cherenkov telescopes (IACT), so that no systematic and unbiased survey exists. Sources are often detected when they are in flaring states, and the optimization of the observing strategy naturally carries on a selection bias in favour of sources physically similar to those already detected. Moreover, it is still true that all the classified extragalactic VHE sources are RL AGNs, yet their radio properties are far less extreme, with low-luminosity sources, and in particular HSP BL Lacs, being more frequently detected than luminous ones, such as LSP BL Lacs and FSRQs.

In this sense, the first *Fermi*-LAT catalogue of sources above 10 GeV [1FHL, 6] represents an ideal resource to try and connect the HE and VHE domains. The 2FHL and the surveys that will be carried out with the Cherenkov Telescope Array (CTA) in the coming years will allow us to extend our understanding. The 1FHL is based on three years of *Fermi*-LAT survey data and it is as uniform and unbiased as possible. It contains 514 sources, 76% of which are AGN and 13% un-associated; the AGN fraction is larger than in 3FGL and the census leans towards extreme spectral type blazars (HSP, 41%). The fraction of UGS is lower, thanks to a generally smaller positional uncertainty ellipses when higher energy photons are considered.

We are carrying out a project aimed at studying all the sources in the 1FHL with very long baseline interferometry (VLBI), in order to characterize the parsec scale properties of this population, which several works have highlighted to be rather peculiar [14,15,21,22,36]. Lico et al. [23] have presented new VLBI images for the less studied objects, proposing also associations for some of the UGS. Thanks to these new images, it is now possible to discuss the radio-gamma correlation for the sample (in particular, the northern subset) without any observational bias. If we consider the VLBI flux density and the 3FGL energy flux (i.e. the gamma-ray energy flux in the entire  $E > 100$  MeV band), the sample shows a strong and highly significant correlation, with  $r = 0.73$  and  $P_{\text{chance}} < 10^{-6}$ , even stronger than what presented in Sect. 2. This is likely due to the fact that VLBI observations filter out extended emission and provide a measurement of a jet region much closer to the gamma-ray zone.

However, when we consider gamma rays of energy  $E > 10$  GeV, the correlation essentially vanishes, with  $r = -0.02$ . Both weak (typically, BL Lacs) and bright (typically, FSRQs) radio sources have lower gamma-ray energy flux in this band, which is obvious as the band itself is narrower. However, the brightest sources have softer photon indexes, which results in a more pronounced



**Figure 2.** Gamma-ray energy flux at  $E > 100$  MeV (a) or  $E > 10$  GeV (b) vs 5 GHz VLBI flux density for the 1FHL sample. Blue: BL Lacs; red: FSRQs; green: BCUs; black: misaligned AGNs.

decrease of the gamma-ray energy flux for these objects (Figure 2). As a consequence, no significant trend is observed. Only if we focus on the sub-class of HSP blazars, we do find evidence of a correlation between the two bands ( $r = 0.60$ ,  $P_{\text{chance}} = 0.0032$ ).


#### 4. Radio-gamma ray connection between $\sim 100$ MHz and $E > 100$ MeV data

Although somehow counter-intuitively, low frequency observations have been shown to be useful in studying blazars and in classifying UGS [29,30,33]. The main reason is that UGSs are in general faint, and on the basis of the correlation demonstrated in [5], we can expect them to be associated to weak radio sources. Since weak radio sources are much more numerous than the bright ones, it is difficult to pinpoint the correct counterpart of a given UGS. It is therefore mandatory to add some physical information about the nature of the known gamma-ray sources and low frequency surveys are a relatively cheap way to obtain this information, e.g. by studying the spectral index.

The MWA [41] is the first operational Square Kilometer Array (SKA) precursor. It paves the way towards low frequency ( $< 200$  MHz) radio astronomy. A commissioning survey catalogue (MWACS, presented in [19]) includes  $\sim 14,000$  sources over  $6,100 \text{ deg}^2$  in the southern sky, with a  $3\sigma$  sensitivity of  $\sim 120$  mJy, a positional accuracy of  $\sim 3'$ , flux density values at 120, 150, 180 MHz, and the corresponding spectral index  $\alpha_{\text{low}}$ . We cross-correlated the MWACS with the fifth edition of the Roma BZCat [28], finding low frequency matches for 36% of all known blazars in the MWACS sky area: the detection rate is higher for FSRQs than for BL Lacs, and it is higher for gamma-ray blazars than for non-gamma-ray ones (see Table 1).

In terms of low frequency spectral indexes, blazars are flatter than the rest of the sources in MWACS, with  $\alpha_{\text{low, blazars}} = 0.51 \pm 0.05$  and  $\alpha_{\text{low, all}} = 0.81 \pm 0.01$  (Figure 3, left panel). Moreover, many sources whose  $\sim$  GHz flux density is above 120 mJy are not detected in MWA, indicating that they must have inverted spectra between 120 MHz and  $\sim 1$  GHz. Taken together, these results indicate that the flat-spectrum core remains prominent also at low frequency.

On the other hand, a linear fit to the gamma-ray energy flux vs MWA flux density at 180 MHz shows only a weak ( $r = 0.26$ ), non significant ( $P_{\text{chance}} = 0.27$ ) correlation (Figure 3, right panel). This indicates that a substantial contribution from the non-beamed lobes enters in the total radio flux density. Assuming the lobes to have a spectral index  $\alpha_1 = 0.81$  (the MWACS mean spectral index for the non-blazar population) and the core to have  $\alpha_c = 0.096$  (as indicated by the  $\sim 1$  GHz to 20



**Figure 3.** (a): cumulative distribution function for the low frequency spectral index of the entire MWACS catalogue (red line) and the blazar subset (blue line); (b): Gamma-ray energy flux at  $E > 100$  MeV vs 120 MHz flux density (main panel) or upper limits (side histogram). Adapted from [16].

GHz spectral index of the MWACS blazars), we determined that the flux density ratio of the two components is  $S_c/S_1 \sim 0.5, 3.5, 27$  at 120 MHz, 1 GHz, and 20 GHz, respectively.

A detailed presentation of this work can be found in [16]. Future deeper all-sky surveys with MWA and LOFAR [42] will further improve our detection rates and provide a clearer picture of the relative contribution of these components.


**Table 1.** Detection rates of blazars in the MWACS.

Class	All blazars		Gamma-ray blazars	
	Fraction	%	Fraction	%
Total	186/517	36%	79/174	45%
FSRQ	147/327	45%	52/71	73%
BL Lacs	23/153	15%	19/87	22%
BCU	16/37	43%	8/16	50%

## 5. Radio-gamma ray connection between 230 GHz and $E > 100$ MeV data

Deep observations at high frequency are certainly more challenging to obtain for large samples, as the atmospheric conditions are more problematic and the instrument field of view smaller. Since they provide crucial information about the most compact regions, it is however important to try to obtain millimetre data for as many sources as possible. For this reason, we selected a representative subsample of the 3LAC to be observed at 230 GHz with ALMA. This sample contains 77 sources, including 26 BL Lacs, 31 FSRQs, 16 blazars of uncertain type, 1 narrow-line Seyfert 1 galaxy, and 2 radio galaxies. In terms of SED-defined sub-classes, it contains 43, 14, and 18 blazars of the LSP, ISP, and HSP types. The gamma-ray flux and photon index distribution are well overlapped to those of the entire 3LAC, so that this sub-sample is suitable to represent the entire gamma-ray blazar population (Figure 4).

The sources were observed in ALMA Cycle 3; all targets were detected, with rms noise generally in agreement with the predicted values of 0.3–0.5 mJy beam $^{-1}$ ; some exceptions were found for the brightest sources, due to dynamic range limitations. Flux densities range between just above 1 mJy and 6.5 Jy, for the core of Cen A.



**Figure 4.** Gamma-ray photon flux and index distribution for the entire 3LAC (solid histogram) and the subset observed with ALMA (dashed histogram).

A preliminary comparison of the ALMA and *Fermi*-LAT data already shows some interesting findings. The 230 GHz flux density and the  $E > 100$  MeV energy flux are correlated with a coefficient  $r_{\text{ALMA-LAT}} = 0.55$ ; the correlation coefficient is significantly larger than what we find if we consider 1.4 GHz data, i.e.  $r_{\text{NVSS-LAT}} = 0.39$ . A more detailed analysis and discussion is in progress, including an assessment of the importance of comparing simultaneous data.

**Acknowledgments:** We thank the local organising committee for the great hospitality and for providing generous support to our participation.

We acknowledge financial contribution from grant PRIN-INAF-2014.

The *Fermi* LAT Collaboration acknowledges generous ongoing support from a number of agencies and institutes that have supported both the development and the operation of the LAT as well as scientific data analysis. These include the National Aeronautics and Space Administration and the Department of Energy in the United States, the Commissariat à l'Energie Atomique and the Centre National de la Recherche Scientifique / Institut National de Physique Nucléaire et de Physique des Particules in France, the Agenzia Spaziale Italiana and the Istituto Nazionale di Fisica Nucleare in Italy, the Ministry of Education, Culture, Sports, Science and Technology (MEXT), High Energy Accelerator Research Organization (KEK) and Japan Aerospace Exploration Agency (JAXA) in Japan, and the K. A. Wallenberg Foundation, the Swedish Research Council and the Swedish National Space Board in Sweden.

Additional support for science analysis during the operations phase is gratefully acknowledged from the Istituto Nazionale di Astrofisica in Italy and the Centre National d'Études Spatiales in France.

This scientific work makes use of the Murchison Radio-astronomy Observatory, operated by CSIRO. We acknowledge the Wajarri Yamatji people as the traditional owners of the Observatory site. Support for the operation of the MWA is provided by the Australian Government Department of Industry and Science and Department of Education (National Collaborative Research Infrastructure Strategy: NCRIS), under a contract to Curtin University administered by Astronomy Australia Limited. We acknowledge the iVEC Petabyte Data Store and the Initiative in Innovative Computing and the CUDA Center for Excellence sponsored by NVIDIA at Harvard University.

This paper makes use of the following ALMA data: ADS/JAO.ALMA#2013.1.01342.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada) and NSC and ASIAA (Taiwan), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ.

This work is based on observations obtained through the S6340 VLBA project, in the framework of the *Fermi*-NRAO cooperative agreement. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Abdo, A. A., Ackermann, M., Ajello, M., et al. The First Catalog of Active Galactic Nuclei Detected by the Fermi Large Area Telescope. *ApJ* **2010a**, *715*, 429–457.
2. Abdo, A. A., Ackermann, M., Agudo, I., et al. Fermi Large Area Telescope and Multi-wavelength Observations of the Flaring Activity of PKS 1510-089 between 2008 September and 2009 June. *ApJ* **2010b**, *721*, 1425–1447.
3. Acero, F., Ackermann, M., Ajello, M., et al. Fermi Large Area Telescope Third Source Catalog. *ApJS* **2015**, *218*, 23.
4. Ackermann, M., Ajello, M., Baldini, L., et al. Fermi Gamma-ray Space Telescope Observations of Gamma-ray Outbursts from 3C 454.3 in 2009 December and 2010 April. *ApJ* **2010**, *721*, 1383–1396.
5. Ackermann, M., Ajello, M., Allafort, A., et al. The Radio/Gamma-Ray Connection in Active Galactic Nuclei in the Era of the Fermi Large Area Telescope. *ApJ* **2011**, *741*, 30.
6. Ackermann, M., Ajello, M., Allafort, A., et al. The First Fermi-LAT Catalog of Sources above 10 GeV. *ApJS* **2013**, *209*, 34.
7. Ackermann, M., Ajello, M., Atwood, W. B., et al. The Third Catalog of Active Galactic Nuclei Detected by the Fermi Large Area Telescope. *ApJ* **2015**, *810*, 14.
8. Dondi, L., & Ghisellini, G. Gamma-ray-loud blazars and beaming. *MNRAS* **1995**, *273*, 583–595.
9. Foschini, L., Bonnoli, G., Ghisellini, G., et al. Fermi/LAT detection of extraordinary variability in the gamma-ray emission of the blazar PKS 1510-089. *A&A* **2013**, *555*, A138.
10. Fossati, G., Maraschi, L., Celotti, A., Comastri, A., & Ghisellini, G. A unifying view of the spectral energy distributions of blazars. *MNRAS* **1998**, *299*, 433–448.
11. Ghirlanda, G., Ghisellini, G., Tavecchio, F., & Foschini, L. Correlation of Fermi Large Area Telescope sources with the 20-GHz Australia Telescope Compact Array radio survey. *MNRAS* **2010**, *407*, 791–803.
12. Ghirlanda, G., Ghisellini, G., Tavecchio, F., Foschini, L., & Bonnoli, G. The radio- $\gamma$ -ray connection in Fermi blazars. *MNRAS* **2011**, *413*, 852–862.
13. Ghisellini, G., Celotti, A., Fossati, G., Maraschi, L., & Comastri, A. A theoretical unifying scheme for gamma-ray bright blazars. *MNRAS* **1998**, *301*, 451–468.
14. Giroletti, M., Giovannini, G., Feretti, L., et al. Parsec-Scale Properties of Markarian 501. *ApJ* **2004**, *600*, 127–140.
15. Giroletti, M., Giovannini, G., Taylor, G. B., & Falomo, R. A Sample of Low-Redshift BL Lacertae Objects. II. EVN and MERLIN Data and Multiwavelength Analysis. *ApJ* **2006**, *646*, 801–814.
16. Giroletti, M., Massaro, F., D'Abrusco, R., et al. High-energy sources at low radio frequency: the Murchison Widefield Array view of Fermi blazars. *A&A* **2016**, *588*, A141.
17. Hartman, R. C., Bertsch, D. L., Bloom, S. D., et al. The Third EGRET Catalog of High-Energy Gamma-Ray Sources. *ApJS* **1999**, *123*, 79–202.
18. Hayashida, M., Nalewajko, K., Madejski, G. M., et al. Rapid Variability of Blazar 3C 279 during Flaring States in 2013–2014 with Joint Fermi-LAT, NuSTAR, Swift, and Ground-Based Multiwavelength Observations. *ApJ* **2015**, *807*, 79.
19. Hurley-Walker, N., Morgan, J., Wayth, R. B., et al. The Murchison Widefield Array Commissioning Survey: A Low-Frequency Catalogue of 14 110 Compact Radio Sources over 6 100 Square Degrees. *PASA* **2014**, *31*, e045.
20. Kovalev, Y. Y., Aller, H. D., Aller, M. F., et al. The Relation Between AGN Gamma-Ray Emission and Parsec-Scale Radio Jets. *ApJ* **2009**, *696*, L17–L21.
21. Lico, R., Giroletti, M., Orienti, M., et al. VLBA monitoring of Mrk 421 at 15 GHz and 24 GHz during 2011. *A&A* **2012**, *545*, A117.
22. Lico, R., Giroletti, M., Orienti, M., et al. Very Long Baseline polarimetry and the  $\gamma$ -ray connection in Markarian 421 during the broadband campaign in 2011. *A&A* **2014**, *571*, A54.
23. Lico, R., Giroletti, M., Orienti, M., & D'Ammando, F. VLBA observations of radio faint Fermi-LAT sources above 10 GeV. *A&A* **2016**, *in press* (arXiv:1607.03415).
24. Linford, J. D., Taylor, G. B., Romani, R. W., et al. Characteristics of Gamma-ray Loud Blazars in the VLBA Imaging and Polarimetry Survey. *ApJ* **2011**, *726*, 16.
25. Lister, M. L., Aller, M. F., Aller, H. D., et al. Why Have Many of the Brightest Radio-loud Blazars Not Been Detected in Gamma-Rays by Fermi?. *ApJ* **2015**, *810*, L9.

26. Mahony, E. K., Sadler, E. M., Murphy, T., et al. High-frequency Radio Properties of Sources in the Fermi-LAT 1 year Point Source Catalog. *ApJ* **2010**, *718*, 587–595.
27. Maraschi, L., Fossati, G., Tavecchio, F., et al. Simultaneous X-Ray and TEV Observations of a Rapid Flare from Markarian 421. *ApJ* **1999**, *526*, L81–L84.
28. Massaro, E., Maselli, A., Leto, C., et al. The 5th edition of the Roma-BZCAT. A short presentation. *Ap&SS* **2015**, *357*, 75
29. Massaro, F., D'Abrusco, R., Giroletti, M., et al. Unveiling the Nature of the Unidentified Gamma-Ray Sources. III. Gamma-Ray Blazar-like Counterparts at Low Radio Frequencies. *ApJS* **2013a**, *207*, 4
30. Massaro, F., Giroletti, M., Paggi, A., et al. Blazar Spectral Properties at 74 MHz. *ApJS* **2013b**, *208*, 15
31. Max-Moerbeck, W., Hovatta, T., Richards, J. L., et al. Time correlation between the radio and gamma-ray activity in blazars and the production site of the gamma-ray emission. *MNRAS* **2014**, *445*, 428–436.
32. Muecke, A., Pohl, M., Reich, P., et al. On the correlation between radio and gamma ray luminosities of active galactic nuclei. *A&A* **1997**, *320*, 33–40.
33. Nori, M., Giroletti, M., Massaro, F., et al. Unveiling the Nature of Unidentified  $\gamma$ -Ray Sources. VI.  $\gamma$ -Ray Blazar Candidates in the WISH Survey and their Radio Properties. *ApJS* **2014**, *212*, 3
34. Padovani, P., Ghisellini, G., Fabian, A. C., & Celotti, A. Radio-loud AGN and the extragalactic gamma-ray background. *MNRAS* **1993**, *260*, L21–L24.
35. Pavlidou, V., Richards, J. L., Max-Moerbeck, W., et al. Assessing the Significance of Apparent Correlations between Radio and Gamma-Ray Blazar Fluxes. *ApJ* **2012**, *751*, 149
36. Piner, B. G., & Edwards, P. G. First-epoch VLBA Imaging of 20 New TeV Blazars. *ApJ* **2014**, *797*, 25
37. Richards, J. L., Max-Moerbeck, W., Pavlidou, V., et al. Blazars in the Fermi Era: The OVRO 40 m Telescope Monitoring Program. *ApJS* **2011**, *194*, 29
38. Saito, S., Stawarz, Ł., Tanaka, Y. T., et al. Very Rapid High-amplitude Gamma-Ray Variability in Luminous Blazar PKS 1510-089 Studied with Fermi-LAT. *ApJ* **2013**, *766*, L11
39. Spingola, C., Dallacasa, D., Orienti, M., et al. Radio follow-up of the  $\gamma$ -ray flaring gravitational lens JVAS B0218+357. *MNRAS* **2016**, *457*, 2263–2271.
40. Tavani, M., Vittorini, V., & Cavalieri, A. An Emerging Class of Gamma-ray Flares from Blazars: Beyond One-zone Models. *ApJ* **2015**, *814*, 51
41. Tingay, S. J., Goeke, R., Bowman, J. D., et al. The Murchison Widefield Array: The Square Kilometre Array Precursor at Low Radio Frequencies. *PASA* **2013**, *30*, e007
42. van Haarlem, M. P., Wise, M. W., Gunst, A. W., et al. LOFAR: The Low-Frequency ARray. *A&A* **2013**, *556*, A2



© 2016 by the authors; licensee *Preprints*, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).