THE MODIFIED KUDRYASHOV METHOD FOR THE CONFORMAL TIME FRACTIONAL (3+1)- DIMENSIONAL KADOMTSEV-PETVIASHVILI AND THE MODIFIED KAWAHARA EQUATIONS

ALPER KORKMAZ

ABSTRACT. The three dimensional conformal time fractional Kadomtsev- Petviashvili and the conformal time fractional modified Kawahara equations are solved by implementing the Kudryashov's procedure. The corresponding wave transformation reduces both equations to some ODEs. Balancing the nonlinear and the highest order derivative terms gives the structure of the solutions in the finite series form. The useful symbolic tools are used to solve the resultant algebraic systems. The solutions are expressed in explicit forms.

1. INTRODUCTION

The Kadomtsev-Petviashvili (KP) equation (sometimes it is called as the two-dimensional Korteweg- de Vries equation) appears in the earlier 70s to study the stability of solitary waves in weakly dispersive media covering fluids or plasma[1]. It is an integrable equation owing to the fact that there exists a connection with a linear spectral problem to a particular Schrödinger equation[2, 3]. Modulation equations of the two-dimensional KP are developed by examining Riemann invariants problem[4]. The two- dimensional form of the KP equation passes the Painlevé(P-) test for the integrability and has soliton-type solutions[5]. The evolution of water wave packages traveling in one direction strongly despite the slowly modulated amplitudes in both directions are studied in details by Ablowitz and Segur[6].

The three- dimensional form of the KP equation is a model to describe rapidly propagating magnetosonic waves of small amplitudes in a low β -magnetized plasma of super fluid helium[7]. The properties of soliton dynamics are also examined deeply for slightly perturbations along z direction in the same study. Bouard and Saut[8] perform a classification of the existence of the localized solitary-type waves by the sign of the transverse dispersion term and by the nonlinearity. The solitary waves of the three- dimensional KP equation both are in cylindrical form for the transverse variables and decay with algebraically optimal rate[9]. Beside the existence of the solitary wave solutions, the three- dimensional KP equation has cnoidal wave solutions to be written as infinite sum of solitons[10] and traveling wave solutions to be expressed in the form of hyperbolic, rational and trigonometric functions[11].

2010 *Mathematics Subject Classification.* Primary 35C07;35R11;35Q53.

Key words and phrases. Conformal time fractional (3+1)- dimensional Kadomtsev-Petviashvili equation, Conformal time fractional modified Kawahara equation, Modified Kudryashov method, wave solution.

The modified Kawahara (mKawahara) equation is a cubic non-linear equation with a fifth-order derivative term. The quadratic nonlinear form of the equation is suggested by Kawahara [12] with its steady solutions. Such solutions may exist due to the sign of the dispersive term and these solutions can be in an oscillatory form because of the dominant fifth-ordered term. The mKawahara equation has implicit doubly periodic solutions and they can be determined by using the method of auxiliary equation [13, 14]. Some soliton and periodic solutions are derived by using a set of forecasting methods having some trigonometric and hyperbolic functions inside [15]. The mKawahara equation has also non-constant meromorphic solutions in explicit form [16]. The existence of traveling wave type solutions in hyperbolic or trigonometric forms are discussed by Al-Ali [17]. Tanh and exp-function methods are also capable to derive the exact solutions to the mKawahara equation [18]. So far, some methods belonging to different categories have been proposed for the exact solutions of both integer or fractional ordered PDEs [19, 20, 21, 22]. In this study, the main aim is to derive some exact solutions to the three-dimensional conformal time fractional KP and the conformal time fractional mKawahara equations. The solutions are obtained explicitly by using the modified Kudryashov method. The existence of the chain rule and other required properties in the definition of the conformal derivative enable some wave transformations to generate the solutions.

2. CONFORMAL DERIVATIVE AND SOME SIGNIFICANT PROPERTIES

Let α be $\in (0, 1]$. Then, the conformal derivative of $f = f(\tau)$ defined in the positive half space $\tau > 0$ is given as

$$D_{\tau}^{\alpha}(f(\tau)) = \lim_{h \rightarrow 0} \frac{f(\tau + h\tau^{1-\alpha}) - f(\tau)}{h}, \quad \tau > 0, \alpha \in (0, 1] \quad (2.1)$$

for $f : [0, \infty) \rightarrow \mathbb{R}$ [23]. Even though it has just been defined, some significant properties covering derivative of multiplication or division of two functions. The following two theorems give a brief summary of those properties.

Theorem 1. *The conformal derivative of order $\alpha \in (0, 1]$ for the α -differentiable functions $u = u(\tau)$ and $w = w(\tau)$ for all positive τ satisfies*

- $D_{\tau}^{\alpha}(c_1 u + c_2 w) = c_1 D_{\tau}^{\alpha}(u) + c_2 D_{\tau}^{\alpha}(w)$
- $D_{\tau}^{\alpha}(\tau^p) = p\tau^{p-\alpha}, \forall p \in \mathbb{R}$
- $D_{\tau}^{\alpha}(u(\tau)) = 0$, when $u(\tau) = c_3$ is a constant function
- $D_{\tau}^{\alpha}(uw) = uD_{\tau}^{\alpha}(w) + wD_{\tau}^{\alpha}(u)$
- $D_{\tau}^{\alpha}\left(\frac{u}{w}\right) = \frac{uD_{\tau}^{\alpha}(w) - wD_{\tau}^{\alpha}(u)}{w^2}$
- $D_{\tau}^{\alpha}(u)(\tau) = \tau^{1-\alpha} \frac{du}{d\tau}$

for all real c_1, c_2, c_3 [24, 25].

Moreover, the definitions of many further properties of the conformal derivative (2.1) are discussed in details in [26]. The Gronwall's inequality, integration by parts, Laplace transform, the conformal derivative of the composite function and more are described in that study.

3. THE MODIFIED KUDRYASHOV METHOD

Let P be

$$P(u, u_{\tau}^{\alpha}, u_x, u_y, u_z, u_{\tau\tau}^{\alpha}, u_{xx}, \dots) = 0 \quad (3.1)$$

where $u = u(x, y, z, \dots, \tau)$ and $\alpha \in (0, 1]$ be the fractional derivative order. The transformation

$$u(x, y, z, \dots, \tau) = u(\xi), \xi = x + y + z + \dots - \frac{c}{\alpha} \tau^\alpha \quad (3.2)$$

converts (3.1) to an ODE for new variable ξ

$$R(u, u', u'', \dots) = 0 \quad (3.3)$$

where the prime ($'$) indicates the derivative operator $\frac{d}{d\xi}$ of u with respect to ξ [27]. Let

$$u(\xi) = a_0 + a_1 U(\xi) + a_2 U^2(\xi) + \dots a_n U^n(\xi) \quad (3.4)$$

be predicted solution of the equation (3.3) for a finite n with all $a_i, 0 \leq i \leq n$ are constants satisfying $a_n \neq 0$. In fact, the n is determined by balancing nonlinear term and the maximal derivative order. Furthermore, this finite series of $U(\xi)$ satisfies the first-order ODE

$$U'(\xi) = U(\xi)(U(\xi) - 1) \ln A \quad (3.5)$$

Accordingly, $U(\xi)$ is of the form

$$U(\xi) = \frac{1}{1 + dA\xi}$$

where d and A are non-zero constants with the conditions $A > 0$ and $A \neq 1$. The balance between the non-linear term and the term having the maximal order derivative in (3.3) enables to determine the positive integer n , if exists. Ultimately, since a solution $u(\xi)$ is sought for (3.3), it must satisfy (3.3). Thus, it is substituted into (3.3) and the resultant equation is rearranged for the powers $U(\xi)$. All the coefficients of the powers of $U(\xi)$ including the remaining part including constants and other parameters are equated to zero. Finally, $a_0, a_1, a_2, \dots, a_n$ are obtained explicitly in terms of constants and coefficients used in the equation (3.1) or originated from the (3.2).

4. THE CONFORMAL TIME FRACTIONAL (3+1) -DIMENSIONAL KP EQUATION

Consider the (3+1)- dimensional KP equation of the form

$$(D_\tau^\alpha(u) + \theta uu_x + \beta u_{xxx})_x - \varepsilon u_{yy} - \delta u_{zz} = 0 \quad (4.1)$$

where $u = u(x, y, z, \tau)$, θ , β , ε and δ are parameters. The transformation (3.2) reduces the KP equation (4.1) to

$$\left(-cu' + \theta uu' + \beta u''' \right)' - (\varepsilon + \delta) u'' = 0 \quad (4.2)$$

where $\left('\right) = \frac{d}{d\xi}$ derivative operator. Integrating the last equation twice gives

$$-cu + \frac{\theta}{2} u^2 + \beta u'' - (\varepsilon + \delta) u = K_1 \xi + K_2 \quad (4.3)$$

where K_1 and K_2 are constants of integration. The balance between u^2 and u'' occurs when $n = 2$. Assume that K_1 is zero. Substitution of the predicted solution

$u(\xi) = a_0 + a_1 U(\xi) + a_2 U^2(\xi)$ into (4.3) and assuming $K_1 = 0$ gives

$$\begin{aligned} & \left(\frac{1}{2} \theta a_2^2 + 6 \beta a_2 (\ln(A))^2 \right) U^4(\xi) \\ & + \left(2 \beta a_1 (\ln(A))^2 - 10 \beta a_2 (\ln(A))^2 + \theta a_1 a_2 \right) U^3(\xi) \\ & + \left(\frac{1}{2} \theta a_1^2 - \delta a_2 - c a_2 - \varepsilon a_2 + \theta a_0 a_2 - 3 \beta a_1 (\ln(A))^2 + 4 \beta a_2 (\ln(A))^2 \right) U^2(\xi) \\ & + \left(-c a_1 - \delta a_1 - \varepsilon a_1 + \beta a_1 (\ln(A))^2 + \theta a_0 a_1 \right) U(\xi) \\ & - K_2 - \varepsilon a_0 - \delta a_0 - c a_0 + \frac{1}{2} \theta a_0^2 = 0 \end{aligned} \quad (4.4)$$

Thus, equating the coefficients of the powers of each $U(\xi)$ and the remaining constants to zero gives

$$\begin{aligned} U^4 : & \frac{1}{2} \theta a_2^2 + 6 \beta a_2 (\ln(A))^2 = 0 \\ U^3 : & 2 \beta a_1 (\ln(A))^2 - 10 \beta a_2 (\ln(A))^2 + \theta a_1 a_2 = 0 \\ U^2 : & \frac{1}{2} \theta a_1^2 - \delta a_2 - c a_2 - \varepsilon a_2 + \theta a_0 a_2 - 3 \beta a_1 (\ln(A))^2 + 4 \beta a_2 (\ln(A))^2 = 0 \\ U^1 : & -c a_1 - \delta a_1 - \varepsilon a_1 + \beta a_1 (\ln(A))^2 + \theta a_0 a_1 = 0 \\ U^0 : & -K_2 - \varepsilon a_0 - \delta a_0 - c a_0 + \frac{1}{2} \theta a_0^2 = 0 \end{aligned} \quad (4.5)$$

The first solution set of the last system constituting five algebraic equations for $\{a_0, a_1, a_2, c\}$ gives

$$\begin{aligned} a_0 &= \frac{-\beta (\ln(A))^2 + \sqrt{\beta^2 (\ln(A))^4 - 2 \theta K_2}}{\theta} \\ a_1 &= 12 \frac{\beta (\ln(A))^2}{\theta} \\ a_2 &= -12 \frac{\beta (\ln(A))^2}{\theta} \\ c &= -\delta - \varepsilon + \sqrt{\beta^2 (\ln(A))^4 - 2 \theta K_2} \end{aligned} \quad (4.6)$$

where $\beta^2 (\ln(A))^4 - 2 \theta K_2 \geq 0$ and $\theta \neq 0$. The solution of the (4.3) takes the form

$$\begin{aligned} U(\xi) &= \frac{-\beta (\ln(A))^2 + \sqrt{\beta^2 (\ln(A))^4 - 2 \theta K_2}}{\theta} + \frac{\beta (\ln(A))^2}{\theta} \frac{1}{1 + dA^\xi} \\ & - 12 \frac{\beta (\ln(A))^2}{\theta} \frac{1}{(1 + dA^\xi)^2} \end{aligned} \quad (4.7)$$

Returning the original variables $\{x, y, z, \tau\}$ gives the solution for the conformal time fractional KP equation (4.1) as

$$\begin{aligned}
u_1(x, y, z, \tau) = & \frac{-\beta (\ln(A))^2 + \sqrt{\beta^2 (\ln(A))^4 - 2\theta K_2}}{\theta} \\
& + \frac{\beta (\ln(A))^2}{\theta} \frac{1}{(-\delta - \varepsilon + \sqrt{\beta^2 (\ln(A))^4 - 2\theta K_2})\tau^\alpha} \\
& - 12 \frac{\beta (\ln(A))^2}{\theta} \frac{1 + dA^{x+y+z-\frac{(-\delta - \varepsilon + \sqrt{\beta^2 (\ln(A))^4 - 2\theta K_2})\tau^\alpha}{\alpha}}}{\left(1 + dA^{x+y+z-\frac{(-\delta - \varepsilon + \sqrt{\beta^2 (\ln(A))^4 - 2\theta K_2})\tau^\alpha}{\alpha}}\right)^2}
\end{aligned} \tag{4.8}$$

The second solution of the system (4.5) for $\{a_0, a_1, a_2, c\}$ gives

$$\begin{aligned}
a_0 &= -\frac{\beta (\ln(A))^2 + \sqrt{\beta^2 (\ln(A))^4 - 2\theta K_2}}{\theta} \\
a_1 &= 12 \frac{\beta (\ln(A))^2}{\theta} \\
a_2 &= -12 \frac{\beta (\ln(A))^2}{\theta} \\
c &= -\delta - \varepsilon - \sqrt{\beta^2 (\ln(A))^4 - 2\theta K_2}
\end{aligned} \tag{4.9}$$

where $\beta^2 (\ln(A))^4 - 2\theta K_2 \geq 0$ and $\theta \neq 0$. Thus, the solution of the (4.3) is determined as

$$\begin{aligned}
U(\xi) = & -\frac{\beta (\ln(A))^2 + \sqrt{\beta^2 (\ln(A))^4 - 2\theta K_2}}{\theta} + 12 \frac{\beta (\ln(A))^2}{\theta} \frac{1}{1 + dA^\xi} \\
& - 12 \frac{\beta (\ln(A))^2}{\theta} \frac{1}{(1 + dA^\xi)^2}
\end{aligned} \tag{4.10}$$

Ultimately, the solution of (4.1) is written in original variables as

$$\begin{aligned}
u_2(x, y, z, \tau) = & -\frac{\beta (\ln(A))^2 + \sqrt{\beta^2 (\ln(A))^4 - 2\theta K_2}}{\theta} \\
& + 12 \frac{\beta (\ln(A))^2}{\theta} \frac{1}{1 + dA^{x+y+z} - \frac{(-\delta - \varepsilon - \sqrt{\beta^2 (\ln(A))^4 - 2\theta K_2} \tau^\alpha)}{\alpha}} \\
& - 12 \frac{\beta (\ln(A))^2}{\theta} \frac{1}{\left(1 + dA^{x+y+z} - \frac{(-\delta - \varepsilon - \sqrt{\beta^2 (\ln(A))^4 - 2\theta K_2} \tau^\alpha)}{\alpha}\right)^2}
\end{aligned} \tag{4.11}$$

5. THE CONFORMAL TIME FRACTIONAL mKAWAHARA EQUATION

The conformal time fractional mKawahara equation

$$D_\tau^\alpha(u) + pu^2u_x + qu_{xxx} + ru_{xxxx} = 0 \tag{5.1}$$

where $u = u(x, \tau)$ and p, q and r are constant parameters can be reduced to a nonlinear ODE of the form

$$-cu' + pu^2u' + qu''' + ru'''' = 0 \tag{5.2}$$

by using the simplest form of the transformation (3.2) that is $u(\xi) = u(x, \tau)$, $\xi = x - c\frac{\tau^\alpha}{\alpha}$. Integrating the last equation once converts it to

$$-cu + \frac{p}{3}u^3 + qu'' + ru'''' = K \tag{5.3}$$

where K denotes the constant of integration. The balance between u^3 and u'''' determines the degree n of the solution as 2. Thus, the predicted solution is constructed in the form $u(\xi) = a_0 + a_1U(\xi) + a_2U^2(\xi)$ with the condition $a_2 \neq 0$. Substituting this solution into the equation (5.3) results

$$\begin{aligned}
& \left(1/3pa_2^3 + 120ra_2(\ln(A))^4\right)U^6(\xi) \\
& + \left(pa_1a_2^2 - 336ra_2(\ln(A))^4 + 24ra_1(\ln(A))^4\right)U^5(\xi) \\
& + \left(pa_1^2a_2 + 6qa_2(\ln(A))^2 + 330ra_2(\ln(A))^4 + pa_0a_2^2 - 60ra_1(\ln(A))^4\right)U^4(\xi) \\
& + \left(-130ra_2(\ln(A))^4 + 2pa_0a_1a_2 + 2qa_1(\ln(A))^2 + 1/3pa_1^3 - 10qa_2(\ln(A))^2 + 50ra_1(\ln(A))^4\right)U^3(\xi) \\
& + \left(-15ra_1(\ln(A))^4 - ca_2 + pa_0a_1^2 + 4qa_2(\ln(A))^2 + 16ra_2(\ln(A))^4 + pa_0^2a_2 - 3qa_1(\ln(A))^2\right)U^2(\xi) \\
& + \left(ra_1(\ln(A))^4 - ca_1 + qa_1(\ln(A))^2 + pa_0^2a_1\right)U(\xi) \\
& - ca_0 + 1/3pa_0^3 - K = 0
\end{aligned} \tag{5.4}$$

Equating the coefficients of each power $U(\xi)$ gives

$$\begin{aligned}
 U^6 : & 1/3 p a_2^3 + 120 r a_2 (\ln(A))^4 = 0 \\
 U^5 : & p a_1 a_2^2 - 336 r a_2 (\ln(A))^4 + 24 r a_1 (\ln(A))^4 = 0 \\
 U^4 : & p a_1^2 a_2 + 6 q a_2 (\ln(A))^2 + 330 r a_2 (\ln(A))^4 + p a_0 a_2^2 - 60 r a_1 (\ln(A))^4 = 0 \\
 U^3 : & -130 r a_2 (\ln(A))^4 + 2 p a_0 a_1 a_2 + 2 q a_1 (\ln(A))^2 + 1/3 p a_1^3 - 10 q a_2 (\ln(A))^2 + 50 r a_1 (\ln(A))^4 = 0 \\
 U^2 : & -15 r a_1 (\ln(A))^4 - c a_2 + p a_0 a_1^2 + 4 q a_2 (\ln(A))^2 + 16 r a_2 (\ln(A))^4 + p a_0^2 a_2 - 3 q a_1 (\ln(A))^2 = 0 \\
 U^1 : & r a_1 (\ln(A))^4 - c a_1 + q a_1 (\ln(A))^2 + p a_0^2 a_1 = 0 \\
 U^0 : & -c a_0 + 1/3 p a_0^3 - K = 0
 \end{aligned} \tag{5.5}$$

The solution of this algebraic system for $\{a_0, a_1, a_2, c, K\}$ gives

$$\begin{aligned}
 a_0 &= \frac{1}{10r} \sqrt{-10 \frac{r}{p}} \left(q + 5 r (\ln(A))^2 \right) \\
 a_1 &= -6 \sqrt{-10 \frac{r}{p}} (\ln(A))^2 \\
 a_2 &= 6 \sqrt{-10 \frac{r}{p}} (\ln(A))^2 \\
 c &= -\frac{1}{10r} \left(15 r^2 (\ln(A))^4 + q^2 \right) \\
 K &= \frac{1}{150r^2} \left(-15 q r^2 (\ln(A))^4 + 50 r^3 (\ln(A))^6 + q^3 \right) \sqrt{-10 \frac{r}{p}}
 \end{aligned} \tag{5.6}$$

with the condition $rp < 0$. Thus, the solution of (5.2) can be written as

$$\begin{aligned}
 U(\xi) &= \frac{1}{10r} \sqrt{-10 \frac{r}{p}} \left(q + 5 r (\ln(A))^2 \right) - 6 \sqrt{-10 \frac{r}{p}} (\ln(A))^2 \frac{1}{1 + dA\xi} \\
 &+ 6 \sqrt{-10 \frac{r}{p}} (\ln(A))^2 \frac{1}{(1 + dA\xi)^2}
 \end{aligned} \tag{5.7}$$

Returning the original variables x and τ gives the solution of (5.1) as

$$\begin{aligned}
 u_3(x, \tau) &= \frac{1}{10r} \sqrt{-10 \frac{r}{p}} \left(q + 5 r (\ln(A))^2 \right) - 6 \sqrt{-10 \frac{r}{p}} (\ln(A))^2 \frac{1}{1 + dA^{x + \frac{1}{10r}} (15 r^2 (\ln(A))^4 + q^2) \frac{t^\alpha}{\alpha}} \\
 &+ 6 \sqrt{-10 \frac{r}{p}} (\ln(A))^2 \frac{1}{\left(1 + dA^{x + \frac{1}{10r}} (15 r^2 (\ln(A))^4 + q^2) \frac{t^\alpha}{\alpha} \right)^2}
 \end{aligned} \tag{5.8}$$

for $pr < 0$.

A second solution to (5.5) for $\{a_0, a_1, a_2, c, K\}$ is of the form

$$\begin{aligned} a_0 &= -\frac{1}{10r} \sqrt{-10 \frac{r}{p}} \left(q + 5r (\ln(A))^2 \right) \\ a_1 &= 6 \sqrt{-10 \frac{r}{p}} (\ln(A))^2 \\ a_2 &= -6 \sqrt{-10 \frac{r}{p}} (\ln(A))^2 \\ c &= -\frac{1}{10r} \left(15r^2 (\ln(A))^4 + q^2 \right) \\ K &= -\frac{1}{150r^2} \left(-15qr^2 (\ln(A))^4 + 50r^3 (\ln(A))^6 + q^3 \right) \sqrt{-10 \frac{r}{p}} \end{aligned} \quad (5.9)$$

and generates the solution to (5.2) as

$$\begin{aligned} U(\xi) &= -\frac{1}{10r} \sqrt{-10 \frac{r}{p}} \left(q + 5r (\ln(A))^2 \right) + 6 \sqrt{-10 \frac{r}{p}} (\ln(A))^2 \frac{1}{1 + dA^\xi} \\ &\quad - 6 \sqrt{-10 \frac{r}{p}} (\ln(A))^2 \frac{1}{(1 + dA^\xi)^2} \end{aligned} \quad (5.10)$$

Similarly, the solution to the conformal time fractional mKawahara equation (5.1) is obtained as

$$\begin{aligned} u_4(x, \tau) &= -\frac{1}{10r} \sqrt{-10 \frac{r}{p}} \left(q + 5r (\ln(A))^2 \right) + 6 \sqrt{-10 \frac{r}{p}} (\ln(A))^2 \frac{1}{1 + dA^{x + \left(\frac{1}{10r} 15r^2 (\ln(A))^4 + q^2 \right) \frac{\tau^\alpha}{\alpha}}} \\ &\quad - 6 \sqrt{-10 \frac{r}{p}} (\ln(A))^2 \frac{1}{\left(1 + dA^{x + \left(\frac{1}{10r} 15r^2 (\ln(A))^4 + q^2 \right) \frac{\tau^\alpha}{\alpha}} \right)^2} \end{aligned} \quad (5.11)$$

where pr is negative.

6. CONCLUSION

The modified form of the Kudryashov method is implemented for the exact solutions of the conformal time fractional $(3+1)$ - dimensional KP and the modified Kawahara equations. The corresponding wave transformations reduce the number of independent variables to one. Thus, both equations are converted some nonlinear ODEs. Balancing the highest ordered derivative term and the nonlinear term leads to determine the degree of the solution in the form of finite series. Substituting the predicted solution into the resultant ODE and equating all coefficients of the powers of the predicted solution gives an algebraic system of equations. Thus, the coefficients of the predicted solution are determined in explicit form by computer aided algebra.

In the study, a couple of explicit solutions to each conformal fractional $(3+1)$ KP and the modified Kawahara equations are determined in the rational form containing exponential function by the modified Kudryashov method.

REFERENCES

- [1] Kadomtsev, B. B., & Petviashvili, V. I., (1970), On the stability of solitary waves in weakly dispersing media. In Soviet Physics Doklady, 15, 539-541.
- [2] Dryuma, V. S. (1974). Analytic solution of the two-dimensional Korteweg-de Vries (KdV) equation. Soviet Journal of Experimental and Theoretical Physics Letters, 19, 387.
- [3] Boiti, M., Pempinelli, F., & Pogrebkov, A., (1994), Solutions of the KPI equation with smooth initial data. Inverse Problems, 10(3), 505.
- [4] Bogayevskii, V. N. (1990). The Korteweg-de Vries, Kadomtsev-Petviashvili and Boussinesq equations. Modulation theory. USSR Computational Mathematics and Mathematical Physics, 30(5), 148-159.
- [5] Alagesan, T., Uthayakumar, A., & Porsezian, K. (1997). Painlevé analysis and Bäcklund transformation for a three-dimensional Kadomtsev-Petviashvili equation. Chaos, Solitons & Fractals, 8(6), 893-895.
- [6] Ablowitz, M. J., & Segur, H. (1979). On the evolution of packets of water waves. Journal of Fluid Mechanics, 92(04), 691-715.
- [7] Senatorski, A., & Infeld, E. (1996). Simulations of two-dimensional Kadomtsev-Petviashvili soliton dynamics in three-dimensional space. Physical review letters, 77(14), 2855.
- [8] De Bouard, A., & Saut, J. C. (1997). Solitary waves of generalized Kadomtsev-Petviashvili equations. In Annales de l'IHP Analyse non linéaire, 14, 2, 211-236.
- [9] De Bouard, A., & Saut, J. C. (1997). Symmetries and Decay of the Generalized Kadomtsev-Petviashvili Solitary Waves. SIAM Journal on Mathematical Analysis, 28(5), 1064-1085.
- [10] Jones, K. L. (2000). Three-dimensional Korteweg-de Vries equation and traveling wave solutions. International Journal of Mathematics and Mathematical Sciences, 24(6), 379-384.
- [11] Song, M., & Ge, Y. (2010). Application of the $(\frac{G'}{G})$ -expansion method to $(3+1)$ -dimensional nonlinear evolution equations. Computers & Mathematics with Applications, 60(5), 1220-1227.
- [12] Kawahara, T. (1972). Oscillatory solitary waves in dispersive media. Journal of the physical society of Japan, 33(1), 260-264.
- [13] Zhang, D. (2005). Doubly periodic solutions of the modified Kawahara equation. Chaos, Solitons & Fractals, 25(5), 1155-1160.
- [14] Sirendaoreji. (2004). New exact travelling wave solutions for the Kawahara and modified Kawahara equations. Chaos Solitons & Fractals, 19(1), 147-150.
- [15] Wazwaz, A. M. (2007). New solitary wave solutions to the modified Kawahara equation. Physics Letters A, 360(4), 588-592.
- [16] Demina, M. V., & Kudryashov, N. A. (2010). From Laurent series to exact meromorphic solutions: the Kawahara equation. Physics Letters A, 374(39), 4023-4029.
- [17] Al-Ali, E. M. (2013). Traveling Wave Solutions for a Generalized Kawahara and Hunter-Saxton Equations. International Journal of Mathematical Analysis, 7, 1647-1666.
- [18] Yusufoglu, E., & Bekir, A. (2008). Symbolic computation and new families of exact travelling solutions for the Kawahara and modified Kawahara equations. Computers & Mathematics with Applications, 55(6), 1113-1121.
- [19] Ray, S. S. (2006). Exact solutions for time-fractional diffusion-wave equations by decomposition method. Physica Scripta, 75(1), 53.
- [20] Alam, M. N., Akbar, M. A., & Mohyud-Din, S. T. (2013). A novel (G'/G) -expansion method and its application to the Boussinesq equation. Chinese Physics B, 23(2), 020203.
- [21] Younis, M. (2013). The first integral method for time-space fractional differential equations. Journal of Advanced Physics, 2(3), 220-223.
- [22] Bhrawy, A. H., Biswas, A., Javidi, M., Ma, W. X., Pinar, Z., & Yildirim, A. (2013). New solutions for $(1+1)$ -dimensional and $(2+1)$ -dimensional Kaup-Kupershmidt equations. Results in Mathematics, 63(1-2), 675-686.
- [23] Khalil, R., Al Horani, M., Yousef, A., & Sababheh, M., A new definition of fractional derivative, Journal of Computational and Applied Mathematics (2014), 264, 65-70.
- [24] Atangana, A., Baleanu, D., & Alsaedi, A., New properties of conformable derivative, Open Mathematics (2015), 13(1), 1-10.
- [25] Çenesiz, Y., Baleanu, D., Kurt, A., & Tasbozan, O., New exact solutions of Burgers' type equations with conformable derivative, Waves in Random and Complex Media (2016), 1-14.
- [26] Abdeljawad, T., On conformable fractional calculus, Journal of computational and Applied Mathematics (2015), 279, 57-66.

- [27] Eslami, M., & Rezazadeh, H., The first integral method for Wu-Zhang system with conformable time-fractional derivative, *Calcolo* (2015), 1-11.
- [28] Kudryashov, N. A., One method for finding exact solutions of nonlinear differential equations, *Communications in Nonlinear Science and Numerical Simulation* (2012), 17(6), 2248-2253.
- [29] Kabir, M. M., Khajeh, A., Abdi Aghdam, E., & Yousefi Koma, A., Modified Kudryashov method for finding exact solitary wave solutions of higher-order nonlinear equations, *Mathematical methods in the Applied Sciences* (2011), 34(2), 213-219.
- [30] Tandogan, Y. A., Pandir, Y., & Gurefe, Y., Solutions of the nonlinear differential equations by use of modified Kudryashov method, *Turkish Journal of Mathematics and Computer Science* (2013), 1, 1-7.

ALPER KORKMAZ: ÇANKIRI KARATEKİN UNIVERSITY, DEPARTMENT OF MATHEMATICS, ÇANKIRI, TURKEY.

E-mail address: akorkmaz@karatekin.edu.tr, alperkorkmaz7@gmail.com

© 2016 by the author; licensee *Preprints*, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons by Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).