

Article

Neutrosophic Modal Logic

Florentin Smarandache

Mathematics & Science Department, University of New Mexico, 705 Gurley Ave., Gallup, NM 87301, USA;
fsmarandache@unm.edu

Abstract: I introduce now for the first time the neutrosophic modal logic. The Neutrosophic Modal Logic includes the neutrosophic operators that express the modalities. It is an extension of neutrosophic predicate logic, and of neutrosophic propositional logic. In order for the paper to be self-contained, I also recall the etymology and definition of neutrosophy and of neutrosophic logic. Several examples are presented as well.

Keywords: neutrosophy; neutrosophic logic; neutrosophic alethic modalities; neutrosophic possibility; neutrosophic necessity; neutrosophic impossibility; neutrosophic temporal modalities; neutrosophic epistemic modalities; neutrosophic doxastic modalities; neutrosophic deontic modalities

1. Introduction

The word “*neutrosophy*” [9, 10, 11] etymologically comes from French *neutre* < Latin *neuter*, neutral, and Greek *sophia*, skill/wisdom and means knowledge of neutral thought.

Neutrosophy is a new branch of philosophy, which studies the origin, nature, and scope of neutralities, as well as their interactions with different ideational spectra.

Neutrosophy is a generalization of dialectics, because while in dialectics one deals with the dynamics of opposites [let’s note them by $\langle A \rangle$ and $\langle \text{anti}A \rangle$, where $\langle A \rangle$ is an item (i.e. idea, notion, proposition, theory etc.) and $\langle \text{anti} \rangle$ is the opposite of $\langle A \rangle$], in neutrosophy one deals with the dynamics of opposites and their neutral [i.e. $\langle A \rangle$, $\langle \text{anti}A \rangle$, and their neutralities $\langle \text{neut}A \rangle$].

For example, if two countries C_1 and C_2 go to war against each other (this is the dynamics of opposites $\langle A \rangle$ and $\langle \text{anti}A \rangle$), some neutral countries ($\langle \text{neut}A \rangle$) interfere into the war in one side or another. So, the neutrals, which were ignored in dialectics, have to be taking into consideration as in neutrosophy.

Neutrosophy is the base of neutrosophic set, neutrosophic logic, neutrosophic probability, neutrosophic statistics [9] and so on.

Neutrosophic logic is a generalization of the intuitionistic fuzzy logic. In neutrosophic logic, a proposition has a degree of truth (t), a degree of indeterminacy/neutrality (neither true, not false) (i), and a degree of falsehood (f), where t, i, f are subsets of the unit interval $[0, 1]$.

The Modal Logic, from classical Boolean logic, I now extend to Neutrosophic Modal Logic.

I introduce therefore the Neutrosophic Modal Logic and the Refined Neutrosophic Modal Logic. Then I can extend them to Symbolic Neutrosophic Modal Logic and Refined Symbolic Neutrosophic Modal Logic. I use labels instead of numerical values.

Neutrosophic Modal Logic is a logic where some neutrosophic modalities have been included.

There is a large variety of neutrosophic modal logics, as actually happens in classical modal logic too. Similarly, the neutrosophic accessibility relation and possible neutrosophic worlds have many interpretations, depending on each application. Several neutrosophic modal applications are also listed.

2. Neutrosophic Modalities

Let \mathcal{P} be a neutrosophic proposition.

One has the following types of **neutrosophic modalities**:

A) Neutrosophic Alethic Modalities (related to *truth*) has three neutrosophic operators:

- i. **Neutrosophic Possibility**: It is neutrosophically possible that \mathcal{P} .
- ii. **Neutrosophic Necessity**: It is neutrosophically necessary that \mathcal{P} .
- iii. **Neutrosophic Impossibility**: It is neutrosophically impossible that \mathcal{P} .

B) Neutrosophic Temporal Modalities (related to *time*)

It was the neutrosophic case that \mathcal{P} .

It will neutrosophically be that \mathcal{P} .

And similarly:

It has always neutrosophically been that \mathcal{P} .

It will always neutrosophically be that \mathcal{P} .

C) Neutrosophic Epistemic Modalities (related to *knowledge*):

It is neutrosophically known that \mathcal{P} .

D) Neutrosophic Doxastic Modalities (related to *belief*):

It is neutrosophically believed that \mathcal{P} .

E) Neutrosophic Deontic Modalities:

It is neutrosophically obligatory that \mathcal{P} .

It is neutrosophically permissible that \mathcal{P} .

2.1. Neutrosophic Alethic Modal Operators

The modalities used in classical (alethic) modal logic can be neutrosophicated by inserting the indeterminacy. I insert the **degrees of possibility** and **degrees of necessity**, as refinement of classical modal operators.

2.2. Neutrosophic Possibility Operator

The classical Possibility Modal Operator « $\diamond P$ » meaning «It is possible that P » is extended to **Neutrosophic Possibility Operator**: $\diamond_N \mathcal{P}$ meaning «It is (t, i, f) -possible that \mathcal{P} », using Neutrosophic Probability, where « (t, i, f) -possible» means t % possible (chance that \mathcal{P} occurs), i % indeterminate (indeterminate-chance that \mathcal{P} occurs), and f % impossible (chance that \mathcal{P} does not occur).

If $\mathcal{P}(t_p, i_p, f_p)$ is a neutrosophic proposition, with t_p, i_p, f_p subsets of $[0, 1]$, then the neutrosophic truth-value of the neutrosophic possibility operator is:

$$\diamond_N \mathcal{P} = (\sup(t_p), \inf(i_p), \inf(f_p)), \quad (1)$$

which means that if a proposition P is t_p true, i_p indeterminate, and f_p false, then the value of the neutrosophic possibility operator $\diamond_N \mathcal{P}$ is: $\sup(t_p)$ possibility, $\inf(i_p)$ indeterminate-possibility, and $\inf(f_p)$ impossibility.

For *example*, let $P =$ «It will be snowing tomorrow».

According to the meteorological center, the neutrosophic truth-value of \mathcal{P} is:

$$\mathcal{P}([0.5, 0.6], (0.2, 0.4), \{0.3, 0.5\}),$$

i.e. $[0.5, 0.6]$ true, $(0.2, 0.4)$ indeterminate, and $\{0.3, 0.5\}$ false.

Then the neutrosophic possibility operator is:

$$\diamond_N \mathcal{P} = (\sup[0.5, 0.6], \inf(0.2, 0.4), \inf\{0.3, 0.5\}) = (0.6, 0.2, 0.3),$$

i.e. 0.6 possible, 0.2 indeterminate-possibility, and 0.3 impossible.

2.3. Neutrosophic Necessity Operator

The classical Necessity Modal Operator « $\square P$ » meaning «It is necessary that P » is extended to **Neutrosophic Necessity Operator**: $\square_N \mathcal{P}$ meaning «It is (t, i, f) -necessary that \mathcal{P} », using again the Neutrosophic Probability, where similarly « (t, i, f) -necessity» means t % necessary (surety that \mathcal{P}

occurs), i % indeterminate (indeterminate-surety that \mathcal{P} occurs), and f % unnecessary (unsafety that \mathcal{P} occurs).

If $\mathcal{P}(t_p, i_p, f_p)$ is a neutrosophic proposition, with t_p, i_p, f_p subsets of $[0, 1]$, then the neutrosophic truth value of the neutrosophic necessity operator is:

$$\square_N \mathcal{P} = (\inf(t_p), \sup(i_p), \sup(f_p)), \quad (2)$$

which means that if a proposition \mathcal{P} is t_p true, i_p indeterminate, and f_p false, then the value of the neutrosophic necessity operator $\square_N \mathcal{P}$ is: $\inf(t_p)$ necessary, $\sup(i_p)$ indeterminate-necessity, and $\sup(f_p)$ unnecessary.

Taking the previous *example*, $\mathcal{P} = \langle \text{It will be snowing tomorrow} \rangle$, with

$$\mathcal{P}([0.5, 0.6], (0.2, 0.4), \{0.3, 0.5\}),$$

then the neutrosophic necessity operator is:

$$\square_N \mathcal{P} = (\inf[0.5, 0.6], \sup(0.2, 0.4), \sup\{0.3, 0.5\}) = (0.5, 0.4, 0.5),$$

i.e. 0.5 necessary, 0.4 indeterminate-necessity, and 0.5 unnecessary.

2.4. Connection between Neutrosophic Possibility Operator and Neutrosophic Necessity Operator

In classical modal logic, a modal operator is equivalent to the negation of the other:

$$\diamond P \leftrightarrow \neg \square \neg P,$$

$$\square P \leftrightarrow \neg \diamond \neg P.$$

In neutrosophic logic one has a class of neutrosophic negation operators. The most used one is:

$$\overline{N}P(t, i, f) = \overline{P}(f, 1 - i, t), \quad (3)$$

where t, i, f are real subsets of the interval $[0, 1]$, and \overline{P} is the negation of P .

Let's check what's happening in the neutrosophic modal logic, using the previous *example*.

One had:

$$\mathcal{P}([0.5, 0.6], (0.2, 0.4), \{0.3, 0.5\}),$$

then

$$\overline{N} \mathcal{P} = \overline{P}(\{0.3, 0.5\}, 1 - (0.2, 0.4), [0.5, 0.6]) = \overline{P}(\{0.3, 0.5\}, (0.6, 0.8), [0.5, 0.6]) = \overline{P}(\{0.3, 0.5\}, (0.6, 0.8), [0.5, 0.6]).$$

Therefore, denoting by \overleftrightarrow{N} the neutrosophic equivalence, one has:

$$\overline{N} \square \overline{N} \mathcal{P}([0.5, 0.6], (0.2, 0.4), \{0.3, 0.5\}) \overleftrightarrow{N}.$$

\overleftrightarrow{N} It is not neutrosophically necessary that $\langle \text{It will not be snowing tomorrow} \rangle$

\overleftrightarrow{N} It is not neutrosophically necessary that $\overline{P}(\{0.3, 0.5\}, (0.6, 0.8), [0.5, 0.6])$

\overleftrightarrow{N} It is neutrosophically possible that $\overline{N} \overline{P}(\{0.3, 0.5\}, (0.6, 0.8), [0.5, 0.6])$

\overleftrightarrow{N} It is neutrosophically possible that $\mathcal{P}([0.5, 0.6], 1 - (0.6, 0.8), \{0.3, 0.5\})$

\overleftrightarrow{N} It is neutrosophically possible that $\mathcal{P}([0.5, 0.6], (0.2, 0.4), \{0.3, 0.5\})$

$\overleftrightarrow{N} \diamond \mathcal{P}([0.5, 0.6], (0.2, 0.4), \{0.3, 0.5\}) = (0.6, 0.2, 0.3).$

Let's check the second neutrosophic equivalence.

$$\overline{N} \diamond \overline{N} \mathcal{P}([0.5, 0.6], (0.2, 0.4), \{0.3, 0.5\}) \overleftrightarrow{N}.$$

\overleftrightarrow{N} It is not neutrosophically possible that $\langle \text{It will not be snowing tomorrow} \rangle$

\overleftrightarrow{N} It is not neutrosophically possible that $\overline{P}(\{0.3, 0.5\}, (0.6, 0.8), [0.5, 0.6])$

\overleftrightarrow{N} It is neutrosophically necessary that $\overline{N} \overline{P}(\{0.3, 0.5\}, (0.6, 0.8), [0.5, 0.6])$

\overleftrightarrow{N} It is neutrosophically necessary that $\mathcal{P}([0.5, 0.6], 1 - (0.6, 0.8), \{0.3, 0.5\})$

\overleftrightarrow{N} It is neutrosophically necessary that $\mathcal{P}([0.5, 0.6], (0.2, 0.4), \{0.3, 0.5\})$

$\overleftrightarrow{N} \square \mathcal{P}([0.5, 0.6], (0.2, 0.4), \{0.3, 0.5\}) = (0.6, 0.2, 0.3).$

2.5. Neutrosophic Modal Equivalences

Neutrosophic Modal Equivalences hold within a certain accuracy, depending on the definitions of neutrosophic possibility operator and neutrosophic necessity operator, as well as on the definition of the neutrosophic negation – employed by the experts depending on each application. Under these conditions, one may have the following neutrosophic modal equivalences:

$$\diamond_N \mathcal{P}(t_p, i_p, f_p) \stackrel{\leftrightarrow}{\sim} \neg \square_N \neg \mathcal{P}(t_p, i_p, f_p), \quad (4)$$

$$\square_N \mathcal{P}(t_p, i_p, f_p) \stackrel{\leftrightarrow}{\sim} \neg \diamond_N \neg \mathcal{P}(t_p, i_p, f_p), \quad (5)$$

For example, other definitions for the **neutrosophic modal operators** may be:

$$\diamond_N \mathcal{P}(t_p, i_p, f_p) = (\sup(t_p), \sup(i_p), \inf(f_p)), \quad (6)$$

or

$$\diamond_N \mathcal{P}(t_p, i_p, f_p) = (\sup(t_p), \frac{i_p}{2}, \inf(f_p)), \quad (7)$$

etc., while

$$\square_N \mathcal{P}(t_p, i_p, f_p) = (\inf(t_p), \inf(i_p), \sup(f_p)), \quad (8)$$

or

$$\square_N \mathcal{P}(t_p, i_p, f_p) = (\inf(t_p), 2i_p \cap [0,1], \sup(f_p)), \quad (9)$$

etc.

2.6. Neutrosophic Truth Threshold

In neutrosophic logic, first I have to introduce a **neutrosophic truth threshold**,

$$TH = \langle T_{th}, I_{th}, F_{th} \rangle, \quad (10)$$

where T_{th}, I_{th}, F_{th} are subsets of $[0, 1]$. I use upper-case letters (T, I, F) in order to distinguish the neutrosophic components of the threshold from those of a proposition in general.

I can say that the proposition $\mathcal{P}(t_p, i_p, f_p)$ is **neutrosophically true** if:

$$\begin{aligned} \inf(t_p) &\geq \inf(T_{th}) \text{ and } \sup(t_p) \geq \sup(T_{th}); \\ \inf(i_p) &\leq \inf(I_{th}) \text{ and } \sup(i_p) \leq \sup(I_{th}); \\ \inf(f_p) &\leq \inf(F_{th}) \text{ and } \sup(f_p) \leq \sup(F_{th}). \end{aligned}$$

For the particular case when all T_{th}, I_{th}, F_{th} and t_p, i_p, f_p are single-valued numbers from the interval $[0, 1]$, then one has:

The proposition $\mathcal{P}(t_p, i_p, f_p)$ is **neutrosophically true** if:

$$\begin{aligned} t_p &\geq T_{th}; \\ i_p &\leq I_{th}; \\ f_p &\leq F_{th}. \end{aligned}$$

The neutrosophic truth threshold is established by experts in accordance to each applications.

2.7. Neutrosophic Semantics

Neutrosophic Semantics of the Neutrosophic Modal Logic is formed by a **neutrosophic frame** G_N , which is a non-empty neutrosophic set, whose elements are called **possible neutrosophic worlds**, and a **neutrosophic binary relation** \mathcal{R}_N , called **neutrosophic accesibility relation**, between the possible neutrosophic worlds. By notation, one has:

$$\langle G_N, \mathcal{R}_N \rangle.$$

A neutrosophic world w'_N that is neutrosophically accessible from the neutrosophic world w_N is symbolized as:

$$w_N \mathcal{R}_N w'_N.$$

In a **neutrosophic model** each neutrosophic proposition \mathcal{P} has a **neutrosophic truth-value** $(t_{w_N}, i_{w_N}, f_{w_N})$ respectively to each neutrosophic world $w_N \in G_N$, where $t_{w_N}, i_{w_N}, f_{w_N}$ are subsets of $[0, 1]$.

A **neutrosophic actual world** can be similarly noted as in classical modal logic as $w_N * .$

2.8. Neutrosophic Formulas

Formalization: Let S_N be a set of neutrosophic propositional variables.

1. Every neutrosophic propositional variable $\mathcal{P} \in S_N$ is a neutrosophic formula.
2. If A, B are neutrosophic formulas, then $\neg_N A, A \wedge_N B, A \vee_N B, A \rightarrow_N B, A \leftrightarrow_N B$, and $\diamond_N A, \square_N A$, are also neutrosophic formulas, where $\neg_N, \wedge_N, \vee_N, \rightarrow_N, \leftrightarrow_N$ and \diamond_N, \square_N represent the neutrosophic negation, neutrosophic intersection, neutrosophic union, neutrosophic implication, neutrosophic equivalence, and neutrosophic possibility operator, neutrosophic necessity operator respectively.

2.9. Accesibility Relation in a Neutrosophic Theory

Let G_N be a set of neutrosophic worlds w_N such that each w_N characterizes the propositions (formulas) of a given neutrosophic theory τ .

One says that the neutrosophic world w'_N is accesible from the neutrosophic world w_N , and one writes: $w_N \mathcal{R}_N w'_N$ or $\mathcal{R}_N(w_N, w'_N)$, if for any proposition (formula) $\mathcal{P} \in w_N$, meaning the neutrosophic truth-value of \mathcal{P} with respect to w_N is

$$\mathcal{P}(t_p^{w_N}, i_p^{w_N}, f_p^{w_N}),$$

one has the neutrosophic truth-value of \mathcal{P} with respect to w'_N

$$\mathcal{P}(t_p^{w'_N}, i_p^{w'_N}, f_p^{w'_N}),$$

where

$$\begin{aligned} \inf(t_p^{w'_N}) &\geq \inf(t_p^{w_N}) \text{ and } \sup(t_p^{w'_N}) \geq \sup(t_p^{w_N}); \\ \inf(i_p^{w'_N}) &\leq \inf(i_p^{w_N}) \text{ and } \sup(i_p^{w'_N}) \leq \sup(i_p^{w_N}); \\ \inf(f_p^{w'_N}) &\leq \inf(f_p^{w_N}) \text{ and } \sup(f_p^{w'_N}) \leq \sup(f_p^{w_N}) \end{aligned}$$

(in the general case when $t_p^{w_N}, i_p^{w_N}, f_p^{w_N}$ and $t_p^{w'_N}, i_p^{w'_N}, f_p^{w'_N}$ are subsets of the interval $[0, 1]$).

But in the instant of $t_p^{w_N}, i_p^{w_N}, f_p^{w_N}$ and $t_p^{w'_N}, i_p^{w'_N}, f_p^{w'_N}$ as single-values in $[0, 1]$, the above inequalities become:

$$\begin{aligned} t_p^{w'_N} &\geq t_p^{w_N}, \\ i_p^{w'_N} &\leq i_p^{w_N}, \\ f_p^{w'_N} &\leq f_p^{w_N}. \end{aligned}$$

2.10. Applications

If the neutrosophic theory τ is the Neutrosophic Mereology, or Neutrosophic Gnosisology, or Neutrosophic Epistemology etc., the neutrosophic accesibility relation is defined as above.

2.11. Neutrosophic n-ary Accesibility Relation

I can also extend the classical **binary** accesibility relation \mathcal{R} to a **neutrosophic n-ary accesibility relation**

$$\mathcal{R}_N^{(n)}, \text{ for } n \text{ integer } \geq 2.$$

Instead of the classical $R(w, w')$, which means that the world w' is accesible from the world w , I generalize it to:

$$\mathcal{R}_N^{(n)}(w_{1N}, w_{2N}, \dots, w_{nN}; w'_N),$$

which means that the neutrosophic world w'_N is accesible from the neutrosophic worlds $w_{1N}, w_{2N}, \dots, w_{nN}$ all together.

2.12. Neutrosophic Kripke Frame

$k_N = \langle G_N, R_N \rangle$ is a neutrosophic Kripke frame, since:

i. G_N is an arbitrary non-empty neutrosophic set of **neutrosophic worlds**, or **neutrosophic states**, or **neutrosophic situations**.

ii. $R_N \subseteq G_N \times G_N$ is a **neutrosophic accessibility relation** of the neutrosophic Kripke frame.

Actually, one has a degree of accesibility, degree of indeterminacy, and a degree of non-accesibility.

2.13. Neutrosophic (t, i, f)-Assignment

The Neutrosophic (t, i, f)-Assignment is a neutrosophic mapping

$$v_N: S_N \times G_N \rightarrow [0,1] \times [0,1] \times [0,1], \quad (11)$$

where, for any neutrosophic proposition $\mathcal{P} \in S_N$ and for any neutrosophic world w_N , one defines:

$$v_N(\mathcal{P}, w_N) = (t_p^{w_N}, i_p^{w_N}, f_p^{w_N}) \in [0,1] \times [0,1] \times [0,1], \quad (12)$$

which is the neutrosophical logical truth value of the neutrosophic proposition \mathcal{P} in the neutrosophic world w_N .

2.14. Neutrosophic Deducibility

One says that the neutrosophic formula \mathcal{P} is neutrosophically deducible from the neutrosophic Kripke frame k_N , the neutrosophic (t, i, f) – assignment v_N , and the neutrosophic world w_N , and one writes as:

$$k_N, v_N, w_N \stackrel{F}{=} \mathcal{P}.$$

Let's make the notation:

$$\alpha_N(\mathcal{P}; k_N, v_N, w_N)$$

that denotes the neutrosophic logical value that the formula \mathcal{P} takes with respect to the neutrosophic Kripke frame k_N , the neutrosophic (t, i, f)-assignment v_N , and the neutrosophic world w_N .

One defines α_N by neutrosophic induction:

1. $\alpha_N(\mathcal{P}; k_N, v_N, w_N) \stackrel{def}{=} v_N(\mathcal{P}, w_N)$ if $\mathcal{P} \in S_N$ and $w_N \in G_N$.
2. $\alpha_N(\neg \mathcal{P}; k_N, v_N, w_N) \stackrel{def}{=} \neg_N[\alpha_N(\mathcal{P}; k_N, v_N, w_N)]$.
3. $\alpha_N(\mathcal{P} \wedge \mathcal{Q}; k_N, v_N, w_N) \stackrel{def}{=} [\alpha_N(\mathcal{P}; k_N, v_N, w_N)] \wedge_N [\alpha_N(\mathcal{Q}; k_N, v_N, w_N)]$
4. $\alpha_N(\mathcal{P} \vee \mathcal{Q}; k_N, v_N, w_N) \stackrel{def}{=} [\alpha_N(\mathcal{P}; k_N, v_N, w_N)] \vee_N [\alpha_N(\mathcal{Q}; k_N, v_N, w_N)]$
5. $\alpha_N(\mathcal{P} \rightarrow \mathcal{Q}; k_N, v_N, w_N) \stackrel{def}{=} [\alpha_N(\mathcal{P}; k_N, v_N, w_N)] \rightarrow_N [\alpha_N(\mathcal{Q}; k_N, v_N, w_N)]$
6. $\alpha_N(\diamond \mathcal{P}; k_N, v_N, w_N) \stackrel{def}{=} \langle \sup, \inf, \inf \rangle \{ \alpha_N(\mathcal{P}; k_N, v_N, w'_N), w'_N \in G_N \text{ and } w_N R_N w'_N \}$.
7. $\alpha_N(\square \mathcal{P}; k_N, v_N, w_N) \stackrel{def}{=} \langle \inf, \sup, \sup \rangle \{ \alpha_N(\mathcal{P}; k_N, v_N, w'_N), w'_N \in G_N \text{ and } w_N R_N w'_N \}$.
8. $\stackrel{F}{=} \mathcal{P}$ if and only if $w_N * \models \mathcal{P}$

(a formula \mathcal{P} is neutrosophically deducible if and only if \mathcal{P} is neutrosophically deducible in the actual neutrosophic world).

One should remark that α_N has a degree of truth (t_{α_N}), a degree of indeterminacy (i_{α_N}), and a degree of falsehood (f_{α_N}), which are in the general case subsets of the interval [0, 1].

Applying $\langle \sup, \inf, \inf \rangle$ to α_N is equivalent to calculating:

$$\langle \sup(t_{\alpha_N}), \inf(i_{\alpha_N}), \inf(f_{\alpha_N}) \rangle,$$

and similarly

$$\langle \inf, \sup, \sup \rangle \alpha_N = \langle \inf(t_{\alpha_N}), \sup(i_{\alpha_N}), \sup(f_{\alpha_N}) \rangle.$$

2.15. Refined Neutrosophic Modal Single-Valued Logic

Using neutrosophic (t, i, f) - thresholds, I refine for the first time the neutrosophic modal logic as:

a) Refined Neutrosophic Possibility Operator.

$\diamond_{1N} \mathcal{P}_{(t,i,f)} = \langle \text{It is very little possible (degree of possibility } t_1) \text{ that } \mathcal{P} \rangle$, corresponding to the threshold (t_1, i_1, f_1) , i.e. $0 \leq t \leq t_1, i \geq i_1, f \geq f_1$, for t_1 a very little number in $[0, 1]$;

$\diamond_{2N} \mathcal{P}_{(t,i,f)} = \langle \text{It is little possible (degree of possibility } t_2) \text{ that } \mathcal{P} \rangle$, corresponding to the threshold (t_2, i_2, f_2) , i.e. $t_1 < t \leq t_2, i \geq i_2 > i_1, f \geq f_2 > f_1$;

... ..

and so on;

$\diamond_m \mathcal{P}_{(t,i,f)} = \langle \text{It is possible (with a degree of possibility } t_m) \text{ that } \mathcal{P} \rangle$, corresponding to the threshold (t_m, i_m, f_m) , i.e. $t_{m-1} < t \leq t_m, i \geq i_m > i_{m-1}, f \geq f_m > f_{m-1}$.

b) Refined Neutrosophic Necessity Operator.

$\square_{1N} \mathcal{P}_{(t,i,f)} = \langle \text{It is a small necessity (degree of necessity } t_{m+1}) \text{ that } \mathcal{P} \rangle$, i.e. $t_m < t \leq t_{m+1}, i \geq i_{m+1}, f \geq f_{m+1} > f_m$;

$\square_{2N} \mathcal{P}_{(t,i,f)} = \langle \text{It is a little bigger necessity (degree of necessity } t_{m+2}) \text{ that } \mathcal{P} \rangle$, i.e. $t_{m+1} < t \leq t_{m+2}, i \geq i_{m+2} > i_{m+1}, f \geq f_{m+2} > f_{m+1}$;

... ..

and so on;

$\square_k \mathcal{P}_{(t,i,f)} = \langle \text{It is a very high necessity (degree of necessity } t_{m+k}) \text{ that } \mathcal{P} \rangle$, i.e. $t_{m+k-1} < t \leq t_{m+k} = 1, i \geq i_{m+k} > i_{m+k-1}, f \geq f_{m+k} > f_{m+k-1}$.

2.16. Application of the Neutrosophic Threshold

I have introduced the term of (t, i, f) -physical law, meaning that a physical law has a degree of truth (t), a degree of indeterminacy (i), and a degree of falsehood (f). A physical law is 100% true, 0% indeterminate, and 0% false in perfect (ideal) conditions only, maybe in laboratory.

But our actual world (w_N^*) is not perfect and not steady, but continuously changing, varying, fluctuating.

For example, there are physicists that have proved a universal constant (c) is not quite universal (i.e. there are special conditions where it does not apply, or its value varies between $(c - \varepsilon, c + \varepsilon)$, for $\varepsilon > 0$ that can be a tiny or even a bigger number).

Thus, one can say that a proposition \mathcal{P} is **neutrosophically nomological necessary**, if \mathcal{P} is neutrosophically true at all possible neutrosophic worlds that obey the (t, i, f) -physical laws of the actual neutrosophic world w_N^* .

In other words, at each possible neutrosophic world w_N , neutrosophically accesible from w_N^* , one has:

$$\mathcal{P}(t_p^{w_N}, i_p^{w_N}, f_p^{w_N}) \geq TH(T_{th}, I_{th}, F_{th}), \quad (13)$$

i.e. $t_p^{w_N} \geq T_{th}, i_p^{w_N} \leq I_{th}, \text{ and } f_p^{w_N} \geq F_{th}$.

2.17. Neutrosophic Mereology

Neutrosophic Mereology means the theory of the neutrosophic relations among the parts of a whole, and the neutrosophic relations between the parts and the whole.

A neutrosophic relation between two parts, and similarly a neutrosophic relation between a part and the whole, has a degree of connection (t), a degree of indeterminacy (i), and a degree of disconnection (f).

2.18. Neutrosophic Mereological Threshold

Neutrosophic Mereological Threshold is defined as:

$$TH_M = (\min(t_M), \max(i_G), \max(f_M)), \quad (14)$$

where t_M is the set of all degrees of connection between the parts, and between the parts and the whole; i_M is the set of all degrees of indeterminacy between the parts, and between the parts and the whole; f_M is the set of all degrees of disconnection between the parts, and between the parts and the whole.

I have considered all degrees as single-valued numbers.

2.19. Neutrosophic Gnosisology

Neutrosophic Gnosisology is the theory of (t, i, f) -knowledge, because in many cases we are not able to completely (100%) find whole knowledge, but only a part of it ($t\%$), another part remaining unknown ($f\%$), and a third part indeterminate (unclear, vague, contradictory) ($i\%$), where t, i, f are subsets of the interval $[0, 1]$.

2.20. Neutrosophic Gnosisological Threshold

Neutrosophic Gnosisological Threshold is defined, similarly, as:

$$TH_G = (\min(t_G), \max(i_G), \max(f_G)), \quad (15)$$

where t_G is the set of all degrees of knowledge of all theories, ideas, propositions etc., i_G is the set of all degrees of indeterminate-knowledge of all theories, ideas, propositions etc., f_G is the set of all degrees of non-knowledge of all theories, ideas, propositions etc.

I have considered all degrees as single-valued numbers.

2.21. Neutrosophic Epistemology

Neutrosophic Epistemology, as part of the Neutrosophic Gnosisology, is the theory of (t, i, f) -scientific knowledge.

Science is infinite. We know only a small part of it ($t\%$), another big part is yet to be discovered ($f\%$), and a third part indeterminate (unclear, vague, contradictory) ($i\%$).

Of course, t, i, f are subsets of $[0, 1]$.

2.22. Neutrosophic Epistemological Threshold

Neutrosophic Epistemological Threshold is defined as:

$$TH_E = (\min(t_E), \max(i_E), \max(f_E)), \quad (16)$$

where t_E is the set of all degrees of scientific knowledge of all scientific theories, ideas, propositions etc., i_E is the set of all degrees of indeterminate scientific knowledge of all scientific theories, ideas, propositions etc., f_E is the set of all degrees of non-scientific knowledge of all scientific theories, ideas, propositions etc.

I have considered all degrees as single-valued numbers.

3. Discussion & Conclusions

I have introduced for the first time the Neutrosophic Modal Logic and the Refined Neutrosophic Modal Logic.

Symbolic Neutrosophic Logic can be connected to the neutrosophic modal logic too, where instead of numbers we may use labels, or instead of quantitative neutrosophic logic we may have a quantitative neutrosophic logic. As an extension, I may introduce **Symbolic Neutrosophic Modal Logic** and **Refined Symbolic Neutrosophic Modal Logic**, where the symbolic neutrosophic modal operators (and the symbolic neutrosophic accessibility relation) have qualitative values (labels) instead on numerical values (subsets of the interval $[0, 1]$).

Applications of neutrosophic modal logic are to neutrosophic modal metaphysics. Similarly to classical modal logic, there is a plethora of neutrosophic modal logics. Neutrosophic modal logics is governed by a set of neutrosophic axioms and neutrosophic rules. The neutrosophic accessibility relation has various interpretations, depending on the applications. Similarly, the notion of possible neutrosophic worlds has many interpretations, as part of possible neutrosophic semantics.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Girle, Rod. *Modal Logics and Philosophy*, 2nd ed.; McGill-Queen's University Press, 2010.
2. Liao, C. J.; Lin, B. I-Pen. Quantitative Modal Logic and Possibilistic Reasoning. 10th European Conference on Artificial Intelligence, pp. 43-47, 1992.
3. Hájek, P.; Harmancová, D. A comparative fuzzy modal logic. *Fuzzy Logic in Artificial Intelligence*, Lecture Notes in AI 695, pp. 27-34, 1993.
4. Smarandache, F. (t, i, f)-Physical Laws and (t, i, f)-Physical Constants. 47th Annual Meeting of the APS Division of Atomic, Molecular and Optical Physics, Volume 61, Number 8, Monday-Friday, May 23-27, 2016; Providence, Rhode Island. <http://meetings.aps.org/Meeting/DAMOP16/Session/Q1.197>
5. Smarandache, F.; Ali, M. Neutrosophic Triplet as extension of Matter Plasma, Unmatter Plasma, and Antimatter Plasma. 69th Annual Gaseous Electronics Conference, Volume 61, Number 9, Monday-Friday, October 10-14, 2016; Bochum, Germany. <http://meetings.aps.org/Meeting/GEC16/Session/HT6.111>
6. Smarandache, Florentin. *Neutrosophic Overset, Neutrosophic Underset, and Neutrosophic Offset. Similarly for Neutrosophic Over-/Under-/Off- Logic, Probability, and Statistics*. Pons Editions, Brussels, Belgium, 2016; Available online: <https://hal.archives-ouvertes.fr/hal-01340830>.
7. Smarandache, Florentin. *Symbolic Neutrosophic Theory*, Europa Nova, Brussels, 2015. Available online: <https://arxiv.org/ftp/arxiv/papers/1512/1512.00047.pdf>
8. José Luis Usó-Doménech, Josué Antonio Nescolarde-Selva and Lorena Segura-Abad. Proposal for the Formalization of Dialectical Logic, *Mathematics* 2016, 4(4), 69; doi:10.3390/math4040069.
9. Florentin Smarandache, *Neutrosophy. Neutrosophic Probability, Set, and Logic*, ProQuest Information & Learning, Ann Arbor, Michigan, USA, 105 p., 1998; <http://fs.gallup.unm.edu/eBook-neutrosophics6.pdf> (last edition online);
10. Neutrosophy, A New Branch of Philosophy, Multiple Valued Logic / An International Journal, USA, Vol. 8, No. 3, pp. 297-384, 2002.
11. University of New Mexico, USA, website: <http://fs.gallup.unm.edu/neutrosophy.htm>



© 2017 by the authors; licensee *Preprints*, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons by Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).