

CLOSED EXPRESSIONS OF THE FIBONACCI POLYNOMIALS IN TERMS OF TRIDIAGONAL DETERMINANTS

FENG QI

Institute of Mathematics, Henan Polytechnic University, Jiaozuo City, Henan Province, 454010, China; College of Mathematics, Inner Mongolia University for Nationalities, Tongliao City, Inner Mongolia Autonomous Region, 028043, China; Department of Mathematics, College of Science, Tianjin Polytechnic University, Tianjin City, 300387, China

JING-LIN WANG

Department of Mathematics, College of Science, Tianjin Polytechnic University, Tianjin City, 300387, China

BAI-NI GUO

School of Mathematics and Informatics, Henan Polytechnic University, Jiaozuo City, Henan Province, 454010, China

ABSTRACT. In the paper, the authors find a new closed expression for the Fibonacci polynomials and, consequently, for the Fibonacci numbers, in terms of a tridiagonal determinant.

1. MAIN RESULTS

A tridiagonal matrix (determinant) is a square matrix (determinant) with nonzero elements only on the diagonal and slots horizontally or vertically adjacent the diagonal. See the paper [14] and closely-related references therein.

A square matrix $H = (h_{ij})_{n \times n}$ is called a tridiagonal matrix if $h_{ij} = 0$ for all pairs (i, j) such that $|i - j| > 1$. On the other hand, a matrix $H = (h_{ij})_{n \times n}$ is called a lower (or an upper, respectively) Hessenberg matrix if $h_{ij} = 0$ for all pairs (i, j) such that $i + 1 < j$ (or $j + 1 < i$, respectively). See the paper [22] and closely-related references therein.

E-mail addresses: qifeng618@gmail.com, qifeng618@hotmail.com, jing-lin.wang@hotmail.com, bai.ni.guo@gmail.com, bai.ni.guo@hotmail.com.

2010 *Mathematics Subject Classification.* 11B39, 11B83, 11C20, 11Y55, 15A15, 65F40.

Key words and phrases. closed expression; Fibonacci number; Fibonacci polynomial; tridiagonal determinant; Hessenberg determinant.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

In mathematics, a closed form is a mathematical expression that can be evaluated in a finite number of operations. It may contain constants, variables, four arithmetic operations, and elementary functions, but usually no limit.

It is general knowledge that the Bernoulli numbers and polynomials B_k and $B_k(u)$ for $k \geq 0$ satisfy $B_k(0) = B_k$ and can be generated respectively by

$$\frac{z}{e^z - 1} = \sum_{k=0}^{\infty} B_k \frac{z^k}{k!} = 1 - \frac{z}{2} + \sum_{k=1}^{\infty} B_{2k} \frac{z^{2k}}{(2k)!} \quad \text{and} \quad \frac{ze^{uz}}{e^z - 1} = \sum_{k=0}^{\infty} B_k(u) \frac{z^k}{k!}$$

for $|z| < 2\pi$. Because the function $\frac{x}{e^x - 1} - 1 + \frac{x}{2}$ is odd in $x \in \mathbb{R}$, all of the Bernoulli numbers B_{2k+1} for $k \in \mathbb{N}$ equal 0. In [26, Theorem 1.2], the Bernoulli polynomials $B_k(u)$ for $k \in \mathbb{N}$ were expressed by lower Hessenberg determinants

$$B_k(u) = (-1)^k \left| \frac{1}{\ell+1} \binom{\ell+1}{m} \left[(1-u)^{\ell-m+1} - (-u)^{\ell-m+1} \right] \right|_{1 \leq \ell \leq k, 0 \leq m \leq k-1}.$$

Consequently, the Bernoulli numbers B_k for $k \in \mathbb{N}$ can be represented by lower Hessenberg determinants

$$B_k = (-1)^k \left| \frac{1}{\ell+1} \binom{\ell+1}{m} \right|_{1 \leq \ell \leq k, 0 \leq m \leq k-1}.$$

For $k \in \{0\} \cup \mathbb{N}$ and $x \in \mathbb{R}$, the Euler numbers E_k and the Euler polynomials $E_k(x)$ can be generated respectively by

$$\frac{2e^{t/2}}{e^t + 1} = \sum_{k=0}^{\infty} \frac{E_k}{k!} \left(\frac{t}{2}\right)^k \quad \text{and} \quad \frac{2e^{xt}}{e^t + 1} = \sum_{k=0}^{\infty} \frac{E_k(x)}{k!} t^k$$

for $t \in (-\pi, \pi)$. Since the generating function $\frac{2e^{t/2}}{e^t + 1}$ of the Euler numbers E_k is even on $(-\pi, \pi)$, then $E_{2k-1} = 0$ for all $k \in \mathbb{N}$. At the website [35], the Euler numbers E_{2k} were represented by lower Hessenberg determinants

$$E_{2k} = (-1)^k (2k)! \begin{vmatrix} \frac{1}{2!} & 1 & 0 & \cdots & 0 & 0 \\ \frac{1}{4!} & \frac{1}{2!} & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \frac{1}{(2k-2)!} & \frac{1}{(2k-4)!} & \frac{1}{(2k-6)!} & \cdots & \frac{1}{2!} & 1 \\ \frac{1}{(2k)!} & \frac{1}{(2k-2)!} & \frac{1}{(2k-4)!} & \cdots & \frac{1}{4!} & \frac{1}{2!} \end{vmatrix}, \quad k \in \mathbb{N}.$$

In [34, Theorem 1.1], the Euler numbers E_{2k} for $k \in \mathbb{N}$ were represented by lower Hessenberg and sparse determinants

$$E_{2k} = (-1)^k \left| \binom{i}{j-1} \cos\left((i-j+1)\frac{\pi}{2}\right) \right|_{(2k) \times (2k)}.$$

Recently, the first author represented the Euler polynomials $E_k(x)$ for $k \geq 0$ by lower Hessenberg determinants

$$E_k(x) = \frac{(-1)^k}{2^k} \begin{vmatrix} 1 & 2 & 0 & 0 & \cdots & 0 & 0 & 0 \\ x & \binom{1}{0} & 2 & 0 & \cdots & 0 & 0 & 0 \\ x^2 & \binom{2}{0} & \binom{2}{1} & 2 & \cdots & 0 & 0 & 0 \\ x^3 & \binom{3}{0} & \binom{3}{1} & \binom{3}{2} & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ x^{k-2} & \binom{k-2}{0} & \binom{k-2}{1} & \binom{k-2}{2} & \cdots & \binom{k-2}{k-3} & 2 & 0 \\ x^{k-1} & \binom{k-1}{0} & \binom{k-1}{1} & \binom{k-1}{2} & \cdots & \binom{k-1}{k-3} & \binom{k-1}{k-2} & 2 \\ x^k & \binom{k}{0} & \binom{k}{1} & \binom{k}{2} & \cdots & \binom{k}{k-3} & \binom{k}{k-2} & \binom{k}{k-1} \end{vmatrix}.$$

Consequently, the Euler numbers E_k for $k \geq 0$ can be expressed by lower Hessenberg determinants

$$E_k = (-1)^k \begin{vmatrix} 1 & 2 & 0 & 0 & \cdots & 0 & 0 & 0 \\ \frac{1}{2} & \binom{1}{0} & 2 & 0 & \cdots & 0 & 0 & 0 \\ \frac{1}{2^2} & \binom{2}{0} & \binom{2}{1} & 2 & \cdots & 0 & 0 & 0 \\ \frac{1}{2^3} & \binom{3}{0} & \binom{3}{1} & \binom{3}{2} & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ \frac{1}{2^{k-2}} & \binom{k-2}{0} & \binom{k-2}{1} & \binom{k-2}{2} & \cdots & \binom{k-2}{k-3} & 2 & 0 \\ \frac{1}{2^{k-1}} & \binom{k-1}{0} & \binom{k-1}{1} & \binom{k-1}{2} & \cdots & \binom{k-1}{k-3} & \binom{k-1}{k-2} & 2 \\ \frac{1}{2^k} & \binom{k}{0} & \binom{k}{1} & \binom{k}{2} & \cdots & \binom{k}{k-3} & \binom{k}{k-2} & \binom{k}{k-1} \end{vmatrix}.$$

For more and detailed information on the Bernoulli numbers B_k , the Bernoulli polynomials $B_k(u)$, the Euler numbers E_k , and the Euler polynomials $E_k(x)$, please refer to recently published papers such as [2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 18, 19, 20, 21, 23, 25, 26, 27, 28, 29, 31, 34] and plenty of closely-related references therein.

It is well-known that the Fibonacci numbers

$$F_n = \frac{(1 + \sqrt{5})^n - (1 - \sqrt{5})^n}{2^n \sqrt{5}}$$

for $n \in \mathbb{N}$ form a sequence of integers and satisfy the linear recurrence relation

$$F_n = F_{n-1} + F_{n-2} \tag{1}$$

with $F_1 = F_2 = 1$. The first fourteen Fibonacci numbers F_n for $1 \leq n \leq 14$ are

$$1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, 233, 377.$$

The Fibonacci numbers F_n can be viewed as a particular case $F_n(1)$ of the Fibonacci polynomials

$$F_n(s) = \frac{1}{2^n} \frac{(s + \sqrt{4 + s^2})^n - (s - \sqrt{4 + s^2})^n}{\sqrt{4 + s^2}}$$

which can be generated by

$$\frac{t}{1 - ts - t^2} = \sum_{n=1}^{\infty} F_n(s)t^n = t + st^2 + (s^2 + 1)t^3 + (s^3 + 2s)t^4 + \cdots \tag{2}$$

See the monograph [5], the websites [32, 33], and related references therein. In [17, p. 215, Example 1], it was deduced that

$$F_n = \begin{pmatrix} 1 & -1 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 1 & -1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 1 & -1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 1 & -1 \\ 0 & 0 & 0 & 0 & \cdots & 1 & 1 \end{pmatrix}_{(n-1) \times (n-1)}, \quad n \in \mathbb{N}.$$

In [15, 16], among other things, it was listed that

$$F_{r+1}(x) = \begin{pmatrix} x & 1 & 0 & \cdots & 0 & 0 \\ -1 & x & 1 & \cdots & 0 & 0 \\ 0 & -1 & x & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & x & 1 \\ 0 & 0 & 0 & \cdots & -1 & x \end{pmatrix}_{r \times r}, \quad F_{r-1} = \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 & 1 \\ -1 & 0 & 1 & \cdots & 1 & 1 \\ 0 & -1 & 0 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & \cdots & -1 & 0 \end{pmatrix}_{r \times r},$$

$$F_{r-1} = |q_{ij}|_{r \times r}, \quad F_{2r} = \begin{pmatrix} 1 & 2 & 3 & \cdots & r-1 & r \\ -1 & 1 & 2 & \cdots & r-2 & r-1 \\ 0 & -1 & 1 & \cdots & r-3 & r-2 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 2 \\ 0 & 0 & 0 & \cdots & -1 & 1 \end{pmatrix}_{r \times r}, \quad F_{2r+1} = |Q_{ij}|_{r \times r}$$

for $r \in \mathbb{N}$, where

$$q_{ij} = \begin{cases} -1, & i - j + 1; \\ i + j + 1 \pmod{2}, & i \leq j; \\ 0, & \text{otherwise} \end{cases} \quad \text{and} \quad Q_{ij} = \begin{cases} -1, & i = j + 1; \\ 2, & i = j; \\ 1, & i < j; \\ 0, & \text{otherwise.} \end{cases}$$

In this paper, motivated by the above closed expressions for the Bernoulli numbers B_k , the Bernoulli polynomials $B_k(u)$, the Euler numbers E_k , the Euler polynomials $E_k(x)$, the Fibonacci numbers F_n , and the Fibonacci polynomials $F_n(s)$, we will find a new closed expression for the Fibonacci polynomials $F_n(s)$ and, consequently, for the Fibonacci numbers F_n , in terms of a tridiagonal determinant.

The main results of this paper can be stated as the following theorem.

Theorem 1. For $n \in \mathbb{N}$, the Fibonacci polynomials $F_n(s)$ can be expressed as

$$F_n(s) = \frac{1}{n!} \begin{vmatrix} 0 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 1 & s & -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 2 \cdot 1 & 2s & -1 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 3 \cdot 2 & 3s & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & (n-2)s & -1 & 0 \\ 0 & 0 & 0 & 0 & \cdots & (n-1)(n-2) & (n-1)s & -1 \\ 0 & 0 & 0 & 0 & \cdots & 0 & n(n-1) & ns \end{vmatrix} \quad (3)$$

and, consequently,

$$F_n = \frac{1}{n!} \begin{vmatrix} 0 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 1 & 1 & -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 2 \cdot 1 & 2 & -1 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 3 \cdot 2 & 3 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & n-2 & -1 & 0 \\ 0 & 0 & 0 & 0 & \cdots & (n-1)(n-2) & n-1 & -1 \\ 0 & 0 & 0 & 0 & \cdots & 0 & n(n-1) & n \end{vmatrix}. \quad (4)$$

2. A LEMMA

In order to supply a concise proof of Theorem 1, we need the following lemma which can be found in [1, p. 40, Exercise 5], [24, Section 2.2, p. 849], [26, p. 94], [30, Remark 6], and [34, Lemma 2.1].

Lemma 1. Let $u(x)$ and $v(x) \neq 0$ be two differentiable functions, let $U_{(n+1) \times 1}(x)$ be an $(n+1) \times 1$ matrix whose elements $u_{k,1}(x) = u^{(k-1)}(x)$ for $1 \leq k \leq n+1$, let $V_{(n+1) \times n}(x)$ be an $(n+1) \times n$ matrix whose elements

$$v_{i,j}(x) = \begin{cases} \binom{i-1}{j-1} v^{(i-j)}(x), & i-j \geq 0; \\ 0, & i-j < 0 \end{cases}$$

for $1 \leq i \leq n+1$ and $1 \leq j \leq n$, and let $|W_{(n+1) \times (n+1)}(x)|$ denote the lower Hessenberg determinant of the $(n+1) \times (n+1)$ lower Hessenberg matrix

$$W_{(n+1) \times (n+1)}(x) = [U_{(n+1) \times 1}(x) \quad V_{(n+1) \times n}(x)].$$

Then the n -th derivative of the ratio $\frac{u(x)}{v(x)}$ can be computed by

$$\frac{d^n}{dx^n} \left[\frac{u(x)}{v(x)} \right] = (-1)^n \frac{|W_{(n+1) \times (n+1)}(x)|}{v^{n+1}(x)}. \quad (5)$$

3. PROOF OF THEOREM 1

Now we start out to concisely prove Theorem 1 by virtue of the formula (5).

Applying $u(t) = t$ and $v(t) = 1 - ts - t^2$ to the formula (5) yields

$$\frac{d^n}{dt^n} \left(\frac{t}{1 - ts - t^2} \right) = \frac{(-1)^n}{(1 - ts - t^2)^{n+1}} \begin{vmatrix} t & 1 - ts - t^2 & 0 & 0 & \dots \\ 1 & -(2t + s) \binom{1}{0} & 1 - ts - t^2 & 0 & \dots \\ 0 & -2 \binom{2}{0} & -(2t + s) \binom{2}{1} & 1 - ts - t^2 & \dots \\ 0 & 0 & -2 \binom{3}{1} & -(2t + s) \binom{3}{2} & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \\ 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \\ - (2t + s) \binom{n-2}{n-3} & 1 - ts - t^2 & 0 & & \\ -2 \binom{n-1}{n-3} & - (2t + s) \binom{n-1}{n-2} & 1 - ts - t^2 & & \\ 0 & -2 \binom{n}{n-2} & - (2t + s) \binom{n}{n-1} & & \end{vmatrix} \\ \rightarrow (-1)^n \begin{vmatrix} 0 & 1 & 0 & 0 & \dots & 0 & 0 & 0 \\ 1 & -s \binom{1}{0} & 1 & 0 & \dots & 0 & 0 & 0 \\ 0 & -2 \binom{2}{0} & -s \binom{2}{1} & 1 & \dots & 0 & 0 & 0 \\ 0 & 0 & -2 \binom{3}{1} & -s \binom{3}{2} & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & -s \binom{n-2}{n-3} & 1 & 0 \\ 0 & 0 & 0 & 0 & \dots & -2 \binom{n-1}{n-3} & -s \binom{n-1}{n-2} & 1 \\ 0 & 0 & 0 & 0 & \dots & 0 & -2 \binom{n}{n-2} & -s \binom{n}{n-1} \end{vmatrix}$$

as $t \rightarrow 0$ for $n \in \mathbb{N}$. By the generating function in (2), we obtain that

$$F_n(s) = \frac{1}{n!} \lim_{t \rightarrow 0} \frac{d^n}{dt^n} \left(\frac{t}{1 - ts - t^2} \right) \\ = \frac{(-1)^n}{n!} \begin{vmatrix} 0 & 1 & 0 & 0 & \dots & 0 & 0 & 0 \\ 1 & -s \binom{1}{0} & 1 & 0 & \dots & 0 & 0 & 0 \\ 0 & -2 \binom{2}{0} & -s \binom{2}{1} & 1 & \dots & 0 & 0 & 0 \\ 0 & 0 & -2 \binom{3}{1} & -s \binom{3}{2} & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & -s \binom{n-2}{n-3} & 1 & 0 \\ 0 & 0 & 0 & 0 & \dots & -2 \binom{n-1}{n-3} & -s \binom{n-1}{n-2} & 1 \\ 0 & 0 & 0 & 0 & \dots & 0 & -2 \binom{n}{n-2} & -s \binom{n}{n-1} \end{vmatrix}$$

which can be rewritten as the expression (3).

The expression (4) follows from taking the limit $s \rightarrow 1$ on both sides of the expression (3). The proof of Theorem 1 is complete.

4. REMARKS

Finally, we list several remarks on Lemma 1, Theorem 1, and others as follows.

Remark 1. As showed, for example, in recently published papers [24, 26, 30, 34] and in this paper, the formula (5) in Lemma 1 is an effectual tool to express some quantities in mathematics as lower Hessenberg determinants.

Remark 2. It is easy to see that, from the expressions (3) and (4), we can recover the recurrence relation

$$F_n(s) = sF_{n-1}(s) + F_{n-2}(s)$$

in [33] and the recurrence relation (1) for $n \geq 3$.

Remark 3. The expressions (3) and (4) can be rearranged as

$$F_n(s) = \frac{1}{n!} \begin{vmatrix} \binom{2}{1}s & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 2\binom{3}{1} & \binom{3}{2}s & -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 2\binom{4}{2} & \binom{4}{3}s & -1 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 2\binom{5}{3} & \binom{5}{4}s & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & \binom{n-2}{n-3}s & -1 & 0 \\ 0 & 0 & 0 & 0 & \cdots & 2\binom{n-1}{n-3} & \binom{n-1}{n-2}s & -1 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 2\binom{n}{n-2} & \binom{n}{n-1}s \end{vmatrix}$$

and

$$F_n = \frac{1}{n!} \begin{vmatrix} \binom{2}{1} & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 2\binom{3}{1} & \binom{3}{2} & -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 2\binom{4}{2} & \binom{4}{3} & -1 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 2\binom{5}{3} & \binom{5}{4} & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & \binom{n-2}{n-3} & -1 & 0 \\ 0 & 0 & 0 & 0 & \cdots & 2\binom{n-1}{n-3} & \binom{n-1}{n-2} & -1 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 2\binom{n}{n-2} & \binom{n}{n-1} \end{vmatrix}$$

for $n \geq 2$.

Remark 4. The Bernoulli numbers B_n , the Bernoulli polynomials $B_n(u)$, the Euler numbers E_n , and the Euler polynomials $E_n(x)$ can be expressed as different closed forms. Let $S(n, m)$ stand for the Stirling numbers of the second kind which can be computed by

$$S(n, m) = \frac{1}{m!} \sum_{\ell=1}^m (-1)^{m-\ell} \binom{m}{\ell} \ell^n, \quad 1 \leq m \leq n.$$

In [26, Theorem 1.1], the Bernoulli polynomials $B_n(u)$ were expressed as

$$B_n(u) = \sum_{k=1}^n k! \sum_{r+s=k} \sum_{\ell+m=n} (-1)^m \binom{n}{\ell} \frac{\ell!}{(\ell+r)!} \frac{m!}{(m+s)!} \\ \times \left[\sum_{i=0}^r \sum_{j=0}^s (-1)^{i+j} \binom{\ell+r}{r-i} \binom{m+s}{s-j} S(\ell+i, i) S(m+j, j) \right] u^{m+s} (1-u)^{\ell+r}$$

for $n \in \mathbb{N}$. Consequently, the Bernoulli numbers B_n can be represented as

$$B_n = \sum_{i=1}^n (-1)^i \frac{\binom{n+1}{i+1}}{\binom{n+i}{i}} S(n+i, i), \quad n \in \mathbb{N}.$$

In [34, Theorem 1.3], the expressions

$$E_n = 1 + \sum_{k=1}^n \frac{(k+1)!}{2^k} S(n, k) \sum_{\ell=1}^k (-1)^\ell \frac{2^\ell}{\ell+1} \binom{\ell+1}{k-\ell}$$

and

$$E_n = 1 + \sum_{\ell=1}^n (-1)^\ell \frac{1}{\ell+1} \sum_{k=0}^{n-\ell} \frac{(k+\ell+1)!}{2^k} \binom{\ell+1}{k} S(n, k+\ell)$$

for $n \in \mathbb{N}$ were acquired. In [10, Theorem 3.1], the Euler polynomials $E_n(x)$ were represented by

$$E_n(x) = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \left[\sum_{\ell=1}^{n-k+1} \frac{(-1)^{\ell-1} (\ell-1)!}{2^{\ell-1}} S(n-k+1, \ell) \right] x^k, \quad n \in \mathbb{N}.$$

Consequently, the Euler numbers E_{2n} for $n \in \mathbb{N}$ can be calculated by

$$E_{2n} = 4^n \sum_{k=0}^{2n} \left[\sum_{\ell=1}^{2n-k+1} \frac{(-1)^{\ell-1} (\ell-1)!}{2^{\ell-1}} S(2n-k+1, \ell) \right] \frac{(-1)^k}{2^k} \binom{2n}{k}.$$

In mathematics, different forms have different implications. In other words, from different forms of mathematical quantities, one can read out different meanings and significance. Therefore, expressing the Fibonacci numbers F_n and the Fibonacci polynomials $F_n(s)$ in terms of tridiagonal determinants is meaningful and significant.

REFERENCES

- [1] N. Bourbaki, *Elements of Mathematics: Functions of a Real Variable: Elementary Theory*, Translated from the 1976 French original by Philip Spain. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2004; Available online at <http://dx.doi.org/10.1007/978-3-642-59315-4>.
- [2] C.-P. Chen and F. Qi, *Three improper integrals relating to the generating function of Bernoulli numbers*, Octagon Math. Mag. **11** (2003), no. 2, 408–409.
- [3] D. V. Dolgy, T. Kim, F. Qi, and J. J. Seo, *A note on three variable symmetric identities for modified q -Bernoulli polynomials arising from bosonic p -adic integral on \mathbb{Z}_p* , Appl. Math. Sci. **9** (2015), no. 92, 4575–4582; Available online at <http://dx.doi.org/10.12988/ams.2015.5339>.
- [4] D. V. Dolgy, T. Kim, F. Qi, and J. J. Seo, *A note on three variable symmetric identities for q -Euler polynomials arising from fermionic p -adic integral on \mathbb{Z}_p* , Appl. Math. Sci. **9** (2015), no. 77, 3819–3826; Available online at <http://dx.doi.org/10.12988/ams.2015.53229>.
- [5] R. A. Dunlap, *The Golden Ratio and Fibonacci Numbers*, World Scientific Publishing Co., Inc., River Edge, NJ, 1997; Available online at <http://dx.doi.org/10.1142/9789812386304>.
- [6] B.-N. Guo, I. Mezö, and F. Qi, *An explicit formula for the Bernoulli polynomials in terms of the r -Stirling numbers of the second kind*, Rocky Mountain J. Math. **46** (2016), no. 6, 1919–1923; Available online at <http://dx.doi.org/10.1216/RMJ-2016-46-6-1919>.
- [7] B.-N. Guo and F. Qi, *A new explicit formula for the Bernoulli and Genocchi numbers in terms of the Stirling numbers*, Glob. J. Math. Anal. **3** (2015), no. 1, 33–36; Available online at <http://dx.doi.org/10.14419/gjma.v3i1.4168>.
- [8] B.-N. Guo and F. Qi, *Alternative proofs of a formula for Bernoulli numbers in terms of Stirling numbers*, Analysis (Berlin) **34** (2014), no. 2, 187–193; Available online at <http://dx.doi.org/10.1515/analy-2012-1238>.
- [9] B.-N. Guo and F. Qi, *An explicit formula for Bernoulli numbers in terms of Stirling numbers of the second kind*, J. Anal. Number Theory **3** (2015), no. 1, 27–30; Available online at <http://dx.doi.org/10.12785/jant/030105>.

- [10] B.-N. Guo and F. Qi, *Explicit formulae for computing Euler polynomials in terms of Stirling numbers of the second kind*, J. Comput. Appl. Math. **272** (2014), 251–257; Available online at <http://dx.doi.org/10.1016/j.cam.2014.05.018>.
- [11] B.-N. Guo and F. Qi, *Explicit formulas for special values of the Bell polynomials of the second kind and the Euler numbers*, ResearchGate Technical Report (2015), available online at <http://dx.doi.org/10.13140/2.1.3794.8808>.
- [12] B.-N. Guo and F. Qi, *Generalization of Bernoulli polynomials*, Internat. J. Math. Ed. Sci. Tech. **33** (2002), no. 3, 428–431; Available online at <http://dx.doi.org/10.1080/002073902760047913>.
- [13] B.-N. Guo and F. Qi, *Some identities and an explicit formula for Bernoulli and Stirling numbers*, J. Comput. Appl. Math. **255** (2014), 568–579; Available online at <http://dx.doi.org/10.1016/j.cam.2013.06.020>.
- [14] V. Higgins and C. Johnson, *Inverse spectral problems for collections of leading principal submatrices of tridiagonal matrices*, Linear Algebra Appl. **489** (2016), 104–122; Available online at <http://dx.doi.org/10.1016/j.laa.2015.10.004>.
- [15] M. Janjić, *Determinants and recurrence sequences*, J. Integer Seq. **15** (2012), no. 3, Article 12.3.5, 21 pages.
- [16] M. Janjić, *Recurrence relations and determinants*, arXiv preprint (2011), available online at <http://arxiv.org/abs/1112.2466>.
- [17] R. K. Kittappa, *A representation of the solution of the n th order linear difference equation with variable coefficients*, Linear Algebra Appl. **193** (1993), 211–222; Available online at [http://dx.doi.org/10.1016/0024-3795\(93\)90278-V](http://dx.doi.org/10.1016/0024-3795(93)90278-V).
- [18] Q.-M. Luo, B.-N. Guo, F. Qi, and L. Debnath, *Generalizations of Bernoulli numbers and polynomials*, Int. J. Math. Math. Sci. **2003** (2003), no. 59, 3769–3776; Available online at <http://dx.doi.org/10.1155/S0161171203112070>.
- [19] Q.-M. Luo and F. Qi, *Generalizations of Euler numbers and Euler numbers of higher order*, Chinese Quart. J. Math. **20** (2005), no. 1, 54–58.
- [20] Q.-M. Luo and F. Qi, *Relationships between generalized Bernoulli numbers and polynomials and generalized Euler numbers and polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **7** (2003), no. 1, 11–18.
- [21] Q.-M. Luo, F. Qi, and L. Debnath, *Generalizations of Euler numbers and polynomials*, Int. J. Math. Math. Sci. **2003** (2003), no. 61, 3893–3901; Available online at <http://dx.doi.org/10.1155/S016117120321108X>.
- [22] R. S. Martin and J. H. Wilkinson, *Handbook Series Linear Algebra: Similarity reduction of a general matrix to Hessenberg form*, Numer. Math. **12** (1968), no. 5, 349–368; Available online at <http://dx.doi.org/10.1007/BF02161358>.
- [23] F. Qi, *A new formula for the Bernoulli numbers of the second kind in terms of the Stirling numbers of the first kind*, Publ. Inst. Math. (Beograd) (N.S.) **100 (114)** (2016), 243–249; Available online at <http://dx.doi.org/10.2298/PIM150501028Q>.
- [24] F. Qi, *Derivatives of tangent function and tangent numbers*, Appl. Math. Comput. **268** (2015), 844–858; Available online at <http://dx.doi.org/10.1016/j.amc.2015.06.123>.
- [25] F. Qi, *Explicit formulas for computing Bernoulli numbers of the second kind and Stirling numbers of the first kind*, Filomat **28** (2014), no. 2, 319–327; Available online at <http://dx.doi.org/10.2298/FIL14023190>.
- [26] F. Qi and R. J. Chapman, *Two closed forms for the Bernoulli polynomials*, J. Number Theory **159** (2016), 89–100; Available online at <http://dx.doi.org/10.1016/j.jnt.2015.07.021>.
- [27] F. Qi, D. V. Dolgy, T. Kim, and C. S. Ryoo, *On the partially degenerate Bernoulli polynomials of the first kind*, Glob. J. Pure Appl. Math. **11** (2015), no. 4, 2407–2412.
- [28] F. Qi and B.-N. Guo, *Alternative proofs of a formula for Bernoulli numbers in terms of Stirling numbers*, Analysis (Berlin) **34** (2014), no. 3, 311–317; Available online at <http://dx.doi.org/10.1515/analy-2014-0003>.
- [29] F. Qi, D. S. Kim, T.-K. Kim, and D. V. Dolgy, *Multiple-poly-Bernoulli polynomials of the second kind*, Adv. Stud. Contemp. Math. (Kyungshang) **25** (2015), no. 1, 1–7.
- [30] F. Qi, X.-T. Shi, and F.-F. Liu, *Several identities involving the falling and rising factorials and the Cauchy, Lah, and Stirling numbers*, Acta Univ. Sapientiae Math. **8** (2016), no. 2, 282–297; Available online at <http://dx.doi.org/10.1515/ausm-2016-0019>.
- [31] F. Qi and X.-J. Zhang, *An integral representation, some inequalities, and complete monotonicity of the Bernoulli numbers of the second kind*, Bull. Korean Math. Soc. **52** (2015), no. 3, 987–998; Available online at <http://dx.doi.org/10.4134/BKMS.2015.52.3.987>.
- [32] R. Stanley and E. W. Weisstein, *Fibonacci Number*, From MathWorld—A Wolfram Web Resource; Available online at <http://mathworld.wolfram.com/FibonacciNumber.html>.

- [33] R. Stanley and E. W. Weisstein, *Fibonacci Polynomial*, From MathWorld—A Wolfram Web Resource; Available online at <http://mathworld.wolfram.com/FibonacciPolynomial.html>.
- [34] C.-F. Wei and F. Qi, *Several closed expressions for the Euler numbers*, J. Inequal. Appl. 2015, **2015**:219, 8 pages; Available online at <http://dx.doi.org/10.1186/s13660-015-0738-9>.
- [35] Wikipedia, *Euler number*, The Free Encyclopedia; Available online at https://en.wikipedia.org/wiki/Euler_number.

URL: <https://qifeng618.wordpress.com>

URL: <http://orcid.org/0000-0001-6725-533X>

URL: <http://orcid.org/0000-0001-6156-2590>



© 2017 by the authors. Licensee *Preprints*, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons by Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).