

IDENTITIES OF THE CHEBYSHEV POLYNOMIALS, THE INVERSE OF A TRIANGULAR MATRIX, AND IDENTITIES OF THE CATALAN NUMBERS

FENG QI AND BAI-NI GUO

ABSTRACT. In the paper, the authors establish two identities to express the generating function of the Chebyshev polynomials of the second kind and its higher order derivatives in terms of the generating function and its derivatives each other, deduce an explicit formula and an identities for the Chebyshev polynomials of the second kind, derive the inverse of an integer, unit, and lower triangular matrix, present several identities of the Catalan numbers, and give some remarks on the closely related results including connections of the Catalan numbers respectively with the Chebyshev polynomials, the central Delannoy numbers, and the Fibonacci polynomials.

1. PRELIMINARIES

It is common knowledge [8, 15, 52] that the generalized hypergeometric series

$${}_pF_q(a_1, \dots, a_p; b_1, \dots, b_q; z) = \sum_{n=0}^{\infty} \frac{(a_1)_n \cdots (a_p)_n}{(b_1)_n \cdots (b_q)_n} \frac{z^n}{n!}$$

is defined for complex numbers $a_i \in \mathbb{C}$ and $b_i \in \mathbb{C} \setminus \{0, -1, -2, \dots\}$, for positive integers $p, q \in \mathbb{N}$, and in terms of the rising factorials $(x)_n$ defined by

$$(x)_n = \prod_{\ell=0}^{n-1} (x + \ell) = \begin{cases} x(x+1) \cdots (x+n-1), & n \geq 1; \\ 1, & n = 0. \end{cases}$$

Specially, one calls ${}_2F_1(a, b; c; z)$ the classical hypergeometric function.

It is well known [12, 47, 55] that the Catalan numbers C_n for $n \geq 0$ form a sequence of natural numbers that occur in tree enumeration problems such as “In how many ways can a regular n -gon be divided into $n-2$ triangles if different orientations are counted separately? whose solution is the Catalan number C_{n-2} ”. The Catalan numbers C_n can be generated by

$$\frac{2}{1 + \sqrt{1-4x}} = \frac{1 - \sqrt{1-4x}}{2x} = \sum_{n=0}^{\infty} C_n x^n = 1 + x + 2x^2 + 5x^3 + \cdots$$

and explicitly expressed as

$$C_n = \frac{1}{n+1} \binom{2n}{n} = {}_2F_1(1-n, -n; 2; 1) = \frac{4^n \Gamma(n+1/2)}{\sqrt{\pi} \Gamma(n+2)},$$

2010 *Mathematics Subject Classification.* Primary 11B83; Secondary 05A15, 05A19, 11C08, 11C20, 11Y35, 15A09, 15B36, 33C05, 34A34.

Key words and phrases. identity; inverse matrix; explicit formula; generating function; Chebyshev polynomials of the second kind; Catalan number; triangular matrix; classical hypergeometric function; integral representation.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

where the classical Euler gamma function can be defined [8, 15, 24, 30, 52] by

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt, \quad \Re(z) > 0$$

or by

$$\Gamma(z) = \lim_{n \rightarrow \infty} \frac{n! n^z}{\prod_{k=0}^n (z+k)}, \quad z \in \mathbb{C} \setminus \{0, -1, -2, \dots\}.$$

For more information on the Catalan numbers C_k and their recent developments, please refer to the monographs [2, 12, 55], the papers [13, 25, 34, 40, 46, 47, 54, 59, 61, 62, 63] and the closely related references therein.

The first six Chebyshev polynomials of the second kind $U_k(x)$ for $0 \leq k \leq 5$ are

$$\begin{aligned} U_0(x) &= 1, & U_1(x) &= 2x, & U_2(x) &= 4x^2 - 1, & U_3(x) &= 8x^3 - 4x, \\ U_4(x) &= 16x^4 - 12x^2 + 1, & U_5(x) &= 32x^5 - 32x^3 + 6x. \end{aligned}$$

They can be generated by

$$F(t) = F(t, x) = \frac{1}{1 - 2xt + t^2} = \sum_{n=0}^{\infty} U_n(x) t^n$$

for $|x| < 1$ and $|t| < 1$. For more information on the Chebyshev polynomials of the second kind $U_k(x)$, please refer to [28, Section 7], the monographs [8, 15, 52] and the closely related references therein.

Let $\lfloor x \rfloor$ denote the floor function whose value is the largest integer less than or equal to x and let $\lceil x \rceil$ stand for the ceiling function which gives the smallest integer not less than x . When $n \in \mathbb{Z}$, it is easy to see that

$$\left\lfloor \frac{n}{2} \right\rfloor = \frac{1}{2} \left[n - \frac{1 - (-1)^n}{2} \right] \quad \text{and} \quad \left\lceil \frac{n}{2} \right\rceil = \frac{1}{2} \left[n + \frac{1 - (-1)^n}{2} \right].$$

In this paper, we will establish two identities to express the generating function $F(t)$ of the Chebyshev polynomials of the second kind $U_k(x)$ and its higher order derivatives $F^{(k)}(t)$ in terms of $F(t)$ and $F^{(k)}(t)$ each other, deduce an explicit formula and an identities for the Chebyshev polynomials of the second kind $U_k(x)$, derive the inverse of an integer, unit, and lower triangular matrix, present several identities of the Catalan numbers C_k , and give some remarks on the closely related results including connections of the Catalan numbers C_k respectively with the Chebyshev polynomials $U_k(x)$, the central Delannoy numbers, and the Fibonacci polynomials.

2. LEMMAS

In order to prove our main results, we recall several lemmas below.

Lemma 2.1 ([2, p. 134, Theorem A] and [2, p. 139, Theorem C]). *For $n \geq k \geq 0$, the Bell polynomials of the second kind, denoted by $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$, are defined by*

$$B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{1 \leq i \leq n, \ell_i \in \{0\} \cup \mathbb{N} \\ \sum_{i=1}^n i \ell_i = n \\ \sum_{i=1}^n \ell_i = k}} \frac{n!}{\prod_{i=1}^{n-k+1} \ell_i!} \prod_{i=1}^{n-k+1} \left(\frac{x_i}{i!} \right)^{\ell_i}.$$

The Faà di Bruno formula can be described in terms of the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ by

$$\frac{d^n}{dt^n} f \circ h(t) = \sum_{k=1}^n f^{(k)}(h(t)) B_{n,k}(h'(t), h''(t), \dots, h^{(n-k+1)}(t)), \quad n \in \mathbb{N}. \quad (2.1)$$

Lemma 2.2 ([2, p. 135]). For complex numbers a and b , we have

$$B_{n,k}(abx_1, ab^2x_2, \dots, ab^{n-k+1}x_{n-k+1}) = a^k b^n B_{n,k}(x_1, x_2, \dots, x_{n-k+1}). \quad (2.2)$$

Lemma 2.3 ([9, Theorem 4.1], [45, Eq. (2.8)], and [56, Lemma 2.5]). For $0 \leq k \leq n$, the Bell polynomials of the second kind $B_{n,k}$ satisfy

$$B_{n,k}(x, 1, 0, \dots, 0) = \frac{1}{2^{n-k}} \frac{n!}{k!} \binom{k}{n-k} x^{2k-n}, \quad (2.3)$$

where $\binom{p}{q} = 0$ for $q > p \geq 0$.

Lemma 2.4 ([7] and [12, pp. 112–114]). Let $T(r, 1) = 1$ and

$$T(r, c) = \sum_{i=c-1}^r T(i, c-1), \quad c \geq 2,$$

or, equivalently,

$$T(r, c) = \sum_{j=1}^c T(r-1, j), \quad r, c \in \mathbb{N}.$$

Then

$$T(r, c) = \frac{r-c+2}{r+1} \binom{r+c-1}{r}, \quad r, c \in \mathbb{N}$$

and $T(n, n) = C_n$ for $n \in \mathbb{N}$.

Lemma 2.5 ([16, p. 2, Eq. (10)] and [3]). For $n \in \mathbb{N}$, the Catalan numbers C_n have the integral representation

$$C_n = \frac{1}{2\pi} \int_0^4 \sqrt{\frac{4-x}{x}} x^n dx. \quad (2.4)$$

Lemma 2.6. For $t \neq 0$ and $j \in \mathbb{N}$, we have

$${}_2F_1\left(\frac{1-j}{2}, \frac{2-j}{2}; 1-j; \frac{1}{t^2}\right) = \frac{1}{2^j} \frac{t}{\sqrt{t^2-1}} \left[\left(1 + \frac{\sqrt{t^2-1}}{t}\right)^j - \left(1 - \frac{\sqrt{t^2-1}}{t}\right)^j \right]. \quad (2.5)$$

Proof. In [8, pp. 999–1000], it was listed that

$$G_n^\lambda(t) = \frac{1}{\sqrt{\pi}} \frac{\Gamma(2\lambda+n)}{n!\Gamma(2\lambda)} \frac{\Gamma(\frac{2\lambda+1}{2})}{\Gamma(\lambda)} \int_0^\pi \left(t + \sqrt{t^2-1} \cos \phi\right)^n \sin^{2\lambda-1} \phi d\phi \quad (2.6)$$

and

$$G_n^\lambda(t) = \frac{2^n \Gamma(\lambda+n)}{n! \Gamma(\lambda)} t^n {}_2F_1\left(-\frac{n}{2}, \frac{1-n}{2}; 1-\lambda-n; \frac{1}{t^2}\right), \quad (2.7)$$

where $G_n^\lambda(t)$ stands for the Gegenbauer polynomials which are the coefficients of α^n in the power-series expansion

$$\frac{1}{(1-2t\alpha+\alpha^2)^\lambda} = \sum_{k=0}^{\infty} G_k^\lambda(t) \alpha^k.$$

Taking $n = j - 1$ and $\lambda = 1$ in equalities (2.6) and (2.7), combining them, and simplifying give

$$\begin{aligned}
 {}_2F_1\left(\frac{1-j}{2}, \frac{2-j}{2}; 1-j; \frac{1}{t^2}\right) &= \frac{j}{2^j} \frac{1}{t^{j-1}} \int_0^\pi \left(t + \sqrt{t^2 - 1} \cos \phi\right)^{j-1} \sin \phi \, d\phi \\
 &= \frac{j}{2^j} \frac{(t^2 - 1)^{(j-1)/2}}{t^{j-1}} \int_0^\pi \left(\frac{t}{\sqrt{t^2 - 1}} + \cos \phi\right)^{j-1} \sin \phi \, d\phi \\
 &= \frac{j}{2^j} \frac{(t^2 - 1)^{(j-1)/2}}{t^{j-1}} \int_0^\pi \sum_{\ell=0}^{j-1} \binom{j-1}{\ell} \left(\frac{t}{\sqrt{t^2 - 1}}\right)^{j-1-\ell} \cos^\ell \phi \sin \phi \, d\phi \\
 &= \frac{j}{2^j} \frac{(t^2 - 1)^{(j-1)/2}}{t^{j-1}} \sum_{\ell=0}^{j-1} \binom{j-1}{\ell} \left(\frac{t}{\sqrt{t^2 - 1}}\right)^{j-1-\ell} \int_0^\pi \cos^\ell \phi \sin \phi \, d\phi \\
 &= \frac{j}{2^j} \frac{(t^2 - 1)^{(j-1)/2}}{t^{j-1}} \left(\frac{t}{\sqrt{t^2 - 1}}\right)^{j-1} \sum_{\ell=0}^{j-1} \binom{j-1}{\ell} \left(\frac{\sqrt{t^2 - 1}}{t}\right)^\ell \frac{(-1)^\ell + 1}{\ell + 1} \\
 &= \frac{j}{2^j} \sum_{\ell=0}^{j-1} \binom{j-1}{\ell} \left(\frac{\sqrt{t^2 - 1}}{t}\right)^\ell \frac{(-1)^\ell + 1}{\ell + 1} \\
 &= \frac{1}{2^j} \frac{t}{\sqrt{t^2 - 1}} \left[\left(1 + \frac{\sqrt{t^2 - 1}}{t}\right)^j - \left(1 - \frac{\sqrt{t^2 - 1}}{t}\right)^j \right].
 \end{aligned}$$

The formula (2.5) is thus proved. The proof of Lemma 2.5 is complete. \square

Lemma 2.7 ([8, p. 399]). *If $\Re(\nu) > 0$, then*

$$\int_0^{\pi/2} \cos^{\nu-1} x \cos(ax) \, dx = \frac{\pi}{2^\nu \nu B\left(\frac{\nu+a+1}{2}, \frac{\nu-a+1}{2}\right)}, \quad (2.8)$$

where $B(\alpha, \beta)$ stands for the classical beta function satisfying

$$B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha + \beta)} = B(\beta, \alpha), \quad \Re(\alpha), \Re(\beta) > 0.$$

3. IDENTITIES OF THE CHEBYSHEV POLYNOMIALS OF THE SECOND KIND

In this section, we establish three identities and an explicit formula for the Chebyshev polynomials of the second kind $U_k(x)$, their generating function $F(t)$, and higher order derivatives $F^{(k)}(t)$. Why do we start our research in this paper from here? Please read Remark 6.1 in Section 6 below.

Theorem 3.1. *For $n \in \mathbb{N}$, the n th derivatives of the generating function $F(t)$ of the Chebyshev polynomials of the second kind $U_k(x)$ satisfy*

$$F^{(n)}(t) = \frac{n!}{[2(t-x)]^n} \sum_{k=\lceil n/2 \rceil}^n (-1)^k \binom{k}{n-k} [2(t-x)]^{2k} F^{k+1}(t) \quad (3.1)$$

and

$$F^{n+1}(t) = \frac{1}{n} \frac{1}{[2(t-x)]^{2n}} \sum_{k=1}^n \frac{(-1)^k}{(k-1)!} \binom{2n-k-1}{n-1} [2(t-x)]^k F^{(k)}(t). \quad (3.2)$$

Consequently, the Chebyshev polynomials of the second kind $U_n(x)$ satisfy

$$U_n(x) = \frac{(-1)^n}{(2x)^n} \sum_{k=\lceil n/2 \rceil}^n (-1)^k \binom{k}{n-k} (2x)^{2k} \quad (3.3)$$

and

$$\sum_{k=1}^n k \binom{2n-k-1}{n-1} (2x)^k U_k(x) = n(2x)^{2n}. \quad (3.4)$$

Proof. By the formulas (2.1), (2.2), and (2.3) in sequence, we have

$$\begin{aligned} F^{(n)}(t) &= \frac{d^n}{dt^n} \left(\frac{1}{1-2tx+t^2} \right) \\ &= \sum_{k=1}^n \left(\frac{1}{u} \right)^{(k)} B_{n,k}(-2x+2t, 2, 0, \dots, 0) \\ &= \sum_{k=1}^n \frac{(-1)^k k!}{u^{k+1}} 2^k B_{n,k}(t-x, 1, 0, \dots, 0) \\ &= \sum_{k=1}^n \frac{(-1)^k k!}{u^{k+1}} 2^k \frac{1}{2^{n-k}} \frac{n!}{k!} \binom{k}{n-k} (t-x)^{2k-n} \\ &= (-1)^n n! \sum_{k=1}^n (-1)^k 2^{2k-n} \binom{k}{n-k} \frac{(x-t)^{2k-n}}{(1-2tx+t^2)^{k+1}} \\ &= (-1)^n n! \sum_{k=1}^n (-1)^k 2^{2k-n} \binom{k}{n-k} (x-t)^{2k-n} F^{k+1}(t) \end{aligned}$$

for $n \in \mathbb{N}$, where $u = u(t, x) = 1 - 2tx + t^2$. This can be rewritten as the formula (3.1).

We can reformulate the formula (3.1) as

$$\begin{pmatrix} \frac{[2(t-x)]^1}{1!} F'(t) \\ \frac{[2(t-x)]^2}{2!} F''(t) \\ \frac{[2(t-x)]^3}{3!} F^{(3)}(t) \\ \vdots \\ \frac{[2(t-x)]^{n-2}}{(n-2)!} F^{(n-2)}(t) \\ \frac{[2(t-x)]^{n-1}}{(n-1)!} F^{(n-1)}(t) \\ \frac{[2(t-x)]^n}{n!} F^{(n)}(t) \end{pmatrix} = A_n \begin{pmatrix} (-1)^1 [2(x-t)]^2 F^2(t) \\ (-1)^2 [2(x-t)]^4 F^3(t) \\ (-1)^3 [2(x-t)]^6 F^4(t) \\ \vdots \\ (-1)^{n-2} [2(x-t)]^{2(n-2)} F^{n-1}(t) \\ (-1)^{n-1} [2(x-t)]^{2(n-1)} F^n(t) \\ (-1)^n [2(x-t)]^{2n} F^{n+1}(t) \end{pmatrix}$$

for $n \in \mathbb{N}$, where $A_n = (a_{i,j})_{n \times n}$ with

$$a_{i,j} = \begin{cases} 0, & i < j \\ \binom{j}{i-j}, & j \leq i \leq 2j \\ 0, & i > 2j \end{cases}$$

for $i, j \in \mathbb{N}$. This means that

$$\begin{pmatrix} (-1)^1 [2(x-t)]^2 F^2(t) \\ (-1)^2 [2(x-t)]^4 F^3(t) \\ (-1)^3 [2(x-t)]^6 F^4(t) \\ \vdots \\ (-1)^{n-2} [2(x-t)]^{2(n-2)} F^{n-1}(t) \\ (-1)^{n-1} [2(x-t)]^{2(n-1)} F^n(t) \\ (-1)^n [2(x-t)]^{2n} F^{n+1}(t) \end{pmatrix} = A_n^{-1} \begin{pmatrix} \frac{[2(t-x)]^1}{1!} F'(t) \\ \frac{[2(t-x)]^2}{2!} F''(t) \\ \frac{[2(t-x)]^3}{3!} F^{(3)}(t) \\ \vdots \\ \frac{[2(t-x)]^{n-2}}{(n-2)!} F^{(n-2)}(t) \\ \frac{[2(t-x)]^{n-1}}{(n-1)!} F^{(n-1)}(t) \\ \frac{[2(t-x)]^n}{n!} F^{(n)}(t) \end{pmatrix} \quad (3.5)$$

for $n \in \mathbb{N}$, where $A_n^{-1} = (b_{i,j})_{n \times n}$ denotes the inverse matrix of A_n .

By the software MATHEMATICA, we can obtain immediately that

$$A_7^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 2 & 1 & 0 & 0 & 0 \\ 0 & 1 & 3 & 1 & 0 & 0 \\ 0 & 0 & 3 & 4 & 1 & 0 \\ 0 & 0 & 1 & 6 & 5 & 1 \end{pmatrix}^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 & 0 & 0 \\ 2 & -2 & 1 & 0 & 0 & 0 \\ -5 & 5 & -3 & 1 & 0 & 0 \\ 14 & -14 & 9 & -4 & 1 & 0 \\ -42 & 42 & -28 & 14 & -5 & 1 \end{pmatrix}. \quad (3.6)$$

The first few values of the sequence $T(r, c)$ can be listed as Table 1, where $T(r, c)$ denote the r th element in column c for $r, c \geq 1$, see [12, p. 113]. Comparing Table 1 and the inverse matrix (3.6),

TABLE 1. Definition of $T(r, c)$

	1	2	3	4	5
1	1				
2	1	2			
3	1	3	5		
4	1	4	9	14	
5	1	5	14	28	42

we should infer that

$$T(k+m, k) = (-1)^{k+1} b_{k+m+1, m+2}, \quad k \geq 1, \quad m \geq 0.$$

Hence, by Lemma 2.4, we should obtain

$$b_{p,q} = (-1)^{p-q} T(p-1, p-q+1) = (-1)^{p-q} \frac{q}{p} \binom{2p-q-1}{p-1}, \quad p \geq q \geq 2.$$

It is easy to see that the formula

$$b_{p,q} = (-1)^{p-q} \frac{q}{p} \binom{2p-q-1}{p-1}$$

should be valid for all $p \geq q \geq 1$. This should imply that

$$(-1)^n [2(x-t)]^{2n} F^{n+1}(t) = \sum_{k=1}^n b_{n,k} \frac{[2(t-x)]^k}{k!} F^{(k)}(t), \quad n \in \mathbb{N}. \quad (3.7)$$

We now start out to inductively verify the equation (3.7). When $n = 1, 2$, the equation (3.7) are

$$-[2(x-t)]^2 F^2(t) = b_{1,1} \frac{2(t-x)}{1!} F'(t) = b_{1,1} \frac{2(t-x)}{1!} \frac{2x-2t}{(1-2tx+t^2)^2}$$

and

$$\begin{aligned} [2(x-t)]^4 F^3(t) &= \sum_{k=1}^2 b_{2,k} \frac{[2(t-x)]^k}{k!} F^{(k)}(t) \\ &= b_{2,1} \frac{2(t-x)}{1!} F'(t) + b_{2,2} \frac{[2(t-x)]^2}{2!} F''(t) \\ &= b_{2,1} \frac{2(t-x)}{1!} \frac{2x-2t}{(1-2tx+t^2)^2} + b_{2,2} \frac{[2(t-x)]^2}{2!} \frac{2(3t^2-6tx+4x^2-1)}{(t^2-2tx+1)^3} \end{aligned}$$

which are clearly valid. When $n \geq 3$, we rewrite (3.7) as

$$(-1)^n F^{n+1}(t) = \sum_{k=1}^n b_{n,k} \frac{[2(t-x)]^{k-2n}}{k!} F^{(k)}(t). \quad (3.8)$$

Differentiating with respect to t on both sides of (3.8) yields

$$\begin{aligned} (-1)^n (n+1) F^n(t) F'(t) &= \sum_{k=1}^n \frac{b_{n,k}}{k!} \{2(k-2n)[2(t-x)]^{k-2n-1} F^{(k)}(t) + [2(t-x)]^{k-2n} F^{(k+1)}(t)\} \\ &= \sum_{k=1}^n \frac{b_{n,k}}{k!} 2(k-2n)[2(t-x)]^{k-2n-1} F^{(k)}(t) + \sum_{k=1}^n \frac{b_{n,k}}{k!} [2(t-x)]^{k-2n} F^{(k+1)}(t) \\ &= \sum_{k=1}^n \frac{2(k-2n)b_{n,k}}{k!} [2(t-x)]^{k-2n-1} F^{(k)}(t) + \sum_{k=2}^{n+1} \frac{b_{n,k-1}}{(k-1)!} [2(t-x)]^{k-1-2n} F^{(k)}(t) \\ &= \frac{b_{n,1}}{1!} \frac{2(1-2n)}{[2(t-x)]^{2n}} F'(t) + \frac{b_{n,n}}{n!} \frac{1}{[2(t-x)]^n} F^{(n+1)}(t) \\ &\quad + \sum_{k=2}^n \left[\frac{b_{n,k}}{k!} 2(k-2n) + \frac{b_{n,k-1}}{(k-1)!} \right] [2(t-x)]^{k-2n-1} F^{(k)}(t) \end{aligned}$$

which can be rearranged as

$$\begin{aligned} (-1)^{n+1} F^{n+2}(t) &= \frac{2(1-2n)b_{n,1}}{n+1} \frac{[2(t-x)]^{1-2(n+1)}}{1!} F'(t) \\ &\quad + b_{n,n} \frac{[2(t-x)]^{(n+1)-2(n+1)}}{(n+1)!} F^{(n+1)}(t) \\ &\quad + \sum_{k=2}^n \frac{2(k-2n)b_{n,k} + kb_{n,k-1}}{n+1} \frac{[2(t-x)]^{k-2(n+1)}}{k!} F^{(k)}(t). \end{aligned}$$

It is easy to see that

$$\frac{2(1-2n)b_{n,1}}{n+1} = \frac{2(1-2n)}{n+1} (-1)^{n-1} \frac{1}{n} \binom{2n-2}{n-1} = (-1)^n \frac{1}{n+1} \binom{2n}{n} = b_{n+1,1}.$$

Since $b_{k,k} = 1$ for all $1 \leq k \leq n \in \mathbb{N}$, it is sufficient to show

$$\frac{2(k-2n)b_{n,k} + kb_{n,k-1}}{n+1} = b_{n+1,k} \quad (3.9)$$

for $2 \leq k \leq n$. This is equivalent to

$$\begin{aligned} \frac{2(k-2n)}{n+1}(-1)^{n-k} \frac{k}{n} \binom{2n-k-1}{n-1} + \frac{k}{n+1}(-1)^{n-k+1} \frac{k-1}{n} \binom{2n-k}{n-1} \\ = (-1)^{n+1-k} \frac{k}{n+1} \binom{2n-k+1}{n} \end{aligned}$$

which can be verified straightforwardly. The equation (3.7), which can be reformulated as (3.2) for $n \in \mathbb{N}$, is thus proved.

The formulas (3.3) and (3.4) follow readily from taking $t \rightarrow 0$ on both sides of (3.1) and (3.2) respectively. The proof of Theorem 3.1 is complete. \square

4. INVERSE OF A TRIANGULAR MATRIX

In this section, basing on equations (3.1) and (3.2), we derive the inverse of an integer, unit, and lower triangular matrix.

Theorem 4.1. For $n \in \mathbb{N}$, let

$$A_n = (a_{i,j})_{n \times n} = \begin{pmatrix} \binom{1}{0} & 0 & 0 & 0 & \cdots & 0 & 0 & 0 & 0 \\ \binom{1}{1} & \binom{2}{0} & 0 & 0 & \cdots & 0 & 0 & 0 & 0 \\ 0 & \binom{2}{1} & \binom{3}{0} & 0 & \cdots & 0 & 0 & 0 & 0 \\ 0 & \binom{2}{2} & \binom{3}{1} & \binom{4}{0} & \cdots & 0 & 0 & 0 & 0 \\ 0 & 0 & \binom{3}{2} & \binom{4}{1} & \cdots & 0 & 0 & 0 & 0 \\ 0 & 0 & \binom{3}{3} & \binom{4}{2} & \cdots & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \binom{4}{3} & \cdots & 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & \binom{n-3}{0} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \cdots & \binom{n-3}{1} & \binom{n-2}{0} & 0 & 0 \\ 0 & 0 & 0 & 0 & \cdots & \binom{n-3}{2} & \binom{n-2}{1} & \binom{n-1}{0} & 0 \\ 0 & 0 & 0 & 0 & \cdots & \binom{n-3}{3} & \binom{n-2}{2} & \binom{n-1}{1} & \binom{n}{0} \end{pmatrix},$$

where

$$a_{i,j} = \begin{cases} 0, & i < j \\ \binom{j}{i-j}, & j \leq i \leq 2j \\ 0, & i > 2j \end{cases}$$

for $1 \leq i, j \leq n$. Then

$$A_n^{-1} = (b_{i,j})_{n \times n} = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ -1 & 1 & 0 & \cdots & 0 & 0 & 0 \\ 2 & -2 & 1 & \cdots & 0 & 0 & 0 \\ -5 & 5 & -3 & \cdots & 0 & 0 & 0 \\ 14 & -14 & 9 & \cdots & 0 & 0 & 0 \\ -42 & 42 & -28 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ \frac{(-1)^{n-1}}{n-2} \binom{2n-6}{n-3} & \frac{(-1)^{n-2}}{n-2} \binom{2n-7}{n-3} & \frac{(-1)^{n-3}}{n-2} \binom{2n-8}{n-3} & \cdots & 1 & 0 & 0 \\ \frac{(-1)^n}{n-1} \binom{2n-4}{n-2} & \frac{(-1)^{n-1}}{n-1} \binom{2n-5}{n-2} & \frac{(-1)^{n-2}}{n-1} \binom{2n-6}{n-2} & \cdots & -(n-2) & 1 & 0 \\ \frac{(-1)^{n-1}}{n} \binom{2n-2}{n-1} & \frac{(-1)^{n-2}}{n} \binom{2n-3}{n-1} & \frac{(-1)^{n-3}}{n} \binom{2n-4}{n-1} & \cdots & \frac{n-2}{n} \binom{n+1}{n-1} & -(n-1) & 1 \end{pmatrix},$$

where

$$b_{i,j} = \begin{cases} 0, & 1 \leq i < j \leq n; \\ (-1)^{i-j} \frac{j}{i} \binom{2i-j-1}{i-1}, & n \geq i > j \geq 1. \end{cases} \quad (4.1)$$

Proof. This follows straightforwardly from combining (3.5) with (3.2). The proof of Theorem 4.1 is complete. \square

5. IDENTITIES OF THE CATALAN NUMBERS

In this section, we present several identities of the Catalan numbers C_k .

Theorem 5.1. For $i \geq j \geq 1$, we have

$$\sum_{\ell=0}^{\lfloor (j-1)/2 \rfloor} (-1)^\ell \binom{j-\ell-1}{\ell} C_{i-\ell-1} = \frac{j}{i} \binom{2i-j-1}{i-1}. \quad (5.1)$$

Proof. Observing the special result (3.6), we guess that the elements $b_{i,j}$ of the inverse of the triangular matrix A_n should satisfy the following relations:

- (1) for $i < j$, the elements in the upper triangle are $b_{i,j} = 0$;
- (2) for all $i \in \mathbb{N}$, the elements on the main diagonal are $b_{i,i} = 1$;
- (3) the elements in the first two columns satisfy $b_{i,1} = -b_{i,2}$ for $i \geq 2$;
- (4) the elements in the first column are $b_{i,1} = (-1)^{i-1} C_{i-1}$;
- (5) for $1 \leq i \leq n-1$ and $1 \leq j \leq n-2$,

$$b_{i+1,j+2} = b_{i,j} - b_{i+1,j+1};$$

- (6) for $i \geq j \geq 2$,

$$b_{i,j} = \sum_{k=-1}^{i-j-1} (-1)^{k+1} b_{i-1,j+k}.$$

Basing on these observations, we guess out that the elements $b_{i,j}$ should alternatively satisfy

$$b_{i,j} = (-1)^{i-j} \sum_{\ell=0}^{\lfloor (j-1)/2 \rfloor} (-1)^\ell \binom{j-\ell-1}{\ell} C_{i-\ell-1}, \quad i \geq j \geq 1. \quad (5.2)$$

Combining this with (4.1) and simplifying yield the identity (5.1).

We now start off to verify the identity (5.1). By virtue of the integral representation (2.4), the formula (2.5) in Lemma 2.6, and the integral (2.8) in Lemma 2.7, we acquire

$$\begin{aligned}
& \sum_{\ell=0}^{\lfloor (j-1)/2 \rfloor} (-1)^\ell \binom{j-\ell-1}{\ell} C_{i-\ell-1} \\
&= \frac{1}{2\pi} \int_0^4 \sqrt{\frac{4-x}{x}} \left[\sum_{\ell=0}^{\lfloor (j-1)/2 \rfloor} (-1)^\ell \binom{j-\ell-1}{\ell} x^{i-\ell-1} \right] dx \\
&= \frac{1}{2\pi} \int_0^4 x^{i-3/2} (4-x)^{1/2} \left[\sum_{\ell=0}^{\lfloor (j-1)/2 \rfloor} \frac{(j-1-\ell)!}{(j-1-2\ell)! \ell!} \left(-\frac{1}{x}\right)^\ell \right] dx \\
&= \frac{1}{2\pi} \int_0^4 x^{i-3/2} (4-x)^{1/2} \left[\sum_{\ell=0}^{\lfloor (j-1)/2 \rfloor} \frac{\left(\frac{1-j}{2}\right)_\ell \left(\frac{2-j}{2}\right)_\ell}{(1-j)_\ell} \frac{1}{\ell!} \left(\frac{4}{x}\right)^\ell \right] dx \\
&= \frac{1}{2\pi} \int_0^4 x^{i-3/2} (4-x)^{1/2} \left[\sum_{\ell=0}^{\infty} \frac{\left(\frac{1-j}{2}\right)_\ell \left(\frac{2-j}{2}\right)_\ell}{(1-j)_\ell} \frac{1}{\ell!} \left(\frac{4}{x}\right)^\ell \right] dx \\
&= \frac{1}{2\pi} \int_0^4 x^{i-3/2} (4-x)^{1/2} {}_2F_1\left(\frac{1-j}{2}, \frac{2-j}{2}; 1-j; \frac{4}{x}\right) dx \\
&= \frac{4^i}{2\pi} \int_0^1 t^{i-3/2} (1-t)^{1/2} {}_2F_1\left(\frac{1-j}{2}, \frac{2-j}{2}; 1-j; \frac{1}{t}\right) dt \\
&= \frac{4^i}{2\pi} \int_0^1 t^{i-3/2} (1-t)^{1/2} \frac{1}{2^j} \frac{\sqrt{t}}{\sqrt{t-1}} \left[\left(1 + \frac{\sqrt{t-1}}{\sqrt{t}}\right)^j - \left(1 - \frac{\sqrt{t-1}}{\sqrt{t}}\right)^j \right] dt \\
&= \frac{2^{2i-j}}{2\pi} i \int_0^1 t^{i-1} \left[\left(1 + \sqrt{1-\frac{1}{t}}\right)^j - \left(1 - \sqrt{1-\frac{1}{t}}\right)^j \right] dt \quad (i = \sqrt{-1}) \\
&= \frac{2^{2i-j}}{\pi} i \int_0^\infty \frac{s}{(1+s^2)^{i+1}} [(1-is)^j - (1+is)^j] ds \\
&= \frac{2^{2i-j}}{\pi} i \int_0^\infty \frac{s}{(1+s^2)^{i+1}} \left[\left(\sqrt{1+s^2} e^{-i \arctan s}\right)^j - \left(\sqrt{1+s^2} e^{i \arctan s}\right)^j \right] ds \\
&= \frac{2^{2i-j}}{\pi} i \int_0^\infty \frac{s}{(1+s^2)^{i-j/2+1}} (e^{-ij \arctan s} - e^{ij \arctan s}) ds \\
&= \frac{2^{2i-j}}{\pi} \int_0^\infty \frac{s}{(1+s^2)^{i-j/2+1}} \sin(j \arctan s) ds \\
&= \frac{2^{2i-j}}{\pi} \int_0^{\pi/2} \frac{\tan t}{(1+\tan^2 t)^{i-j/2+1}} \sin(jt) \sec^2 t dt \\
&= \frac{2^{2i-j}}{\pi} \int_0^{\pi/2} \frac{\tan t}{\sec^{2i-j} t} \sin(jt) dt \\
&= \frac{2^{2i-j}}{\pi} \int_0^{\pi/2} \sin t \cos^{2i-j-1} t \sin(jt) dt
\end{aligned}$$

$$\begin{aligned}
&= \frac{2^{2i-j}}{\pi} \int_0^{\pi/2} [\cos((j-1)t) - \cos((j+1)t)] \cos^{2i-j-1} t \, dt \\
&= \frac{2^{2i-j}}{\pi} \left[\frac{\pi}{2^{2i-j}(2i-j)B(i, i-j+1)} - \frac{\pi}{2^{2i-j}(2i-j)B(i+1, i-j)} \right] \\
&= \frac{1}{2i-j} \left[\frac{1}{B(i, i-j+1)} - \frac{1}{B(i+1, i-j)} \right] \\
&= \frac{1}{2i-j} \left[\frac{\Gamma(2i-j+1)}{\Gamma(i)\Gamma(i-j+1)} - \frac{\Gamma(2i-j+1)}{\Gamma(i+1)\Gamma(i-j)} \right] \\
&= (2i-j-1)! \left[\frac{1}{\Gamma(i)\Gamma(i-j+1)} - \frac{1}{\Gamma(i+1)\Gamma(i-j)} \right] \\
&= (2i-j-1)! \left[\frac{1}{(i-1)!(i-j)!} - \frac{1}{i!(i-j-1)!} \right] \\
&= \frac{j}{i} \binom{2i-j-1}{i-1}.
\end{aligned}$$

The identity (5.1) is thus proved. The proof of Theorem 5.1 is complete. \square

Theorem 5.2. For $i, j, n \in \mathbb{N}$, the Catalan numbers C_n satisfy

$$\sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \binom{n-k}{k} C_{n-k} = 1, \quad (5.3)$$

$$\sum_{\substack{i \leq 2\ell \leq 2i \\ \ell \geq j}} \sum_{k=0}^{\lfloor (j-1)/2 \rfloor} (-1)^{\ell-k} \binom{\ell}{i-\ell} \binom{j-k-1}{k} C_{\ell-k-1} = 0, \quad (5.4)$$

and

$$\sum_{\substack{i \geq \ell \geq j \\ \ell \leq 2j}} \sum_{k=0}^{\lfloor (\ell-1)/2 \rfloor} (-1)^{\ell-k} \binom{j}{\ell-j} \binom{\ell-k-1}{k} C_{i-k-1} = 0. \quad (5.5)$$

Proof. This follows from expanding the matrix equation

$$A_n A_n^{-1} = A_n^{-1} A_n = I_n \quad (5.6)$$

and utilizing the expression (5.2) in Theorem 4.1, where I_n stands for the identity matrix of n orders. This can be written in details as follows.

The matrix equation (5.6) is equivalent to

$$\sum_{\ell=1}^n a_{i,\ell} b_{\ell,j} = \begin{cases} 0, & i < j \\ \sum_{\ell=j}^i a_{i,\ell} b_{\ell,j}, & i \geq j \end{cases} = \begin{cases} 0, & i \neq j \\ 1, & i = j \end{cases}$$

and

$$\sum_{\ell=1}^n b_{i,\ell} a_{\ell,j} = \begin{cases} 0, & i < j \\ \sum_{\ell=j}^i b_{i,\ell} a_{\ell,j}, & i \geq j \end{cases} = \begin{cases} 0, & i \neq j \\ 1, & i = j \end{cases}$$

which can be rearranged as

$$\sum_{\ell=j}^i a_{i,\ell} b_{\ell,j} = \begin{cases} 0, & i > j \\ 1, & i = j \end{cases} \quad \text{or} \quad \sum_{\ell=j}^i b_{i,\ell} a_{\ell,j} = \begin{cases} 0, & i > j \\ 1, & i = j \end{cases}$$

for $1 \leq i, j \leq n$.

When $1 \leq i = j \leq n$, it follows that

$$\begin{aligned} 1 &= \sum_{\ell=j}^i a_{i,\ell} b_{\ell,j} = \sum_{\ell=j}^i b_{i,\ell} a_{\ell,j} = a_{i,i} b_{i,i} = b_{i,i} \\ &= \sum_{k=0}^{i-1} (-1)^k \binom{i-k-1}{k} C_{i-k-1} = \sum_{k=0}^{\lfloor (i-1)/2 \rfloor} (-1)^k \binom{i-k-1}{k} C_{i-k-1}. \end{aligned}$$

The identity (5.3) is thus concluded.

When $1 \leq j < i \leq n$, it follows that

$$\begin{aligned} 0 &= \sum_{\ell=j}^i a_{i,\ell} b_{\ell,j} = \sum_{\substack{i/2 \leq \ell \leq i \\ \ell \geq j}} a_{i,\ell} b_{\ell,j} \\ &= \sum_{\substack{i/2 \leq \ell \leq i \\ \ell \geq j}} \binom{\ell}{i-\ell} (-1)^{\ell-j} \sum_{k=0}^{\lfloor (j-1)/2 \rfloor} (-1)^k \binom{j-k-1}{k} C_{\ell-k-1} \\ &= (-1)^j \sum_{\substack{i/2 \leq \ell \leq i \\ \ell \geq j}} \sum_{k=0}^{\lfloor (j-1)/2 \rfloor} (-1)^{\ell-k} \binom{\ell}{i-\ell} \binom{j-k-1}{k} C_{\ell-k-1} \end{aligned}$$

and

$$\begin{aligned} 0 &= \sum_{\ell=j}^i b_{i,\ell} a_{\ell,j} = \sum_{\substack{i \geq \ell \geq j \\ \ell \leq 2j}} b_{i,\ell} a_{\ell,j} \\ &= \sum_{\substack{i \geq \ell \geq j \\ \ell \leq 2j}} (-1)^{i-\ell} \sum_{k=0}^{\lfloor (\ell-1)/2 \rfloor} (-1)^k \binom{\ell-k-1}{k} C_{i-k-1} \binom{j}{\ell-j} \\ &= (-1)^i \sum_{\substack{i \geq \ell \geq j \\ \ell \leq 2j}} \sum_{k=0}^{\lfloor (\ell-1)/2 \rfloor} (-1)^{\ell-k} \binom{j}{\ell-j} \binom{\ell-k-1}{k} C_{i-k-1}. \end{aligned}$$

The identities (5.4) and (5.5) are thus derived. The proof of Theorem 5.2 is complete. \square

Theorem 5.3. Let $m, n \in \mathbb{N}$. If $n \geq 2m \geq 2$, then

$$\frac{\sum_{\ell=0}^{m-1} (-1)^\ell \binom{2m-\ell-1}{\ell} \frac{n+2\ell+1}{n-\ell+1} C_{n-\ell-1}}{\sum_{\ell=0}^{m-1} (-1)^\ell \binom{2m-\ell-2}{\ell} \frac{1}{2m-2\ell-1} C_{n-\ell-1}} = m(2m-1). \quad (5.7)$$

Proof. Employing the expression (5.2) and making use of Theorem 5.1, we can write the recursive equation (3.9) as

$$\begin{aligned}
 & 2(k-2n)(-1)^{n-k} \sum_{\ell=0}^{\lfloor (k-1)/2 \rfloor} (-1)^\ell \binom{k-\ell-1}{\ell} C_{n-\ell-1} \\
 & + k(-1)^{n-k+1} \sum_{\ell=0}^{\lfloor (k-2)/2 \rfloor} (-1)^\ell \binom{k-\ell-2}{\ell} C_{n-\ell-1} \\
 & = (-1)^{n-k+1} \left\{ k \sum_{\ell=0}^{\lfloor (k-2)/2 \rfloor} (-1)^\ell \binom{k-\ell-2}{\ell} C_{n-\ell-1} \right. \\
 & \quad \left. - 2(k-2n) \sum_{\ell=0}^{\lfloor (k-1)/2 \rfloor} (-1)^\ell \binom{k-\ell-1}{\ell} C_{n-\ell-1} \right\} \\
 & = (-1)^{n-k+1} (n+1) \sum_{\ell=0}^{\lfloor (k-1)/2 \rfloor} (-1)^\ell \binom{k-\ell-1}{\ell} C_{n-\ell}
 \end{aligned}$$

for $n \geq 2$, that is,

$$\begin{aligned}
 k \sum_{\ell=0}^{\lfloor (k-2)/2 \rfloor} (-1)^\ell \binom{k-\ell-2}{\ell} C_{n-\ell-1} - 2(k-2n) \sum_{\ell=0}^{\lfloor (k-1)/2 \rfloor} (-1)^\ell \binom{k-\ell-1}{\ell} C_{n-\ell-1} \\
 = (n+1) \sum_{\ell=0}^{\lfloor (k-1)/2 \rfloor} (-1)^\ell \binom{k-\ell-1}{\ell} C_{n-\ell}, \quad n \geq 2. \quad (5.8)
 \end{aligned}$$

When $k = 2m$ and $m \in \mathbb{N}$, the equation (5.8) is equivalent to

$$\begin{aligned}
 & 2m \sum_{\ell=0}^{m-1} (-1)^\ell \binom{2m-\ell-2}{\ell} C_{n-\ell-1} - 4(m-n) \sum_{\ell=0}^{m-1} (-1)^\ell \binom{2m-\ell-1}{\ell} C_{n-\ell-1} \\
 & = (n+1) \sum_{\ell=0}^{m-1} (-1)^\ell \binom{2m-\ell-1}{\ell} C_{n-\ell}, \\
 & 2m \sum_{\ell=0}^{m-1} (-1)^\ell \binom{2m-\ell-2}{\ell} C_{n-\ell-1} - 4m \sum_{\ell=0}^{m-1} (-1)^\ell \binom{2m-\ell-1}{\ell} C_{n-\ell-1} \\
 & = (n+1) \sum_{\ell=0}^{m-1} (-1)^\ell \binom{2m-\ell-1}{\ell} C_{n-\ell} - 4n \sum_{\ell=0}^{m-1} (-1)^\ell \binom{2m-\ell-1}{\ell} C_{n-\ell-1}, \\
 & 2m \sum_{\ell=0}^{m-1} (-1)^\ell \left[\binom{2m-\ell-2}{\ell} - 2 \binom{2m-\ell-1}{\ell} \right] C_{n-\ell-1} \\
 & = \sum_{\ell=0}^{m-1} (-1)^\ell \binom{2m-\ell-1}{\ell} [(n+1)C_{n-\ell} - 4nC_{n-\ell-1}],
 \end{aligned}$$

and

$$m(2m-1) \sum_{\ell=0}^{m-1} (-1)^\ell \frac{(2m-\ell-2)!}{\ell!(2m-2\ell-1)!} C_{n-\ell-1} = \sum_{\ell=0}^{m-1} (-1)^\ell \binom{2m-\ell-1}{\ell} \frac{n+2\ell+1}{n-\ell+1} C_{n-\ell-1}$$

which can be rearranged as

$$\sum_{\ell=0}^{m-1} (-1)^\ell \left[m(2m-1) - \frac{(2m-\ell-1)(n+2\ell+1)}{n-\ell+1} \right] \frac{(2m-\ell-2)!}{\ell!(2m-2\ell-1)!} C_{n-\ell-1} = 0$$

for $n \geq 2m \geq 2$. This can be further rewritten as (5.7). The proof of Theorem 5.3 is complete. \square

6. REMARKS

Finally, we give some remarks on the closely related results stated in previous sections.

Remark 6.1. Now we explain the motivation of Theorem 3.1 as follows. In [11], the following results were inductively and recursively obtained.

(1) The nonlinear differential equations

$$2^n n! F^{n+1}(t) = \sum_{i=1}^n a_i(n) (x-t)^{i-2n} F^{(i)}(t), \quad n \in \mathbb{N}$$

has a solution

$$F(t) = F(t, x) = \frac{1}{1-2tx+t^2},$$

where $a_1(n) = (2n-3)!!$ and

$$\begin{aligned} a_i(n) = & \sum_{k_{i-1}=0}^{n-i} \sum_{k_{i-2}=0}^{n-i-k_{i-1}} \cdots \sum_{k_1=0}^{n-i-k_{i-1}-\cdots-k_2} 2^{\sum_{j=1}^{i-1} k_j} \\ & \times \prod_{j=2}^i \left\langle n - \sum_{\ell=j}^{i-1} k_\ell - \frac{2i+2-j}{2} \right\rangle_{k_{j-1}} \left(2 \left(n-i - \sum_{j=1}^{i-1} k_j \right) - 1 \right)!! \end{aligned} \quad (6.1)$$

for $2 \leq i \leq n$, with the notation that

$$\langle x \rangle_n = \prod_{k=0}^{n-1} (x-k) = \begin{cases} x(x-1) \cdots (x-n+1), & n \geq 1 \\ 1, & n = 0 \end{cases}$$

is the falling factorial and that the double factorial of negative odd integers $-2n-1$ is defined by

$$(-2n-1)!! = \frac{(-1)^n}{(2n-1)!!} = (-1)^n \frac{2^n n!}{(2n)!}$$

for $n \geq 0$. See [11, Theorem 1].

(2) The higher order Chebyshev polynomials of the second kind $U_n^{(\alpha)}(x)$ generated by

$$\left(\frac{1}{1-2xt+t^2} \right)^\alpha = \sum_{n=0}^{\infty} U_n^{(\alpha)}(x) t^n$$

satisfy

$$U_n^{(k+1)}(x) = \frac{1}{2^k k!} \sum_{i=1}^k a_i(k) \sum_{\ell=0}^n \binom{2k+n-\ell-i-1}{n-\ell} U_{\ell+i}(x) x^{i+\ell-2k-n} \langle \ell+i \rangle_i$$

for $k \in \mathbb{N}$, where $U_n^{(1)}(x) = U_n(x)$. See [11, Theorem 2].

(3) The higher order Legendre polynomials $p_n^{(\alpha)}(x)$ generated by

$$\left(\frac{1}{\sqrt{1-2xt+t^2}} \right)^\alpha = \sum_{n=0}^{\infty} p_n^{(\alpha)}(x) t^n$$

satisfy

$$\sum_{\ell=0}^n p_\ell^{(k+1)}(x) p_{n-\ell}^{(k+1)}(x) = \frac{1}{2^k k!} \sum_{i=1}^k a_i(k) \sum_{\ell=0}^n \binom{2k+n-\ell-i-1}{n-\ell} U_{\ell+i}(x) \langle \ell+i \rangle_i x^{i+\ell-2k-n}$$

for $k \in \mathbb{N}$ and $n \geq 0$ and

$$U_n^{(k+1)}(x) = \frac{1}{2^k k!} \sum_{i=1}^k a_i(k) \sum_{\ell=0}^n \sum_{j=0}^{\ell+i} \binom{2k+n-\ell-i-1}{n-\ell} x^{i+\ell-2k-n} \langle \ell+i \rangle_i p_{\ell+i-j}(x)$$

for $k, n \in \mathbb{N}$, where $p_n^{(1)}(x) = p_n(x)$. See [11, Corollaries 3 and 4].

(4) The higher order Chebyshev polynomials of the third kind $V_n^{(\alpha)}(x)$ generated by

$$\left(\frac{1-t}{\sqrt{1-2xt+t^2}} \right)^\alpha = \sum_{n=0}^{\infty} V_n^{(\alpha)}(x) t^n$$

satisfy

$$\begin{aligned} \sum_{\ell=0}^n \binom{k+n-\ell}{n-\ell} V_\ell^{(k+1)}(x) &= \frac{1}{2^k k!} \sum_{i=1}^k \sum_{\ell=0}^i a_i(k) \frac{i!}{\ell!} \\ &\times \sum_{m+s+p=n} \binom{2k+m-i-1}{m} \binom{i-\ell+s}{s} \langle \ell+p \rangle_\ell x^{i-2k-m} V_{\ell+p}(x) \end{aligned}$$

for $k \in \mathbb{N}$ and $n \geq 0$, where $V_n^{(1)}(x) = V_n(x)$. See [11, Theorem 5].

(5) The higher order Chebyshev polynomials of the fourth kind $W_n^{(\alpha)}(x)$ generated by

$$\left(\frac{1+t}{\sqrt{1-2xt+t^2}} \right)^\alpha = \sum_{n=0}^{\infty} W_n^{(\alpha)}(x) t^n$$

satisfy

$$\begin{aligned} \sum_{\ell=0}^n (-1)^{n-\ell} \binom{k+n-\ell}{n-\ell} W_\ell^{(k+1)}(x) &= \frac{1}{2^k k!} \sum_{i=1}^k \sum_{\ell=0}^i (-1)^{i-\ell} a_i(k) \frac{i!}{\ell!} \\ &\times \sum_{m+s+p=n} (-1)^s \binom{2k+m-i-1}{m} \binom{i-\ell+s}{s} \langle \ell+p \rangle_\ell x^{i-2k-m} W_{\ell+p}(x) \end{aligned}$$

for $k \in \mathbb{N}$ and $n \geq 0$, where $W_n^{(1)}(x) = W_n(x)$. See [11, Theorem 6].

(6) The higher order Chebyshev polynomials of the first kind $T_n^{(\alpha)}(x)$ generated by

$$\left(\frac{1-t^2}{\sqrt{1-2xt+t^2}} \right)^\alpha = \sum_{n=0}^{\infty} T_n^{(\alpha)}(x) t^n$$

satisfy

$$\begin{aligned} & 2^{k+1} k! \sum_{s+m+p=n} \binom{k+s}{s} \binom{m+k}{m} (-1)^m T_p^{(k+1)}(x) \\ &= \sum_{i=1}^k \sum_{\ell=0}^i a_i(k) \frac{i!}{\ell!} \sum_{m+s+p=n} \binom{2k+m-i-1}{m} \binom{i+s-\ell}{s} \langle \ell+p \rangle_\ell x^{i-2k-m} T_{p+\ell}(x) \\ &+ \sum_{i=1}^k \sum_{\ell=0}^i a_i(k) \frac{i!}{\ell!} (-1)^{i-\ell} \sum_{m+s+p=n} (-1)^s \binom{2k+m-i-1}{m} \binom{i+s-\ell}{s} \langle \ell+p \rangle_\ell x^{i-2k-m} T_{p+\ell}(x) \end{aligned}$$

for $k \in \mathbb{N}$ and $n \geq 0$. See [11, Theorem 7].

It is clear that the quantities $a_i(n)$ defined by (6.1) play a key role in the above-mentioned conclusions obtained in the paper [11]. However, the quantities $a_i(n)$ are expressed complicatedly and can not be computed easily. Can one find a simple expression for the quantities $a_i(n)$? Theorem 3.1 answers this question by

$$a_k(n) = \frac{(-1)^{n-k}}{2^{n-k}} \frac{n!}{k!} b_{n,k} = \frac{1}{2^{n-k}} \frac{(n-1)!}{(k-1)!} \binom{2n-k-1}{n-1} \quad (6.2)$$

for $n \geq k \geq 1$. By this much simpler expression for $a_k(n)$, we can reformulate all the above-mentioned main results in the paper [11] in terms of the quantities defined in (6.2). For saving time of the authors and space of this paper, we do not write down them in details.

Due to the same motivation and reason as Theorem 3.1, the authors composed and published the papers [10, 19, 20, 21, 35, 36, 37, 38, 48, 49, 50, 57, 58], for examples.

Remark 6.2. The identity (5.3) recovers [61, p. 2187, Theorem 2, Eq. (15b)]. It can also be verified alternatively and directly by the same method used in the proof of the identity (5.1).

Actually, the identity (5.3) is a special case $i = j \in \mathbb{N}$ of the identity (5.1). In other words, the identity (5.1) generalizes, or say, extends (5.3).

It is clear that the proof of the identity (5.3) is simpler than the one of (5.3) adopted in [61] and related references therein.

Remark 6.3. The integral representation (2.4) for the Catalan numbers C_k and its variant forms can be found in [3, 4, 5, 6, 16, 17, 44] and the closely related references therein.

In recent years, there are plenty of literature, such as [14, 18, 25, 27, 31, 32, 40, 41, 44, 46, 47, 59, 62, 63], dedicated to generalizations of the Catalan numbers C_n and to investigating their properties.

Remark 6.4. The formula (2.3) in Lemma 2.3 has also been applied many times in some papers such as [9, 26, 29, 33, 36, 38, 39, 42, 43, 45, 51, 56, 60] and the closely related references therein.

Remark 6.5. Let $A_n = I_n + M_n$ and I_n be the identity matrix of order n . By linear algebra, it is easy to see that $M_n^n = 0$ and

$$(I_n + M_n)(I_n - M_n + M_n^2 - M_n^3 + \cdots + (-1)^{n-1} M_n^{n-1}) = I_n - M_n^n = I_n.$$

This means that

$$A_n^{-1} = (I_n + M_n)^{-1} = I_n + \sum_{k=1}^{n-1} (-1)^k M_n^k.$$

In theory, this formula is useful for computing the inverse A_n^{-1} . But, in practice, it is too difficult to acquire the simple form in (4.1).

Can one conclude a general and concrete formula for computing M_n^k from Theorem 4.1?

Remark 6.6. Motivated by the proof of the identity 5.1, we naturally ask a question: can one explicitly compute integrals of the type

$$\int_0^1 z^{\alpha-1} (1-z)^{\beta-1} {}_pF_q(a, b; c; xz^\sigma) dz?$$

In [53, p. 340, Remark], it was given that

$$\int_0^1 z^{\alpha-1} (1-z)^{\beta-1} {}_2F_1(a, b; c; xz^\sigma) dz = \frac{\Gamma(c)\Gamma(\beta)}{\Gamma(a)\Gamma(b)} {}_3\Phi_2((a, 1), (b, 1), (\alpha, \sigma); (c, 1), (\alpha + \beta, \sigma); x),$$

where

$${}_p\Phi_q((\alpha_1, \beta_1), \dots, (\alpha_p, \beta_p); (\rho_1, \mu_1), \dots, (\rho_p, \mu_q); z) = \sum_{n=0}^{\infty} \frac{\Gamma(\alpha_1 + \beta_1 n) \cdots \Gamma(\alpha_p + \beta_p n)}{\Gamma(\rho_1 + \mu_1 n) \cdots \Gamma(\rho_q + \mu_q n)} \frac{z^n}{n!}$$

and β_r, μ_t are real positive numbers such that

$$1 + \sum_{t=1}^q \mu_t - \sum_{r=1}^p \beta_r > 0.$$

Making use of this result, we can supply an alternative proof of the identity 5.1 in Theorem 5.1.

There is a similar formula in [52, p. 104, Theorem 38].

This question has also been considered in [1, 23] and the closely related references therein.

Remark 6.7. In [15, p. 387, 15.4.18], it was listed that the formula

$${}_2F_1\left(a, a + \frac{1}{2}; 2a; z\right) = \frac{1}{\sqrt{1-z}} \left(\frac{1}{2} + \frac{\sqrt{1-z}}{2}\right)^{1-2a}$$

holds for the principal branch when $|z| < 1$, and by analytic continuation elsewhere. Straightforwardly letting $a = \frac{1-j}{2}$ results in

$${}_2F_1\left(\frac{1-j}{2}, \frac{2-j}{2}; 1-j; t\right) = \frac{1}{\sqrt{1-t}} \left(\frac{1}{2} + \frac{\sqrt{1-t}}{2}\right)^j, \quad |t| < 1.$$

Replacing t by $\frac{1}{t}$ leads to

$${}_2F_1\left(\frac{1-j}{2}, \frac{2-j}{2}; 1-j; \frac{1}{t}\right) = \frac{1}{2^j} \sqrt{\frac{t}{t-1}} \left(1 + \sqrt{\frac{t-1}{t}}\right)^j, \quad |t| > 1.$$

This expression for ${}_2F_1\left(\frac{1-j}{2}, \frac{2-j}{2}; 1-j; \frac{1}{t}\right)$ is slightly different from (2.5) in Lemma 2.6.

Remark 6.8. Comparing main results of this paper with those in [28], we can see that there exist some close connections among the Chebyshev polynomials of the second kind U_n , the Catalan numbers C_n , the central Delannoy numbers D_n , the Fibonacci polynomials $F_n(x)$, and triangular and tridiagonal matrices.

Comparing Theorem 3.1 with Theorem 5.1 reveals that the equality (3.4) can be reformulated in terms of the Catalan numbers C_n as

$$\sum_{k=1}^n \left[\sum_{\ell=0}^{\lfloor (k-1)/2 \rfloor} (-1)^\ell \binom{k-\ell-1}{\ell} C_{n-\ell-1} \right] (2x)^k U_k(x) = (2x)^{2n}. \quad (6.3)$$

Taking $x = 3$ in (6.3) and considering results in [28, Section 10] disclose that

$$\sum_{k=1}^n 6^k \left[\sum_{\ell=0}^{\lfloor (k-1)/2 \rfloor} (-1)^\ell \binom{k-\ell-1}{\ell} C_{n-\ell-1} \right] \left[\sum_{\ell=0}^k D(\ell) D(k-\ell) \right] = 6^{2n},$$

where $D(k)$ denotes the central Delannoy numbers which are combinatorially the numbers of “king walks” from the $(0, 0)$ corner of an $n \times n$ square to the upper right corner (n, n) and can be generated analytically by

$$\frac{1}{\sqrt{1-6x+x^2}} = \sum_{k=0}^{\infty} D(k) x^k = 1 + 3x + 13x^2 + 63x^3 + \cdots.$$

Taking $x = \frac{s}{2}\sqrt{-1}$ in (6.3) and utilizing results in [28, Section 8] expose that

$$\sum_{k=1}^n (-1)^k \left[\sum_{\ell=0}^{\lfloor (k-1)/2 \rfloor} (-1)^\ell \binom{k-\ell-1}{\ell} C_{n-\ell-1} \right] s^k F_{k+1}(s) = (-1)^n s^{2n},$$

where the Fibonacci polynomials

$$F_n(s) = \frac{1}{2^n} \frac{(s + \sqrt{4+s^2})^n - (s - \sqrt{4+s^2})^n}{\sqrt{4+s^2}}$$

can be generated by

$$\frac{t}{1-ts-t^2} = \sum_{n=1}^{\infty} F_n(s) t^n = t + st^2 + (s^2+1)t^3 + (s^3+2s)t^4 + \cdots.$$

Remark 6.9. Now we can see that our main results in this paper stride analysis, special functions, combinatorics, number theory, matrix theory, integral transforms, and the like.

REFERENCES

- [1] P. Agarwal, F. Qi, M. Chand, and S. Jain, *Certain integrals involving the generalized hypergeometric function and the Laguerre polynomials*, J. Comput. Appl. Math. **313** (2017), 307–317; Available online at <http://dx.doi.org/10.1016/j.cam.2016.09.034>.
- [2] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Revised and Enlarged Edition, D. Reidel Publishing Co., Dordrecht and Boston, 1974.
- [3] T. Dana-Picard, *Integral presentations of Catalan numbers*, Internat. J. Math. Ed. Sci. Tech. **41** (2010), no. 1, 63–69; Available online at <http://dx.doi.org/10.1080/00207390902971973>.
- [4] T. Dana-Picard, *Integral presentations of Catalan numbers and Wallis formula*, Internat. J. Math. Ed. Sci. Tech. **42** (2011), no. 1, 122–129; Available online at <http://dx.doi.org/10.1080/0020739X.2010.519792>.
- [5] T. Dana-Picard, *Parametric integrals and Catalan numbers*, Internat. J. Math. Ed. Sci. Tech. **36** (2005), no. 4, 410–414; Available online at <http://dx.doi.org/10.1080/00207390412331321603>.
- [6] T. Dana-Picard and D. G. Zeitoun, *Parametric improper integrals, Wallis formula and Catalan numbers*, Internat. J. Math. Ed. Sci. Tech. **43** (2012), no. 4, 515–520; Available online at <http://dx.doi.org/10.1080/0020739X.2011.599877>.
- [7] O. Dunkel, W. A. Bristol, W.R. Church, and V. F. Ivanoff, *Problems and Solutions: Solutions: 3421*, Amer. Math. Monthly **38** (1931), no. 1, 54–57; Available online at <http://dx.doi.org/10.2307/2301598>.

- [8] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, Translated from the Russian, Translation edited and with a preface by Daniel Zwillinger and Victor Moll, Eighth edition, Revised from the seventh edition, Elsevier/Academic Press, Amsterdam, 2015; Available online at <http://dx.doi.org/10.1016/B978-0-12-384933-5.00013-8>.
- [9] B.-N. Guo and F. Qi, *Explicit formulas for special values of the Bell polynomials of the second kind and the Euler numbers*, ResearchGate Technical Report (2015), available online at <http://dx.doi.org/10.13140/2.1.3794.8808>.
- [10] B.-N. Guo and F. Qi, *Some identities and an explicit formula for Bernoulli and Stirling numbers*, J. Comput. Appl. Math. **255** (2014), 568–579; Available online at <http://dx.doi.org/10.1016/j.cam.2013.06.020>.
- [11] T. Kim, D. S. Kim, J.-J. Seo, and D. V. Dolgy, *Some identities of Chebyshev polynomials arising from non-linear differential equations*, J. Comput. Anal. Appl. **23** (2017), no. 5, 820–832.
- [12] T. Koshy, *Catalan Numbers with Applications*, Oxford University Press, Oxford, 2009.
- [13] F.-F. Liu, X.-T. Shi, and F. Qi, *A logarithmically completely monotonic function involving the gamma function and originating from the Catalan numbers and function*, Glob. J. Math. Anal. **3** (2015), no. 4, 140–144; Available online at <http://dx.doi.org/10.14419/gjma.v3i4.5187>.
- [14] M. Mahmoud and F. Qi, *Three identities of the Catalan–Qi numbers*, Mathematics **4** (2016), no. 2, Article 35, 7 pages; Available online at <http://dx.doi.org/10.3390/math4020035>.
- [15] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark (eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York, 2010; Available online at <http://dlmf.nist.gov/>.
- [16] K. A. Penson and J.-M. Sixdeniers, *Integral representations of Catalan and related numbers*, J. Integer Seq. **4** (2001), no. 2, Article 01.2.5.
- [17] F. Qi, *An improper integral with a square root*, Preprints **2016**, 2016100089, 8 pages; Available online at <http://dx.doi.org/10.20944/preprints201610.0089.v1>.
- [18] F. Qi, *Asymptotic expansions, complete monotonicity, and inequalities of the Catalan numbers*, ResearchGate Technical Report (2015), available online at <http://dx.doi.org/10.13140/RG.2.1.4371.6321>.
- [19] F. Qi, *Explicit formulas for the convolved Fibonacci numbers*, ResearchGate Working Paper (2016), available online at <http://dx.doi.org/10.13140/RG.2.2.36768.17927>.
- [20] F. Qi and B.-N. Guo, *Explicit formulas and nonlinear ODEs of generating functions for Eulerian polynomials*, ResearchGate Working Paper (2016), available online at <http://dx.doi.org/10.13140/RG.2.2.23503.69288>.
- [21] F. Qi and B.-N. Guo, *Explicit formulas for derangement numbers and their generating function*, J. Nonlinear Funct. Anal. **2016**, Article ID 45, 10 pages.
- [22] F. Qi, B.-N. Guo, V. Čerňanová, and X.-T. Shi, *Explicit expressions, Cauchy products, integral representations, convexity, and inequalities of central Delannoy numbers*, ResearchGate Working Paper (2016), available online at <http://dx.doi.org/10.13140/RG.2.1.4889.6886>.
- [23] F. Qi and K. S. Nisar, *Some integral transforms of the generalized k-Mittag-Leffler function*, Preprints **2016**, 2016100020, 8 pages; Available online at <http://dx.doi.org/10.20944/preprints201610.0020.v1>.
- [24] F. Qi, *Limit formulas for ratios between derivatives of the gamma and digamma functions at their singularities*, Filomat **27** (2013), no. 4, 601–604; Available online at <http://dx.doi.org/10.2298/FIL1304601Q>.
- [25] F. Qi, A. Akkurt, and H. Yildirim, *Catalan numbers, k-gamma and k-beta functions, and parametric integrals*, J. Comput. Anal. Appl. **24** (2018), in press. ResearchGate Working Paper (2017), available online at <http://dx.doi.org/10.13140/RG.2.2.19398.06721>.
- [26] F. Qi and V. Čerňanová, *Some discussions on a kind of improper integrals*, Internat. J. Anal. Appl. **11** (2016), no. 2, 101–109.
- [27] F. Qi and P. Cerone, *Several expressions, some properties, and a double inequality of the Fuss–Catalan numbers*, ResearchGate Research (2015), available online at <http://dx.doi.org/10.13140/RG.2.1.1655.6004>.
- [28] F. Qi, V. Čerňanová, and Y. S. Semenov, *On tridiagonal determinants and the Cauchy product of central Delannoy numbers*, ResearchGate Working Paper (2016), available online at <http://dx.doi.org/10.13140/RG.2.1.3772.6967>.
- [29] F. Qi and B.-N. Guo, *Explicit and recursive formulas, integral representations, and properties of the large Schröder numbers*, Kragujevac J. Math. **41** (2017), no. 1, 121–141.
- [30] F. Qi and B.-N. Guo, *Integral representations and complete monotonicity of remainders of the Binet and Stirling formulas for the gamma function*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM **111** (2017), no. 2, 425–434; Available online at <http://dx.doi.org/10.1007/s13398-016-0302-6>.

- [31] F. Qi and B.-N. Guo, *Logarithmically complete monotonicity of a function related to the Catalan–Qi function*, Acta Univ. Sapientiae Math. **8** (2016), no. 1, 93–102; Available online at <http://dx.doi.org/10.1515/ausm-2016-0006>.
- [32] F. Qi and B.-N. Guo, *Logarithmically complete monotonicity of Catalan–Qi function related to Catalan numbers*, Cogent Math. (2016), **3**:1179379, 6 pages; Available online at <http://dx.doi.org/10.1080/23311835.2016.1179379>.
- [33] F. Qi and B.-N. Guo, *Several explicit and recursive formulas for the generalized Motzkin numbers*, Preprints **2017**, 22017030200, 11 pages; Available online at <http://dx.doi.org/10.20944/preprints201703.0200.v1>.
- [34] F. Qi and B.-N. Guo, *Some properties and generalizations of the Catalan, Fuss, and Fuss–Catalan numbers*, Mathematical Analysis and Applications: Selected Topics, 35 pages, edited by H. Dutta, M. Ruzhansky, and R. P. Agarwal, Wiley, October 2017.
- [35] F. Qi and B.-N. Guo, *Some properties of a solution to a family of inhomogeneous linear ordinary differential equations*, Preprints **2016**, 2016110146, 11 pages; Available online at <http://dx.doi.org/10.20944/preprints201611.0146.v1>.
- [36] F. Qi and B.-N. Guo, *Some properties of the Hermite polynomials and their squares and generating functions*, Preprints **2016**, 2016110145, 14 pages; Available online at <http://dx.doi.org/10.20944/preprints201611.0145.v1>.
- [37] F. Qi and B.-N. Guo, *Viewing some nonlinear ODEs and their solutions from the angle of derivative polynomials*, ResearchGate Working Paper (2016), available online at <http://dx.doi.org/10.13140/RG.2.1.4593.1285>.
- [38] F. Qi and B.-N. Guo, *Viewing some ordinary differential equations from the angle of derivative polynomials*, Preprints **2016**, 2016100043, 12 pages; Available online at <http://dx.doi.org/10.20944/preprints201610.0043.v1>.
- [39] F. Qi, B.-N. Guo, V. Čerňanová, and X.-T. Shi, *Explicit expressions, Cauchy products, integral representations, convexity, and inequalities of central Delannoy numbers*, ResearchGate Working Paper (2016), available online at <http://dx.doi.org/10.13140/RG.2.1.4889.6886>.
- [40] F. Qi, M. Mahmoud, X.-T. Shi, and F.-F. Liu, *Some properties of the Catalan–Qi function related to the Catalan numbers*, SpringerPlus (2016), **5**:1126, 20 pages; Available online at <http://dx.doi.org/10.1186/s40064-016-2793-1>.
- [41] F. Qi, X.-T. Shi, and P. Cerone, *A unified generalization of the Catalan, Fuss, and Fuss–Catalan numbers and the Catalan–Qi function*, ResearchGate Working Paper (2015), available online at <http://dx.doi.org/10.13140/RG.2.1.3198.6000>.
- [42] F. Qi, X.-T. Shi, and B.-N. Guo, *Integral representations of the large and little Schröder numbers*, ResearchGate Working Paper (2016), available online at <http://dx.doi.org/10.13140/RG.2.1.1988.3288>.
- [43] F. Qi, X.-T. Shi, and B.-N. Guo, *Two explicit formulas of the Schröder numbers*, Integers **16** (2016), Paper No. A23, 15 pages.
- [44] F. Qi, X.-T. Shi, and F.-F. Liu, *An integral representation, complete monotonicity, and inequalities of the Catalan numbers*, ResearchGate Technical Report (2015), available online at <http://dx.doi.org/10.13140/RG.2.1.3754.4806>.
- [45] F. Qi, X.-T. Shi, F.-F. Liu, and D. V. Kruchinin, *Several formulas for special values of the Bell polynomials of the second kind and applications*, J. Appl. Anal. Comput. (2017), in press. ResearchGate Technical Report (2015), available online at <http://dx.doi.org/10.13140/RG.2.1.3230.1927>.
- [46] F. Qi, X.-T. Shi, M. Mahmoud, and F.-F. Liu, *Schur-convexity of the Catalan–Qi function related to the Catalan numbers*, Tbilisi Math. J. **9** (2016), no. 2, 141–150; Available online at <http://dx.doi.org/10.1515/tmj-2016-0026>.
- [47] F. Qi, X.-T. Shi, M. Mahmoud, and F.-F. Liu, *The Catalan numbers: a generalization, an exponential representation, and some properties*, J. Comput. Anal. Appl. **23** (2017), no. 5, 937–944.
- [48] F. Qi, J.-L. Wang, and B.-N. Guo, *Simplifying two families of nonlinear ordinary differential equations*, Preprints **2017**, 22017030192, 6 pages; Available online at <http://dx.doi.org/10.20944/preprints201703.0192.v1>.
- [49] F. Qi and J.-L. Zhao, *Some properties of the Bernoulli numbers of the second kind and their generating function*, J. Differ. Equ. Appl. (2017), in press. ResearchGate Working Paper (2017), available online at <http://dx.doi.org/10.13140/RG.2.2.13058.27848>.
- [50] F. Qi, J.-L. Zhao, and J.-L. Wang, *Derivative polynomials of a function related to the Apostol–Euler and Frobenius–Euler numbers*, J. Nonlinear Sci. Appl. **10** (2017), in press. ResearchGate Working Paper (2017), available online at <http://dx.doi.org/10.13140/RG.2.2.35572.35204>.

- [51] F. Qi and M.-M. Zheng, *Explicit expressions for a family of the Bell polynomials and applications*, Appl. Math. Comput. **258** (2015), 597–607; Available online at <http://dx.doi.org/10.1016/j.amc.2015.02.027>.
- [52] E. D. Rainville, *Special Functions*, The Macmillan Co., New York, 1960.
- [53] S. B. Rao, J. C. Prajapati, A. K. Shukla, *Wright type hypergeometric function and its properties*, Adv. Pure Math. **3** (2013), 335–342; Available online at <http://dx.doi.org/10.4236/apm.2013.33048>.
- [54] X.-T. Shi, F.-F. Liu, and F. Qi, *An integral representation of the Catalan numbers*, Glob. J. Math. Anal. **3** (2015), no. 3, 130–133; Available online at <http://dx.doi.org/10.14419/gjma.v3i3.5055>.
- [55] R. P. Stanley, *Catalan Numbers*, Cambridge University Press, New York, 2015; Available online at <http://dx.doi.org/10.1017/CB09781139871495>.
- [56] C.-F. Wei and F. Qi, *Several closed expressions for the Euler numbers*, J. Inequal. Appl. **2015**, 2015:219, 8 pages; Available online at <http://dx.doi.org/10.1186/s13660-015-0738-9>.
- [57] A.-M. Xu and G.-D. Cen, *Closed formulas for computing higher-order derivatives of functions involving exponential functions*, Appl. Math. Comput. **270** (2015), 136–141; Available online at <http://dx.doi.org/10.1016/j.amc.2015.08.051>.
- [58] A.-M. Xu and Z.-D. Cen, *Some identities involving exponential functions and Stirling numbers and applications*, J. Comput. Appl. Math. **260** (2014), 201–207; Available online at <http://dx.doi.org/10.1016/j.cam.2013.09.077>.
- [59] L. Yin and F. Qi, *Several series identities involving the Catalan numbers*, Preprints **2017**, 2017030029, 11 pages; Available online at <http://dx.doi.org/10.20944/preprints201703.0029.v1>.
- [60] J.-L. Zhao and F. Qi, *Two explicit formulas for the generalized Motzkin numbers*, J. Inequal. Appl. **2017**, 2017:44, 8 pages; Available online at <http://dx.doi.org/10.1186/s13660-017-1313-3>.
- [61] R. R. Zhou and W. Chu, *Identities on extended Catalan numbers and their q -analogs*, Graphs Combin. **32** (2016), no. 5, 2183–2197; Available online at <http://dx.doi.org/10.1007/s00373-016-1694-y>.
- [62] Q. Zou, *Analogues of several identities and supercongruences for the Catalan–Qi numbers*, J. Inequal. Spec. Funct. **7** (2016), no. 4, 235–241.
- [63] Q. Zou, *The q -binomial inverse formula and a recurrence relation for the q -Catalan–Qi numbers*, J. Math. Anal. **8** (2017), no. 1, 176–182.

(Qi) INSTITUTE OF MATHEMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA; COLLEGE OF MATHEMATICS, INNER MONGOLIA UNIVERSITY FOR NATIONALITIES, TONGLIAO CITY, INNER MONGOLIA AUTONOMOUS REGION, 028043, CHINA; DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN CITY, 300387, CHINA

E-mail address: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com

URL: <https://qifeng618.wordpress.com>

(Guo) SCHOOL OF MATHEMATICS AND INFORMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

E-mail address: bai.ni.guo@gmail.com, bai.ni.guo@hotmail.com

URL: http://www.researchgate.net/profile/Bai-Ni_Guo/



© 2017 by the authors. Licensee Preprints, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons by Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).