

Review

Human MAP tau based targeted cytolytic fusion proteins

Olusiji A. Akinrinmade¹, Sandra Jordaan¹, Dmitrij Hristodorov², Radoslav Mladenov², Neelakshi Mungra¹, Shivan Chetty^{1,*}, Stefan Barth^{1,*,#}

¹South African Research Chair in Cancer Biotechnology, Institute of Infectious Disease and Molecular Medicine (IDM), Department of Integrative Biomedical Sciences, Faculty of Health Sciences, University of Cape Town, South Africa

²Fraunhofer Institute for Molecular Biology and Applied Ecology, Aachen, Germany

*Equal contribution

#Corresponding Author: stefan.barth@uct.ac.za; Tel.: +27 21 406 6938

Abstract: Some of the most promising small molecule toxins used to generate antibody drug conjugates (ADCs) include anti-mitotic agents (e.g. auristatin and its derivatives) which are designed to attack cancerous cells at their most vulnerable state during mitosis. We were interested to identify a human cystostatic protein eventually showing comparable activities and allowing the generation of corresponding targeted fully human cytolytic fusion proteins. Recently, we identified the human microtubule associated protein tau (MAP tau) which binds specifically to tubulin and modulates the stability of microtubules thereby blocking mitosis and presumably vesicular transport. By binding and stabilizing polymerized microtubule filaments, MAP tau-based fusion proteins skew microtubule dynamics towards cell cycle arrest and apoptosis. This biological activity makes rapidly proliferating cells (e.g. cancer and inflammatory cells) an excellent target for MAP tau-based targeted treatments. Their superior selectivity for proliferating cells confers additional selectivity towards upregulated tumor-associated antigens at their surface, thereby preventing off-target related toxicity against normal cells bearing tumor-associated antigens at physiologically normal to low levels. In this review, we highlight recent findings on MAP tau-based targeted cytolytic fusion proteins reported in preclinical immunotherapeutic studies.

Keywords: human cytolytic fusion proteins; immunotherapy; microtubule associated protein tau (MAP); cancer; inflammatory diseases

1. INTRODUCTION

Of all pharmacological agents developed for the treatment of cancer, anti-mitotic drugs have till now remained the most successful [1]. All such drugs are largely defined by their profound ability to compromise cellular dynamics during the crucial period of mitotic cell division and by so doing initiate apoptosis or autophagy related cell death [2]. As a reason of their success, traditional anti-mitotic agents like the vinca alkaloids have remained an integral part of anti-cancer agents available to clinicians since their first use about seven decades ago [3]. Their clinical effectiveness has encouraged the developments of newer therapeutics against several cellular and molecular targets involved in different stages of mitosis (e.g. microtubules, kinases, motor proteins, and multi-protein complexes) [2]. Importantly, those targeting the microtubules have been regarded as the most clinically effective [4]. They work by either stabilizing or destabilizing the polymerization of microtubule, therein disrupting the organization of mitotic spindle necessary for the completion of the M phase of the cell cycle [5]. Normally, microtubules springing from the opposite poles of a dividing cell during metaphase will allow spindle fibres to make productive attachment to the kinetochores and by so doing establish connections necessary for the precise arrangement of chromosomes along the equatorial plate. This is particularly dependent on the dynamic nature of microtubules through its polymerization (grow) and depolymerization (shorten) cycles which allows

the microtubule to search the cytoplasm for the kinetochores [6, 7]. Importantly, the precise alignment of chromosomes at the equatorial plate is a pre-requisite to allow spindle assembly checkpoint-SAC (a cell cycle regulator) permit the commencement of anaphase. This is because any abnormal alignment or absence of a single chromosome at the equatorial plate will prevent the cell from completing mitosis and eventually lead to the induction of apoptosis [8]. Most microtubule targeting drugs (stabilizing or destabilizing) exploit this delicate moment of mitotic cell division to exert their cytotoxic activity, often by the common mechanism of suppressing microtubule dynamics and the chronic activation of spindle assembly checkpoint-SAC which halts cell-division in a prolonged event that signals the induction of apoptosis (figure 1) [9]. Anti-mitotics like the vinca alkaloids (vinflunine, vincristine, vinorelbine, vindesine and eribulin), cryptophycins, halichondrins, estramustine, colchicines, and combretastatins are examples of drugs known to depolymerize / destabilize microtubules by binding to various β -tubulin sites [5, 10, 11]. On the other hand, taxanes paclitaxel (Taxol), Lauimalide, Dictyostatin, the epothilones, etc. are examples of microtubule stabilizing compounds [12, 13] known to enhance tubulin polymerization / stabilization and prevent the ability of microtubules to shorten or separate sister chromatids [10, 14]. Clinically, the vinca alkaloids and taxanes have been integrated in combination chemotherapy regimens for the treatment of breast cancer, ovarian cancer, non-small-cell lung carcinoma and haematological malignancies [15]. Based on their popularity, semi-synthetic and synthetic strategies have been used to produce analogues of these agents for a wider therapeutic application [16]. This is because most of these agents are scarce and only found in minute amounts in their natural sources (mostly marine organisms and plants) herewith slowing down clinical development [17]. Examples include auristatin E (AE) and Monomethyl auristatin E (MMAE), which are derivatives of the tubulin polymerisation inhibitor dolastatin 10 [18]. Interestingly, these analogues are highly effective and have shown more cytotoxicity than their natural counterparts with preclinical studies showing AE to be 200 times more potent than vinblastine [19]. The therapeutic effectiveness of MMAE at the cytotoxic domain of brentuximab vedotin has also resulted in its approval by the US Food and Drug Administration (FDA) for the treatment of refractory Hodgkin lymphoma and systemic anaplastic large-cell lymphoma [20, 21]. Unfortunately, as this is not often the case with other microtubule binding agents, especially the synthetic derivatives of taxanes faced with several clinical drawbacks [17], we were interested to identify a human anti-mitotic protein eventually showing comparable activities and allowing the generation of corresponding fully human cytolytic fusion proteins.

We identified microtubule associated proteins tau (MAP tau) which belongs to a family of proteins (microtubule associated proteins, MAPs) endowed with the ability to bind tubulin and modulate the stability of microtubules. Different classes of MAPs occur in different cell types and are known to have specific functions. Of interest to scientists are MAPs that can stabilize microtubules including MAP2, MAP4 and MAP tau [22]. They act primarily by binding to microtubules and by promoting the formation of crosslinks between tubulin units. By promoting tubulin assembly, the rate of depolymerization is reduced and an overall stabilization effect is observed. In so doing, these proteins promote rapid formation of the mitotic spindle, which allows for the movement of chromosomes to opposite poles during mitosis [23]. As the enzymatic phosphorylation of MAPs is a tightly regulated system for controlling microtubule dynamics and ensuring a successful cell division [24], we postulated that the presence of non-phosphorylated MAPs would considerably slow down mitosis and force the cell to undergo apoptosis [18]. The implementation of MAP tau as an effector protein in fully humanized antibody-based immunotherapy was not accidental but based on its vital role in regulating microtubule behaviour. Thus, the first MAP tau-based fusion protein was engineered, bearing specificity towards the human epidermal growth factor (EGF) receptor [4]. In this review, we provide a summary on the biology of MAP tau, including data from preclinical studies incorporating the selective cytotoxicity of MAP tau towards proliferating cells for the development of targeted human cytolytic fusion proteins.




Figure 1. Mechanistic scenario of events leading to mitotic cell death (MCD) by microtubule targeting drugs. Microtubule stabilizing agents (e.g. paclitaxel, dictyostatin, MAPs) promote the formation of microtubule bundles that cannot disassemble while microtubule destabilizing agents (e.g. vinca alkaloids, colchicines) prevent formation of the mitotic spindle by inhibiting tubulin polymerization. Both results in the activation of SAC eventually leading to mitotic arrest and apoptosis.

2. MAP Tau: DISCOVERY AND STRUCTURE

Tau proteins are found in many animal species, including *Drosophila* [25], rodents [26], goat [27], goldfish [28] and humans [29]. Discovered in 1975 by Weingarten et al. by co-purification with tubulin [30], MAP tau was recognized as a protein factor essential for microtubule assembly [31]. It is most abundantly expressed in the neurons of the human brain [12] and results from a single tau gene present on chromosome 17 [32]. However, evidence shows that MAP tau expression is not solely restricted to neuronal cells. Differential trace levels of MAP tau are also detectable in the heart, muscle, testis, lung and pancreatic tissues [33, 34].

Interestingly, when subjected to electrophoresis on sodium dodecyl sulfate polyacrylamide gels, it was found that purified tau consists of four to six polypeptides of molecular weights ranging from 55-62 kDa [35]. Subsequent discoveries then led to confirm that alternative splicing of tau mRNA indeed results in the formation of six isoforms in the adult mammalian brain tissue [36]. However, they all exhibit a similar structure, differing only in the number of microtubule-binding domains, as well as in the number of N-terminal inserts. In general, tau proteins consist of a C-terminal region that encompasses a proline-rich domain which is capable of interacting with microtubules [37]. On the other hand, the N-terminal domain consists of repeats of amino acids which do not bind to tubulin. By projecting from the microtubule surface, tau's N-terminal domains mediate the

interaction between microtubules and the plasma membrane, thereby promoting neuronal development [38].

Map Tau FUNCTION

The primary role of MAP tau revolves around its ability to stabilize as well as promote the polymerization of microtubules. Indeed, a study conducted in 1992 demonstrated that the addition of tau to pure tubulin solutions, culminates in an increase in the rate of growth of microtubules [39]. Such an observation is also accompanied by hindrance of progression into the shrinking phase of microtubule dynamics. Through new technological approaches, such as cryo-electron microscopy, it was determined that tau binds in a longitudinal fashion to microtubules and that such binding results in the bridging of tubulin interfaces [40].

Several *in vitro* experiments were carried out to characterize the defining role of MAP tau in neuronal development. By selectively inhibiting tau expression in cell culture, it was found that neuronal elongation was severely affected [41]. While these results demonstrated an indispensable role of tau in the neuron, other studies begged to differ. Tau knockout mice did not exhibit any kind of abnormalities in brain development; microtubule stability and neuronal growth were essentially normal [42]. Therefore, this fact lead to consider the role of other nervous system MAPs in compensating for any tau deficiency. Moreover, the important functions of tau are directly related to its peculiar structure. It can be speculated that each tau isoform has different functions based on its structural composition. Such claim is warranted by the varying spatial and temporal expression of MAP tau variants. For instance, only one tau isoform is prominently expressed during fetal stages of development, while all six isoforms are usually present during adulthood [43]. Additionally, fetal tau undergoes a higher level of phosphorylation as compared to adult tau [44]. Together, these studies provide evidence towards the cooperative role of isoforms and phosphorylation in regulating microtubule dynamics.

Pathologies associated with MAP tau

To ensure the normal biological function of tau, a regulatory network of kinases and phosphatases has proved to be highly indispensable. A KXGS motif found within tau's repeat domain [45] is known to be the site of phosphorylation, to which any abnormalities in the degree of phosphorylation can have serious impacts on an organism's well-being. Clinically, aggregation of filamentous tau proteins is a distinctive feature of various neurodegenerative diseases, collectively termed as tauopathies [46]. Some examples include Pick's disease, progressive supranuclear palsy, as well as Alzheimer's disease (AD) [47]. AD remains one of the most common occurrence of such pathology. It is characterized by an excessive phosphorylation at Ser396 [48] and the accumulation of extracellular plaques consisting primarily of β -amyloid peptides and MAP tau [49]. The pathology of such a disease may be in part due to the loss of tau's ability to bind to tubulin. Microtubule destabilisation thereby follows and causes a serious threat to neuronal integrity. By removing the vital phosphorylation sites of tau, we made a *bona fide* attempt towards preventing any possible hyper-phosphorylation of MAP tau-based fusion proteins. Theoretically, it is also not realistic that these fusion proteins will cross a healthy blood-brain-barrier. This could only be possible in patients already suffering from neurodegenerative diseases (e.g. MS or Alzheimer) and having a leaky BBB. Furthermore, the selective nature of human cytolytic fusion proteins towards their target receptors helps prevents the accumulation of MAP tau in the human brain.

Map Tau-based targeted human cytolytic fusion proteins

For the first time, we recently demonstrated the selective delivery and cytotoxicity of a constitutively active human microtubule stabilizing protein towards cancer cells by reporting a new drug format incorporating the human MAP tau protein in the fashion of a targeted human cytolytic fusion protein. The expression construct for this fusion protein comprised of the human MAP tau (isoform 3) protein mutated at two phosphorylation sites and genetically fused to the human anti-

EGFR single chain fragment variable (scFv). This double mutation (S156A and S204A) allows MAP tau to bind microtubules and by so doing interrupt their dynamic “grow” and “shorten” behaviour, ultimately leading to mitotic arrest and induction of apoptosis. To allow nuclear enrichment of the internalised protein, we inserted a nuclear localization signal sequence (NLS) by adding the sequence (5'-CCCAAAAAAAAAAAGGAAAGTG-3') derived from the SV40 T-antigen to the C-terminus of the open reading frame (ORF). A schematic representation of this expression construct is depicted in figure 2 below.



Figure 2: Schematic representation of the expression cassette for the recombinant EGF-MAP human cytolytic fusion protein. The open reading frame (ORF) was expressed in *E. coli* BL21(DE3) using protocol for periplasmic stress expression in the presence of compatible solutes and includes an N-terminal peptidase B (pelB) leader sequence from *Erwinia carotovora* CE which directs the fusion protein to the periplasmic space, a His₁₀ tag to facilitate purification, and enterokinase cleavage site to allow proteolytic separation of the fusion construct from the upstream production sequence. * = C-terminal NLS.

Based on our hypothesis that a constitutively active anti-EGFR-MAP tau fusion protein would trigger apoptosis in a proliferation dependent manner, we tested the specificity of the fusion protein in *in vitro* and *in vivo* experiments. Specific binding of the EGF-MAP Tau construct to EGFR-overexpressing pancreatic cancer cell line (L3.6pl) was revealed by flow cytometry with the absence of non-specific binding to EGFR⁻ HEK293 cells. Cellular cytotoxicity assays revealed that the EGF-MAP fusion protein had an IC₅₀ value of ~1 μM against L3.6pl cells with no cytotoxicity towards EGFR⁻ HEK293 cell line *in vitro*, which importantly demonstrates the selective cytotoxicity of human MAP Tau towards its targeted ligand. Next, we further evaluated the proliferation dependent cytotoxicity of EGF-MAP by halting cell division with a pre-titrated concentration (100 pM for 24 h) of paclitaxel known to arrest mitosis. Importantly, this mitosis blocking reagent is confirmed (paclitaxel, at above concentration) not to induce apoptosis in cell lines [50]. Findings from the study revealed that no cytotoxic activity was evident for EGF-MAP in non-proliferating cells in comparison to the control immunotoxin 425(scFv)-ETA' which was highly cytotoxic (IC₅₀ ~1nM) to L3.6pl cells arrested at the G1 growth phase. Furthermore, apoptosis and tubulin polymerisation assays showed EGF-MAP to induce apoptosis by promoting microtubule polymerisation. Taking advantage of the popularity of the xenograft models to screen and evaluate new anti-cancer agents for their clinical potential [51], we investigated the efficacy of EGF-MAP in BALB/C nu/nu mice subcutaneously injected with Katushka-2 (far-red fluorescent protein) transfected L3.6pl cells. An anti-tumor treatment cycle of 4 mg kg⁻¹ EGF-MAP intravenously administered to mice at 11 days after tumor challenge resulted in a significant reduction in tumor growth when compared to the control group that received PBS [18]. More recently, subsequent evaluation of several MAP tau-based human cytolytic fusion proteins have been described and are summarized in table 1 below.

Construct	Cell line	Disease model	IC ₅₀ value (nmol/l)	Reference
Ki-4(scFv)-MAP	L540cy	HL & sALCL	53	[52]
	L428	HL & sALCL	135	
	Karpas 299	HL & Salcl	220	
anti-EpCAM(scFv)-MAP	L3.6pl	Pancreas Carcinoma	43	[53]
	A431	Epidermoid carcinoma	67	
	22Rv1	Prostate carcinoma	677	

	C4-2	Prostate carcinoma	161	
	SU86.86	Pancreas carcinoma	333	
scFv35-MAP	FL-OH1	Rhabdomyosarcoma	900	[54]
	RD	Rhabdomyosarcoma	950	
α CSPG4(scFv)-MAP	MDA-MB-231	TNBC	219	[55]
	Hs 578T	TNBC	480	
H22(scFv)-MAP	HL-60 cells	M1 macrophage-mediated diseases	0.04	[56]
	CD64+ leukemic blasts	AML / CML		[57]

Table 1: Cytotoxicity of MAP-tau based human cytolytic fusion proteins to different cell lines. HL: Hodgkin lymphoma, TNBC: Triple negative breast cancer, sALCL: systemic anaplastic large cell lymphoma, AML: Acute myeloid leukemia, CML: Chronic myelomonocytic leukemia.

As depicted in the table above, most of the MAP-tau based fusion proteins did demonstrate improved IC_{50} values when compared to the IC_{50} value of 1 μ M from its first use, even though they were constructed on the same Map Tau ORF described earlier. This difference could mainly be linked to the different binding affinity of the targeting moiety and/or internalization frequency of target cell receptors towards the MAP-tau bound ligand, hence the different concentration of the protein available to confer its cytolytic activity. Also, all of the MAP-tau based fusion proteins recorded above were favourably well tolerated *in vivo* at an intravenous dose of 4 mg kg^{-1} . Interestingly, this treatment schedule was sufficient to significantly inhibit tumor growth when compared with the control group. It is also important to note that the cytotoxicity conferred by this new class of therapeutics is comparably able to compete with other effector proteins especially those of human origin. The anti-CD30 single-chain antibody fragment Ki4(scFv) fused in frame to MAP tau for treatment of CD30⁺ malignancies such as Hodgkin lymphoma (HL) and systemic anaplastic large cell lymphoma (sALCL) was shown to induce apoptosis in rapidly proliferating L540cy, L428, and Karpas 299 cells in a dose-dependent manner with IC_{50} values of ~53 nmol/L, which is comparably like to that obtained (IC_{50} 36 nmol/L) for the Gb_R201K-Ki4(scFv), a granzyme B-derived mutant. Also, an anti-EpCAM(scFv)-MAP fusion protein showed promising results with IC_{50} values of 43 nmol/L and 67 nmol/L against L3.6pl and A431 EpCAM⁺ cancer cell lines, which is also comparably better when compared with that obtained for anti-EpCAM(scFv)-ETA' (409 nmol/L on L3.6pl and 91 nmol/L on A431 [53]. As earlier described, it is important to note that differences in IC_{50} values are attributed to different target antigens and as well as their differences in expression on different cells and cell lines. The highly promising serpin B9-resistant Gb_R201K-Ki4(scFv) construct has also been documented in literature to possess single digit IC_{50} values, with picomolar IC_{50} values also published in literature for the bacterial toxin ETA'. On the other hand, the cytotoxicity of MAP-tau based fusion proteins can still be greatly improved by the insertion of adapter sequences that facilitate the translocation of the effector molecule from the endosome to the cytosol [58]. This is because most endogenously secreted human molecules are entirely dependent on transport vesicles to move from one cellular location to the other unlike the bacteria enzymes which possess a natural evolutionary adapted membrane translocation domain which allows a higher concentration of the effector enzyme per target substrate in the cytosol. Hetzel et al., in 2008 reported the significant contribution of small cleavable adapter sequences to enhancing the cytotoxicity of human cytolytic fusion proteins. Indeed the use of these adapter sequence is a major aim in our future MAP-tau based preclinical studies [53].

Another mechanistic study into the apoptosis pathway induced by compromising microtubules dynamics by MAP Tau has also offered a significant advantage to better understanding their underlying mechanism of action (MOA). Amoury and colleagues recently highlighted the possibility of a completely caspase-independent pathway for a fusion protein consisting of the high affinity

CSPG4-specific single-chain antibody fragment fused to MAP-tau when used on MDA-MB-231 cells. Investigation of the cellular apoptotic pathways activated by α CSPG4(scFv)-MAP in MDA-MB-231 cells revealed no signal for caspase-3 and PARP-1 in treated cells, and a moderate activation of caspase 9 which was not sufficient to match the cytotoxic readouts of MAP tau [55]. A closer look confirmed the significant activation of the mitochondrial EndoG apoptotic pathway by α CSPG4(scFv)-MAP which is similar to that reported for taxanes. Our understanding of the apoptosis signalling pathways after induction of mitotic arrest is crucial to better optimizing human MAP tau based proteins for clinical use. Further studies are therefore required to fully demonstrate how MAP tau based human cytolytic fusion proteins induce cell death in targeted cells.

Finally, potential clinical applications of MAP tau based human cytolytic fusion proteins do not end with their anti-cancer properties. Several other diseases including atherosclerosis, rheumatoid arthritis, psoriasis, idiopathic pulmonary fibrosis, scleroderma and cirrhosis of the liver etc. are reportedly driven by cells undergoing excessive or dysregulated proliferation [59]. It is of general agreement that atherogenic proliferation of smooth muscle cells in the inner most layer (intima) of the arterial wall is a principal event that occurs during the early stages of the pathogenesis of atherosclerosis which would result in the formation of new connective tissue and intracellular and extracellular lipid deposit [59, 60]. Also, the chronic inflammatory skin diseases, psoriasis is characterized by the hyper-proliferation and aberrant differentiation of keratinocytes [61]. In a much recent publication by Hristodorov, Mladenov et al., the proliferative state of disease causing pro-inflammatory macrophages (M1) has also been reported in tissue sections from inflamed skins. This discovery became of significant importance in shedding more light to the current debate of whether macrophages proliferate *in situ*. It also allowed the evaluation and documentation of the first MAP tau based fusion protein, H22(scFv)-MAP for the treatment of diseases mediated by M1 pro-inflammatory macrophages. In the study, authors documented the proliferation dependent cytotoxicity of H22(scFv)-MAP towards the pro-monocytic cell line HL-60. And interestingly, H22(scFv)-MAP could not kill peritoneum-derived murine macrophages and human macrophages derived from peripheral blood mononuclear cells (PBMCs) because they are known to stop proliferating when isolated and cultivated *in vitro* [56]. On top of that, *ex vivo* data demonstrated that H22(scFv)-MAP recognizes efficiently CD64⁺ leukemic blasts leading to their elimination, while it spares the homeostasis of healthy CD64⁺ PBMC. Same study shows the ability of the human cytolytic fusion protein to kill leukemic cells independent of the target receptor profile, which suggest the high potency of MAP tau as cytostatic agent [57]. In context of leukaemia, similar results on the efficacy of MAP tau were obtained when evaluating CD89 as a target for immunotherapy (manuscript submitted). Taken together, these findings confirm the versatility and the therapeutic suitability of MAP-based immunotherapeutic agents for the treatment of a wide range of indications including immunological diseases.

3. Conclusion and future remarks

On a much broader landscape, the introduction of MAP tau has culminated into the creation of a wider portfolio of targeted human cytolytic fusion proteins, in addition to human granzymes and RNases. As described above, MAP tau based fusion proteins have demonstrated the preclinical potential and ability to compromise mitosis and inducing apoptosis in proliferating target cells. MAP-tau based fusion proteins offer two levels of protection by a) target cell surface antigen specificity and b) killing of proliferating cells. This would prevent healthy cells expressing tumor or disease markers at physiological levels from being affected. On the other hand, it is important to mention that studies aimed at evaluating the retention ability of MAP tau in target cells are still needed and crucial to the eventual success of MAP tau based fusion proteins. This is because cells in the G1 and S-phase of cell division possess the potential to repopulate cleared fractions once the drug clears since they are generally not affected by anti-mitotic drugs. The ability of paclitaxel to linger for about a week in tumor cells and by so doing confer longer cytotoxic effect in comparison to the ~13 hour half-life of newer drugs has been linked to its popularity and success [62]. Further studies along this line might thus help improve the effectiveness of MAP tau. However, MAP tau also has its own

advantage when compared with other anti-cancer or anti-mitotic agent. It is well tolerated *in vivo* as seen in the above findings, which could allow the administration of repeated doses. The human origin of MAP tau is expected to confer a low immunogenic response when compared with other treatment modalities. With the active removal of vital phosphorylation sites lower off-target toxicities should be associated with this emerging novel class of treatment.

Acknowledgments: R.M. was supported by a scholarship from the Jürgen-Manchot-Foundation, parts of this work were funded by the Fraunhofer project "The markets of the future – SKINHEAL", as well as the ForSaTum project, which was sponsored within the NRW-EU Ziel 2-Programm "Regionale Wettbewerbsfähigkeit und Beschäftigung 2007-20013" (EFRE).

Conflicts of Interest: DH, RM, and SB are inventors on a European Patent Application on MAP based targeted human cytolytic fusion proteins assigned to Fraunhofer. No potential conflicts of interest are disclosed by the other authors

References

1. Nagle A, Hur W, Gray NS. Antimitotic agents of natural origin. *Current drug targets*. 2006;7(3):305-26.
2. Penna LS, Henriques JAP, Bonatto D. Anti-mitotic agents: Are they emerging molecules for cancer treatment? *Pharmacology & Therapeutics*. 2017.
3. Prendergast GC, Jaffee EM. *Cancer immunotherapy: immune suppression and tumor growth*: Academic Press; 2013.
4. Mc Gee MM. Targeting the mitotic catastrophe signaling pathway in cancer. *Mediators of inflammation*. 2015;2015.
5. Fanale D, Bronte G, Passiglia F, Calò V, Castiglia M, Di Piazza F, et al. Stabilizing versus destabilizing the microtubules: a double-edge sword for an effective cancer treatment option? *Analytical cellular pathology*. 2015;2015.
6. Jordan M. Mechanism of action of antitumor drugs that interact with microtubules and tubulin. *Current Medicinal Chemistry-Anti-Cancer Agents*. 2002;2(1):1-17.
7. Sudakin V, Yen TJ. Targeting mitosis for anti-cancer therapy. *BioDrugs*. 2007;21(4):225-33.
8. Stanton RA, Gernert KM, Nettles JH, Aneja R. Drugs that target dynamic microtubules: a new molecular perspective. *Medicinal research reviews*. 2011;31(3):443-81.
9. Gascoigne KE, Taylor SS. How do anti-mitotic drugs kill cancer cells? *Journal of cell science*. 2009;122(15):2579-85.
10. van Vuuren RJ, Visagie MH, Theron AE, Joubert AM. Antimitotic drugs in the treatment of cancer. *Cancer chemotherapy and pharmacology*. 2015;76(6):1101-12.
11. Mukhtar E, Adhami VM, Mukhtar H. Targeting microtubules by natural agents for cancer therapy. *Molecular cancer therapeutics*. 2014;13(2):275-84.
12. Zhang B, Carroll J, Trojanowski JQ, Yao Y, Iba M, Potuzak JS, et al. The microtubule-stabilizing agent, epothilone D, reduces axonal dysfunction, neurotoxicity, cognitive deficits, and Alzheimer-like pathology in an interventional study with aged tau transgenic mice. *Journal of Neuroscience*. 2012;32(11):3601-11.
13. Vollmer LL, Jiménez M, Camarco DP, Zhu W, Daghestani HN, Balachandran R, et al. A Simplified Synthesis of Novel Dictyostatin Analogues with In Vitro Activity against Epothilone B-Resistant Cells and Antiangiogenic Activity in Zebrafish Embryos. *Molecular cancer therapeutics*. 2011;10(6):994-1006.

14. Peters C, Brown S. Antibody–drug conjugates as novel anti-cancer chemotherapeutics. *Bioscience reports*. 2015;35(4):e00225.
15. Norris B, Pritchard K, James K, Myles J, Bennett K, Marlin S, et al. Phase III comparative study of vinorelbine combined with doxorubicin versus doxorubicin alone in disseminated metastatic/recurrent breast cancer: National Cancer Institute of Canada Clinical Trials Group Study MA8. *Journal of clinical oncology*. 2000;18(12):2385-94.
16. Lewis WH, Elvin-Lewis MP. *Medical botany: plants affecting human health*: John Wiley & Sons; 2003.
17. Dumontet C, Jordan MA. Microtubule-binding agents: a dynamic field of cancer therapeutics. *Nature reviews Drug discovery*. 2010;9(10):790-803.
18. Hristodorov D, Mladenov R, Pardo A, Pham A, Huhn M, Fischer R, et al. Microtubule-associated protein tau facilitates the targeted killing of proliferating cancer cells in vitro and in a xenograft mouse tumour model in vivo. *British journal of cancer*. 2013;109(6):1570-8.
19. Dosio F, Brusa P, Cattel L. Immunotoxins and anticancer drug conjugate assemblies: the role of the linkage between components. *Toxins*. 2011;3(7):848-83.
20. Klute K, Nackos E, Tasaki S, Nguyen DP, Bander NH, Tagawa ST. Microtubule inhibitor-based antibody–drug conjugates for cancer therapy. *OncoTargets and therapy*. 2014;7:2227.
21. de Claro RA, McGinn K, Kwitkowski V, Bullock J, Khandelwal A, Habtemariam B, et al. US Food and Drug Administration approval summary: brentuximab vedotin for the treatment of relapsed Hodgkin lymphoma or relapsed systemic anaplastic large-cell lymphoma. *Clinical Cancer Research*. 2012;18(21):5845-9.
22. Mohan R, John A. Microtubule-associated proteins as direct crosslinkers of actin filaments and microtubules. *IUBMB life*. 2015;67(6):395-403.
23. Hayden JH, Bowser SS, Rieder CL. Kinetochores capture astral microtubules during chromosome attachment to the mitotic spindle: direct visualization in live newt lung cells. *J Cell Biol*. 1990;111(3):1039-45.
24. Billingsley ML, Kincaid RL. Regulated phosphorylation and dephosphorylation of tau protein: effects on microtubule interaction, intracellular trafficking and neurodegeneration. *Biochemical Journal*. 1997;323(3):577-91.
25. Cambiazo V, González M, Maccioni RB. DMAP-85: A τ -Like Protein from *Drosophila melanogaster* Larvae. *Journal of neurochemistry*. 1995;64(3):1288-97.
26. Kosik KS, Finch EA. MAP2 and tau segregate into dendritic and axonal domains after the elaboration of morphologically distinct neurites: an immunocytochemical study of cultured rat cerebrum. *Journal of Neuroscience*. 1987;7(10):3142-53.
27. Nelson PT, Stefansson K, Gulcher J, Saper CB. Molecular evolution of τ protein: implications for Alzheimer's disease. *Journal of neurochemistry*. 1996;67(4):1622-32.
28. Liu Y, Xia J, Ma D, Faber DS, Fischer I. Tau-like proteins in the nervous system of goldfish. *Neurochemical research*. 1997;22(12):1511-6.
29. Goedert M, Spillantini M, Jakes R, Rutherford D, Crowther R. Multiple isoforms of human microtubule-associated protein tau: sequences and localization in neurofibrillary tangles of Alzheimer's disease. *Neuron*. 1989;3(4):519-26.

30. Cleveland DW, Hwo S-Y, Kirschner MW. Purification of tau, a microtubule-associated protein that induces assembly of microtubules from purified tubulin. *Journal of molecular biology*. 1977;116(2):207-25.
31. Weingarten MD, Lockwood AH, Hwo S-Y, Kirschner MW. A protein factor essential for microtubule assembly. *Proceedings of the National Academy of Sciences*. 1975;72(5):1858-62.
32. Schoenfeld TA, Obar RA. Diverse distribution and function of fibrous microtubule-associated proteins in the nervous system. *International review of cytology*. 1994;151:67-137.
33. Gu Y, Oyama F, Ihara Y. τ is widely expressed in rat tissues. *Journal of neurochemistry*. 1996;67(3):1235-44.
34. Michalik L, Neuville P, Vanier M, Launay J. Pancreatic tau related maps: biochemical and immunofluorescence analysis in a tumoral cell line. *Molecular and cellular biochemistry*. 1995;143(2):107-12.
35. Cleveland DW, Hwo S-Y, Kirschner MW. Physical and chemical properties of purified tau factor and the role of tau in microtubule assembly. *Journal of molecular biology*. 1977;116(2):227-47.
36. Himmler A, Drechsel D, Kirschner MW, Martin D. Tau consists of a set of proteins with repeated C-terminal microtubule-binding domains and variable N-terminal domains. *Molecular and Cellular Biology*. 1989;9(4):1381-8.
37. Kolarova M, García-Sierra F, Bartos A, Ricny J, Ripova D. Structure and pathology of tau protein in Alzheimer disease. *International journal of Alzheimer's disease*. 2012;2012.
38. Brandt R, Léger J, Lee G. Interaction of tau with the neural plasma membrane mediated by tau's amino-terminal projection domain. *The Journal of cell biology*. 1995;131(5):1327-40.
39. Drechsel DN, Hyman A, Cobb MH, Kirschner M. Modulation of the dynamic instability of tubulin assembly by the microtubule-associated protein tau. *Molecular biology of the cell*. 1992;3(10):1141-54.
40. Al-Bassam J, Ozer RS, Safer D, Halpain S, Milligan RA. MAP2 and tau bind longitudinally along the outer ridges of microtubule protofilaments. *The Journal of cell biology*. 2002;157(7):1187-96.
41. Caceres A, Kosik KS. Inhibition of Neurite Polarity by Tau Antisense Oligonucleotides in Primary Cerebellar Neurons. *Nature*. 1990;343(6257):461.
42. Harada A, Oguchi K, Okabe S, Kuno J, Terada S, Ohshima T, et al. Altered microtubule organization in small-calibre axons of mice lacking tau protein. *Nature*. 1994;369(6480):488.
43. Goedert M, Jakes R. Expression of separate isoforms of human tau protein: correlation with the tau pattern in brain and effects on tubulin polymerization. *The EMBO journal*. 1990;9(13):4225.
44. Kanemaru K, Takio K, Miura R, Titani K, Ihara Y. Fetal-Type Phosphorylation of the τ in Paired Helical Filaments. *Journal of neurochemistry*. 1992;58(5):1667-75.
45. Trinczek B, Biernat J, Baumann K, Mandelkow E-M, Mandelkow E. Domains of tau protein, differential phosphorylation, and dynamic instability of microtubules. *Molecular biology of the cell*. 1995;6(12):1887-902.
46. Garcia ML, Cleveland DW. Going new places using an old MAP: tau, microtubules and human neurodegenerative disease. *Current opinion in cell biology*. 2001;13(1):41-8.
47. Lee VM, Goedert M, Trojanowski JQ. Neurodegenerative tauopathies. *Annual review of neuroscience*. 2001;24(1):1121-59.

48. Bramblett GT, Goedert M, Jakes R, Merrick SE, Trojanowski JQ, Lee VM. Abnormal tau phosphorylation at Ser 396 in Alzheimer's disease recapitulates development and contributes to reduced microtubule binding. *Neuron*. 1993;10(6):1089-99.
49. Wasiak T, Ionov M, Nieznanski K, Nieznanska H, Klementieva O, Granell M, et al. Phosphorus dendrimers affect Alzheimer's (A β 1-28) peptide and MAP-Tau protein aggregation. *Molecular pharmaceutics*. 2012;9(3):458-69.
50. Pasquier E, Carré M, Pourroy B, Camoin L, Rebaï O, Briand C, et al. Antiangiogenic activity of paclitaxel is associated with its cytostatic effect, mediated by the initiation but not completion of a mitochondrial apoptotic signaling pathway. *Molecular Cancer Therapeutics*. 2004;3(10):1301-10.
51. Jung J. Human tumor xenograft models for preclinical assessment of anticancer drug development. *Toxicological Res*. 2014;30(1):1-5.
52. Hristodorov D, Nordlohne J, Mladenov R, Huhn M, Fischer R, Thepen T, et al. Human microtubule-associated protein tau mediates targeted killing of CD30+ lymphoma cells in vitro and inhibits tumour growth in vivo. *British journal of haematology*. 2014;164(2):251-7.
53. Hristodorov D, Amoury M, Mladenov R, Niesen J, Arens K, Berges N, et al. EpCAM-selective elimination of carcinoma cells by a novel MAP-based cytolytic fusion protein. *Molecular cancer therapeutics*. 2014;13(9):2194-202.
54. Brehm H, Hristodorov D, Pardo A, Mladenov R, Niesen J, Fischer R, et al. Targeted killing of rhabdomyosarcoma cells by a MAP-based human cytolytic fusion protein. *Cancer letters*. 2015;365(2):149-55.
55. Amoury M, Mladenov R, Nachreiner T, Pham AT, Hristodorov D, Di Fiore S, et al. A novel approach for targeted elimination of CSPG4-positive triple-negative breast cancer cells using a MAP tau-based fusion protein. *International journal of cancer*. 2016;139(4):916-27.
56. Hristodorov D, Mladenov R, Fischer R, Barth S, Thepen T. Fully human MAP-fusion protein selectively targets and eliminates proliferating CD64+ M1 macrophages. *Immunology and cell biology*. 2016;94(5):470-8.
57. Mladenov R, Hristodorov D, Cremer C, Gresch G, Grieger E, Schenke L, et al. CD64-directed microtubule associated protein tau kills leukemic blasts ex vivo. *leukemia (CML)*. 2016;1:3.
58. Hetzel C, Bachran C, Fischer R, Fuchs H, Barth S, Stöcker M. Small cleavable adapters enhance the specific cytotoxicity of a humanized immunotoxin directed against CD64-positive cells. *Journal of immunotherapy*. 2008;31(4):370-6.
59. Sporn MB, Harris ED. Proliferative diseases. *The American journal of medicine*. 1981;70(6):1231-6.
60. Ross R, Glomset J, Harker L. Response to injury and atherogenesis. *The American journal of pathology*. 1977;86(3):675.
61. Cai Y, Fleming C, Yan J. New insights of T cells in the pathogenesis of psoriasis. *Cellular & molecular immunology*. 2012;9(4):302-9.
62. Chan K, Koh CG, Li H. Mitosis-targeted anti-cancer therapies: where they stand. *Cell death & disease*. 2012;3(10):e411.