

# Subterranean Karst Environments as a Global Sink for Atmospheric Methane

Kevin D. Webster<sup>1,2,\*</sup>, Agnieszka Drobniak<sup>3</sup>, Giuseppe Etiope<sup>4,5</sup>, Maria Mastalerz<sup>3</sup>, Peter E. Sauer<sup>6</sup>, Arndt Schimmelmann<sup>6</sup>

<sup>1</sup>School of Natural Resources and the Environment, University of Arizona, 1064 E Lowell St, Tucson, Arizona 85721, USA

<sup>2</sup>Department of Ecology and Evolutionary Biology, University of Arizona, 1041 E Lowell St, Tucson, Arizona 85721, USA.

<sup>3</sup>Indiana Geological Survey, Indiana University, 611 North Walnut Grove Ave., Bloomington, Indiana 47405, USA.

<sup>4</sup>Istituto Nazionale di Geofisica e Vulcanologia, Sezione Roma 2, Italy.

<sup>5</sup>Faculty of Environmental Science and Engineering, Babes-Bolyai University, Cluj-Napoca, Romania.

<sup>6</sup>Department of Earth and Atmospheric Sciences, Indiana University, 1001 E 10<sup>th</sup> St. Bloomington, IN 47405, USA.

\* corresponding author: kevdwebs@email.arizona.edu; telephone: 520-621-2539.

**Abstract:** The air in subterranean karst cavities is often depleted in methane (CH<sub>4</sub>) relative to the atmosphere. Karst is considered a potential sink for the atmospheric greenhouse gas CH<sub>4</sub> because its subsurface drainage networks and solution-enlarged fractures facilitate atmospheric exchange. Karst landscapes cover about 14 % of earth's continental surface, but observations of CH<sub>4</sub> concentrations in cave air are limited to localized studies in Gibraltar, Spain, Indiana (USA), Vietnam, Australia, and by incomplete isotopic data. To test if karst is systematically acting as a global CH<sub>4</sub> sink, we measured the CH<sub>4</sub> concentrations,  $\delta^{13}\text{C}_{\text{CH}_4}$ , and  $\delta^2\text{H}_{\text{CH}_4}$  values of cave air from 33 caves in the USA and three caves in New Zealand. We also measured CO<sub>2</sub> concentrations,  $\delta^{13}\text{C}_{\text{CO}_2}$ , and radon (Rn) concentrations to support CH<sub>4</sub> data interpretation by assessing cave air residence times and mixing processes. Among these caves, 35 exhibited subatmospheric CH<sub>4</sub> concentrations in at least one location compared to their local atmospheric backgrounds. CH<sub>4</sub> concentrations and  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  values suggest that microbial methanotrophy within caves is the primary CH<sub>4</sub> consumption mechanism as the atmosphere exchanges with subsurface air. The pattern of  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  values along CH<sub>4</sub> concentration gradients in cave air provides evidence for incomplete oxidation by methanotrophy. Only 5 locations from 3 caves showed elevated CH<sub>4</sub> concentrations compared to the atmospheric background and could be ascribed to local CH<sub>4</sub> sources from sewage and outgassing swamp water. Several associated  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  values point to carbonate reduction and acetate fermentation as biochemical pathways of limited methanogenesis in karst environments and suggest that these pathways occur in the environment over large spatial scales. Our data show that karst environments function as a global CH<sub>4</sub> sink. Estimates of CH<sub>4</sub> flux in karst landscapes are needed in order to include the subterranean CH<sub>4</sub> sink in climate models.

**Keywords:** cave; greenhouse gas; karst; methane; methanogenesis; methanotrophy

## 1. Introduction

Atmospheric methane ( $\text{CH}_4$ ) is a greenhouse gas and its concentration is increasing in the atmosphere (Dlugokencky et al., 2011; Sussmann et al., 2012; Ciais et al., 2013). The increase in atmospheric  $\text{CH}_4$  is due to an imbalance between  $\text{CH}_4$  sources and sinks. Anthropogenic and natural sources combine to contribute about 680 Tg of  $\text{CH}_4$  to the atmosphere while reactions with hydroxyl ( $\cdot\text{OH}$ ) and chlorine radicals in the troposphere and stratosphere remove about 600 Tg  $\text{a}^{-1}$  (Kirschke et al., 2013). Methanotrophic consumption in soils removes 30 Tg  $\text{a}^{-1}$  (Kirschke et al., 2013). The present globally averaged  $\text{CH}_4$  concentration is 1.87 ppmv which is 2.5 times higher than preindustrial levels (Nisbet et al., 2016). Despite improvements in estimating individual sources and sinks of atmospheric  $\text{CH}_4$ , the associated errors remain large (Kirschke et al., 2013). Recent studies suggest that caves may act as an additional  $\text{CH}_4$  sink (Mattey et al., 2013; Fernandez-Cortes et al., 2015; McDonough et al., 2016; Webster et al., 2016; Lennon et al., 2017).

Caves and associated karst landscapes may be an important overlooked sink for atmospheric  $\text{CH}_4$  because they are estimated to cover as much as 10 to 20 % of the continental surface with the more precise estimates suggesting about 13.8 % (Palmer, 1991; Ford and Williams, 2007). Karst landscapes are frequently associated with the chemical dissolution of limestones, but can form in any soluble rock body. The resulting caves, solution-enlarged fractures, and internal drainage networks that function to transport mass from high elevations to low elevations also allow for subsurface-surface atmospheric exchange (Kowalczyk and Froelich, 2010; Garcia-Anton et al., 2014). The total volume and surface area of karst conduits able to interact with the atmosphere is unknown, in part due to small fractures and the difficulty of imaging the subsurface with geophysical methods. Karst caves, due to their accessibility, provide opportunities for non-invasive, *in-situ* analyses and sampling.

Cave and karst landscapes form in two common ways, each of which influences karst's capacity to act as a  $\text{CH}_4$  sink. Epigenic karst forms through the interaction of limestone with carbonic acid derived from the dissolution of atmospheric and soil  $\text{CO}_2$  into surface waters, and hypogenic caves form when corrosive water from deep sources migrates into and dissolves limestone bedrock. Epigenic caves are more widespread, and atmospheric to subatmospheric  $\text{CH}_4$  concentrations of 1.8 ppmv to  $<0.1$  ppmv have been observed in these settings (Mattey et al., 2013; Fernandez-Cortes et al., 2015; McDonough et al., 2016; Webster et al., 2016; Lennon et al., 2017). In some hypogenic caves by contrast, elevated  $\text{CH}_4$  concentrations from 2 ppmv to 1 % have been observed in association with  $\text{CH}_4$ -rich springs or seeps related to fluid migration from deep hydrocarbon-bearing sedimentary rocks, i.e. seepage processes that are widespread on Earth (Sarbu et al., 1996; Hutchens et al., 2004; Jones et al., 2012; Webster et al., 2017). The dominance of epigenic karst suggests that karst is likely functioning as a  $\text{CH}_4$  sink at the global scale, but more observations are needed.

Different hypotheses have been put forward to explain the low  $\text{CH}_4$  concentrations observed in epigenic cave air. The combination of subatmospheric  $\text{CH}_4$  concentrations and the stable carbon isotopic ratio of  $\text{CH}_4$  in the air of caves in Gibraltar led to the hypothesis that microorganisms were responsible for the removal of  $\text{CH}_4$  (Mattey et al., 2013). In turn, low  $\text{CH}_4$  concentrations in Spanish caves, in the presumed absence of  $\text{CH}_4$ -consuming (methanotrophic) bacteria, led to the hypothesis that  $\text{CH}_4$  oxidation was induced by ions and  $\cdot\text{OH}$  generated by the radioactive decay of radon and daughter nuclides (Fernandez-Cortes et al., 2015). Since these initial observations, datasets from caves in Australia, the USA, and Vietnam have pointed towards methanotrophic  $\text{CH}_4$  oxidation (McDonough et al., 2016; Webster et al., 2016; Lennon et al., 2017).

The chemical composition of cave air results from the mixing of the atmosphere and air from the overlying soils and epikarst and these processes should influence the  $\text{CH}_4$  concentrations of cave air. Previous studies have shown that  $\text{CH}_4$  concentrations have been inversely correlated with  $\text{CO}_2$  concentrations in cave air (Mattey et al., 2013; Fernandez-Cortes et al., 2015; McDonough et al., 2016;

Webster et al., 2016). Cave air CO<sub>2</sub> concentrations are positively correlated with radon (Rn) concentrations and Rn is known to track cave air residence time (Cunningham and LaRock, 1991; Batiot-Guilhe et al., 2007; Kowalczyk and Froelich, 2010; Matthey et al., 2010; Gregorič et al., 2011, 2014). Additionally, the stable C isotope composition of CO<sub>2</sub> ( $\delta^{13}\text{C}_{\text{CO}_2}$ ), can track the sources of CO<sub>2</sub> in the environment. For example,  $\delta^{13}\text{C}_{\text{CO}_2}$  values of -24 ‰ are associated with soil CO<sub>2</sub>, while atmospheric CO<sub>2</sub> has  $\delta^{13}\text{C}_{\text{CO}_2}$  values ranging from -8.5 ‰ to -10 ‰ (Amundson et al., 1998; Keeling et al., 2010; Peyraube et al., 2013). Thus CO<sub>2</sub>, Rn, and  $\delta^{13}\text{C}_{\text{CO}_2}$  in cave air can help determine the influence of cave air mixing processes on CH<sub>4</sub>.

The stable C and H isotope compositions of CH<sub>4</sub> ( $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$ ) also provide tools for understanding the sources and sinks of CH<sub>4</sub> in caves because different CH<sub>4</sub> sources are associated with characteristic  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  values. For example, CH<sub>4</sub> produced from carbonate reduction has  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  values that range from -112 to -60 ‰ VPDB and from -350 to -100 ‰ VSMOW respectively (Whiticar, 1999), and atmospheric CH<sub>4</sub> has  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  values around -47.5 and -100 ‰ (Miller et al., 2002; Townsend-Small et al., 2012). The  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  values of CH<sub>4</sub> can also be altered through secondary processes such as oxidation and mixing. The oxidation pathways of CH<sub>4</sub> by methanotrophs or ·OH have fractionation factors that result in residual CH<sub>4</sub> having increased in  $\delta^2\text{H}_{\text{CH}_4}$  values of 8.5 ‰ for every 1 ‰ increase in  $\delta^{13}\text{C}_{\text{CH}_4}$  value and increases in  $\delta^2\text{H}_{\text{CH}_4}$  values of 72 ‰ for every 1 ‰ increase in  $\delta^{13}\text{C}_{\text{CH}_4}$  value, respectively (Feisthauer et al., 2011; Saueressig et al., 2001). Mixing between two different CH<sub>4</sub> sources creates a linear trend between the two members. Thus measuring the  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  of cave air should allow for the determination of cave air CH<sub>4</sub> sources.

The objective of the present work is to extend the karst CH<sub>4</sub> dataset and test the hypothesis that karst systems act as a CH<sub>4</sub> sink on a global scale. To this aim, we studied CH<sub>4</sub> concentration,  $\delta^{13}\text{C}_{\text{CH}_4}$ , and  $\delta^2\text{H}_{\text{CH}_4}$  in cave air from 33 epigenic caves in the USA and three epigenic caves in New Zealand. CO<sub>2</sub>,  $\delta^{13}\text{C}_{\text{CO}_2}$ , and Rn were also measured to support CH<sub>4</sub> data interpretation *via* assessing cave air residence times and mixing processes. Data analysis is focused on determining CH<sub>4</sub> concentrations, origin, mixing processes and isotopic fractionations.

## 2. Methods

### 2.1. Sampling and analyses

Air samples from 16 limestone or dolostone caves in the Appalachian fold and thrust belt, 17 limestone caves in gently warped intracratonic basins of the USA, and 3 caves from the North Island of New Zealand were collected over a timespan of roughly four years (Fig. 1; Table 1). Cave air was analyzed using *in-situ* methods and was further sampled for laboratory analysis. *In-situ* CH<sub>4</sub>, CO<sub>2</sub>, and Rn abundance analyses were carried out using a suite of instruments (Table 2). Discrete samples of cave air were collected in pre-evacuated 50-mL serum vials, in 1 to 3-L Tedlar® bags, or in 4-L glass bottles. CH<sub>4</sub> and CO<sub>2</sub> concentrations of discrete samples were measured *via* gas chromatography.

We assessed cave air mixing processes through the following techniques. A qualitative estimate on cave air residence time was obtained by comparing CH<sub>4</sub> to CO<sub>2</sub> concentrations at individual locations in each cave. Additionally, we measured the Rn concentrations of caves 32 through 36 to assess the relationship between cave air residence time, CH<sub>4</sub> concentrations, and CO<sub>2</sub> concentrations.  $\delta^{13}\text{C}_{\text{CO}_2}$  data were used to assess the sources of CO<sub>2</sub> and thus of air entering the caves. We also assessed the distance from each sampling location to cave entrances as another tool to understand the sources and sinks of CH<sub>4</sub> in caves.

CH<sub>4</sub> and CO<sub>2</sub> concentrations from discrete air samples were measured at Indiana University using a Varian 450 gas chromatograph (GC) (Varian – Agilent Technologies, Palo Alto, California). The GC was fitted with a flame ionization detector (FID) for CH<sub>4</sub> and a thermal conductivity detector (TCD) for CO<sub>2</sub>. Standard gas mixtures from Air Liquide America Specialty Gasses LLC (Plumsteadville, Pennsylvania) were used for 3-point calibration curves to convert signals measured on the GC to concentrations. CH<sub>4</sub> standards measured on the GC had errors of  $\pm 5$  to  $\pm 14$  % of the reported concentrations. CH<sub>4</sub> concentrations are reported with the uncertainty associated with the standard curve unless the calculated uncertainty was  $\leq 0.1$  ppmv. Samples with calculated uncertainties  $\leq 0.1$  ppmv were assigned uncertainties of 0.1 ppmv based on replicate measurements. The uncertainty associated with standard curves for CO<sub>2</sub> concentrations varied from  $< \pm 1$  to 5 %. CO<sub>2</sub> concentrations were assigned uncertainties based on their associated standard curve.

The stable carbon isotope ratios of CH<sub>4</sub> and CO<sub>2</sub> and hydrogen stable isotope ratios of CH<sub>4</sub> were measured on a ThermoFinnigan Delta Plus XP mass spectrometer in the Stable Isotope Research Facility at Indiana University. Carbon stable isotope ratios are expressed as conventional  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^{13}\text{C}_{\text{CO}_2}$  values in ‰ along the scale anchored to Vienna Pee Dee Belemnite (VPDB). Hydrogen stable isotope ratios are expressed as  $\delta^2\text{H}_{\text{CH}_4}$  values in ‰ along the scale anchored to Vienna Standard Mean Ocean Water (VSMOW). CH<sub>4</sub> samples were measured in continuous-flow mode using CH<sub>4</sub> preconcentration, cryofocusing (Miller et al., 2002), and a gas chromatography-oxidation/pyrolysis-isotope ratio mass spectrometer (GC-ox/pyr-IRMS) interface. Varying sample extraction times were used to isolate roughly 0.45 and 0.90  $\mu\text{mol}$  of CH<sub>4</sub> prior to the introduction of the sample to the GC-ox/pyr-IRMS for analysis of  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  values, respectively. In-house CH<sub>4</sub> standards methane #3, methane #6, and methane ALM with  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  values of  $[+19.86 \pm 0.05; +2.2 \pm 1.2]$  ‰,  $[-39.40 \pm 0.02; -153 \pm 2]$  ‰, and  $[-58.6 \pm 0.03; -272.2 \pm 3.4]$  ‰ were used for 2-point normalizations. Errors associated with  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  values were calculated using a standard curve that accounted for the peak size of the measurement. Analytical repeatability of internal standards ranged from 0.14 to 0.6 ‰ for  $\delta^{13}\text{C}_{\text{CH}_4}$  and from 7 to 18 ‰ for  $\delta^2\text{H}_{\text{CH}_4}$ .

$\delta^{13}\text{C}_{\text{CO}_2}$  values were measured in continuous-flow mode using a GasBench II inlet (Tu et al., 2001). Measured  $^{13}\text{C}/^{12}\text{C}$  ratios of CO<sub>2</sub> from cave air were converted to the VPDB scale using a single isotopically characterized in-house standard that has a value of  $12.0 \pm 0.2$  ‰.

## 2.2 Data elaboration and quality control

*In-situ* measurements were preferentially used when statistically analyzing gas concentration data. When *in-situ* measurements were not available, concentrations measured on the GC were used in the statistical analyses. Samples and data were screened for quality control by comparing the samples with *in-situ* measurements and visual estimation of the volume of sample bags. If a sample bag had been shown to exhibit a leak for one analyte, data from that sample were discarded. CH<sub>4</sub> and CO<sub>2</sub> concentrations measured by both GC-FID and FTIR showed strong agreements ( $\text{GC}(\text{CO}_2) = 0.92 \pm 0.04 * \text{FTIR}(\text{CO}_2) + 100 \pm 300$ ,  $r^2 = 0.99$ ,  $p = 5 * 10^{-19}$ ;  $\text{GC}(\text{CH}_4) = 0.7 \pm 0.2 * \text{FTIR}(\text{CH}_4) + 0.2 \pm 0.2$ ,  $r^2 = 0.62$ ), and the stable isotopic composition of the samples was not related to their storage time ( $\delta^{13}\text{C}_{\text{CH}_4} = 0.2 \pm 0.3 * \text{day} - 47 \pm 2$ ,  $r^2 = 0.03$ ,  $p = 0.32$ ;  $\delta^2\text{H}_{\text{CH}_4} = 0.03 \pm 0.14 * \text{day} - 96 \pm 11$ ,  $r^2 = 0.005$ ,  $p = 0.72$ ;  $\delta^{13}\text{C}_{\text{CO}_2} = -0.05 \pm 0.16 * \text{day} - 19.9 \pm 1.8$ ,  $r^2 = 0.006$ ,  $p = 0.57$ ). In locations where more than one sample was taken with *in-situ* methods, the values of the samples were averaged. In total, 199 CH<sub>4</sub> concentrations, 192 CO<sub>2</sub> concentrations, 32  $\delta^{13}\text{C}_{\text{CH}_4}$  values, 26  $\delta^2\text{H}_{\text{CH}_4}$  values, and 60  $\delta^{13}\text{C}_{\text{CO}_2}$  values are reported in this study (Supplemental Tables 1, 2). All samples are reported with 95 % confidence intervals.

Two different modeling techniques were used to assess trace gas sources and sinks in the studied caves. Keeling plots were used to assess the possibility of a two end member mixing system affecting  $\delta^{13}\text{C}_{\text{CO}_2}$ . The stable isotopic composition of  $\text{CO}_2$  entering the caves ( $\delta^{13}\text{C}_s$ ) was assessed through equation 1

$$\delta^{13}\text{C}_s = (\delta^{13}\text{C}_m - F_{\text{atm}} * \delta^{13}\text{C}_{\text{atm}}) (1 - F_{\text{atm}})^{-1} \quad (1)$$

where  $\delta^{13}\text{C}_m$  is the  $\delta^{13}\text{C}_{\text{CO}_2}$  of the sample,  $\delta^{13}\text{C}_{\text{atm}}$  is the  $\delta^{13}\text{C}_{\text{CO}_2}$  of the atmosphere, and  $F_{\text{atm}}$  is the fraction of atmospheric  $\text{CO}_2$  in the  $\text{CO}_2$  concentration of the sample (Peyraube et al., 2016). We used values of  $-10\text{‰}$  for  $\delta^{13}\text{C}_{\text{atm}}$  and 400 ppmv for the concentration of atmospheric  $\text{CO}_2$ .

Rayleigh distillation models were used as the theoretical basis to examine changes in the stable isotopic composition of  $\text{CH}_4$  in cave air caused by methanotrophy or  $\cdot\text{OH}$  oxidation. The  $\delta$ -value of an isotope system in a chemical compound of interest (e.g.,  $\text{CH}_4$ ) in cave air can be modeled as

$$\delta_c = (\delta_i + 1000) f^{-(\alpha+1)} - 1000 \quad (2)$$

where  $\delta_c$  is the instantaneous  $\delta$ -value of a particular isotope system in cave air after partial consumption,  $\delta_i$  is the initial  $\delta$ -value of the isotope system in cave air,  $f$  is the fraction of the compound remaining, and  $\alpha$  is the kinetic isotope fractionation factor (Mattey et al., 2013).  $\alpha$  values of 1.018 and 1.1353 were used to model changes in  $\delta^{13}\text{CCH}_4$  and  $\delta^2\text{HCH}_4$  caused by methanotrophy (Coleman et al., 1981; Feisthauer et al., 2011).  $\alpha$  values for changes in  $\delta^{13}\text{CCH}_4$  caused by methanotrophy have been observed to range from 1.003 to 1.039 (Templeton et al., 2006; Feisthauer et al., 2011); 1.018 was selected based on observations of methanotrophy in soils and its similarity to the  $\alpha$  value of 1.012 observed in St. Michael's Cave in Gibraltar (Feisthauer et al., 2011; Mattey et al., 2013).  $\alpha$  values of 1.0039 and 1.294 were used to model changes in  $\delta^{13}\text{CCH}_4$  and  $\delta^2\text{HCH}_4$  values caused by  $\cdot\text{OH}$  oxidation (Saueressig et al., 2001). The initial stable isotopic composition of atmospheric  $\text{CH}_4$  was modeled with a  $\delta^{13}\text{CCH}_4 = -47.5\text{‰}$  (VPDB), and  $\delta^2\text{HCH}_4 = -100\text{‰}$  (VSMOW) based on the work of Townsend-Small et al. (2012).

### 3. Results

Each of the 36 caves showed atmospheric to subatmospheric  $\text{CH}_4$  concentrations in at least one location. Only five locations from three different caves showed elevated  $\text{CH}_4$  concentrations relative to the surface atmosphere (Fig. 2, Supplemental Table 1). The  $\text{CH}_4$  concentration in the surface atmosphere at study sites ranged from  $1.8 \pm 0.3$  to  $2.8 \pm 0.7$  ppmv.  $\text{CH}_4$  concentrations in cave air ranged from  $\leq 0.1 \pm 0.1$  ppmv to  $5 \pm 1$  ppmv, and were generally observed to decrease with the distance from cave entrances (Fig. 3, Supplemental Table 1). Two thirds of the caves where three or more air measurements and distance data were recorded showed decreases in  $\text{CH}_4$  concentration from cave entrances to interiors. For example, caves 7, 8, and 9 from Kentucky all showed progressive decreases in  $\text{CH}_4$  concentration from about 2 ppmv at the entrance of the cave, down to zero or near zero ppmv in the more inner rooms (from 2 to 0 in caves 8 and 9 and from 1.9 to 0.3 ppmv in cave 7). Additionally,  $\text{CH}_4$  concentrations were negatively correlated with  $\text{CO}_2$  concentrations in cave air following an inverse power law relationship ( $[\text{CH}_4] = 17.5[\text{CO}_2]^{-0.41}$ ,  $r^2 = 0.26$ ) (Fig. 2). In the caves where Rn concentrations were measured, the average  $\text{CO}_2$  concentration of cave air was correlated with the average Rn concentration of cave air ( $[\text{CO}_2] = (1.42 \pm 0.09)[\text{Rn}] + 400 \pm 120$ ,  $n = 4$ ,  $r^2 = 0.99$ ,  $p = 0.009$ ).

Values of  $\delta^{13}\text{C}_{\text{CO}_2}$  in cave air ranged from  $-10.7 \pm 0.4$  to  $-23.81 \pm 0.10\text{‰}$ . Analysis of  $\delta^{13}\text{C}_{\text{CO}_2}$  values from samples with  $\text{CO}_2$  concentrations above 600 ppmv showed that  $\delta^{13}\text{C}_s$  ranging from  $-28\text{‰}$  to  $-20\text{‰}$  contributed to the composition of  $\text{CO}_2$  in cave air. Pooled analysis of the  $\text{CO}_2$  dataset shows that the average apparent source  $\delta^{13}\text{C}_{\text{CO}_2}$  value is  $-23.3 \pm 0.5\text{‰}$  ( $\delta^{13}\text{C}_{\text{CO}_2} = 4600 [\text{CO}_2]^{-1} - 23.3 \pm 0.5\text{‰}$ ,  $r^2 = 0.83$ ) (Fig. 4).

Values of  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  in cave air ranged from to  $-57.2 \pm 0.6$  to  $-27.1 \pm 0.2$  ‰ and  $-196 \pm 10$  to  $+2 \pm 18$  ‰, respectively. Keeling plots of  $\delta^{13}\text{C}_{\text{CH}_4}$  did not suggest that a two end member model was an adequate fit for the system ( $\delta^{13}\text{C}_{\text{CH}_4}$  vs.  $[\text{CH}_4]^{-1}$ :  $r^2 = 0.05$ ,  $\delta^2\text{H}_{\text{CH}_4}$  vs.  $[\text{CH}_4]^{-1}$ :  $r^2 = 0.12$ ). Some cave air samples plotted near the theoretical relationship between  $\delta^{13}\text{C}_{\text{CH}_4}$  values and  $\text{CH}_4$  concentrations caused by methanotrophy (Fig. 5). However, many points fell below and to the left of the line representing the theoretical incomplete oxidation of atmospheric  $\text{CH}_4$  (Figs. 5, 6). When  $\delta^2\text{H}_{\text{CH}_4}$  and  $\delta^{13}\text{C}_{\text{CH}_4}$  values were plotted against each other, many samples clustered tightly near the signature of atmospheric  $\text{CH}_4$  (Fig. 7). Some points, like those from caves 25 and 26, plotted near the expected trend of partial atmospheric  $\text{CH}_4$  oxidation by methanotrophy. Other points plotted below or above this trend and indicated that at least two additional sources of  $\text{CH}_4$  had to be entering the caves.

## 4. Discussion

### 4.1. Subsurface-Surface Atmospheric Exchange

The concentrations and stable isotopic compositions of  $\text{CH}_4$ ,  $\text{CO}_2$ ,  $\text{Rn}$  in cave air overlapped and diverged from those of the surface atmosphere. This suggests that atmospheric and cave processes influenced the composition of cave air. The atmospheric  $\text{CH}_4$  concentration at many study locations was above the globally averaged atmospheric background concentrations (1.87 ppmv  $\text{CH}_4$ ; Ciais et al., 2013) probably because they were near roads or pastures with local  $\text{CH}_4$  sources (Gioli et al., 2012; Harper et al., 2014).  $\text{Rn}$  concentrations were positively correlated with  $\text{CO}_2$  concentrations in cave air which agrees with other observations that have shown that  $\text{CO}_2$  concentrations in cave air track cave air residence time. The majority of cave air samples were depleted in  $\text{CH}_4$  and enriched in  $\text{CO}_2$  relative to the atmosphere, pointing to processes like in-situ  $\text{CH}_4$  oxidation and diffusion of air from the epikarst to decrease  $\text{CH}_4$  and increase  $\text{CO}_2$  concentrations (Fig. 2). Additionally, cave air  $\text{CH}_4$  concentrations generally decreased as the distance from an entrance increased (Fig. 3), and departures from this trend can be explained by fast airflow (caves 13, 25), cave air flowing out of the entrance (cave 26), multiple entrances resulting in multiple flow paths (cave 24), distance scales that were too small to observe a decrease in  $\text{CH}_4$  concentration (cave 20), or internal  $\text{CH}_4$  sources (cave 9). The patterns of  $\text{CH}_4$ ,  $\text{CO}_2$ , and  $\text{Rn}$  concentration indicate that  $\text{CH}_4$  concentrations decrease with cave air residence time.

The concentration and stable isotopic composition of  $\text{CO}_2$  in cave air show that in addition to an atmospheric end member,  $\text{CO}_2$  with  $\delta^{13}\text{C}_s$  values ranging from  $-28$  to  $-20$  ‰ were entering the caves and that the average source  $\delta^{13}\text{C}_{\text{CO}_2}$  value was  $-23.3 \pm 0.5$  ‰ (Fig. 4). These observations generally agree with a  $\text{CO}_2$  source from the soils and an atmospheric end-member. The  $\delta^{13}\text{C}_{\text{CO}_2}$  values of many soils is near  $-24$  ‰ (Amundson et al., 1998), and atmospheric  $\delta^{13}\text{C}_{\text{CO}_2}$  values are frequently observed to be near  $-10$  ‰ (Peyraube et al., 2013; Fernandez-Cortes et al., 2015). Due to fast airflow in cave 13 in Kentucky, the  $\text{CO}_2$  concentration of  $350 \pm 6$  ppmv and  $\delta^{13}\text{C}_{\text{CO}_2} = -10.7 \pm 0.4$  ‰ in cave air were similar to the local background atmosphere (i.e.  $400 \pm 90$  ppmv  $\text{CO}_2$ ). It is possible that the observed source values that are more negative than  $-24$  ‰ may be due to dripwater degassing or fast airflow (Spötl et al., 2005; Garcia-Anton et al., 2014). The  $\delta^{13}\text{C}_s$  values that are more positive than  $-24$  ‰ may be caused by differential abundances of C3 and C4 plants above the caves (Breecker et al., 2012). Our  $\text{CO}_2$  data show that the caves in our study are not atypical compared to other caves in the literature and that cave air residence time increased as the  $\delta^{13}\text{C}_{\text{CO}_2}$  values of cave air decreased.

### 4.2. Sources and Stable Isotopic Composition of Methane

Subatmospheric CH<sub>4</sub> concentrations in cave air suggest that CH<sub>4</sub> from the outside atmosphere is consumed in caves, and the addition of these observations to the existing datasets shows that karst is behaving as a sink for atmospheric CH<sub>4</sub> (Mattey et al., 2013; Ferndandez-Cortes et al., 2015; McDonough et al., 2016; Webster et al., 2016; Nguyễn-Thuỳ et al., 2017). Only three caves exhibited CH<sub>4</sub> concentrations above the atmospheric background, namely (i) cave 3, Indiana, (ii) cave 22, Tennessee, and (iii) cave 32, New Zealand. Caves with CH<sub>4</sub> concentrations above the atmospheric background appear to be uncommon, and understanding if there is a systematic change in karst landscapes from smaller order to higher order drainages awaits future work.

In some of the caves where we measured Rn concentrations, cave air flow was relatively fast. The average CO<sub>2</sub> and Rn concentrations from these caves were relatively low (cave 33, [CO<sub>2</sub>] = 540 ± 20 ppmv, [Rn] = 20 ± 100 Bq m<sup>-3</sup>; cave 34, [CO<sub>2</sub>] = 560 ± 30 ppmv, [Rn] = 200 ± 400 Bq m<sup>-3</sup>). Despite the similarity to atmospheric CO<sub>2</sub> concentrations in these caves, average CH<sub>4</sub> concentrations were still depleted relative to the atmosphere (cave 33, [CH<sub>4</sub>] = 1.07 ± 0.06 ppmv; cave 34, [CH<sub>4</sub>] = 1.34 ± 0.06 ppmv). These observations show that an *in-situ* process is removing CH<sub>4</sub> from the cave air because subatmospheric CH<sub>4</sub> concentrations in cave air are still observed in the absence of large increases in CO<sub>2</sub> concentrations as would be expected if CH<sub>4</sub> were only diluted by the arrival of CH<sub>4</sub>-free air into caves from soils.

In several caves, values of δ<sup>13</sup>CCH<sub>4</sub> and δ<sup>2</sup>HCH<sub>4</sub> show that non-atmospheric CH<sub>4</sub> enters caves because they were more negative than those predicted by the oxidation of atmospheric CH<sub>4</sub> via methanotrophy or reaction with ·OH. Other studies have also shown microbially produced CH<sub>4</sub> entering caves (Mattey et al., 2013; Webster et al., 2016). Our data point to the methanogenic sources of acetate fermentation and carbonate reduction (Fig. 7). Mixing between residual atmospheric CH<sub>4</sub> after partial methanotrophic oxidation and CH<sub>4</sub> from acetate fermentation in the soil-epikarst-cave system will cause a decrease in the δ<sup>13</sup>CCH<sub>4</sub> and δ<sup>2</sup>H CH<sub>4</sub> values of cave air compared to the atmospheric oxidation curve. Caves 23, 24, and 26 all appear to be influenced by acetoclastic methanogenesis. CH<sub>4</sub> produced from carbonate reduction is inferred to enter the caves based on samples that had δ<sup>2</sup>HCH<sub>4</sub> values that were roughly equal to, or more positive than atmospheric values (Fig. 7). The stable isotopic compositions of CH<sub>4</sub> from caves 5, 15, 25, and 27 can be explained by partial methanotrophic consumption of CH<sub>4</sub> generated from carbonate reduction with the strongest source signal in cave 5 (Fig. 7). CO<sub>2</sub> reduction is typically observed in lake sediments, but has been observed in oxidizing environments such as biological soil crusts in deserts after rain events (Angel et al., 2011). We hypothesize that karst environments, which are less oxidizing, exhibit similar behavior. Our data show that carbonate reduction and acetate fermentation can occur in similar environments over large spatial scales and are not limited to arctic environments.

Sites of methanogenesis in or near the studied cave systems may include waterlogged soils above caves, cave soils themselves, and the epikarst. It is possible that after rain events anoxic micro niches occur in soil, the epikarst, or caves themselves and that the generated CH<sub>4</sub> is dissolved and later introduced into caves with drip waters. We confirmed that dissolved CH<sub>4</sub> outgasses in drip water of cave 32, which was situated underneath a wetland, by placing our CH<sub>4</sub> detecting probe near the water and measuring increased CH<sub>4</sub> concentrations in its vicinity. We confirmed that *in-situ* CH<sub>4</sub> production can take place in locally anoxic environments within caves by measuring CH<sub>4</sub> concentrations close to a bat guano deposit in cave 25, Tennessee (site 4h; average CH<sub>4</sub> concentration = 0.3 ± 0.5 ppmv). Time series measurements near the large bat guano deposit showed that CH<sub>4</sub> concentrations oscillated between 0.5 ppmv and 0.1 ppmv over the course of seconds, presumably in response to episodic migration of CH<sub>4</sub> bubbles through the moist guano (similar oscillations in ammonia, NH<sub>3</sub>, were also observed). Additionally, we observed circumstantial evidence for local *in-situ* CH<sub>4</sub> production in cave 3 because measured CH<sub>4</sub> concentrations upstream of a restroom in the cave were low, while measured CH<sub>4</sub> concentrations downstream of the restroom were enhanced. Our data show that caves are capable of expressing elevated CH<sub>4</sub> concentrations

due to in-situ CH<sub>4</sub> production when accumulations of organic matter, such as guano or plant material, foster methanogenesis or when dissolved CH<sub>4</sub> from waters outgasses into cave air.

We observed minor amounts of thermogenic CH<sub>4</sub> entering in at least one cave. Locally elevated CH<sub>4</sub> concentrations in cave 9 (Mammoth Cave, Kentucky) were associated with a known hydrocarbon seep that is also transporting sulfide (Olson, 2013). Elevated CH<sub>4</sub> concentrations, thought to derive from thermogenic CH<sub>4</sub>, have also been observed at sulfidic springs in Cueva de Villa Luz (Webster et al., 2017). Some  $\delta^{13}\text{CCH}_4$  and  $\delta^2\text{HCH}_4$  values in cave air, i.e. two of the samples from cave 24, are compatible with thermogenic CH<sub>4</sub> isotopic signatures (e.g., Schoell, 1988; Etiope et al., 2009; Fig. 7). It cannot be excluded that small fluxes of CH<sub>4</sub> from shales or hydrocarbon deposits underlying the limestone are entering caves through natural fractures, but our present isotope data cannot confirm this source for cave 24. The Antes and Utica shales, which contain hydrocarbon gases, are stratigraphically below cave 24 (Coleman et al., 2014), and geologic faults and joints, which are often aligned with caves, may serve as conduits for the flow of hydrocarbons (Powell, 1969; Koša et al., 2003). A confirmation of hydrocarbons entering from deep sources may be obtained through measurements of 'radiocarbon-dead' CO<sub>2</sub>, Rn, and ethane.

#### 4.3. Methane Oxidation Mechanisms

The combination of  $\delta^{13}\text{CCH}_4$  and  $\delta^2\text{HCH}_4$  values allow for inferences to be made about how CH<sub>4</sub> oxidizing reactions in karst environments. We distinguish between two scenarios with distinct sets of assumptions. Many isotopic compositions of CH<sub>4</sub> in this study cannot be accounted for in the first scenario in which we assume that (i) CH<sub>4</sub> enters the caves from the atmosphere, through acetate fermentation, and through carbonate reduction, and that (ii) CH<sub>4</sub> is removed from cave air through reactions involving the ·OH. Conversely, in a second scenario, if it is assumed that (i) sources of CH<sub>4</sub> in cave air include the atmosphere, acetate fermentation, and carbonate reduction, and that (ii) CH<sub>4</sub> is removed from cave air by methanotrophy, all of the points fall within the plausibility envelope of the model, suggesting that methanotrophy is the mechanism responsible for removing CH<sub>4</sub> from cave air (Fig. 7). Consideration of an additional source of thermogenic CH<sub>4</sub> (natural gas) from deep geologic sources enlarges the plausibility fields of both prior scenarios to encompass all of the data. Our  $\delta^{13}\text{CCH}_4$  and  $\delta^2\text{HCH}_4$  data also agree with observations of  $\delta^{13}\text{CCH}_4$  and  $\delta^2\text{HCH}_4$  from a cave in Indiana where it appeared that CH<sub>4</sub> from both acetate fermentation and carbonate reduction influenced cave air geochemistry (Webster et al., 2016). Additionally our data resemble an arctic system characterized by acetoclastic and hydrogenotrophic CH<sub>4</sub> sources and methanotrophy (McCalley et al., 2014). Our isotopic evidence for *in-situ* microbial CH<sub>4</sub> oxidation in caves is corroborated by recent results from *in-situ* mesocosm experiments in Vietnam where cave rocks with live microorganisms were shown to consume CH<sub>4</sub> even in cases where surface soils were very thin to non-existent (Lennon et al., 2017; Nguyễn-Thùy et al., 2017).

## 5. Conclusions

Subterranean karst air generally shows subatmospheric CH<sub>4</sub> concentrations. CH<sub>4</sub> and CO<sub>2</sub> concentrations were negatively correlated in cave air showing that as the residence time of cave air increases the CH<sub>4</sub> concentration of cave air decreases. The stable isotopic composition of CH<sub>4</sub> in studied caves suggests that CH<sub>4</sub> is being oxidized by microbial methanotrophy. This evidence adds to earlier reports that methanotrophy is the mechanism by which CH<sub>4</sub> is removed in cave air (Mattey et al., 2013; McDonough et al., 2016; Webster et al., 2016; Lennon et al., 2017). The observations of sub-atmospheric CH<sub>4</sub>

concentrations in cave air from this study and other studies shows that karst is behaving as a global sink for CH<sub>4</sub>. CH<sub>4</sub> flux data from cave and karst landscapes are needed to estimate the size of the karst sink.

The stable isotopic composition of CH<sub>4</sub> in the studied caves suggests that, in addition to atmospheric CH<sub>4</sub>, at least two additional CH<sub>4</sub> sources are present in some caves. We suggest that the sources include CH<sub>4</sub> produced from acetate fermentation and from CO<sub>2</sub> reduction. These data corroborate recent findings of partially oxidized CH<sub>4</sub> entering cave air from acetate fermentation and CO<sub>2</sub> reduction in Indiana (Webster et al., 2016). These observations of CH<sub>4</sub> production by acetate fermentation and carbonate reduction suggest that both processes happen over a wide scale in the environment.

Cave air CH<sub>4</sub> appears to have been isotopically shifted towards more positive  $\delta^{13}\text{C}_{\text{CH}_4}$  and  $\delta^2\text{H}_{\text{CH}_4}$  values due to methanotrophy in the soil-epikarst-cave system. Even though minor CH<sub>4</sub> production appears to be taking place in karst environments, CH<sub>4</sub> consumption is the dominant process.

## Acknowledgements

This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, Chemical Sciences, Geosciences, and Biosciences Division under Award Number DE-SC0006978. This study was partially funded by NASA ASTEP (Award # NNX11AJ01G) and a National Speleological Society research grant. We thank J. Forsythe of Lost River Cave (cave 7) and R. S. Toomey III of Mammoth Cave National Park for granting access. Members of the Bloomington Indiana Grotto facilitated access to caves in Indiana. The Richard Blenz Nature Conservancy permitted access to cave 35. We thank M. Lefticariu and J. Hayden-Moore of Little Egypt Grotto in Carbondale, Illinois for providing assistance and access to gas sampling in cave 36 (SE Missouri). T.V. Royer in SPEA at Indiana University made analytical gas chromatography available. We are grateful for constructive comments from anonymous reviewers.

## References


1. Amundson, R., Stern, L., Baisden, T., Wang, Y., 1998. The isotopic composition of soil and soil-respired CO<sub>2</sub>. *Geoderma* 82, 83–114. [http://dx.doi.org/10.1016/S0016-7061\(97\)00098-0](http://dx.doi.org/10.1016/S0016-7061(97)00098-0)
2. Angel, R., Matthies D., Conrad, R., 2011. Activation of methanogenesis in arid biological soil crusts despite the presence of oxygen. *PLoS ONE* 6(5), e20453. <http://dx.doi.org/10.1371/journal.pone.0020453>
3. Batiot-Guilhe, C., Seidel, J.-L., Jourde, H., Hébrard, O., Bailly-Comte, V., 2007. Seasonal variations of CO<sub>2</sub> and <sup>222</sup>Rn in a Mediterranean sinkhole – spring (Causse d’Aumelas, SE France). *International Journal of Speleology* 36(1), 51-56.
4. Breecker, D.O., Payne, A.E., Quade, J., Banner, J.L., Ball, C.E., Meyer, K.W., Cowan, B.D., 2012. The sources and sinks of CO<sub>2</sub> in caves under mixed woodland and grassland vegetation. *Geochimica et Cosmochimica Acta* 96, 230-246. <http://dx.doi.org/10.1016/j.gca.2012.08.023>
5. Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R.B., Piao, S., Thornton, P., 2013. Carbon and other biogeochemical cycles. *In* (Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, A., Nauels, A., Xia, Y., Bex, V., Midgley, P.M., eds.) *Climate Change 2013: The Physical Science Basis. Contribution to Working Group I to the Fifth Assessment Report of the Intergovernmental*

- Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
6. Coleman D.D., Risatti J.B., Schoell M., 1981. Fractionation of carbon and hydrogen isotopes by methane-oxidizing bacteria. *Geochimica et Cosmochimica Acta* 45(7), 1033-1037. [http://dx.doi.org/10.1016/0016-7037\(81\)90129-0](http://dx.doi.org/10.1016/0016-7037(81)90129-0)
  7. Coleman, J.L., Jr., Ryder, R.T., Milici, R.C., Brown, S., 2014. Overview of the potential and identified petroleum source rocks of the Appalachian Basin, Eastern United States. *In* (Ruppert, L.F., Ryder, R.T., eds.) *Coal and Petroleum Resources in the Appalachian basin; Distribution, Geologic Framework, and Geochemical Character*: U.S. Geological Survey Professional Paper 1708, <http://dx.doi.org/10.3133/pp1708G.13>
  8. Cunningham, K.I., LaRock, E.J., 1991. Recognition of microclimate zones through radon mapping, Lecheguilla Cave, Carlsbad Caverns National Park, New Mexico. *Health Physics* 61(4), 493-500.
  9. Dlugokencky, E.J., Houweling, S., Bruhwiler, L., Masarie, K.A., Lang, P.M., Miller, J.B., Tans, P.P., 2003. Atmospheric methane levels off: Temporary pause or a new steady state? *Geophysical Research Letters* 30 (1992), <http://dx.doi.org/10.1029/2003GL018126>
  10. Dlugokencky, E.J., Nisbet, E.G., Fisher, R., Lowry, D., 2011. Global atmospheric methane: budget changes, and dangers. *Phil. Trans. R. Soc. A* 369, 2058-2072. <http://dx.doi.org/10.1098/rsta.2010.0341>
  11. Etiopé G., Feyzullayev A., Baciú C.L., 2009. Terrestrial methane seeps and mud volcanoes: a global perspective of gas origin. *Marine and Petroleum Geology* 26, 333-344. <http://dx.doi.org/10.1016/j.marpetgeo.2008.03.001>
  12. Feisthauer, S., Vogt, C., Modrzynski, J., Szlenkier, M., Krüger, M., Siegert, M., Richnow, H.-H., 2011. Different types of methane monooxygenases produce similar carbon and hydrogen isotope fraction patterns during methane oxidation. *Geochimica et Cosmochimica Acta* 75(5), 1173-1184. <http://dx.doi.org/10.1016/j.gca.2010.12.006>
  13. Fernandez-Cortes, A., Cuezva, S., Alvarez-Gallego, M., Garcia-Anton, E., Pla C., Benavente, D., Jurado V., Saiz-Jimenez, C., Sanchez-Moral, S., 2015. Subterranean atmospheres may act as daily methane sinks. *Nature Communications* 6(7003). <http://dx.doi.org/10.1038/ncomms8003>
  14. Garcia-Anton, E., Cuezva, S., Fernandez-Cortes, A., Benavente, D., Sanchez-Moral, S., 2014. Main drivers of diffusive and advective processes of CO<sub>2</sub>-gas exchange between a shallow vadose zone and the atmosphere. *International Journal of Greenhouse Gas Control* 21, 113-129. <http://dx.doi.org/10.1016/j.ijggc.2013.12.006>
  15. Ford, D., Williams, P., 2007. *Karst Hydrogeology and Geomorphology*. John Wiley & Sons Ltd, England.
  16. Gioli, B., Toscano, P., Lugato, E., Matese, A., Miglietta, F., Zaldei, A., Vaccari, F.P., 2012. Methane and carbon dioxide fluxes and source partitioning in urban areas: The case study of Florence, Italy. *Environmental Pollution* 164, 125-131. <http://dx.doi.org/10.1016/j.envpol.2012.01.019>
  17. Gregorič, A., Zidanšek, A., Vaupotič, J., 2011. Dependence of radon levels in Postojna Cave on outside air temperature. *Natural Hazards and Earth System Sciences* 11, 1523-1528. <http://dx.doi.org/10.5194/nhess-11-1523-2011>
  18. Gregorič, A., Vaupotič, J., Šebela, S., 2014. The role of cave ventilation in governing cave air temperature and radon levels (Postojna Cave, Slovenia). *International Journal of Climatology* 34(5), 1488-1500. <http://dx.doi.org/10.1002/joc.3778>
  19. Harper, L.A., Weaver, K.H., De Visscher, A., 2014. Dinitrogen and methane gas production during the anaerobic/anoxic decomposition of animal manure. *Nutrient Cycling in Agroecosystems* 100(1), 53-64. <http://dx.doi.org/10.1007/s10705-014-9626-9>
  20. Hutchens, E., Radajewski, S., Dumont, M.G., McDonald, I.R., Murrell, J.C., 2004. Analysis of methanotrophic bacteria in Movile Cave by stable isotope probing. *Environmental Microbiology* 6(2), 111-120. <http://dx.doi.org/10.1046/j.1462-2920.2003.00543.x>


21. Jones, D.S., Albrecht, H.L., Dawson, K.S., Schaperdoth, I., Freeman, K.H., Pi, Y., Pearson, A., Macalady, J.L., 2012. Community genomic analysis of an extremely acidophilic sulfur oxidizing biofilm. *The ISME Journal* 6, 158-170. <http://dx.doi.org/10.1038/ismej.2011.75>
22. Kirschke, S., Bousquet, P., Ciais, P., Saunois, M., Canadell, J.G., Dlugokencky, E.J., Bergamaschi, P., Bergmann, D., Blake, D.R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser, A., Heimann, M., Hodson, E.L., Houweling, S., Josse, B., Fraser, P.J., Krummel, P.B., Lamarque, J.-F., Langenfelds, R.L., Le Quéré, C., Naik, V., O'Doherty, S., Palmer, P.I., Pison, I., Plummer, D., Poulter, B., Prinn, R.G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Schindell, D.T., Simpson, I.J., Spahni, R., Steele, L.P., Strode, S.A., Sudo, K., Szopa, S., van der Werf, G.R., Voulgarakis, A., van Weele, M., Weiss, R.F., Williams, J.E., Zeng, G., 2014. Three decades of global methane sources and sinks. *Nature Geoscience* 6, 812-8123.
23. Keeling, R.F., Piper, S.C., Bollenbacher, A.F., Walker, S.J., 2010. Monthly atmospheric  $^{13}\text{C}/^{12}\text{C}$  isotopic ratios for 11 SIO stations. *In Trends: A Compendium of Data on Global Change. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tenn., U.S.A.*
24. Koša, E., Hunt, D., Fitchen, W.M., Bockel-Rebelle, M.-O., Roberts, G., 2003. The heterogeneity of paleocavern systems developed along syndepositional fault zones: The Upper Permian Capitan Platform, Guadalupe Mountains, U.S.A. *SEPM Special Publications No. 78 and APGP Memoir 83*, 291-322.
25. Kowalczyk, A.J., Froelich, P.N., 2010. Cave air ventilation and  $\text{CO}_2$  outgassing by radon-222 monitoring: How fast do caves breathe? *Earth and Planetary Science Letters* 289, 209-219. <http://dx.doi.org/10.1016/j.epsl.2009.11.010>
26. Lennon, J.T., Nguyễn-Thùy, D., Phạm, T.M., Drobnik, A., Tạ, P.H., Phạm, N.Đ., Streil, T., Webster, K.D., Schimmelmann, A., 2017. Microbial contributions to subterranean methane sinks. *Geobiology* 15(2), 254-258. <http://dx.doi.org/10.1111/gbi.12214>
27. Matthey, D.P., Fairchild, I.J., Atkinson, T.C., Latin, J.-P., Ainsworth, M., Durrell, R., 2010. Seasonal microclimate control of calcite fabrics, stable isotopes and trace elements in modern speleothem from St. Michaels Cave, Gibraltar. *In (Pedley, H.M., Rogerson, M. eds.) Tufas and Speleothems: Unravelling the Microbial and Physical Controls: Geological Society, London, Special Publications 336*, 323-344. <http://dx.doi.org/10.1144/SP336.17>
28. Matthey, D.P., Fisher, R., Atkinson, T.C., Latin, J.-P., Durrell, R., Ainsworth, M., Lowry, D., Fairchild, I.J., 2013. Methane in underground air in Gibraltar karst. *Earth and Planetary Science Letters* 374, 71-80. <http://dx.doi.org/10.1016/j.epsl.2013.05.011>
29. McCalley, C.K., Woodcroft, B.J., Hodgkins, S.B., Wehr, R.A., Kim, E.-H., Mondav, R., Crill, P.M., Chanton, J.P., Rich, V.I., Tyson, G.W., Saleska, S.R., 2014. Methane dynamics regulated by microbial community response to permafrost thaw. *Nature* 514, 478-481. <http://dx.doi.org/10.1038/nature13798>
30. McDonough, L.K., Iverach, C.P., Beckman, S., Manfield, M., Rau, G.C., Baker, A., Kelly, B.F.J., 2016. Spatial variability of cave-air carbon dioxide and methane concentrations and isotopic compositions in a semi-arid karst environment. *Environmental Earth Sciences* 75(700), <http://dx.doi.org/10.1007/s12665-016-5497-5>
31. Miller, J.B., Mack, K.A., Dissly, R., White, J.W.C., Dlugokencky, E.J., Tans, P.P., 2002. Development of analytical methods and measurements of  $^{13}\text{C}/^{12}\text{C}$  in atmospheric  $\text{CH}_4$  from the NOAA Climate Monitoring and Diagnostics Laboratory Global Air Sampling Network. *Journal of Geophysical Research: Atmospheres* 107(D13), ACH 11-1-ACH 11-15. <http://dx.doi.org/10.1029/2001JD000630>
32. Nisbet, E.G., Dlugokencky, E.J., Manning, M.R., Lowry, D., Fisher, R.E., France, J.L., Michel, M.E., Miller, J.B., White, W.J.C., Vaughn, B., Bousquet, P., Pyle, J.A., Warwick, N.J., Cain, M., Brownlow, R., Zazzeri, G., Lanoisellé, M., Manning, A.C., Gloor, E., Worthy, D.E.J., Brunke, E.-G.,

- Labuschagne, C., Wolff, E.W., Ganesan, A.L., 2016. Rising atmospheric methane: 2007-2014 growth and isotopic shift. *Global Biogeochemical Cycles* 30, 1356-1370. <http://dx.doi.org/10.1002/2016GB005406>
33. Nguyễn-Thùy, D., Schimmelmann, A., Nguyễn-Văn, H., Drobniak, A., Lennon, J.T., Tạ, P.H., Nguyễn, N.T.Á., (2017). Subterranean microbial oxidation of atmospheric methane in cavernous tropical karst. *Chemical Geology*. <https://doi.org/10.1016/j.chemgeo.2017.06.014>
  34. Olson, R., 2013. Potential effects of hydrogen sulfide and hydrocarbon seeps on Mammoth Cave ecosystems. Mammoth Cave Research Symposia. Paper 28. [http://digitalcommons.wku.edu/mc\\_research\\_symp/10th\\_Research\\_Symposium\\_2013/Research\\_Posters/28](http://digitalcommons.wku.edu/mc_research_symp/10th_Research_Symposium_2013/Research_Posters/28)
  35. Palmer, A.N., 1991. Origin and morphology of limestone caves. *Geological Society of America Bulletin* 103(1), 1-21. [http://dx.doi.org/10.1130/0016-7606\(1991\)103<0001:OAMOLC>2.3.CO;2](http://dx.doi.org/10.1130/0016-7606(1991)103<0001:OAMOLC>2.3.CO;2)
  36. Peyraube, N., Lastennet, R., Denis, A., Malaurent, P., 2013. Estimation of epikarst air  $P_{CO_2}$  using measurements of water  $\delta^{13}C_{TDIC}$ , cave air  $P_{CO_2}$  and  $\delta^{13}C_{CO_2}$ . *Geochimica et Cosmochimica Acta* 118, 1-17. <http://dx.doi.org/10.1016/j.gca.2013.03.046>
  37. Peyraube, N., Lastennet, R., Villanueva, J.D., Houillon, N., Malaurent, P., Denis, A., 2016. Effect of diurnal and seasonal temperature variation on Cussac cave ventilation using  $CO_2$  assessment. *Theoretical and Applied Climatology*, <http://dx.doi.org/10.1007/s00704-016-1824-8>
  38. Powell, R.L. 1969. Base level, and climatic controls of karst groundwater zones in South-Central Indiana. *Indiana Academy of Science*, 281-291.
  39. Sarbu, S.M., Kane, T.C., Kinkle, B.K., 1996. A chemoautotrophically based cave ecosystem. *Science* 272(5270), 1953-1955. <http://dx.doi.org/10.1126/science.272.5270.1953>
  40. Saueressig, G., Crowley, J.N., Bergamaschi, P., Brühl, C., Brenninkmeijer, C.A.M., Fischer, H., 2001. Carbon 13 and D kinetic isotope effects in the reactions of  $CH_4$  with  $O(^1D)$  and  $OH$ : New laboratory measurements and their implications for the isotopic composition of stratospheric methane. *Journal of Geophysical Research* 106 (D19), 23127–23138. <http://dx.doi.org/10.1029/2000JD000120>
  41. Schoell, M., 1988. Multiple origins of methane in the Earth, in Schoell, M. ed. *Origins of Methane in the Earth*. *Chemical Geology* 71, 1-10. [http://dx.doi.org/10.1016/0009-2541\(88\)90101-5](http://dx.doi.org/10.1016/0009-2541(88)90101-5)
  42. Spötl, C., Fairchild, I.J., Tooth, A.F., 2005. Cave air control on dripwater geochemistry, Obir Caves (Austria): Implications for speleothem deposition in dynamically ventilated caves. *Geochimica et Cosmochimica Acta* 69(10), 2451-2468. <http://dx.doi.org/10.1016/j.gca.2004.12.009>
  43. Sussman, R., Forster, F., Rettinger, M., Bousquet, P., 2012. Renewed methane increase for five years (2007-2011) observed by solar FTIR spectrometry. *Atmospheric Chemistry and Physics* 12, 4885-4891. <http://dx.doi.org/10.5194/acp-12-4885-2012>
  44. Templeton, A., Chu, K.-H., Alvarez-Cohen, L., Conrad, M.E., 2006. Variable carbon isotope fractionation expressed by aerobic  $CH_4$ -oxidizing bacteria. *Geochimica et Cosmochimica Acta* 70(7), 1739-1752. <http://dx.doi.org/10.1016/j.gca.2005.12.002>
  45. Townsend-Small, A., Tyler, S.C., Pataki, D.E., Xu, X., Christensen, L.E., 2012. Isotopic measurements of atmospheric methane in Los Angeles, California, USA: Influence of “fugitive” fossil fuel emissions. *Journal of Geophysical Research* 117(D07308). <http://dx.doi.org/10.1029/2011JD016826>
  46. Tu, K.P., Brooks, P.D., Dawson, T.E., 2001. Using septum-capped vials with continuous-flow isotope ratio mass spectrometric analysis of atmospheric  $CO_2$  for Keeling plot applications. *Rapid Communications in Mass Spectrometry* 15(12), 952-956. <http://dx.doi.org/10.1002/rcm.320>
  47. Weary, D.J., Doctor, D.H., 2014. Karst in the United States: A digital map compilation and database: U.S. Geological Survey Open-File Report 2014–1156. <http://dx.doi.org/10.3133/ofr20141156>

48. Webster, K.D., Mirza, A., Deli, J.M., Sauer, P.E., Schimmelmann, A., 2016. Consumption of atmospheric methane in a limestone cave in Indiana, USA. *Chemical Geology* 443, 1-9. <http://dx.doi.org/10.1016/j.chemgeo.2016.09.020>
49. Webster, K.D., Rosales Lagarde, L., Sauer, P.E., Schimmelmann, A., Lennon, J.T., Boston, P.J., 2017. Isotopic evidence for the migration of thermogenic methane into a sulfidic cave, Cueva de Villa Luz, Tabasco, Mexico. *Journal of Cave and Karst Studies* 79(1), 24-34. <https://caves.org/pub/journal/PDF/V79/cave-79-01-24.pdf>.
50. Whiticar, M.J., 1999. Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane. *Chemical Geology* 161, 291-314. [http://dx.doi.org/10.1016/S0009-2541\(99\)00092-3](http://dx.doi.org/10.1016/S0009-2541(99)00092-3)




**Figure 1.** Locations of the U.S. study caves in their regional context (A) and within the contiguous USA (B). The location of the New Zealand caves are presented in C. Karst land cover data were obtained from Weary and Doctor (2014) and Ford and Williams (2007).




**Figure 2.** CH<sub>4</sub> concentrations *versus* CO<sub>2</sub> concentrations in studied caves. The majority of samples appear to follow an inverse power law relationship with CH<sub>4</sub> concentrations being inversely related to CO<sub>2</sub> concentrations (with points from caves 3 and 22 removed from the overall trend). Numbers represent individual caves. For clarity some error bars were omitted from the figure. Data associated with error bars are also representative for typical errors of data where no bars are shown.







**Figure 4.** A Keeling plot of  $\delta^{13}\text{C}_{\text{CO}_2}$  versus inverse CO<sub>2</sub> concentration in cave air samples. The data show that atmospheric CO<sub>2</sub> is mixing with apparent isotopic endmembers between  $-33$  and  $-20.4$  ‰. A regression analysis of the entire data set shows that the average  $\delta^{13}\text{C}_{\text{CO}_2}$  value entering the caves is  $-23.3 \pm 0.5$  ‰. Numbers represent individual caves. Error bars are not included for clarity.



**Figure 5.** Relationship between CH<sub>4</sub> concentration and  $\delta^{13}\text{C}_{\text{CH}_4}$  in cave air. Some samples plot along the expected relationship between CH<sub>4</sub> concentration and  $\delta^{13}\text{C}_{\text{CH}_4}$  caused either by methanotrophy modeled with an  $\alpha$  value of 1.018, or by oxidation with ·OH modeled with an  $\alpha$  value of 1.0039. Other samples plot below and to the left of the theoretical shifts of the oxidation trends. Numbers represent individual caves. If error bars are not visible, they are smaller than the data points.



**Figure 6.** Relationship between methane concentration and  $\delta^2\text{H}_{\text{CH}_4}$  in cave air. Some samples plot along the expected relationship between  $\text{CH}_4$  concentration and  $\delta^2\text{H}_{\text{CH}_4}$  caused by methanotrophy modeled with an  $\alpha$  value of 1.1353, or by oxidation with  $\cdot\text{OH}$  modeled with an  $\alpha$  value 1.294. Other samples plot below and to the left of the shift caused by methanotrophy. Numbers represent individual caves. If error bars are not visible, they are smaller than the data points.



**Figure 7.** Stable isotopic composition of CH<sub>4</sub> in cave air samples plotted in δ<sup>2</sup>H<sub>CH<sub>4</sub></sub> versus δ<sup>13</sup>C<sub>CH<sub>4</sub></sub> space.

CH<sub>4</sub> generated by carbonate reduction, acetate fermentation, and thermogenesis are plotted within labeled fields (Whiticar, 1999). Dotted lines indicate the expected shift in δ<sup>2</sup>H<sub>CH<sub>4</sub></sub> and δ<sup>13</sup>C<sub>CH<sub>4</sub></sub> caused by partial aerobic methane oxidation adopting a slope of 8.5. Dashed lines indicate the expected shift in δ<sup>2</sup>H<sub>CH<sub>4</sub></sub> and δ<sup>13</sup>C<sub>CH<sub>4</sub></sub> caused by oxidation with ·OH adopting a slope 75 (Saueressig et al., 2001; Feisthauer et al., 2011). Mixing in δ<sup>2</sup>H<sub>CH<sub>4</sub></sub> vs δ<sup>13</sup>C<sub>CH<sub>4</sub></sub> space plots as a straight line. Numbers represent individual caves. If the cave system can be characterized by sources from the atmosphere (□, -47.5 ‰, -100 ‰), acetate fermentation (S1, -49 ‰, -325 ‰), and carbonate reduction (S2, -63 ‰, -125 ‰), and methanotrophy the

data should plot between the methanotrophic boundaries between S1 and S2, shaded in medium gray. If the same sources are considered to be entering the cave and are oxidized by the  $\cdot\text{OH}$  radical they should plot between the  $\cdot\text{OH}$  oxidized boundary of S1 and the atmosphere, and mixing between S1 and S2 as well as mixing between S2 and the atmosphere, shaded light gray. Error bars are not included with black circles for clarity. In other locations, if error bars are not visible, they are smaller than the data points.

**Table 1:** Overview of collected data. 'Discrete' measurements refer to the laboratory.

| Cave | State           | Collection date <sup>a</sup> | Sites measured | [CH <sub>4</sub> ] min (ppmv) | [CH <sub>4</sub> ] max (ppmv) | [CO <sub>2</sub> ] min (ppmv) | [CO <sub>2</sub> ] max (ppmv) | Methods        |
|------|-----------------|------------------------------|----------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|----------------|
| 1    | AL              | 2015-04-07                   | 1              | NA <sup>b</sup>               | 2.2                           | NA                            | 378                           | <i>In-situ</i> |
| 2    | IN              | 2013-08-10                   | 4              | 0.42                          | 1.89                          | 2140                          | 8000                          | Discrete       |
| 3    | IN              | 2013-08-07                   | 3              | 0.23                          | 4.2                           | 13400                         | 17900                         | Discrete       |
| 4    | IN              | 2013-06-18                   | 2              | 0.3                           | 0.7                           | 3900                          | 5200                          | Discrete       |
| 5    | IN              | 2013-06-29                   | 3              | 0.4                           | 1.18                          | 3300                          | 8600                          | Discrete       |
| 6    | IN              | 2013-06-14                   | 3              | 0.32                          | 1.32                          | 750                           | 3700                          | Discrete       |
| 7    | KY              | 2012-05-05                   | 4              | 0.3                           | 0.9                           | 3500                          | 6020                          | <i>In-situ</i> |
| 8    | KY              | 2012-05-06                   | 5              | 0.1                           | 0.4                           | 5200                          | 6200                          | <i>In-situ</i> |
| 9    | KY              | 2012-05-07                   | 7              | 0.08                          | 0.6                           | 900                           | 1500                          | <i>In-situ</i> |
| 10   | KY              | 2013-07-18                   | 1              | NA                            | 1.79                          | NA                            | 920                           | Discrete       |
| 11   | KY              | 2013-07-18                   | 1              | NA                            | 0.91                          | NA                            | 2790                          | Discrete       |
| 12   | KY              | 2013-07-18                   | 1              | NA                            | 1.37                          | NA                            | 1790                          | Discrete       |
| 13   | KY              | 2015-04-10                   | 4              | 2.3                           | 2.6                           | 349                           | 390                           | Mixed          |
| 14   | KY              | 2015-04-10                   | 3              | 1.82                          | 2.33                          | 351                           | 390                           | <i>In-situ</i> |
| 15   | KY              | 2015-04-10                   | 2              | 0.9                           | 1.8                           | 440                           | 486                           | Mixed          |
| 16   | OH              | 2015-04-10                   | 1              | NA                            | 1.95                          | NA                            | 940                           | <i>In-situ</i> |
| 17   | PA              | 2013-07-15                   | 5              | 1.4                           | 1.9                           | 1430                          | 2400                          | Mixed          |
| 18   | PA              | 2013-07-15                   | 4              | 0.4                           | 1.1                           | 760                           | 4250                          | Mixed          |
| 19   | PA              | 2013-07-15                   | 4              | 0.4                           | 0.5                           | 900                           | 1400                          | Mixed          |
| 20   | PA              | 2013-07-16                   | 3              | 0.4                           | 1.0                           | 3390                          | 4240                          | Mixed          |
| 21   | TN              | 2013-07-18                   | 3              | 0.5                           | 1.25                          | 2210                          | 2430                          | Mixed          |
| 22   | TN              | 2013-07-19                   | 3              | 2.6                           | 3.5                           | 490                           | 1610                          | Mixed          |
| 23   | TN              | 2015-04-07                   | 7              | 0.2                           | 1.02                          | 1100                          | 1600                          | Mixed          |
| 24   | TN              | 2015-04-08                   | 6              | 0.6                           | 1.6                           | 1100                          | 1515                          | Mixed          |
| 25   | TN              | 2015-04-09                   | 7              | 0.2                           | 0.8                           | 1430                          | 2100                          | Mixed          |
| 26   | TN              | 2015-04-09                   | 3              | 0.6                           | 0.79                          | 630                           | 660                           | Mixed          |
| 27   | VA              | 2015-07-17                   | 5              | 0.6                           | 1.0                           | 6300                          | 14700                         | Mixed          |
| 28   | VA              | 2015-07-17                   | 4              | 0.4                           | 0.8                           | 2480                          | 7600                          | Mixed          |
| 29   | VA              | 2015-07-17                   | 5              | 0.4                           | 1.6                           | 1100                          | 8740                          | Mixed          |
| 30   | VA              | 2015-07-18                   | 5              | 0.3                           | 1.5                           | 1090                          | 19900                         | Mixed          |
| 31   | VA              | 2015-07-18                   | 6              | 0.4                           | 2.2                           | 410                           | 8900                          | Mixed          |
| 32   | NZ <sup>c</sup> | 2014-10-04                   | 8              | 1.5                           | 2.8                           | 500                           | 590                           | <i>In-situ</i> |
| 33   | NZ              | 2014-10-04                   | 3              | 1.0                           | 1.7                           | 500                           | 620                           | <i>In-situ</i> |
| 34   | NZ              | 2014-10-04                   | 6              | 1.0                           | 1.8                           | 500                           | 1000                          | <i>In-situ</i> |
| 35   | IN              | 2014-06-11                   | 2              | 0.14                          | 0.3                           | NA <sup>d</sup>               | NA <sup>d</sup>               | <i>In-situ</i> |
| 36   | MO              | 2016-09-24                   | 9              | 0.1                           | 1.5                           | 2590                          | 6190                          | <i>In-situ</i> |

<sup>a</sup> Dates are formatted yyyy-mm-dd<sup>b</sup> NA = not applicable where only 1 sample was obtained.<sup>c</sup> NZ = New Zealand, <sup>d</sup> NA = Only CH<sub>4</sub> was measured

**Table 2:** *In-situ* instrumentation used in this study.

| Instrument    | Maker                      | Location                      | Method                                  | Analytes Measured                                   | Lower detection limits                                                         |
|---------------|----------------------------|-------------------------------|-----------------------------------------|-----------------------------------------------------|--------------------------------------------------------------------------------|
| GasFinder2    | Boreal Laser               | Spruce Grove, Alberta, Canada | Tunable Diode Laser Spectroscopy        | CH <sub>4</sub>                                     | 1 ppmv m <sup>-1</sup>                                                         |
| Inspectra     | Gazomat, WEST Systems      | Pontedera, Italy              | Tunable Diode Laser Spectroscopy        | CH <sub>4</sub>                                     | 0.1 ppmv                                                                       |
| LGD F200      | Axetris, SARAD             | Kägiswil, Switzerland         | Tunable Diode Laser Spectroscopy        | CH <sub>4</sub>                                     | 0.2 ppmv                                                                       |
| madIR-D01 CO2 | Madur, SARAD               | Zgierz, Poland                | Fourier Transform Infrared Spectroscopy | CO <sub>2</sub>                                     | 400 ppmv                                                                       |
| DX4030        | Gasmet LICOR, WEST Systems | Milton Keynes, United Kingdom | Nondispersive Infrared Spectroscopy     | CH <sub>4</sub> , CO <sub>2</sub> , NH <sub>3</sub> | 0.3 ppmv CH <sub>4</sub> , 200 ppmv CO <sub>2</sub> , 0.1 ppbv NH <sub>3</sub> |
| LI820         | WEST Systems               | Pontedera, Italy              | Alpha Spectroscopy                      | CO <sub>2</sub>                                     | 5 ppmv                                                                         |
| RTM 2200      | SARAD                      | Dresden, Germany              | Alpha spectroscopy                      | Rn                                                  | 0 Bq/m <sup>3</sup>                                                            |