

*Review*

## **A Comprehensive Profile of Mature MicroRNA in the Pig**

Xiliang Wang<sup>1,2</sup>, David K.C. Cooper<sup>3</sup>, Zhiming Cai<sup>1</sup>, Lisha Mou<sup>1\*</sup>

1. Shenzhen Xenotransplantation Medical Engineering Research and Development Center,

Institute of Translational Medicine, Shenzhen Second People's Hospital, First Affiliated

Hospital of Shenzhen University, Shenzhen, Guangdong, 518035, China

2. Department of Biochemistry, Zhongshan School of Medicine, Sun Yat-Sen University,

Guangzhou, Guangdong, 510080, China

3. Xenotransplantation Program, Department of Surgery, University of Alabama at

Birmingham, Birmingham, AL, 35233, USA

*\* Corresponding author*

Lisha Mou PhD

Shenzhen Second People's Hospital

NO.3002 Sungang Road

Futian district, Shenzhen, Guangdong 518035, China

Tel: (086) 0755-83366388-3230

Fax: (086) 0755-83366388-3230

E-mail: lishamou@gmail.com

---

**Abstract**

The pig is an important source of meat production and provides a valuable model for certain human diseases. MicroRNA (miRNA), which is non-coding RNA and regulates gene expression at the posttranscriptional level, plays a critical role in various biological processes. Studies on identification and function of mature miRNAs in multiple pig tissues are increasing, yet the literature is limited. Therefore, we reviewed current research to determine the miRNAs expressed in specific pig tissues that are involved in carcass values (including muscle and adipocytes), reproduction (including pituitary, testis, and ovary), and development of some solid organs (e.g., brain, lung, kidney, and liver). We also discuss the possible regulating mechanisms of miRNA. Finally, as pig organs are suitable candidates for xenotransplantation, biomarkers of their miRNA in xenotransplantation were evaluated.

**Keywords:**

microRNA; pig; biomarker; xenotransplantation

---

## Introduction

The pig is one of the most widespread livestock species in the world, providing meat production [1]. Although relatively expensive to breed and with a relatively long gestation period (about 114 days) [2], the pig is still an ideal animal model for biomedical research because of its close phylogenetic proximity and similarities with humans, such as organ size, anatomical features, physiology, and organ development, compared with the mouse [3]. The pig has been used as a model to study various issues, such as reproduction [4], and the neuronal system [5], and is employed as a source of organs and cells in xenotransplantation [6].

Despite its apparent importance, knowledge about the pig is still much less than has been accumulated for the mouse and rat, such as genome organization and gene expression regulation. The latest porcine genome reference (Sscrofa 11.1) was shared in the NCBI website (<https://www.ncbi.nlm.nih.gov/genome/?term=pig>) by the Swine Genome Sequencing Consortium (SGSC) in 2017.

When compared with the human, the pig's genome is of comparable size and contains a similar number of protein-coding genes [7]. One might think that the pig should also have a similar number of microRNAs (miRNAs). However, the number of porcine miRNAs available in public databases is still limited, with only 411 mature miRNAs in the miRBase (v21) compared to human (2,588) and mouse (1,982) [8], partly because only part of the porcine genome is available for study [9]. The common research strategy on miRNA may include three aims: (i) characterization of miRNAs (including identification of novel


---

miRNAs), (ii) determination of target genes, and (iii) illumination of function of miRNAs and target genes. These strategies have been widely used to describe the miRNAome in various pig tissues, such as muscle [9-12], brain [13], fat [14], embryo [3], pituitary [15], intestine [16], ovary [17] and testes [4].

This review is focused on the possible functions and regulating mechanisms of miRNAs in pigs, aiming towards a better understanding of the miRNAome in various tissues. Because of many similarities in morphology and physiology between pig and human, we also evaluate biomarker values of pig miRNA in pig-to-human xenotransplantation.

### **Origins and mechanisms of miRNAs**

The miRNAs, which are typical transcripts of RNA polymerase II, are small non-coding RNAs in animals and plants [18]. They are transcribed from genomic DNA as long hairpins (pri-miRNA) with an imperfectly paired stem of ~33 bp [19]. The pri-miRNA is excised by Drosha to generate a pre-miRNA species in the nucleus, which is the first processing step. In the second processing step, pre-miRNA is exported from the nucleus and processed by Dicer to form the mature miRNA/miRNA\* duplex of ~22bp length. The miRNA is then assembled into RISC (RNA-induced silencing complexes). Generally, only one strand of the duplex is stably associated with an RISC [20] (Fig 1).



**Figure 1.** Biogenesis of miRNA and assembly into miRISC, and possible mechanisms of miRISC-mediated repression. In animals, the pri-miRNA is transcribed by RNA polymerase II from genomic DNA, and is processed by Drosha with the aid of DGCR8 to generate a pre-miRNA species, which is exported from the nucleus and processed by Dicer to form the mature miRNA/miRNA\* duplex. Generally, only one strand of the duplex is then assembled into miRISC. When RISCs bind to mRNAs, they can repress initiation of translation at the stage of cap recognition (I) or 60S recruitment (II). Alternatively, they can induce mRNA deadenylation and thereby inhibit its circularization (III). They can also repress translation at the post-initiation stage through inducing ribosomes to dissociate prematurely (IV). They can also induce deadenylation followed by decapping to facilitate mRNA degradation (V). Without repression, mRNAs recruit initiation factors and ribosomal subunits and form circularized structures (VI).

---

The miRNA acts as an adaptor for RISC to specifically recognize and regulate particular target mRNAs (Fig 1). Recognition involves Watson-Crick base pairing of the 2-8th miRNA nucleotides, which is the so-called seed region [21]. The binding sites of miRNA to mRNAs are located in the 3' UTR (un-translated region) and usually exist in multiple copies. Most animal miRNAs bind imperfectly with mismatches. When RISCs bind to mRNAs, they can repress initiation of translation at the stage of cap recognition or 60S recruitment [22,23]. Alternatively, they can induce mRNA deadenylation and thereby inhibit its circularization [24]. They can also repress translation at post-initiation stage through inducing ribosomes to drop off prematurely [25]. They can also induce deadenylation followed by decapping to facilitate mRNA degradation [26,27] (Fig 1).

### **miRNAs in pig skeletal muscle (Table 1)**

As the pig is an agriculturally important species, miRNAs that affect development and growth of economically important skeletal muscle are of interest. Several miRNAs may promote myogenesis [28-35] (Table 1). Several others have potential function during muscle development [35-51]. Others are involved in the development of the *longissimus doris* and *psaos major* muscles [46-51], whereas others [9,52-55] are expressed in a stage-specific manner across muscle development periods. Among them, the miR-1, miR-133, and miR-206, which are specifically expressed in cardiac and skeletal muscle [32], are frequently listed as the highest expressed miRNAs in porcine muscle [9,10,41,56,57]. These multiple above-mentioned miRNAs provide insights into the regulation of muscle growth, and are potential candidates for further improvement of meat quality and production using molecular approaches.

**Table 1: miRNAs in pig skeletal muscle**

miRNA	Target	Function	Reference
miR-1a	HDAC4	promotes myogenesis during embryonic development and muscle cell differentiation	28,29,46-48
miR-133	SRF	enhances the proliferation of myoblasts	29,46-48
miR-27b		involved in myogenic differentiation, fast-specific and glucocorticoid-dependent myostatin expression	30-32
miR-148a	ROCK1	a novel myogenic miRNA that mediates myogenic differentiation	33
miR-143		controls performance of different fiber types	34,35
miR-378	BMP2 MAPK1	a candidate for myogenesis	36,37
miR-128		regulates adipogenesis, osteogenesis, and myogenesis	38
miR-126		attenuates insulin signaling and governs vascular integrity and angiogenesis	39,40
miR-92a		regulates skeletal muscle growth	35
miR-127		regulate the callipyge muscular hypertrophy phenotype	41,42
miR-432			
miR-136			
miR-10b		regulates myogenesis and muscle development	41,43
let-7 family		key miRNA regulators of development	44,45
miR-103		involved in cellular Acetyl-CoA and lipid levels	46,49
miR-107			
miR-23	PGC-1 $\alpha$	affects the ratio of oxidative red muscle and oxidative white muscle fibers	46,50,51
miR-181	Hox-A11	barely detectable in resting muscle, establishes the muscle phenotype	52,53
miR-206	Cx43	only highly-expressed in newly-formed muscle fibers, promotes myoblast differentiation and development	32,54,55
miR-486		Expressed postnatally, acts as an inhibitor of myogenesis	9
miR-376b		expressed prenatally, plays a role in promotion of myogenesis	9
miR-363		differentially expressed between 33 days post-gestation and adult life, long-term regulation of muscle growth and development	9
miR-365			
miR-422b			

### miRNAs in pig adipose tissue (Table 2)

In addition to skeletal muscle, adipose also affects the carcass value of a pig, including meat quality grade and yield. It also plays an important role in metabolic health. miR-143 was the first miRNA reported to be involved in adipose cell biology [58-60] (Table 2). The miR-210 [14,61] and the miR-27 family [14,62] are involved in adipogenesis. Several are abundant in both visceral and subcutaneous adipose tissues [60,61,63,64], whereas others are subcutaneous adipose tissue-specific enriched miRNAs [64-67]. Some are specifically expressed in the greater omentum [64,68,69]. Research into pig adipose tissue miRNAs may be beneficial in meeting the increasing demand of consumers for improved pork quality, which is a topic of worldwide concern [14].

**Table 2: miRNAs in pig adipose tissue**

miRNA	Target	Function	Reference
miR-143	ERK5	promotes adipocyte differentiation	58-60,63
miR-210		promotes adipogenesis	14,61
miR-27 family	INSR IRS1-4 PDK1/2 CREB S6K1	inhibit adipogenesis	14,62
miR-148a-3p		involved in differentiation of 3T3-L1 pre-adipocytes	61
let-7a-1-5p		play potential housekeeping roles in adipocytes	64
let-7f-5p			
miR-155-5p	C/EBP-b	inhibits adipogenesis	64-67
miR-193b-3p		act as central regulators of brown fat differentiation and adipogenesis	64
miR-365			
miR-374b-5p	C/EBP-b	involved in the effect of maternal dietary protein on lipid metabolism	64
miR-18a-3p		involved in development and production of pro-inflammatory B cells and T cells	64,68,69
miR-20-3p			
miR-19b-1-5p			
miR-181a-2-3p			
miR-181b-2-3p			



### miRNAs in other pig tissues (Table 3)

Although most research has hitherto been dedicated to miRNA's roles in meat quality, emerging research has evaluated miRNA in other solid tissues in pigs, including, but not limited to, reproduction. Many miRNAs are enriched in reproductive tissues [4,15,17,44] (Table 3). Others may play a role in differentiating neurons in brain development [3,13,70-73]. Several are highly-expressed in big solid organs [1,44,74-76]. The miR-200b and miR-214 are key miRNAs in tooth development [77,78].

When a miRNA is predominant, this suggests that it could have a significant role in the tissue, and that it could govern or be implicated in the major constitutive functions carried out by this tissue.

**Table 3: miRNAs in other pig tissues**

miRNA	Tissue	Function	Reference
miR-7	pituitary		15
miR-760	pituitary		44
miR-1296			
miR-137			
miR-362			
miR-153	mature testis		4
miR-205			
miR-196	immature		4
miR-149*	testis		
miR-485-3p			
miR-21-5p	ovary, testis		17
miR-9	head region	miR-9 regulates proliferation and migration of human neural progenitor cells	3,70
miR-30a			
miR-17	neurons and brain	involved in neurons differentiation and brain development by regulating APP	13,71
miR-106a			
miR-29c	adult cortex,	an effective biomarker of	72,73

	cerebellum	radiation-induced brain response	
miR-320	lung		74
miR-375	stomach and lymph nodes		74
miR-23a	kidney		1
miR-125b			
miR-23b			
miR-126			
miR-200b-3p			
miR-122-5p	liver	plays a role in cholesterol, fatty acid, and lipid metabolism	44,75,76
miR-200b	teeth	miR-200b is key in tooth development	77,78
miR-214			

### **Biomarker values of miRNA in xenotransplantation**

The pig has become the most suitable candidate as a source organ for xenotransplantation to overcome the growing gap between the need and availability of human donor organs [6]. Detailed genome information and emerging gene-editing technologies increase the possibility of producing pigs specific for this purpose. The xenotransplantation of organs from gene-modified pigs is associated with longer survival and less rejection [6]. Biopsy is the gold standard for diagnosis of conditions such as acute rejection (AR), disease recurrence, and drug toxicity [79]. However, biopsy often relies on ‘subjective’ measures, with some variability in results and reporting methods among pathologists, or limited diagnostic accuracy associated with sampling error [80]. There is a critical need for biomarkers for early diagnosis, treatment response, and outcome prediction in organ transplantation, with the final goal of an individualized treatment to prevent or reverse graft injury [81].

---

The miRNAs may be ideal candidates as biomarkers of disease [82-86]. Several factors (the relatively consistent changes seen in diseases, reliable analysis methods, tissue-specific expression patterns, less complexity than mRNAs, no post-processing modification, and amplifiable signals) contribute to making miRNAs ideal candidates, especially in the cancer diagnostic field [87-91]. Profiling miRNAs can be used as markers of organ donor quality/ischemia reperfusion injury [92]. The strong association between miRNA expression and allograft function or acute rejection demonstrates that miRNAs may be excellent biomarkers of human allograft status [81,93]. For example, levels of miR-142-5p, miR-155 and miR-223 can each predict acute rejection with >90% sensitivity and specificity in human renal allografts [94,95]. Because miRNAs are stably expressed in serum, plasma, urine, saliva, and other body fluids, this makes them ideal non-invasive biomarkers [83] to accurately monitor graft function in xenotransplantation. There are minor differences in the nucleotide composition of miRNAs among species [74]. Therefore, the circulating pig-specific miRNAs in human body fluids have vast potential to be biomarkers after pig-to-human xenotransplantation. Graft tissue and/or circulating miRNA profiles may be used as new biomarkers in guiding the diagnostic, therapeutic, and prognostic strategies that are associated with over-immunosuppression, organ toxicity, and graft rejection or loss.

## **Discussion**

Knowledge of human development, physiology, and pathology can be obtained from suitable animal models, especially the mouse and rat, but many of their physiological parameters (e.g., size, respiratory rate) are significantly different from those in humans [2].

---

Rodent genomes also have a faster rate of evolution than the human genome [96]. The pig is not only of significant agricultural value, but is also considered a good model for biomedical research [97]. Furthermore, pigs have been identified as the most promising source of organs for xenotransplantation to counteract the shortage of human organs for transplantation [98].

Many miRNAs are highly-conserved among related species [99]. Studies on miRNAs in the pig will be beneficial in understanding their key regulatory roles in humans. To obtain a better insight into the biological functions of miRNAs, it is imperative to identify all miRNAs expressed in the pig genome and their potential mRNA targets [100,101], which is becoming easier using bioinformatic methods, with a growing number of excellent tools becoming available [8,102]. However, the false discovery rate in predictive-results remains high, and experimental validation will be needed after bioinformatic prediction. We suggest, however, that pig miRNA profiles will be used as new biomarkers in pig-to-human xenotransplantation.

### **Acknowledgments**

This work was supported by Science and Technology Planning Project of Guangdong Province, China (2016A020226025), Shenzhen Foundation of Science and Technology (JCYJ20160229204849975, and GCZX2015043017281705), Shenzhen Foundation of Science and Technology grant (SZXJ2017021), and Fund for High Level University's Medical Discipline Construction (2016031638). The costs to publish in open access is covered by Shenzhen Foundation of Science and Technology grant (SZXJ2017021).

---

### **Author Contributions**

X.L.W. wrote the manuscript, D.C. revised the manuscript, Z.M.C editing the manuscript,

L.S.M. directed the work.

### **Conflict of interest**

The authors declare no conflict of interest.

### **Abbreviations**

miRNA microRNA

RISC RNA-induced silencing complexes

---

## References

1. Timoneda, O.; Balcells, I.; Nunez, J.I.; Egea, R.; Vera, G.; Castello, A.; Tomas, A.; Sanchez, A. Mirna expression profile analysis in kidney of different porcine breeds. *PLoS One* **2013**, *8*, e55402.
2. Martini, P.; Sales, G.; Brugiolo, M.; Gandaglia, A.; Naso, F.; De Pitta, C.; Spina, M.; Gerosa, G.; Chemello, F.; Romualdi, C., *et al.* Tissue-specific expression and regulatory networks of pig micrornaome. *PLoS One* **2014**, *9*, e89755.
3. Zhou, Y.; Tang, X.; Song, Q.; Ji, Y.; Wang, H.; Wang, H.; Jiao, H.; Ouyang, H.; Pang, D. Identification and characterization of pig embryo micrnas by solexa sequencing. *Reproduction in Domestic Animals* **2013**, *48*, 112-120.
4. Lian, C.; Sun, B.; Niu, S.; Yang, R.; Liu, B.; Lu, C.; Meng, J.; Qiu, Z.; Zhang, L.; Zhao, Z. A comparative profile of the micrna transcriptome in immature and mature porcine testes using solexa deep sequencing. *The FEBS journal* **2012**, *279*, 964-975.
5. Kragh, P.M.; Nielsen, A.L.; Li, J.; Du, Y.; Lin, L.; Schmidt, M.; Bogh, I.B.; Holm, I.E.; Jakobsen, J.E.; Johansen, M.G., *et al.* Hemizygous minipigs produced by random gene insertion and handmade cloning express the alzheimer's disease-causing dominant mutation appsw. *Transgenic research* **2009**, *18*, 545-558.
6. Cooper, D.K.C.; Ezzelarab, M.B.; Hara, H.; Iwase, H.; Lee, W.; Wijkstrom, M.; Bottino, R. The pathobiology of pig-to-primate xenotransplantation: A historical review. *Xenotransplantation* **2016**, *23*, 83-105.
7. Groenen, M.A.; Archibald, A.L.; Uenishi, H.; Tuggle, C.K.; Takeuchi, Y.; Rothschild, M.F.; Rogel-Gaillard, C.; Park, C.; Milan, D.; Megens, H.J., *et al.* Analyses of pig genomes provide insight into porcine demography and evolution. *Nature* **2012**, *491*, 393-398.
8. Kozomara, A.; Griffiths-Jones, S. Mirbase: Annotating high confidence micrnas using deep sequencing data. *Nucleic Acids Res* **2014**, *42*, D68-73.
9. Huang, T.H.; Zhu, M.J.; Li, X.Y.; Zhao, S.H. Discovery of porcine micrnas and profiling from skeletal muscle tissues during development. *PLoS One* **2008**, *3*, e3225.
10. McDaneld, T.G.; Smith, T.P.; Harhay, G.P.; Wiedmann, R.T. Next-generation sequencing of the porcine skeletal muscle transcriptome for computational prediction of micrna gene targets. *PLoS One* **2012**, *7*, e42039.
11. Xie, S.S.; Huang, T.H.; Shen, Y.; Li, X.Y.; Zhang, X.X.; Zhu, M.J.; Qin, H.Y.; Zhao, S.H. Identification and characterization of micrnas from porcine skeletal muscle. *Animal genetics* **2010**, *41*, 179-190.
12. Maxmen, A. Model pigs face messy path. *Nature* **2012**, *486*, 453.
13. Podolska, A.; Kaczkowski, B.; Kamp Busk, P.; Sokilde, R.; Litman, T.; Fredholm, M.; Cirera, S. Micrna expression profiling of the porcine developing brain. *PLoS One* **2011**, *6*, e14494.
14. Guo, Y.; Mo, D.; Zhang, Y.; Zhang, Y.; Cong, P.; Xiao, S.; He, Z.; Liu, X.; Chen, Y. Micrnaome comparison between intramuscular and subcutaneous vascular stem cell adipogenesis. *Plos One* **2012**, *7*, e45410.
15. Li, H.; Xi, Q.; Xiong, Y.; Cheng, X.; Qi, Q.; Yang, L.; Shu, G.; Wang, S.; Wang, L.;

- 
- Gao, P., *et al.* A comprehensive expression profile of micrnas in porcine pituitary. *Plos One* **2011**, *6*, e24883.
16. Sharbati, S.; Friedlander, M.R.; Sharbati, J.; Hoeke, L.; Chen, W.; Keller, A.; Stahler, P.F.; Rajewsky, N.; Einspanier, R. Deciphering the porcine intestinal micrna transcriptome. *BMC Genomics* **2010**, *11*, 275.
  17. Li, M.; Liu, Y.; Wang, T.; Guan, J.; Luo, Z.; Chen, H.; Wang, X.; Chen, L.; Ma, J.; Mu, Z., *et al.* Repertoire of porcine micrnas in adult ovary and testis by deep sequencing. *International journal of biological sciences* **2011**, *7*, 1045-1055.
  18. Kim, V.N. Micrna biogenesis: Coordinated cropping and dicing. *Nat Rev Mol Cell Biol* **2005**, *6*, 376-385.
  19. Bartel, D.P. Micrnas: Genomics, biogenesis, mechanism, and function. *Cell* **2004**, *116*, 281-297.
  20. Kim, V.N.; Han, J.; Siomi, M.C. Biogenesis of small rnas in animals. *Nat Rev Mol Cell Biol* **2009**, *10*, 126-139.
  21. Pillai, R.S.; Artus, C.G.; Filipowicz, W. Tethering of human ago proteins to mrna mimics the mirna-mediated repression of protein synthesis. *RNA* **2004**, *10*, 1518-1525.
  22. Ding, X.C.; Großhans, H. Repression of c. *Elegans* micrna targets at the initiation level of translation requires gw182 proteins. *EMBO J* **2009**, *28*, 213-222.
  23. Wakiyama, M.; Takimoto, K.; Ohara, O.; Yokoyama, S. Let-7 micrna-mediated mrna deadenylation and translational repression in a mammalian cell-free system. *Genes & development* **2007**, *21*, 1857-1862.
  24. Behm-Ansmant, I.; Rehwinkel, J.; Doerks, T.; Stark, A.; Bork, P.; Izaurralde, E. Mrna degradation by mirnas and gw182 requires both ccr4:Not deadenylase and dcp1:Dcp2 decapping complexes. *Genes & Development* **2006**, *20*, 1885-1898.
  25. Nottrott, S.; Simard, M.J.; Richter, J.D. Human let-7a mirna blocks protein production on actively translating polyribosomes. *Nat Struct Mol Biol* **2006**, *13*, 1108-1114.
  26. Carthew, R.W.; Sontheimer, E.J. Origins and mechanisms of mirnas and sirnas. *Cell* **2009**, *136*, 642-655.
  27. Giraldez, A.J.; Mishima, Y.; Rihel, J.; Grocock, R.J.; Van Dongen, S.; Inoue, K.; Enright, A.J.; Schier, A.F. Zebrafish mir-430 promotes deadenylation and clearance of maternal mrnas. *Science* **2006**, *312*, 75-79.
  28. Liu, N.; Williams, A.H.; Kim, Y.; McAnally, J.; Bezprozvannaya, S.; Sutherland, L.B.; Richardson, J.A.; Bassel-Duby, R.; Olson, E.N. An intragenic mef2-dependent enhancer directs muscle-specific expression of micrnas 1 and 133. *Proceedings of the National Academy of Sciences of the United States of America* **2007**, *104*, 20844-20849.
  29. Chen, J.F.; Mandel, E.M.; Thomson, J.M.; Wu, Q.; Callis, T.E.; Hammond, S.M.; Conlon, F.L.; Wang, D.Z. The role of micrna-1 and micrna-133 in skeletal muscle proliferation and differentiation. *Nat Genet* **2006**, *38*, 228-233.
  30. Lozano-Velasco, E.; Contreras, A.; Crist, C.; Hernandez-Torres, F.; Franco, D.; Aranega, A.E. Pitx2c modulates pax3+/pax7+ cell populations and regulates pax3 expression by repressing mir27 expression during myogenesis. *Dev Biol* **2011**, *357*, 165-178.

31. Allen, D.L.; Loh, A.S. Posttranscriptional mechanisms involving microRNA-27a and b contribute to fast-specific and glucocorticoid-mediated myostatin expression in skeletal muscle. *American journal of physiology. Cell physiology* **2011**, *300*, C124-C137.
32. Qin, L.; Chen, Y.; Liu, X.; Ye, S.; Yu, K.; Huang, Z.; Yu, J.; Zhou, X.; Chen, H.; Mo, D. Integrative analysis of porcine microRNAome during skeletal muscle development. *PLoS One* **2013**, *8*, e72418.
33. Zhang, J.; Ying, Z.Z.; Tang, Z.L.; Long, L.Q.; Li, K. MicroRNA-148a promotes myogenic differentiation by targeting the rock1 gene. *J Biol Chem* **2012**, *287*, 21093-21101.
34. Chen, L.; Wu, P.; Guo, X.H.; Hu, Y.; Li, Y.L.; Shi, J.; Wang, K.Z.; Chu, W.Y.; Zhang, J.S. Mir-143: A novel regulator of myod expression in fast and slow muscles of siniperca chuatsi. *Current molecular medicine* **2014**, *14*, 370-375.
35. Zuo, J.; Wu, F.; Liu, Y.; Xiao, J.; Xu, M.; Yu, Q.; Xia, M.; He, X.; Zou, S.; Tan, H., et al. MicroRNA transcriptome profile analysis in porcine muscle and the effect of mir-143 on the myh7 gene and protein. *PLoS One* **2015**, *10*, e0124873.
36. Gagan, J.; Dey, B.K.; Layer, R.; Yan, Z.; Dutta, A. MicroRNA-378 targets the myogenic repressor myor during myoblast differentiation. *J Biol Chem* **2011**, *286*, 19431-19438.
37. Hou, X.; Tang, Z.; Liu, H.; Wang, N.; Ju, H.; Li, K. Discovery of microRNAs associated with myogenesis by deep sequencing of serial developmental skeletal muscles in pigs. *PLoS One* **2012**, *7*, e52123.
38. Motohashi, N.; Alexander, M.S.; Casar, J.C.; Kunkel, L.M. Identification of a novel microRNA that regulates the proliferation and differentiation in muscle side population cells. *Stem Cells and Development* **2012**, *21*, 3031-3043.
39. Ryu, H.S.; Park, S.Y.; Ma, D.; Zhang, J.; Lee, W. The induction of microRNA targeting irs-1 is involved in the development of insulin resistance under conditions of mitochondrial dysfunction in hepatocytes. *PLoS One* **2011**, *6*, e17343.
40. Wang, S.; Aurora, A.B.; Johnson, B.A.; Qi, X.; McAnally, J.; Hill, J.A.; Richardson, J.A.; Bassel-Duby, R.; Olson, E.N. The endothelial-specific microRNA mir-126 governs vascular integrity and angiogenesis. *Developmental cell* **2008**, *15*, 261-271.
41. Nielsen, M.; Hansen, J.H.; Hedegaard, J.; Nielsen, R.O.; Panitz, F.; Bendixen, C.; Thomsen, B. MicroRNA identity and abundance in porcine skeletal muscles determined by deep sequencing. *Animal genetics* **2010**, *41*, 159-168.
42. Davis, E.; Caiment, F.; Tordoir, X.; Cavaille, J.; Ferguson-Smith, A.; Cockett, N.; Georges, M.; Charlier, C. Rnai-mediated allelic trans-interaction at the imprinted rtl1/peg11 locus. *Curr Biol* **2005**, *15*, 743-749.
43. Li, T.; Wu, R.; Zhang, Y.; Zhu, D. A systematic analysis of the skeletal muscle mirna transcriptome of chicken varieties with divergent skeletal muscle growth identifies novel mirnas and differentially expressed mirnas. *BMC Genomics* **2011**, *12*, 186.
44. Li, M.; Xia, Y.; Gu, Y.; Zhang, K.; Lang, Q.; Chen, L.; Guan, J.; Luo, Z.; Chen, H.; Li, Y., et al. MicroRNAome of porcine pre- and postnatal development. *PLoS One* **2010**, *5*, e11541.
45. Roush, S.; Slack, F.J. The let-7 family of microRNAs. *Trends in cell biology* **2008**, *18*, 505-516.



46. Liu, Y.; Li, M.; Ma, J.; Jie, Z.; Zhou, C.; Tao, W.; Gao, X.; Li, X. Identification of differences in miRNA transcriptomes between porcine oxidative and glycolytic skeletal muscles. *BMC Molecular Biology* **2013**, *14*, 1-8.
47. Niu, Z.; Li, A.; Zhang, S.X.; Schwartz, R.J. Serum response factor micromanaging cardiogenesis. *Current opinion in cell biology* **2007**, *19*, 618-627.
48. Boutz, P.L.; Chawla, G.; Stoilov, P.; Black, D.L. MicroRNAs regulate the expression of the alternative splicing factor nptb during muscle development. *Genes Dev* **2007**, *21*, 71-84.
49. Mahoney, D.J.; Parise, G.; Melov, S.; Safdar, A.; Tarnopolsky, M.A. Analysis of global mRNA expression in human skeletal muscle during recovery from endurance exercise. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* **2005**, *19*, 1498-1500.
50. Handschin, C.; Spiegelman, B.M. The role of exercise and PGC1 $\alpha$  in inflammation and chronic disease. *Nature* **2008**, *454*, 463-469.
51. Safdar, A.; Abadi, A.; Akhtar, M.; Hettinga, B.P.; Tarnopolsky, M.A. MiRNA in the regulation of skeletal muscle adaptation to acute endurance exercise in c57bl/6j male mice. *PLoS One* **2009**, *4*, e5610.
52. Naguibneva, I.; Ameyar-Zazoua, M.; Poleskaya, A.; Ait-Si-Ali, S.; Groisman, R.; Souidi, M.; Cuvellier, S.; Harel-Bellan, A. The miRNA mir-181 targets the homeobox protein hox-a11 during mammalian myoblast differentiation. *Nature cell biology* **2006**, *8*, 278-284.
53. Townley-Tilson, W.H.; Callis, T.E.; Wang, D. MicroRNAs 1, 133, and 206: Critical factors of skeletal and cardiac muscle development, function, and disease. *The international journal of biochemistry & cell biology* **2010**, *42*, 1252-1255.
54. Yuasa, K.; Hagiwara, Y.; Ando, M.; Nakamura, A.; Takeda, S.i.; Hijikata, T. MicroRNA-206 is highly expressed in newly formed muscle fibers: Implications regarding potential for muscle regeneration and maturation in muscular dystrophy. *Cell Structure and Function* **2008**, *33*, 163-169.
55. Anderson, C.; Catoe, H.; Werner, R. Mir-206 regulates connexin43 expression during skeletal muscle development. *Nucleic Acids Res* **2006**, *34*, 5863-5871.
56. McDanel, T.G.; Smith, T.P.; Doumit, M.E.; Miles, J.R.; Coutinho, L.L.; Sonstegard, T.S.; Matukumalli, L.K.; Nonneman, D.J.; Wiedmann, R.T. MicroRNA transcriptome profiles during swine skeletal muscle development. *BMC Genomics* **2009**, *10*, 77.
57. Brennecke, J.; Stark, A.; Cohen, S.M. Not mir-ly muscular: MicroRNAs and muscle development. *Genes & Development* **2005**, *19*, 2261-2264.
58. Esau, C.; Kang, X.; Peralta, E.; Hanson, E.; Marcusson, E.G.; Ravichandran, L.V.; Sun, Y.; Koo, S.; Perera, R.J.; Jain, R., *et al.* MicroRNA-143 regulates adipocyte differentiation. *J Biol Chem* **2004**, *279*, 52361-52365.
59. Cordes, K.R.; Sheehy, N.T.; White, M.P.; Berry, E.C.; Morton, S.U.; Muth, A.N.; Lee, T.H.; Miano, J.M.; Ivey, K.N.; Srivastava, D. Mir-145 and mir-143 regulate smooth muscle cell fate and plasticity. *Nature* **2009**, *460*, 705-710.
60. Li, G.; Li, Y.; Li, X.; Ning, X.; Li, M.; Yang, G. MicroRNA identity and abundance in developing swine adipose tissue as determined by solexa sequencing. *Journal of cellular biochemistry* **2011**, *112*, 1318-1328.
61. Qin, L.; Chen, Y.; Niu, Y.; Chen, W.; Wang, Q.; Xiao, S.; Li, A.; Xie, Y.; Li, J.; Zhao,

- X., *et al.* A deep investigation into the adipogenesis mechanism: Profile of micrnas regulating adipogenesis by modulating the canonical wnt/ $\beta$ -catenin signaling pathway. *BMC Genomics* **2010**, *11*, 320-320.
62. Karbiener, M.; Fischer, C.; Nowitsch, S.; Opriessnig, P.; Papak, C.; Ailhaud, G.; Dani, C.; Amri, E.Z.; Scheideler, M. Microna mir-27b impairs human adipocyte differentiation and targets ppargamma. *Biochem Biophys Res Commun* **2009**, *390*, 247-251.
63. Alexander, R.; Lodish, H.; Sun, L. Micrnas in adipogenesis and as therapeutic targets for obesity. *Expert opinion on therapeutic targets* **2011**, *15*, 623-636.
64. Ma, J.; Jiang, Z.; He, S.; Liu, Y.; Chen, L.; Long, K.; Jin, L.; Jiang, A.; Zhu, L.; Wang, J., *et al.* Intrinsic features in micrna transcriptomes link porcine visceral rather than subcutaneous adipose tissues to metabolic risk. *PLoS One* **2013**, *8*, e80041.
65. Liu, S.; Yang, Y.; Wu, J. Tnf $\alpha$ -induced up-regulation of mir-155 inhibits adipogenesis by down-regulating early adipogenic transcription factors. *Biochemical and Biophysical Research Communications* **2011**, *414*, 618-624.
66. Sun, L.; Xie, H.; Mori, M.A.; Alexander, R.; Yuan, B.; Hattangadi, S.M.; Liu, Q.; Kahn, C.R.; Lodish, H.F. Mir193b-365 is essential for brown fat differentiation. *Nature cell biology* **2011**, *13*, 958-965.
67. Pan, S.; Zheng, Y.; Zhao, R.; Yang, X. Microna-130b and microna-374b mediate the effect of maternal dietary protein on offspring lipid metabolism in meishan pigs. *The British journal of nutrition* **2013**, *109*, 1731-1738.
68. Xiao, C.; Srinivasan, L.; Calado, D.P.; Patterson, H.C.; Zhang, B.; Wang, J.; Henderson, J.M.; Kutok, J.L.; Rajewsky, K. Lymphoproliferative disease and autoimmunity in mice with increased mir-17-92 expression in lymphocytes. *Nature immunology* **2008**, *9*, 405-414.
69. Ventura, A.; Young, A.G.; Winslow, M.M.; Lintault, L.; Meissner, A.; Erkeland, S.J.; Newman, J.; Bronson, R.T.; Crowley, D.; Stone, J.R., *et al.* Targeted deletion reveals essential and overlapping functions of the mir-17 through 92 family of mirna clusters. *Cell* **2008**, *132*, 875-886.
70. Uchida, N. Microna-9 controls a migratory mechanism in human neural progenitor cells. *Cell Stem Cell* **2010**, *6*, 294-296.
71. Hebert, S.S.; Horre, K.; Nicolai, L.; Bergmans, B.; Papadopoulou, A.S.; Delacourte, A.; De Strooper, B. Microna regulation of alzheimer's amyloid precursor protein expression. *Neurobiology of disease* **2009**, *33*, 422-428.
72. Park, S.Y.; Lee, J.H.; Ha, M.; Nam, J.W.; Kim, V.N. Mir-29 mirnas activate p53 by targeting p85 alpha and cdc42. *Nat Struct Mol Biol* **2009**, *16*, 23-29.
73. Koturbash, I.; Zemp, F.; Kolb, B.; Kovalchuk, O. Sex-specific radiation-induced micrnaome responses in the hippocampus, cerebellum and frontal cortex in a mouse model. *Mutation research* **2011**, *722*, 114-118.
74. Xie, S.S.; Li, X.Y.; Liu, T.; Cao, J.H.; Zhong, Q.; Zhao, S.H. Discovery of porcine micrnas in multiple tissues by a solexa deep sequencing approach. *PLoS One* **2011**, *6*, e16235.
75. Gatfield, D.; Le Martelot, G.; Vejnar, C.E.; Gerlach, D.; Schaad, O.; Fleury-Olela, F.; Ruskeepaa, A.L.; Oresic, M.; Esau, C.C.; Zdobnov, E.M., *et al.* Integration of

- microrna mir-122 in hepatic circadian gene expression. *Genes Dev* **2009**, *23*, 1313-1326.
76. Esau, C.; Davis, S.; Murray, S.F.; Yu, X.X.; Pandey, S.K.; Pear, M.; Watts, L.; Booten, S.L.; Graham, M.; McKay, R., *et al.* Mir-122 regulation of lipid metabolism revealed by in vivo antisense targeting. *Cell metabolism* **2006**, *3*, 87-98.
77. Li, A.; Song, T.; Wang, F.; Liu, D.; Fan, Z.; Zhang, C.; He, J.; Wang, S. Micrornaome and expression profile of developing tooth germ in miniature pigs. *PLoS One* **2012**, *7*, e52256.
78. Sehic, A.; Risnes, S.; Khuu, C.; Khan, Q.-E.S.; Osmundsen, H. Effects of in vivo transfection with anti-mir-214 on gene expression in murine molar tooth germ. *Physiological Genomics* **2011**, *43*, 488-498.
79. Williams, W.W.; Taheri, D.; Tolckoff-Rubin, N.; Colvin, R.B. Clinical role of the renal transplant biopsy. *Nature reviews. Nephrology* **2012**, *8*, 110-121.
80. Kozakowski, N.; Regele, H. Biopsy diagnostics in renal allograft rejection: From histomorphology to biological function. *Transplant international : official journal of the European Society for Organ Transplantation* **2009**, *22*, 945-953.
81. Mas, V.R.; Dumur, C.I.; Scian, M.J.; Gehrau, R.C.; Maluf, D.G. Micrnas as biomarkers in solid organ transplantation. *American Journal of Transplantation Official Journal of the American Society of Transplantation & the American Society of Transplant Surgeons* **2013**, *13*, 11-19.
82. Maluf, D.G.; Dumur, C.I.; Suh, J.L.; Scian, M.J.; King, A.L.; Cathro, H.; Lee, J.K.; Gehrau, R.C.; Brayman, K.L.; Gallon, L., *et al.* The urine microrna profile may help monitor post-transplant renal graft function. *Kidney international* **2014**, *85*, 439-449.
83. Bhatt, K.; Mi, Q.S.; Dong, Z. Micrnas in kidneys: Biogenesis, regulation, and pathophysiological roles. *American journal of physiology. Renal physiology* **2011**, *300*, F602-610.
84. Chen, X.; Ba, Y.; Ma, L.; Cai, X.; Yin, Y.; Wang, K.; Guo, J.; Zhang, Y.; Chen, J.; Guo, X., *et al.* Characterization of micrnas in serum: A novel class of biomarkers for diagnosis of cancer and other diseases. *Cell Res* **2008**, *18*, 997-1006.
85. Godwin, J.G.; Ge, X.; Stephan, K.; Jurisch, A.; Tullius, S.G.; Iacomini, J. Identification of a microrna signature of renal ischemia reperfusion injury. *Proceedings of the National Academy of Sciences of the United States of America* **2010**, *107*, 14339-14344.
86. Kanak, M.A.; Takita, M.; Shahbazov, R.; Lawrence, M.C.; Chung, W.Y.; Dennison, A.R.; Levy, M.F.; Naziruddin, B. Evaluation of microrna375 as a novel biomarker for graft damage in clinical islet transplantation. *Transplantation* **2015**, *99*, 1568-1573.
87. Chawla, J.P.S.; Nageshwar, I.; Kanwaldeep Singh, S.; Atul, S.; Sunpreet Kaur, K.; Pratiksha, P. Role of mirna in cancer diagnosis, prognosis, therapy and regulation of its expression by epstein-barr virus and human papillomaviruses: With special reference to oral cancer. *Oral Oncology* **2015**, *51*, 731-737.
88. Sun, L.; Chen, Y.; Su, Q.; Tang, X.; Liang, Y.; Che, G.; Luo, F. Increased plasma mirna-30a as a biomarker for non-small cell lung cancer. *Medical Science Monitor International Medical Journal of Experimental & Clinical Research* **2016**, *22*,

- 
- 647-655.
89. Wang, J.; Ye, H.; Zhang, D.; Hu, Y.; Yu, X.; Long, W.; Zuo, C.; Yu, Y.; Xu, G.; Liu, S. MicroRNA-410-5p as a potential serum biomarker for the diagnosis of prostate cancer. *Cancer Cell International* **2016**, *16*, 1-6.
  90. Guo, S.; Guo, W.; Li, S.; Dai, W.; Zhang, N.; Zhao, T.; Wang, H.; Ma, J.; Yi, X.; Ge, R. Serum mir-16: A potential biomarker for predicting melanoma prognosis. *Journal of Investigative Dermatology* **2016**.
  91. Widodo; Djati, M.S.; Rifa'i, M. Role of microRNAs in carcinogenesis that potential for biomarker of endometrial cancer. *Annals of Medicine and Surgery* **2016**, *7*, 9-13.
  92. Shapiro, M.D.; Bagley, J.; Latz, J.; Godwin, J.G.; Ge, X.; Tullius, S.G.; Iacomini, J. MicroRNA expression data reveals a signature of kidney damage following ischemia reperfusion injury. *PLoS One* **2011**, *6*, e23011.
  93. Shan, J.; Feng, L.; Luo, L.; Wu, W.; Li, C.; Li, S.; Li, Y. MicroRNAs: Potential biomarker in organ transplantation. *Transplant immunology* **2011**, *24*, 210-215.
  94. Anglicheau, D.; Sharma, V.K.; Ding, R.; Hummel, A.; Snopkowski, C.; Dadhania, D.; Seshan, S.V.; Suthanthiran, M. MicroRNA expression profiles predictive of human renal allograft status. *Proceedings of the National Academy of Sciences of the United States of America* **2009**, *106*, 5330-5335.
  95. Tao, J.; Yang, X.; Han, Z.; Lu, P.; Wang, J.; Liu, X.; Wu, B.; Wang, Z.; Huang, Z.; Lu, Q., *et al.* Serum microRNA-99a helps detect acute rejection in renal transplantation. *Transplantation proceedings* **2015**, *47*, 1683-1687.
  96. Wu, C.I.; Li, W.H. Evidence for higher rates of nucleotide substitution in rodents than in man. *Proceedings of the National Academy of Sciences of the United States of America* **1985**, *82*, 1741-1745.
  97. Rocha, D.; Plastow, G. Commercial pigs: An untapped resource for human obesity research? *Drug Discov Today* **2006**, *11*, 475-477.
  98. Cooper, D.K.; Gollackner, B.; Sachs, D.H. Will the pig solve the transplantation backlog? *Annu Rev Med* **2002**, *53*, 133-147.
  99. Leung, A.K.; Sharp, P.A. MicroRNAs: A safeguard against turmoil? *Cell* **2007**, *130*, 581-585.
  100. Martini, P.; Sales, G.; Calura, E.; Brugiolo, M.; Lanfranchi, G.; Romualdi, C.; Cagnin, S. Systems biology approach to the dissection of the complexity of regulatory networks in the s. Scrofa cardiocirculatory system. *International journal of molecular sciences* **2013**, *14*, 23160-23187.
  101. Chen, C.; Ai, H.; Ren, J.; Li, W.; Li, P.; Qiao, R.; Ouyang, J.; Yang, M.; Ma, J.; Huang, L. A global view of porcine transcriptome in three tissues from a full-sib pair with extreme phenotypes in growth and fat deposition by paired-end rna sequencing. *BMC Genomics* **2011**, *12*, 448.
  102. Zhao, W.; Liu, W.; Tian, D.; Tang, B.; Wang, Y.; Yu, C.; Li, R.; Ling, Y.; Wu, J.; Song, S., *et al.* Waprna: A web-based application for the processing of rna sequences. *Bioinformatics* **2011**, *27*, 3076-3077.