

Article

Assembly of the *Boechera retrofracta* genome and evolutionary analysis of apomixis-associated genes

Sergei Kliver^{1,2}, Mike Rayko¹, Alexey Komissarov¹, Evgeny Bakin¹, Daria Zhernakova¹, Kasavajhala V. S. K. Prasad³, Catherine Rushworth⁴, R. Baskar⁵, Dmitry Smetanin⁶, Jeremy Schmutz^{7,8}, Daniel S. Rokhsar⁷, Thomas Mitchell-Olds⁹, Ueli Grossniklaus⁶, Vladimir Brukhin^{1,10} *

¹ Dobzhansky Center for Genome Bioinformatics, St. Petersburg State University, 41 Sredniy Prospekt, Vasilievsky Island, St. Petersburg 199004, Russia; mahajrod@gmail.com (S.K.), ad3002@gmail.com (A.K.), dashzhernakova@gmail.com (D. Z.), mikerayko@gmail.com (M. R.), eugene.bakin@gmail.com (E. B.), vbrukhin@gmail.com (V. B.)
² All-Russia Research Institute for Agricultural Microbiology, 196608, Podbelskogo sh., 3, Pushkin, St. Petersburg, Russia
³ Department of Biology, Colorado State University, Fort Collins, Colorado 80523, USA; e-mail: kasavajhalaprasad@gmail.com
⁴ University and Jepson Herbaria, University of California, Berkeley, USA; crushworth@berkeley.edu
⁵ Department of Biotechnology, Bhupat and Jyoti Mehta School of Biosciences, Indian Institute of Technology Madras, Chennai 600036, India; rbaskar@iitm.ac.in
⁶ Department of Plant & Microbial Biology Zurich-Basel Plant Science Center, University of Zurich, Zollikerstrasse 107, 8008 Zurich, Switzerland; dmitry.smetanin@botinst.uzh.ch (D. S.), grossnik@botinst.uzh.ch (U. G.)
⁷ Department of Energy Joint Genome Institute, Walnut Creek, California 94598, USA; dsrokhsar@gmail.com (D. S. R.)
⁸ HudsonAlpha Institute of Biotechnology, Huntsville, AL, USA; jschmutz@hudsonalpha.org (J. S.)
⁹ Department of Biology, Duke University, Durham NC 27708-0338 NC, USA; tmo1@duke.edu
¹⁰ Department of Plant Embryology & Reproductive Biology, Komarov Botanical Institute RAS, St. Petersburg, 197376 Russia.
* Correspondence: vbrukhin@gmail.com; Tel.: +7-965-046-5605

Abstract: Closely related to the model plant *Arabidopsis thaliana*, the genus *Boechera* is known to contain both sexual and apomictic species or accessions. *Boechera retrofracta* is a diploid sexually reproducing species and is thought to be an ancestral parent species of the apomictic species *Boechera divaricarpa*. Here we report the de novo assembly of the *B. retrofracta* genome using short Illumina and Roche reads from 1 paired-end and 3 mate pair libraries. The distribution of 23-mers from the paired end library has indicated a low level of heterozygosity and the presence of detectable duplications and triplications. The genome size was estimated to be equal 227 Mb. N50 of the assembled scaffolds was 2.3 Mb. 27048 protein-coding genes were predicted using a hybrid approach that combines homology-based and *de novo* methods. Also repeats, tRNA and rRNA genes were annotated. Finally, genes of *B. retrofracta* and 6 other *Brassicaceae* species were used for phylogenetic tree reconstruction. Also, a detailed analysis of evolution of the *APOLLO* apomixis-associated locus was performed. An assembled genome of *B. retrofracta* will help in the challenging assembly of the highly heterozygous genomes of hybrid apomictic species such as *B. divaricarpa*.

Keywords: *Boechera*, Brassicaceae, genome, assembly, annotation, apomixis

1. Introduction

Among over the 370 genera belonging to the family Brassicaceae (Cruciferae), only the genus *Boecheira* shows asexual reproduction by seeds [1-4]. Apomixis is defined as asexual reproduction through seeds that results in progeny identical to the maternal plant. The harnessing of apomixis is widely considered as a key enabling technology for crop improvement because it allows the fixation of any heterozygous genotype, leading to simpler and faster breeding schemes [5-7]. The *Boecheira* genus includes 110 sexual and apomictic species, widely distributed in North America. Plants from the *Boecheira* genus are represented by biannual and perennial herbs with a chromosome base number of $n = 7$ [8, 9].

Apomixis in the *Boecheira* genus is of special interest because it can occur at the diploid level, which is very rare. Furthermore, the phylogenetic proximity of *Boecheira* to the model plant *Arabidopsis thaliana* is attractive for potential functional studies. Although the genus *Boecheira* includes both sexual and apomictic species and accessions that are of variable ploidy and geographical origin, search for homologous sequences are feasible across the genus [10]. The sexual accessions of *Boecheira* are self-compatible and largely self-pollinating [11], unlike the sexual ancestors of most other apomicts, which are typically self-incompatible and cross-pollinating [12]. This inbreeding causes low heterozygosity in sexual *Boecheira* species. Apomictic *Boecheira* accessions have likely arisen through independent hybridisation events [13]. Their hybridogenic origin is supported by the aberrant structure of their chromosomes, as they are often observed as a consequence of hybridization, leading to allopolyploidy, aneuploidy, the replacement of homeologous chromosomes, and aberrant chromosomes [13, 14].

Certain apomictic *Boecheira* accessions are hypothesized to have arisen through hybridization between sexual *B. stricta* and *B. retrofracta* (Figure 1). *B. retrofracta* was previously included within *B. holboellii* (*sensu lato*) [15]. Up to now only the genome sequence of *B. stricta* was available [16], while the genome of *B. retrofracta* has not been assembled yet.

In this paper, we present the assembly and annotation of the *B. retrofracta* genome. The availability of the *B. retrofracta* genome sequence together with the previously assembled *B. stricta* genome will greatly help in the assembly and annotation of related apomictic hybrid species and provide the basis to investigate the peculiarities of hybridisation events, chromosomal organisation, the stability of apomictic genomes, and the genetic factors underlying apomixis. The performed assembly and annotation allowed to make a detailed analysis of *APOLLO* (APOmixis-Linked LOcus) genes, which may be involved in enucleation of sieve elements and, consequently, apomixis.

The following abbreviations will be used further: BAC – bacterial artificial chromosome BES – BAC end sequencing, PE – paired end, MP – mate pairs. An information about all the initial data and the final results availability see in Appendix B.

2. Materials and Methods

2.1. Sample Information

The reference *B. retrofracta* genotype was collected in Panther Creek, Lemhi County, Idaho, at 45°18'11.9"N 114°22'35.9"W, 1610 m elevation (see Figure 1). Plant growth, DNA extraction, and library construction were the same as with *B. stricta* [16]. Briefly, seedlings were germinated in aseptic culture in half-strength MS liquid medium. Cell nuclei were used for isolation of clean high-molecular-size nuclear DNA in TE buffer.



Figure 1. Plant and flower of *B. retrofracta*

2.2. Sequencing Strategy

The *B. retrofracta* genome was sequenced within the JGI Community Sequencing Project to produce sequence data for the *Boechera* genus [17]. Six libraries were constructed and sequenced using three platforms including Illumina, Roche, and Sanger: 1 PE (paired-end) library, 4 MP (mate pairs) libraries and 1 Sanger BAC end library. Read length and actual insert sizes for each library are given in Appendix A (see Table A1). This sequencing scheme was specially developed for the initial contig assembly by the DISCOVAR assembler [18], followed by scaffolding. Construction of genomic libraries and sequencing were performed following Lee et al [16].

2.2. Raw Data Filtration and pre-processing

Filtration of the PE library LIB400 was performed in two stages. First, reads containing long adapter fragments were removed using Cookiecutter [19]. Then Trimmomatic [20] was used to filter out reads with short adapter fragments. However, according to the DISCOVAR requirements no trimming or quality filtration was performed at those two stages. Only whole reads contaminated by adapters were discarded.

To process Illumina MP libraries LIB5000 and LIB7000 the NextClip [21] tool was modified to handle Cre-Lox libraries. It is important to note that original NextClip uses a very simple algorithm to align linker sequences to reads. It takes into account only the number of matching bases. As the CreLox linker is significantly longer than the Nextera linker, the number of false hits may significantly increase. To mitigate this effect, a requirement for the presence of a continuous 9-bp core alignment was added. The modified tool was named CreClip and can be found in [22].

Reads from Roche MP Libraries LIB4000R and LIB24000R were split into “forward” and “reverse” segments separated by linker. Then low quality ends were trimmed from “reverse” segments by Trimmomatic. Finally, reverse segments were reverse complemented to mimic to Illumina MP libraries.

2.2. Genome Size Estimation

Estimation of the genome size based on the 23-mer distribution (as well as other *k*-mer based statistics) was performed using the KrATER software [23] on the LIB400 library and further compared with the previous estimations of *Boechera* genus [24].

2.3. Genome Assembly and Quality Evaluation

At the assembly stage initial contigs were constructed from the filtered LIB400 reads by DISCOVAR. Then, the obtained contigs were extended using a BAC end sequencing (BES) library and the SSPACE scaffolder [25].

Before scaffolding the assessment of the actual (mean) insert size was performed. Filtered reads from all libraries were aligned to initial contigs by BWA [26]. For each library, only alignments to contigs with 3x length of the target insert size were used in the estimation (Table 1) to minimize alignment artifacts. Next, the extended contigs were scaffolded by SSPACE in two stages: at the first stage, all four MP libraries (LIB4000R, LIB5000, LIB7000, LIB24000R) were used to produce raw scaffolds, at the second stage, raw scaffolds were linked to the intermediate scaffolds using the BES library only. Scaffolding was carried out in several stages because different options were required to utilize the BES data. Gap closing in the intermediate scaffolds was performed using GapCloser (a module for SOAPdenovo2) [27] on the LIB400 library only. Finally, all scaffolds with a length of less than 250 bp (i.e. less than read length of LIB400, the library used for initial contig construction) were filtered out, as the corresponding short fragments most likely are the assembly artifacts. Integrity of the assembly was verified by CEGMA [28] and BUSCO [29]. A schematic diagram of the assembly pipeline is shown in Figure A1 in Appendix A.

2.4. Repeats Analysis

A *de novo* repeat identification in the *B. retrofracta* genome was performed using RepeatModeler [30] with default parameters. The obtained repeat library was combined with *A. thaliana* repeats from RepBase [31], and the merged library was used to annotate repeats by RepeatMasker [32]. Then repeats in the *B. retrofracta* genome were softmasked by Bedtools [33] for the prediction of protein coding genes. Also, masking of tandem and interspersed repeats by TRF [34] and Windowmasker [35], respectively, were performed.

2.5. Variants Calling and Genotyping

For variant calling and genotyping filtered reads were aligned to the assembled genome using BWA mem with default options. Next, the GATK pipeline [36] for variant calling was applied in the following way: duplicates were marked using Picard MarkDuplicates, realigned reads at indels, and, finally, HaplotypeCaller was used to call variants. Only Single Nucleotide Polymorphisms (SNPs) and Indels passing the following filtering criteria were kept: QualByDepth (QD) > 2.0, FisherStrand (FS) < 20.0, RMSMappingQuality (MQ) > 40.0, MappingQualityRankSumTest (MQRankSum) > -12.5, ReadPosRankSumTest (ReadPosRankSum) > -8.0 for SNPs, and QualByDepth (QD) > 2.0, FisherStrand (FS) < 20.0, ReadPosRankSumTest (ReadPosRankSum) > -20.0 for indels, respectively. Finally, the variants falling into the repeats masked by RepeatMasker were excluded.

2.6. Prediction of Protein-coding Genes and ncRNA

The prediction of protein-coding genes was performed using a combined approach that synthesizes both homology-based and *de novo* predictions, where *de novo* predictions are used only to fill gaps and to extend the homology-based predictions. Pure *de novo* predictions were filtered out.

As homology-based evidence for gene presence, we have used proteins and transcripts of 5 closely-related species. Proteins of the four reference species – *Arabidopsis thaliana* (assembly TAIR10), *Brassica rapa* (Brapa_1.0), *Capsella rubella* (Caprub1_0), and *Eutrema salsugineum* (Eutsalg1_0) were aligned to the *B. retrofracta* assembly by Exonerate [37], using the protein2genome model with a maximum of three hits per protein. The obtained alignments were classified into the top (primary) and secondary hits; the coding sequence (CDS) fragments were cut from each side by 3 bp for the top hits and by 9 bp for the secondary hits. Transcripts of *B. stricta* (assembly v1.2, [16]) with marked CDS regions were also aligned to the *B. retrofracta* genome by Exonerate using the

cdna2genome model leaving the other options unchanged. Alignments of CDS segments were not cut for top hits, but cut by 3bp for secondary hits.

These truncated fragments were clustered and supplied as hints to the AUGUSTUS software package [38], and the CDS segments of genes were predicted in the soft-masked *B. retrofracta* assembly using *A. thaliana* gene models. Proteins were translated from the predicted genes and aligned by HMMER 3.1 [39] and BLAST [40] to the Pfam [41] and Swiss-Prot [42] databases, respectively. Only genes supported by the both hints and hits to one of the protein databases were retained; the rest were discarded. tRNA and rRNA genes were predicted by tRNAscan-SE v1.3.1 [43] and Barrnap v0.6 [44], respectively.

2.7. Phylogenetic Analysis

The longest proteins corresponding to each predicted gene of *B. retrofracta* and six other *Brassicaceae* species - *B. stricta* (assembly v1.2), *A. thaliana* (TAIR10), *A. lyrata* (v.1.0), *C. rubella* (*Caprub1_0*), *Cardamine hirsuta* (v1.0), and *E. salsugineum* (Eutsalg1_0) - were aligned to profile HMM models of the braNOG subset from the eggNOG database [45] using HMMER. The top hits from the alignments were extracted and used for assignment of the corresponding proteins to orthologous groups, followed by extraction of single-copy orthologs.

To verify topology concordance and get a basis for future studies of positive selection, a species tree reconstruction was performed. Single-copy orthologous proteins of the seven species included in the analysis were aligned by MAFFT [46]. Based on the obtained protein alignments, a maximum likelihood tree was reconstructed by RAxML v8.2 [47] with the PROTGAMMAAUTO option, and the JTT fitting model was tested with 1,000 bootstrap replications. The tree was rooted with *E. salsugineum* as an outgroup. The resulting tree was drawn with FigTree software [48].

2.7. APOLLO Evolution Analysis

The evolutionary history of APOLLO gene was inferred by using the Maximum Likelihood method based on the Tamura-Nei model [51]. Initial alignment of corresponding CDS was performed using prank v.140110 [49] in codon-aware mode. The tree with the highest log likelihood (-12153.79) was selected (see Fig. 4). Initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using the Maximum Composite Likelihood (MCL) approach, and then selecting the topology with superior log likelihood value. All positions containing gaps and missing data were eliminated. There were a total of 1158 positions in the final dataset. Evolutionary analyses were conducted in MEGA7 [52].

3. Results

3.1. k-mer Based Statistics

k-mer spectrum built by KrATER is shown in Figure 2. The 23-mer distribution has a peak of erroneous 23-mers at 1x coverage corresponding to sequencing errors and one major peak at 371x coverage corresponding to diploid 23-mers (shared between homologous chromosomes), but no significant peak related to heterozygous genome positions was detected (Figure 2). However, we detected several small additional peaks at double (737x) and triple (1120x) depth, which are probably related to duplications and triplications, respectively.

The genome size of *B. retrofracta* was estimated to be close to 227 Mbp, which is close to the previous estimations of a minimal genome size of 200 Mbp in the *Boechera* genus [24].




Figure 2. Distribution of 23-mers for PE LIB400 library. Only one major peak at 371x coverage is present, suggesting low heterozygosity (bottom plot), however there are detectable duplications and triplications at 737x and 1120x coverage (upper plot, Y axis is on a logarithmic scale).

3.2. Genome Assembly and Evaluation

We have achieved N50 of 2,297,899 bp, L50 of 25, and a total assembly length of 222.25 Mbp for the final scaffolds, which is very close to our 23-mer based estimation. Detailed statistics including N50 and total assembly values for every stage of the assembly pipeline are listed in Tables 1 and 2. It is important to note that the final assembly (Table 1, column final scaffolds) have smaller size than previous intermediate assemblies due to the last filtration step. All scaffolds shorter than 250 bp (a read length of LIB400) were treated as artifacts of assembly and were removed. However, size of final assembly(222.25 Mbp) is closer to estimated genome size (226.87 Mbp) than the size of intermediate assemblies.

Evaluation of the assembly completeness was performed using CEGMA and BUSCO. 242 (97.58%) complete CEGs were identified in the assembled genome. Out of 1440 BUSCO genes from the Embryophyta, set only 12 (0.8%) genes were not found, 6 were fragmented, 36 (2.5%) were duplicated and 1422 (98.8 %) were complete. This high fraction of complete BUSCO genes suggests high completeness of the assembly and its integrity at least in gene-coding regions.

Table 1. General statistics for all stages of the assembly pipeline.

Parameter	Contigs	Extended contigs	Raw scaffolds	Intermediate scaffolds	Gap closed scaffolds	Final scaffolds
Longest contig	791 985	792 340	8 101 256	9 045 706	9 049 080	9 049 080
Ns	28 100	28 100	11 890 519	16 366 994	12 409 189	12 409 189
Total length	225 649 216	226 402 628	236 469 041	240 945 496	241 014 839	222 253 471

Table 2. N50 values for all stages of the assembly pipeline and several different cutoffs for minimal scaffold length.

scaffold length cutoff	Contigs	Extended contigs	Raw scaffolds	Intermediate scaffolds	Gap closed scaffolds	Final scaffolds
all	85 286	84 648	1 256 534	1 898 006	1 898 985	2 297 899
>=100	85 286	84 648	1 256 534	1 898 006	1 898 985	2 297 899
>=250	101 388	100 393	1 442 421	2 296 484	2 297 899	2 297 899
>=500	115 732	115 486	1 538 795	2 678 857	2 680 941	2 680 941
>=1000	122 300	121 678	1 704 064	2 678 857	2 680 941	2 680 941

3.3. Repeats Annotation

In total approximately 85 Mbp (38.13 %) of the assembly were masked. The detailed description of the annotated repeat types is listed in Table 3. It is important to note that a large number (10.96% of the assembly size) of interspersed repeats was not classified. The results are shown in Table 4.

Table 3. Repeats found by RepeatMasker.

Class	Number of elements	Total length, bp	Fraction of assembly, %
SINEs	577	125 298	0.06
LINEs	7 075	4 351 241	1.96
LTR elements	51 040	40 608 195	18.27
DNA elements	31 638	12 868 684	5.79
Unclassified	82 693	24 363 135	10.96

Total interspersed repeats	-	82 316 553	37.04
Small RNA	5 461	1599354	0.72
Satellites	1 541	573 026	0.26
Simple repeats	2044	363 642	0.16
Low complexity	56	7 456	0

Table 4. Results of repeat masking performed by three different tools: RepeatMasker, TRF, Windowmasker.

Tool	Number of repeats	Total length, Mbp
RepeatMasker	173 023	82.31
TRF	100 593	17.41
Windowmasker	1 104 650	64.20

3.4. Variant Calling and Genotyping

In the genome 3341 SNPs and 1317 indels were detected. Among these, 103 (3.08 %) SNPs and 97 (7.37%) indels were homozygous and, therefore, most likely artifacts of alignment or SNP calling. Mean heterozygous SNP and indel densities in non-masked regions (138 Mbp in total) are 0.0235 SNP and 0.0089 indel per kbp, respectively, suggesting a very low heterozygosity of the *B. retrofracta* genome.

3.5. Prediction of Protein-coding Genes and ncRNAs

In total 27,048 genes with 28,269 transcripts were predicted. tRNA and rRNA genes predicted by tRNAscan-SE and Barrnap are given in Tables 5 and 6 respectively.

Table 5. Annotated tRNAs.

tRNA type	Number
tRNAs decoding standard 20 AA	1126
Selenocysteine tRNAs (TCA)	0
Possible suppressor tRNAs (CTA,TTA)	3
tRNAs with undetermined isotypes	5
Resolution of Brassicaceae Phylogeny Using Nuclear Genes	32
Uncovers Nested Radiations and Supports Convergent	
Morphological Evolution Predicted pseudogenes	
Total tRNAs	1 166

Table 6. Annotated rRNAs.

rRNA	Complete (>= 80% of expected length)	Partial (<80% of expected length)
5.8S	178	53
5S	601	104
28S	0	1 782
18S	1	1 458
12S	0	173
16S	0	607

3.5. Species tree reconstruction

In course of the assignment of proteins to orthologous groups 8,959 single-copy orthologs were identified among the seven species (*B. retrofracta*, *B. stricta*, *A. thaliana*, *A. lyrata*, *C. rubella*, *Cardamine hirsuta*, and *E. salsugineum*).

The corresponding phylogenetic tree was rooted with *E. salsugineum* as an outgroup (see Figure 3). All nodes have a high support and no topology discordance was found with the tree reconstructed previously by Huang et al, 2015 [52].




Figure 3. Phylogenetic tree of seven Brassicaceae species used for analysis. Maximum likelihood tree was reconstructed by RAxML using 8959 single copy orthologs and was tested with 1000 bootstrap replicates. Numbers near nodes represent corresponding bootstrap support.

3.6. Analysis of Evolution of the APOLLO locus

Results from Corral et. al. [53] suggest that *APOLLO* (Aspartate Glutamate Aspartate Aspartate histidine exonuclease) is one of the important apomixis-related genes in *Boechera*. It was shown that the *APOLLO* locus has several alleles with apomixis-associated polymorphisms. All studied apomictic plants carry at least one of the “apoalleles”, while both copies in sexual genotypes were “sexalleles”.

In this study we decided to take a closer look to this locus in our assembly and other *Brassicaceae* species in this study. Along with an exact copy of the *APOLLO* locus, we also found two other, more distant copies, which may indicate past duplication events. We searched for these orthologs in other species, and reconstructed phylogenetic tree (see Figure 4). All *Brassicaceae* genomes in the study also carried these three copies, related to the clusters of orthologous genes ENOG410BURN (*APOLLO* locus), ENOG410BUTR, and ENOG410C333 in the EggNOG database.

We observed that branches in the tree were grouped by genes rather than by species, suggesting that the triplication event took place before the separation of the *Brassicaceae* species in this study. It worth noting that in *Populus trichocarpa* genome there is only one copy of these locus, which gives an upper-bound time estimate of the series of duplication events.

We also examined *APOLLO* alleles (both apo- and sex-alleles) described in *Boechnera* spp. by Corral et al. We can see that these alleles arise after the separation of the *Boechnera* genus, and compose two separate clades. Given the fact that *B.retrofracta* and *B.stricta* are the sexual species, it was not surprising that in both cases all corresponding polymorphic sites were in the “sexallele”-state, and clustered with sex-alleles.

We calculated the *Ka/Ks* ratio for the internal branches in this tree and found that branch leading to apo-alleles is under positive selection (*Ka/Ks* 1.4646, the branch is shown in red in Figure 3), which is typical for paralogues that are required to serve a novel function.

The *APOLLO* gene was initially described in *A. thaliana* as an exonuclease, protein NEN3, Q9CA74 in Uniprot database, probably involved in enucleation of sieve elements, whereas two other copies were described as NEN1 (Q9FLR0) and NEN2 (Q0V842). Given that, we may suggest an evolutionary scenario where, after the series of duplications, one of the NEN protein copies in the common ancestor of *Boechnera* spp might have acquired alter regulation, and might induce development of the apomictic reproduction from the ancestral “sexual” state, following by separation of the apomictic lineages.

That could explain the phenomena of the diploid apomictic *Boechnera*, emerged as a result of duplication events rather than polyploidy.




Figure 4. Phylogenetic tree of the isoforms of *APOLLO* locus (exonuclease NEN) in seven species of interest and alleles of *APOLLO* locus of apomictic *Boechnera* species from Corral et al (2013). Sequences of *Populus trichocarpa*, *Vitus vinifera* and *Glycine max* were used as outgroup. The clade related to the *APOLLO* locus is shown in blue, with apo-alleles shown in red.

Table 7. Comparison of genome characteristics of *B.retrofracta* with previously sequenced *B.stricta* and *A.thaliana* genomes. Source for *B.retrofracta* - this paper, *B.stricta*, *A.lyrata* and *A.thaliana* - Phytozome v12.1 database [54]

	<i>B. retrofracta</i>	<i>B. stricta</i> v.1.2	<i>A.lyrata</i> v2.1	<i>A. thaliana</i> TAIR10
Total length	227 M	184 M	207 Mb	135 Mb
Chromosomes	<i>n</i> = 7	<i>n</i> = 7	<i>n</i> = 8	<i>n</i> = 5
Protein-coding loci	27,048	27,416	31,073	27,416
Transcripts	28,269	29,812	33,132	35,386

4. Discussion

In this study we present a *de novo* assembly and annotation of the genome of *Boechnera retrofracta*, a perennial flowering plant belonging to Brassicaceae family that is native to North America. The genome of *B. retrofracta* demonstrated a very low level of heterozygosity. Notably, repeats in the genome of *B. retrofracta* occupied almost 40% of the genome space. Nearly half of them were LTRs (18.27%). The genome size was found to be 227 Mb, nearly two-fold larger than the *Arabidopsis thaliana* genome, and one quarter greater than *Boechnera stricta*, and almost equal to *Arabidopsis lyrata* genome size (Table 7). At the same time the amount of protein-coding genes in the genome of *B. retrofracta* is slightly less than in the *B. stricta* and *A. thaliana* genomes and much less than that in the *A. lyrata* genome (Table 1). Despite the largest genome size, the number of predicted transcripts in *B. retrofracta* is the smallest among the four Brassicaceae species compared (Table 1). The presence of a slightly greater number of genes in *B. stricta* compared with *B. retrofracta*, despite a smaller genome size, may be associated with aneuploidy of the chromosomal fragments, or genome rearrangements occurred as a result of interhybridization, which is characteristic of many *Boechnera* species and accessions.

As an example of how the genome of the sexual species *B. retrofracta* could be used to study evolution and origin of apomixis, we performed an evolutionary analysis of the three alleles of the *APOLLO* (APOmixis-Linked LOcus) gene (apo- and sex-alleles) described by Corral et al [53]. We examined this gene in more detail in our assembly and in other Brassicaceae species. Along with the described copy of *APOLLO*, we also found two other, more distant copies, which evidently arose by two sequential gene duplications (triplication). The *APOLLO* phylogenetic tree may indicate that triplication event occurred before the separation of Brassicaceae species under study (Figure 3). We also analyzed the *APOLLO* alleles described in *Boechnera ssp.* It was clear that these alleles arose after separation of the *Boechnera* genus. In sexual *B.retrofracta* and *B.stricta* polymorphic sites corresponded to the “sexallele”-state and clustered with sex-alleles of the other species.

These results are compatible with an evolutionary scenario where, after the series of duplications, one of the NEN exonuclease protein (ancestor of *APOLLO*) copies in the common ancestor of *Boechnera spp* experiencing relaxed selection might be deregulated, promoting development of the apomictic reproduction from the ancestral “sexual” state, following by separation of the apomictic lineages. This model of evolution of *APOLLO* alleles might explain the phenomena of apomictic development in *Boechnera* in the diploid condition, emerged as a result of duplication events rather than polyploidy.

In conclusion, increasing number of sequenced genomes from the economically important Brassicaceae family will facilitate future genetic, genomic, evolutionary, and domestication studies in this family. *B. retrofracta* is thought to be an ancestor of certain hybrids including apomictic species, for example *B. divaricarpa*. Consequently, the genome assembly presented in this report may help with the challenging genome assembly of highly heterozygous hybrid *Boechnera* species that are apomictic. Thus, the *B. retrofracta* genome reported here will provide a basis to decipher the hybridogenesis events that led to the formation of apomictic *Boechnera* accessions.

Acknowledgments: This work was supported by grants 16-54-21014 and 15-54-45001 from the Russian Foundation for Basic Research and by grant 1.52.1647.2016 from St. Petersburg State University (to VB), grant IZLRZ3_163885 from the Swiss National Science Foundation (to UG) through the Scientific & Technological Cooperation Programme Switzerland-Russia (STCPSR). US National Science Foundation Doctoral Dissertation

Improvement Grant 1311269 (to CR) and grant R01 GM086496 from the National Institutes of Health (USA) (to TMO). Work conducted by the US Department of Energy Joint Genome Institute was supported by the Office of Science of the US Department of Energy under Contract No. DE-AC02-05CH11231.

Author Contributions: TMO, VB and UG designed the study. TMO, CR, JS, KVS KP gathered samples, extracted and sequenced genomic DNA. SK, AK, DZ and MR performed genome assembly and annotation. SK, VB, MR, EB and RB wrote manuscript with revision by all other authors. All authors read and approved the final manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Table A1. Sequencing scheme of *B. retrofracta* genome

ID	Library type	Platform	Read length	Mean insert size, bp
LIB400	paired ends	Illumina	250	402
LIB4000R	mate pairs	Roche	-	4 014
LIB5000	mate pairs	Illumina	150	4 877
LIB7000	mate pairs	Illumina	150	6 882
LIB24000R	mate pairs	Roche	-	24 332
BES	BAC end sequencing	Sanger	-	147 708



Figure A1. Pipeline used to assembly genome of *B. retrofracta*.

Appendix B

The original data could be found at:
<http://public.dobzhanskycenter.ru/ad89dedc8b4674276c9b0760f29b07af/>
http://garfield.dobzhanskycenter.org/boechera_retrofracta/
http://garfield.dobzhanskycenter.org/cgi-bin/hgTracks?db=hub_65_b_retrofracta&lastVirtModeType=e=default&lastVirtModeExtraState=&virtModeType=default&virtMode=0&nonVirtPosition=&position=scaffold100%3A1-247471&hgsid=135_4PIU5GWT5fPBAsQNvjXcOsZU2vBG
http://garfield.dobzhanskycenter.org/cgi-bin/hgGateway?hgsid=136_aTuM2vMkx2XM8rnOkblRRRQZDuJA
or at NCBI, BioProject ID: PRJNA418376.

References

1. Sharbel TF and Mitchell-Olds T. Recurrent polyploid origins and chloroplast phylogeography in the *Arabis holboellii* complex (Brassicaceae). *Heredity*, 2001, 87: 59. DOI: 10.1046/j.1365-2540.2001.00908.x
2. Schranz ME, Dobes C, Koch MA and Mitchell-Olds T. Sexual reproduction, hybridization, apomixis and polyploidization in the genus *Boechera* (Brassicaceae). *American J Botany*, 2005, 92: 1797 - 810. DOI: 10.3732/ajb.92.11.1797
3. Naumova TN. Apomixis and amphimixis in flowering plants. *Cyt.Genetics*, 2008, 3: 51-63. DOI: 10.3103/S0095452708030055

4. Aliyu OM, Schranz ME and Sharbel TF Quantitative variation for apomictic reproduction in the genus Boechera (Brassicaceae). *Am. J. Bot.*, 2010, 97: 1719. DOI: 10.3732/ajb.1000188

5. Koltunow AM and Grossniklaus U. Apomixis: a developmental perspective. *Annu. Rev. Plant Biol.*, 2003, 54: 547–574. DOI: 10.1146/annurev.arplant.54.110901.160842

6. Rodriguez-Leal D, Vielle-Calza da J-P. Regulation of apomixis: learning from sexual experience. *Curr Opin Plant Biol.*, 2012, 15: 549–555. DOI: 10.1016/j.pbi.2012.09.005

7. Brukhin V. Molecular and genetic regulation of apomixis. *Russ. J. Genetics*, 2017, 53: 943–964. DOI: 10.1134/s1022795417090046

8. Windham MD, and Al-Shehbaz IA. New and noteworthy species of Boechera (Brassicaceae) II: apomictic hybrids. *Harvard Pap. Bot.*, 2007, 11: 257–274. DOI: 10.3100/1043-4534(2007)11[257:nansob]2.0.co;2

9. Windham MD, and Al-Shehbaz IA. New and noteworthy species of Boechera (Brassicaceae) III: additional sexual diploids and apomictic hybrids. *Harvard Pap. Bot.*, 2007b, 12: 235–257. DOI: 10.3100/1043-4534(2007)12[235:nansob]2.0.co;2

10. Lovell, J. T., et al. Mutation Accumulation in an Asexual Relative of Arabidopsis. *PLoS Genetics*, 2017, 13(1): e1006550. DOI: 10.1371/journal.pgen.1006550

11. Rushworth, C. A., et al. Boechera, a model system for ecological genomics. *Molecular Ecology*, 2011, 20(23): 4843–4857. DOI: 10.1111/j.1365-294x.2011.05340.x

12. Beck JB, Alexander PJ, Allphin L, Al-Shehbaz IA, Rushworth C, Bailey CD et al. Does hybridization drive the transition to asexuality in diploid Boechera? *Evolution*, 2011, 66: 985–995. DOI: 10.1111/j.1558-5646.2011.01507.x

13. Kantama L, Sharbel TF, Schranz ME, Mitchell-Olds ., de Vries Sde Jong H. Diploid apomicts of the Boechera holboellii complex display large-scale chromosome substitutions and aberrant chromosomes. *Proc. Natl. Acad. Sci. USA*, 2007, 104: 14026–14031. DOI: 10.1073/pnas.0706647104

14. Mandáková T., Schranz M.E., Sharbel T.F., de Jong H., Lysak M.A. Karyotype evolution in apomictic Boechera and the origin of the aberrant chromosomes. *The Plant Journal*, 2015, 82: 785–793. DOI: 10.1111/tpj.12849

15. Windham MD, and Al-Shehbaz IA. New and noteworthy species of Boechera I: sexual diploids. *Harvard Pap. Bot.*, 2006, 11: 61–88. DOI: 10.3100/1043-4534(2006)11[61:nansob]2.0.co;2

16. Lee CR, Wang B, Mojica JP, Mandáková T, Prasad KVSK, Goicoechea JL, et al. Young inversion with multiple linked QTLs under selection in a hybrid zone. *Nat. Ecol. Evol.*, 2017, 1: 119. DOI: 10.1038/s41559-017-0119

17. Why sequence Boechera holboellii? <http://jgi.doe.gov/why-sequence-boechera-holboellii/> Accessed 01 Nov 2017

18. Weisenfeld NI, Yin S, Sharpe T, Lau B, Hegarty R, Holmes L, et al. Comprehensive variation discovery in single human genomes. *Nat. Genetics*, 2014, 46: 1350–1355. DOI: 10.1038/ng.3121

19. Starostina E. Tamazian G, Dobrynin P, O'Brien S and Komissarov A. Cookiecutter: a tool for kmer-based read filtering and extraction. *bioRxiv*, 2015, 024679. DOI: 10.1101/024679

20. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, 2014, 30: 2114–2120. DOI: 10.1093/bioinformatics/btu170

21. Leggett R M, Clavijo BJ, Clissold L, Clark MD and Caccamo M. NextClip: an analysis and read preparation tool for Nextera Long Mate Pair libraries. *Bioinformatics*, 2013, 30: 566–568. DOI: 10.1093/bioinformatics/btt702

22. Kliver S. CreClip. <https://github.com/mahajrod/CreClip> Accessed on 01 Jul 2016.

23. Kliver S, Tamazian G, O'Brien SJ, Brukhin V, Komissarov A. KrATER (K-mer Analysis Tool Easy to Run). <https://github.com/mahajrod/KrATER> Accessed on 01 Nov 2017

24. Anderson J T, J Willis JH and Mitchell-Olds T. Evolutionary genetics of plant adaptation. *Trends Genetics*, 2011, 27: 258–266. DOI: 10.1016/j.tig.2011.04.001 Actions

25. Boetzer M, Henkel CV, Jansen HJ, Butler D and Pirovano W. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics*, 2010, 27: 578–579. DOI: 10.1093/bioinformatics/btq683

26. Li H, and Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics*, 2009, 25: 1754–1760. DOI: 10.1093/bioinformatics/btp324

27. Luo, R., Liu, B., Xie, Y., Li, Z., Huang, W., Yuan, J. and Tang, J. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. *Gigascience*, 2012, 1(1), 18. DOI: 10.1186/2047-217x-1-18

28. Parra, G., Bradnam, K., & Korf, I. CEGMA: a pipeline to accurately annotate core genes in eukaryotic genomes. *Bioinformatics*, 2007, 23(9), 1061-1067. DOI: 10.1093/bioinformatics/btm071
29. Simão F A, Waterhouse RM. Ioannidis P, Kriventseva EV and Zdobnov EM. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, 2015, 31: 3210-3212. DOI: 10.1093/bioinformatics/btv351
30. Smit, AFA, Hubley, R. RepeatModeler Open-1.0. 2008-2015 <http://www.repeatmasker.org>
31. Bao W., Kenji K. Kojima, Kohany O. Repbase Update a database of repetitive elements in eukaryotic genomes. *Mobile DNA*, 2015, 6, no. 1: 11. DOI: 10.1186/s13100-015-0041-9
32. Smit AFA, Hubley R and Green P. RepeatMasker Open-4.0. <http://www.repeatmasker.org>.
33. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics*, 2010, 26: 841-842. DOI: 10.1093/bioinformatics/btq033
34. Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic acids research*, 1999, 27(2), 573. DOI: 10.1093/nar/27.2.573
35. Morgulis A, Gertz EM, Schäffer AA and Agarwala R. WindowMasker: window-based masker for sequenced genomes. *Bioinformatics*, 2005, 22: 134-141. DOI: 10.1093/bioinformatics/bti774
36. Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, et al. From FastQ data to high-confidence variant calls: the genome analysis toolkit best practices pipeline. *Current Prot. Bioinformatics*, 2013, 11-10. DOI: 10.1002/0471250953.bi1110s43
37. Slater G., and Birney E. Automated generation of heuristics for biological sequence comparison. *BMC Bioinformatics*, 2005, 6: 31. DOI: 10.1186/1471-2105-6-31
38. Stanke M, Keller O, Gunduz I, Hayes A, Waack S and Morgenstern B. AUGUSTUS: ab initio prediction of alternative transcripts. *Nucl. Acids Res*, 2006, 34, Suppl_2 : W435-W439. DOI: 10.1093/nar/gkl200
39. Johnson LS, Eddy SR and Portugaly E. Hidden Markov model speed heuristic and iterative HMM search procedure. *BMC Bioinformatics*, 2010, 11: 431. DOI: 10.1186/1471-2105-11-431
40. Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ. Basic local alignment search tool. *J.Mol. Biol.*, 1990, 215: 403-410. DOI: 10.1006/jmbi.1990.9999
41. Bateman A, Coin L, Durbin R, Finn RD, Hollich V, Griffiths-Jones S, et al. The Pfam protein families database. *Nucl. Acids Res.*, 2004, 32, Suppl_1: D138-D141. DOI: 10.1093/nar/28.1.263
42. UniProt Consortium. UniProt: the universal protein knowledgebase. *Nucl. Acids Res.*, 2017, 45: D158-D169. DOI: 10.1093/nar/gkw1099
43. Lowe TM and Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucl. Acids Res.*, 1997, 25: 955-964. DOI: 10.1093/nar/25.5.955
44. Victorian Bioinformatics Consortium. Barrnap. <http://www.vicbioinformatics.com/software/barrnap.shtml> Accessed on 01 Nov 2017.
45. Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, et al. eggNOG 4.5: a hierarchical orthology framework with improved functional annotations for eukaryotic, prokaryotic and viral sequences. *Nucl. Acids Res.*, 2015, 44: D286-D293. DOI: 10.1093/nar/gkv1248
46. Katoh, K., & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. biol. and evol.*, 2013, 30(4), 772-780. DOI: 10.1093/molbev/mst010
47. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*, 2014, 30: 1312-1313. DOI: 10.1093/bioinformatics/btu033
48. FigTree. <http://tree.bio.ed.ac.uk/software/figtree/> Accessed on 01 Nov 2017
49. Löytynoja A Phylogeny-aware alignment with PRANK, *Methods Mol Biol*, 2014; 1079:155-70. doi: 10.1007/978-1-62703-646-7_10.
50. Talavera, G., and Castresana, J. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Sys. Biol.*, 2007, 56, 564-577. DOI: 10.1080/10635150701472164
51. Tamura K. and Nei M. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol. Biol. and Evol.*, 1993, 10:512-526. DOI: 10.1093/oxfordjournals.molbev.a040023
52. Kumar S., Stecher G., and Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Mol. Biol. and Evol.*, 2016, 33:1870-1874. DOI: 10.1093/molbev/msw054

- 486 53. Corral, José M., et al. A conserved apomixis-specific polymorphism is correlated with exclusive
487 exonuclease expression in premeiotic ovules of apomictic *Boechera* species. *Plant physiology*, 2013, 163.4:
488 1660-1672. DOI: 10.1104/pp.113.222430
489 54. Phytozome v12.1 database. <https://phytozome.jgi.doe.gov/pz/portal.html> Accessed on 30 Jan 2018