

SOME INEQUALITIES OF HERMITE-HADAMARD TYPE FOR HYPERBOLIC p -CONVEX FUNCTIONS

SILVESTRU SEVER DRAGOMIR^{1,2}

ABSTRACT. In this paper we establish several Hermite-Hadamard type integral inequalities for hyperbolic p -convex functions. Applications for special means are provided as well.

1. INTRODUCTION

The following integral inequality

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) dt \leq \frac{f(a)+f(b)}{2},$$

which holds for any convex function $f : [a, b] \rightarrow \mathbb{R}$, is well known in the literature as the Hermite-Hadamard inequality.

There is an extensive amount of literature devoted to this simple and nice result which has many applications in the Theory of Special Means and in Information Theory for divergence measures, for which we would like to refer the reader to the monograph [9], the recent survey paper [8] and the references therein.

Let I be a finite or infinite open interval of real numbers and $p \in \mathbb{R}$, $p \neq 0$.

In the following we present the basic definitions and results concerning the class of hyperbolic p -convex function, see [3]. For other concepts of modified convex functions see for example [11], [12] and [4], [6], [7], [10], [13], [14] and [15].

We consider the hyperbolic functions of a real argument $x \in \mathbb{R}$ defined by

$$\sinh x := \frac{e^x - e^{-x}}{2} = \frac{e^{2x} - 1}{2e^x}, \quad \cosh x := \frac{e^x + e^{-x}}{2} = \frac{e^{2x} + 1}{2e^x},$$

$$\tanh x := \frac{\sinh x}{\cosh x} \quad \text{and} \quad \coth x := \frac{\cosh x}{\sinh x}.$$

We say that a function $f : I \rightarrow \mathbb{R}$ is *hyperbolic p -convex* (or sub H -function, according with [3]) on I , if for any closed subinterval $[a, b]$ of I we have

$$(1.2) \quad f(x) \leq \frac{\sinh[p(b-x)]}{\sinh[p(b-a)]} f(a) + \frac{\sinh[p(x-a)]}{\sinh[p(b-a)]} f(b)$$

for all $x \in [a, b]$.

If the inequality (1.2) holds with " \geq ", then the function will be called *hyperbolic p -concave* on I .

1991 *Mathematics Subject Classification.* 26D15, 26D10, 26D07, 26A33.

Key words and phrases. Convex functions, Hyperbolic p -convex Functions, Hermite-Hadamard inequalities, Integral inequalities.

Geometrically speaking, this means that the graph of f on $[a, b]$ lies nowhere above the p -hyperbolic function determined by the equation

$$H(x) = H(x; a, b, f) := A \cosh(px) + B \sinh(px)$$

where A and B are chosen such that $H(a) = f(a)$ and $H(b) = f(b)$.

If we take $x = (1-t)a + tb \in [a, b]$, $t \in [0, 1]$, then the condition (1.2) becomes

$$(1.3) \quad f((1-t)a + tb) \leq \frac{\sinh[p(1-t)(b-a)]}{\sinh[p(b-a)]} f(a) + \frac{\sinh[pt(b-a)]}{\sinh[p(b-a)]} f(b)$$

for any $t \in [0, 1]$.

We have the following properties of hyperbolic p -convex on I , [3].

- (i) A hyperbolic p -convex function $f : I \rightarrow \mathbb{R}$ has finite right and left derivatives $f'_+(x)$ and $f'_-(x)$ at every point $x \in I$ and $f'_-(x) \leq f'_+(x)$. The function f is differentiable on I with the exception of an at most countable set.
- (ii) A necessary and sufficient condition for the function $f : I \rightarrow \mathbb{R}$ to be hyperbolic p -convex function on I is that it satisfies the gradient inequality

$$(1.4) \quad f(y) \geq f(x) \cosh[p(y-x)] + K_{x,f} \sinh[p(y-x)]$$

for any $x, y \in I$ where $K_{x,f} \in [f'_-(x), f'_+(x)]$. If f is differentiable at the point x then $K_{x,f} = f'(x)$.

- (iii) A necessary and sufficient condition for the function f to be a hyperbolic p -convex in I , is that the function

$$\varphi(x) = f'(x) - p^2 \int_a^x f(t) dt$$

is nondecreasing on I , where $a \in I$.

- (iv) Let $f : I \rightarrow \mathbb{R}$ be a two times continuously differentiable function on I . Then f is hyperbolic p -convex on I if and only if for all $x \in I$ we have

$$(1.5) \quad f''(x) - p^2 f(x) \geq 0.$$

For other properties of hyperbolic p -convex functions, see [3].

Consider the function $f_r : (0, \infty) \rightarrow (0, \infty)$, $f_r(x) = x^r$ with $p \in \mathbb{R} \setminus \{0\}$. If $r \in (-\infty, 0) \cup [1, \infty)$ the function is convex and if $r \in (0, 1)$ it is concave. We have for $r \in (-\infty, 0) \cup [1, \infty)$

$$f_r''(x) - p^2 f_r(x) = r(r-1)x^{r-2} - p^2 x^r = p^2 x^{r-2} \left(\frac{r(r-1)}{p^2} - x^2 \right), \quad x > 0.$$

We observe that $f_r''(x) - p^2 f_r(x) > 0$ for $x \in \left(0, \frac{\sqrt{r(r-1)}}{|p|}\right)$ and $f_r''(x) - p^2 f_r(x) < 0$ for $x \in \left(\frac{\sqrt{r(r-1)}}{|p|}, \infty\right)$, which shows that the power function f_r for $r \in (-\infty, 0) \cup [1, \infty)$ is hyperbolic p -convex on $\left(0, \frac{\sqrt{r(r-1)}}{|p|}\right)$ and hyperbolic p -concave on $\left(\frac{\sqrt{r(r-1)}}{|p|}, \infty\right)$.

If $r \in (0, 1)$, then $f_r''(x) - p^2 f_r(x) < 0$ for any $x > 0$, which shows that f_r is hyperbolic p -concave on $(0, \infty)$ for any $r \in (0, 1)$.

Consider the exponential function $f_\alpha(x) = \exp(\alpha x)$ for $\alpha \neq 0$ and $x \in \mathbb{R}$. Then

$$f_\alpha''(x) - p^2 f_\alpha(x) = \alpha^2 e^{\alpha x} - p^2 e^{\alpha x} = (\alpha^2 - p^2) e^{\alpha x}, \quad x > 0.$$

If $|\alpha| > |p|$, then f_α is hyperbolic p -convex on \mathbb{R} and if $|\alpha| < |p|$ then f_α is hyperbolic p -concave on \mathbb{R} .

In this paper we establish several Hermite-Hadamard type integral inequalities for hyperbolic p -convex functions. Applications for special means are provided as well.

2. SOME HERMITE-HADAMARD TYPE INEQUALITIES

We start with the following lemma of interest in itself:

Lemma 1. *Assume that the function $f : I \rightarrow \mathbb{R}$ is hyperbolic p -convex on I . Then for any $a, b \in I$ with $a < b$ and $x \in [a, b]$ we have*

$$(2.1) \quad f\left(\frac{a+b}{2}\right) \cosh\left[p\left(x - \frac{a+b}{2}\right)\right] \leq \frac{1}{2} [f(x) + f(a+b-x)] \\ \leq \frac{f(a) + f(b)}{2} \frac{\cosh\left[p\left(x - \frac{a+b}{2}\right)\right]}{\cosh\left[\frac{p(b-a)}{2}\right]}.$$

Proof. From (1.2) we have by replacing x with $a+b-x$ that

$$(2.2) \quad f(a+b-x) \leq \frac{\sinh[p(x-a)]}{\sinh[p(b-a)]} f(a) + \frac{\sinh[p(b-x)]}{\sinh[p(b-a)]} f(b)$$

for any $x \in [a, b]$.

If we add (1.2) with (2.2) we get

$$(2.3) \quad f(x) + f(a+b-x) \\ \leq \frac{\sinh[p(b-x)]}{\sinh[p(b-a)]} f(a) + \frac{\sinh[p(x-a)]}{\sinh[p(b-a)]} f(b) \\ + \frac{\sinh[p(x-a)]}{\sinh[p(b-a)]} f(a) + \frac{\sinh[p(b-x)]}{\sinh[p(b-a)]} f(b) \\ = \frac{\sinh[p(b-x)] + \sinh[p(x-a)]}{\sinh[p(b-a)]} f(a) \\ + \frac{\sinh[p(b-x)] + \sinh[p(x-a)]}{\sinh[p(b-a)]} f(b) \\ = \frac{\sinh[p(b-x)] + \sinh[p(x-a)]}{\sinh[p(b-a)]} [f(a) + f(b)]$$

for any $x \in [a, b]$.

Observe that

$$(2.4) \quad \frac{\sinh[p(b-x)] + \sinh[p(x-a)]}{\sinh[p(b-a)]} \\ = \frac{2 \sinh\left[\frac{p(b-a)}{2}\right] \cosh\left[p\left(x - \frac{a+b}{2}\right)\right]}{2 \sinh\left[\frac{p(b-a)}{2}\right] \cosh\left[\frac{p(b-a)}{2}\right]} = \frac{\cosh\left[p\left(x - \frac{a+b}{2}\right)\right]}{\cosh\left[\frac{p(b-a)}{2}\right]}$$

for any $x \in [a, b]$.

Using the equality (2.4) and dividing by 2 in (2.3) we get the second inequality in (2.1).

From (1.3) for $t = \frac{1}{2}$ and $a = u, b = v$ we get

$$\begin{aligned} f\left(\frac{u+v}{2}\right) &\leq \frac{\sinh\left[p\left(\frac{v-u}{2}\right)\right]}{\sinh[p(v-u)]}f(u) + \frac{\sinh\left[p\left(\frac{v-u}{2}\right)\right]}{\sinh[p(v-u)]}f(v) \\ &= \frac{\sinh\left[p\left(\frac{v-u}{2}\right)\right]}{\sinh[p(v-u)]}[f(u) + f(v)] \\ &= \frac{\sinh\left[p\left(\frac{v-u}{2}\right)\right]}{2\sinh\left[p\left(\frac{v-u}{2}\right)\right]\cosh\left[p\left(\frac{v-u}{2}\right)\right]}[f(u) + f(v)] \\ &= \frac{1}{\cosh\left[p\left(\frac{v-u}{2}\right)\right]}\frac{f(u) + f(v)}{2}, \end{aligned}$$

which implies that

$$(2.5) \quad f\left(\frac{u+v}{2}\right)\cosh\left[p\left(\frac{v-u}{2}\right)\right] \leq \frac{f(u) + f(v)}{2}$$

for any $u, v \in I$.

Now, if in (2.5) we take $v = x$ and $u = a + b - x$, then we get the first inequality in (2.1). \square

Remark 1. By taking $x = (1-t)a + tb$ in (2.1) we get the equivalent double inequality

$$(2.6) \quad \begin{aligned} f\left(\frac{a+b}{2}\right)\cosh\left[p\left(t - \frac{1}{2}\right)(b-a)\right] \\ \leq \frac{1}{2}[f((1-t)a + tb) + f(ta + (1-t)b)] \\ \leq \frac{f(a) + f(b)}{2} \frac{\cosh\left[p\left(t - \frac{1}{2}\right)(b-a)\right]}{\cosh\left[\frac{p(b-a)}{2}\right]} \end{aligned}$$

for any $a, b \in I$ and $t \in [0, 1]$.

We have the following Hermite-Hadamard type inequality that was obtained in a different and slightly more difficult manner in [2]

Theorem 1. Assume that the function $f : I \rightarrow \mathbb{R}$ is hyperbolic p -convex on I . Then for any $a, b \in I$ with $0 < b - a < \frac{\pi}{p}$ we have

$$(2.7) \quad \frac{2}{p}f\left(\frac{a+b}{2}\right)\sinh\left[\frac{p(b-a)}{2}\right] \leq \int_a^b f(x)dx \leq \frac{f(a) + f(b)}{p}\tanh\left[\frac{p(b-a)}{2}\right].$$

Proof. Integrating the inequality (2.1) over x on $[a, b]$ we get

$$\begin{aligned}
 (2.8) \quad f\left(\frac{a+b}{2}\right) \int_a^b \cosh\left[p\left(x - \frac{a+b}{2}\right)\right] dx \\
 \leq \frac{1}{2} \left[\int_a^b f(x) dx + \int_a^b f(a+b-x) dx \right] \\
 \leq \frac{f(a) + f(b)}{2} \frac{\int_a^b \cosh\left[p\left(x - \frac{a+b}{2}\right)\right] dx}{\cosh\left[\frac{p(b-a)}{2}\right]}.
 \end{aligned}$$

Since

$$\begin{aligned}
 \int_a^b \cosh\left[p\left(x - \frac{a+b}{2}\right)\right] dx &= \frac{1}{p} \sinh\left[p\left(x - \frac{a+b}{2}\right)\right] \Big|_a^b \\
 &= \frac{1}{p} \sinh\left[p\left(b - \frac{a+b}{2}\right)\right] - \frac{1}{p} \sinh\left[p\left(a - \frac{a+b}{2}\right)\right] \\
 &= \frac{2}{p} \sinh\left[\frac{p(b-a)}{2}\right]
 \end{aligned}$$

and by the change of variable $y = a + b - x$

$$\int_a^b f(a+b-x) dx = \int_a^b f(y) dy,$$

hence by (2.8) we get (2.7). □

Remark 2. The inequality (2.7) for $p = 1$ was obtained in 2004 by M. Bessenyei in his Ph.D. Thesis [5, Corollary 2.12] in the context of Chebyshev system (cosh, sinh).

We use the notation $\operatorname{sech} t = \frac{1}{\cosh t}$ for to state the following weighted integral inequality of Hermite-Hadamard type:

Theorem 2. Assume that the function $f : I \rightarrow \mathbb{R}$ is hyperbolic p -convex on I . Then for any $a, b \in I$ with $a < b$

$$(2.9) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) \operatorname{sech}\left[p\left(x - \frac{a+b}{2}\right)\right] dx \leq \frac{f(a) + f(b)}{2}.$$

Proof. From (2.1) we have

$$\begin{aligned}
 (2.10) \quad f\left(\frac{a+b}{2}\right) &\leq \frac{1}{2} [f(x) + f(a+b-x)] \operatorname{sech}\left[p\left(x - \frac{a+b}{2}\right)\right] \\
 &\leq \frac{f(a) + f(b)}{2} \operatorname{sech}\left[\frac{p(b-a)}{2}\right]
 \end{aligned}$$

for any $a, b \in I$ with $a < b$ and $x \in [a, b]$.

If we take the integral mean in (2.10) we get

$$\begin{aligned}
 (2.11) \quad f\left(\frac{a+b}{2}\right) &\leq \frac{1}{2(b-a)} \int_a^b [f(x) + f(a+b-x)] \operatorname{sech}\left[p\left(x - \frac{a+b}{2}\right)\right] dx \\
 &\leq \frac{f(a) + f(b)}{2} \operatorname{sech}\left[\frac{p(b-a)}{2}\right].
 \end{aligned}$$

Using the change of variable $y = a + b - x$ we get

$$\begin{aligned} \int_a^b f(a+b-x) \operatorname{sech} \left[p \left(x - \frac{a+b}{2} \right) \right] dx &= \int_a^b f(y) \operatorname{sech} \left[p \left(\frac{a+b}{2} - y \right) \right] dy \\ &= \int_a^b f(y) \operatorname{sech} \left[p \left(y - \frac{a+b}{2} \right) \right] dy \end{aligned}$$

and by (2.11) we obtain the desired result (2.9). \square

The following weighted inequality also holds:

Theorem 3. Assume that the function $f : I \rightarrow \mathbb{R}$ is hyperbolic p -convex on I . Then for any $a, b \in I$ with $0 < b - a < \frac{\pi}{p}$ we have

$$\begin{aligned} (2.12) \quad \frac{1}{2} \left[b - a + \frac{1}{p} \sinh [p(b-a)] \right] f \left(\frac{a+b}{2} \right) \\ \leq \int_a^b f(x) \cosh \left[p \left(x - \frac{a+b}{2} \right) \right] dx \\ \leq \frac{b - a + \frac{1}{p} \sinh [p(b-a)]}{2 \cosh \left[\frac{p(b-a)}{2} \right]} \frac{f(a) + f(b)}{2}. \end{aligned}$$

Proof. If we multiply the inequality (2.1) by $\cosh \left[p \left(x - \frac{a+b}{2} \right) \right] \geq 0$ we get

$$\begin{aligned} f \left(\frac{a+b}{2} \right) \cosh^2 \left[p \left(x - \frac{a+b}{2} \right) \right] \\ \leq \frac{1}{2} [f(x) + f(a+b-x)] \cosh \left[p \left(x - \frac{a+b}{2} \right) \right] \\ \leq \frac{f(a) + f(b)}{2} \frac{\cosh^2 \left[p \left(x - \frac{a+b}{2} \right) \right]}{\cosh \left[\frac{p(b-a)}{2} \right]} \end{aligned}$$

for any $x \in [a, b]$.

If we integrate this inequality over $x \in [a, b]$ we get

$$\begin{aligned} (2.13) \quad f \left(\frac{a+b}{2} \right) \int_a^b \cosh^2 \left[p \left(x - \frac{a+b}{2} \right) \right] dx \\ \leq \frac{1}{2} \int_a^b [f(x) + f(a+b-x)] \cosh \left[p \left(x - \frac{a+b}{2} \right) \right] dx \\ \leq \frac{f(a) + f(b)}{2} \frac{\int_a^b \cosh^2 \left[p \left(x - \frac{a+b}{2} \right) \right] dx}{\cosh \left[\frac{p(b-a)}{2} \right]}. \end{aligned}$$

Since

$$\int_a^b \cosh^2 \left[p \left(x - \frac{a+b}{2} \right) \right] dx = \frac{1}{2} \left[(b-a) + \int_a^b \cosh \left[2p \left(x - \frac{a+b}{2} \right) \right] dx \right]$$

and

$$\begin{aligned} \int_a^b \cosh \left[2p \left(x - \frac{a+b}{2} \right) \right] dx &= \frac{1}{2p} \sinh \left[2p \left(x - \frac{a+b}{2} \right) \right] \Big|_a^b \\ &= \frac{1}{2p} \sinh \left[2p \left(b - \frac{a+b}{2} \right) \right] - \frac{1}{2p} \sinh \left[2p \left(a - \frac{a+b}{2} \right) \right] \\ &= \frac{1}{p} \sinh [p(b-a)], \end{aligned}$$

hence

$$\int_a^b \cosh^2 \left[p \left(x - \frac{a+b}{2} \right) \right] dx = \frac{1}{2} \left[(b-a) + \frac{1}{p} \sinh [p(b-a)] \right].$$

Also, by the change of variable $y = a + b - x$ we have

$$\int_a^b f(a+b-x) \cosh \left[p \left(x - \frac{a+b}{2} \right) \right] dx = \int_a^b f(y) \cosh \left[p \left(y - \frac{a+b}{2} \right) \right] dx$$

and by (2.13) we get (2.12). \square

3. RELATED RESULTS

Theorem 4. Assume that the function $f : I \rightarrow \mathbb{R}$ is hyperbolic p -convex on I . Then for any $a, b \in I$ with $0 < b - a < \frac{\pi}{p}$ we have

$$(3.1) \quad \int_a^b f(t) dt \geq \frac{2}{p} \sinh \left[p \left(\frac{b-a}{2} \right) \right] \times \left\{ f(x) \cosh \left[p \left(x - \frac{a+b}{2} \right) \right] + K_{x,f} \sinh \left[p \left(\frac{a+b}{2} - x \right) \right] \right\}$$

for any $x \in [a, b]$, where $K_{x,f} \in [f'_-(x), f'_+(x)]$.

If f is differentiable in x , then

$$(3.2) \quad \int_a^b f(t) dt \geq \frac{2}{p} \sinh \left[p \left(\frac{b-a}{2} \right) \right] \times \left\{ f(x) \cosh \left[p \left(x - \frac{a+b}{2} \right) \right] + f'(x) \sinh \left[p \left(\frac{a+b}{2} - x \right) \right] \right\}.$$

Proof. If we take the integral over $y \in [a, b]$ in the gradient inequality (1.4) we get

$$(3.3) \quad \int_a^b f(y) dy \geq f(x) \int_a^b \cosh [p(y-x)] dy + K_{x,f} \int_a^b \sinh [p(y-x)] dy$$

where $K_{x,f} \in [f'_-(x), f'_+(x)]$.

Observe that

$$\begin{aligned} \int_a^b \cosh [p(y-x)] dy &= \frac{1}{p} \sinh [p(y-x)] \Big|_a^b = \frac{1}{p} \sinh [p(b-x)] + \frac{1}{p} \sinh [p(x-a)] \\ &= \frac{2}{p} \sinh \left[p \left(\frac{b-a}{2} \right) \right] \cosh \left[p \left(x - \frac{a+b}{2} \right) \right] \end{aligned}$$

and

$$\begin{aligned} \int_a^b \sinh [p(y-x)] dy &= \frac{1}{p} \cosh [p(y-x)] \Big|_a^b = \frac{1}{p} \cosh [p(b-x)] - \frac{1}{p} \cosh [p(a-x)] \\ &= \frac{2}{p} \sinh \left[p \left(\frac{b-a}{2} \right) \right] \sinh \left[p \left(\frac{a+b}{2} - x \right) \right] \end{aligned}$$

or any $x \in [a, b]$.

By utilising the inequality (3.3) we then get (3.1). \square

Remark 3. We observe that, if we take $x = \frac{a+b}{2}$ in (3.1), then we recapture the first inequality in (2.7).

Theorem 5. Assume that the function $f : I \rightarrow \mathbb{R}$ is hyperbolic p -convex on I . Then for any $a, b \in I$ with $0 < b-a < \frac{\pi}{p}$ we have

$$(3.4) \quad \int_a^b f(t) \operatorname{sech} [p(t-x)] dt \geq f(x)(b-a) + \frac{1}{p} K_{x,f} \ln \left(\frac{\cosh [p(b-x)]}{\cosh [p(x-a)]} \right)$$

for any $x \in [a, b]$, where $K_{x,f} \in [f'_-(x), f'_+(x)]$.

If f is differentiable in x , then

$$(3.5) \quad \int_a^b f(t) \operatorname{sech} [p(t-x)] dt \geq f(x)(b-a) + \frac{1}{p} f'(x) \ln \left(\frac{\cosh [p(b-x)]}{\cosh [p(x-a)]} \right)$$

for any $x \in [a, b]$.

Proof. We have by the gradient inequality (1.4) that

$$(3.6) \quad f(y) \operatorname{sech} [p(y-x)] \geq f(x) + K_{x,f} \tanh [p(y-x)]$$

for any $x, y \in [a, b]$.

If we integrate (3.6) over $y \in [a, b]$, then we get

$$(3.7) \quad \int_a^b f(y) \operatorname{sech} [p(y-x)] dy \geq f(x)(b-a) + K_{x,f} \int_a^b \tanh [p(y-x)] dy$$

for any $x \in [a, b]$.

Since

$$\begin{aligned} \int_a^b \tanh [p(y-x)] dy &= \int_a^b \frac{\sinh [p(y-x)]}{\cosh [p(y-x)]} dy \\ &= \frac{1}{p} \int_a^b \frac{1}{\cosh [p(y-x)]} d(\cosh [p(y-x)]) \\ &= \frac{1}{p} \ln (\cosh [p(y-x)]) \Big|_a^b \\ &= \frac{1}{p} \ln (\cosh [p(b-x)]) - \frac{1}{p} \ln (\cosh [p(a-x)]) \\ &= \frac{1}{p} \ln \left(\frac{\cosh [p(b-x)]}{\cosh [p(x-a)]} \right) \end{aligned}$$

for any $x \in [a, b]$, hence by (3.7) we get (3.4). \square

Remark 4. We observe that, if we take $x = \frac{a+b}{2}$ in (3.4), then we recapture the first inequality in (2.9).

From a different perspective, we have:

Theorem 6. Assume that the function $f : I \rightarrow \mathbb{R}$ is hyperbolic p -convex on I . Then for any $a, b \in I$ with $0 < b - a < \frac{\pi}{p}$ we have

$$(3.8) \quad \frac{1}{p+1} \{f(y)(b-a) + f(b) \sinh [p(b-y)] + f(a) \sinh [p(y-a)]\} \\ \geq \int_a^b f(x) \cosh [p(x-y)] dx$$

for any $y \in [a, b]$.

In particular, for $y = \frac{a+b}{2}$, we get

$$(3.9) \quad \frac{1}{p+1} \left\{ f \left(\frac{a+b}{2} \right) (b-a) + [f(b) + f(a)] \sinh \left[p \left(\frac{b-a}{2} \right) \right] \right\} \\ \geq \int_a^b f(x) \cosh \left[p \left(x - \frac{a+b}{2} \right) \right] dx.$$

Proof. The function $f : I \rightarrow \mathbb{R}$ is differentiable almost everywhere and we have

$$(3.10) \quad f(y) \geq f(x) \cosh [p(y-x)] + f'(x) \sinh [p(y-x)]$$

for almost every $x \in [a, b]$ where $y \in [a, b]$.

If we integrate the inequality over $x \in [a, b]$ we get

$$(3.11) \quad f(y)(b-a) \geq \int_a^b f(x) \cosh [p(y-x)] dx + \int_a^b f'(x) \sinh [p(y-x)] dx$$

for any $y \in [a, b]$.

Using the integration by parts, we have

$$\int_a^b f'(x) \sinh [p(y-x)] dx = f(x) \sinh [p(y-x)] \Big|_a^b + p \int_a^b f(x) \cosh [p(y-x)] dx \\ = -f(b) \sinh [p(b-y)] - f(a) \sinh [p(y-a)] \\ + p \int_a^b f(x) \cosh [p(y-x)] dx$$

and by (3.11) we get

$$f(y)(b-a) \geq -f(b) \sinh [p(b-y)] - f(a) \sinh [p(y-a)] \\ + (p+1) \int_a^b f(x) \cosh [p(y-x)] dx,$$

for any $y \in [a, b]$, which is equivalent to the desired result (3.8). \square

4. SOME INEQUALITIES FOR SPECIAL MEANS

Recall the following special means:

(1) The arithmetic mean

$$A = A(a, b) := \frac{a+b}{2}, \quad a, b \geq 0;$$

(2) The geometric mean:

$$G = G(a, b) := \sqrt{ab}, \quad a, b \geq 0;$$

(3) The harmonic mean:

$$H = H(a, b) := \frac{2}{\frac{1}{a} + \frac{1}{b}}, \quad a, b \geq 0;$$

(4) The logarithmic mean:

$$L = L(a, b) := \begin{cases} a & \text{if } a = b \\ \frac{b-a}{\ln b - \ln a} & \text{if } a \neq b \end{cases} \quad a, b > 0;$$

(5) The identric mean:

$$I := I(a, b) = \begin{cases} a & \text{if } a = b \\ \frac{1}{e} \left(\frac{b^b}{a^a} \right)^{\frac{1}{b-a}} & \text{if } a \neq b \end{cases} \quad a, b > 0;$$

(6) The p -logarithmic mean:

$$L_p = L_p(a, b) := \begin{cases} \left[\frac{b^{p+1} - a^{p+1}}{(p+1)(b-a)} \right]^{\frac{1}{p}} & \text{if } a \neq b; \\ a & \text{if } a = b \end{cases}$$

where $p \in \mathbb{R} \setminus \{-1, 0\}$ and $a, b > 0$.

It is well known that L_p is monotonic nondecreasing over $p \in \mathbb{R}$ with $L_{-1} := L$ and $L_0 := I$.

For $r \in \mathbb{R} \setminus \{-1, 0\}$ and $0 < a < b$ we have

$$\frac{1}{b-a} \int_a^b x^r dx = L_r^r(a, b).$$

We know, from Introduction, that the power function $f_r(x) = x^r$, $x > 0$, for $r \in (-\infty, 0) \cup [1, \infty)$ is hyperbolic p -convex on $\left(0, \frac{\sqrt{r(r-1)}}{|p|}\right)$ and hyperbolic p -concave on $\left(\frac{\sqrt{r(r-1)}}{|p|}, \infty\right)$. Also f_r is hyperbolic p -concave on $(0, \infty)$ for any $r \in (0, 1)$.

By using the inequality (2.7) we have for $p \neq 0$, $r \in (-\infty, 0) \cup [1, \infty)$

$$(4.1) \quad \frac{2}{p} A^r(a, b) \sinh \left[\frac{p(b-a)}{2} \right] \leq (b-a) L_r^r(a, b) \leq \frac{2}{p} A(a^r, b^r) \tanh \left[\frac{p(b-a)}{2} \right]$$

provided $[a, b] \subset \left(0, \frac{\sqrt{r(r-1)}}{|p|}\right)$.

If either $[a, b] \subset \left(\frac{\sqrt{r(r-1)}}{|p|}, \infty\right)$ or $r \in (0, 1)$ and $[a, b] \subset (0, \infty)$, then the inequality reverses in (4.1).

Now, by utilising the inequality (2.9) for the function f_r we get

$$(4.2) \quad A^r(a, b) \leq \frac{1}{b-a} \int_a^b x^r \operatorname{sech} \left[p \left(x - \frac{a+b}{2} \right) \right] dx \leq A(a^r, b^r),$$

where $p \neq 0$, $r \in (-\infty, 0) \cup [1, \infty)$ and $[a, b] \subset \left(0, \frac{\sqrt{r(r-1)}}{|p|}\right)$. If either $r \in (-\infty, 0) \cup [1, \infty)$ and $[a, b] \subset \left(\frac{\sqrt{r(r-1)}}{|p|}, \infty\right)$ or $r \in (0, 1)$ and $[a, b] \subset (0, \infty)$, then the sign of inequality reverses in (4.2).

On making use of the other general inequalities obtained above, further similar Hermite-Hadamard type inequalities may be stated, however the details are left to the interested reader.

REFERENCES

- [1] M. S. S. Ali, On certain properties of trigonometrically ρ -convex functions, *Advances in Pure Mathematics*, 2012, 2, 337-340 <http://dx.doi.org/10.4236/apm.2012.25047>.
- [2] M. S. S. Ali, On Hadamard's inequality for trigonometrically ρ -convex functions," accepted to appear in *Theoretical Mathematics & Applications.*, March, 2013.
- [3] M. S. S. Ali, On certain properties for two classes of generalized convex functions, *Abstract and Applied Analysis* Volume 2016, Article ID 4652038, 7 pages <http://dx.doi.org/10.1155/2016/4652038>
- [4] E. F. Beckenbach, Convex functions, *Bulletin of the American Mathematical Society*, Vol. 54, No. 5, 1948, pp. 439-460. doi:10.1090/S0002-9904-1948-08994-7
- [5] M. Bessenyei, Hermite-Hadamard-type inequalities for generalized convex functions, *PhD Thesis*, Univ. Debrecen, Hungary, 2004. RGMIA Monographs, [<http://rgmia.org/papers/monographs/dissertatio.pdf>]
- [6] F. F. Bonsall, The characterization of generalized convex functions, *The Quarterly Journal of Mathematics Oxford Series*, Vol. 1, 1950, pp. 100-111. doi:10.1093/qmath/1.1.100
- [7] A. M. Bruckner and E. Ostrow, Some functions classes related to the class of convex functions," *Pacific Journal of Mathematics*, Vol. 12, 1962, pp. 1203-1215.
- [8] S. S. Dragomir, Ostrowski type inequalities for Lebesgue integral: a survey of recent results, *Australian J. Math. Anal. Appl.*, Volume 14, Issue 1, Article 1, pp. 1-287, 2017. [Online <http://ajmaa.org/cgi-bin/paper.pl?string=v14n1/V14I1P1.tex>].
- [9] S. S. Dragomir and C. E. M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, 2000. [Online http://rgmia.org/monographs/hermite_hadamard.html].
- [10] J. W. Green, Support, Convergence, and Differentiability Properties of Generalized Convex Functions, *Proceedings of the American Mathematical Society*, Vol. 4, No. 3, 1953, pp. 391-396. doi:10.1090/S0002-9939-1953-0056039-2
- [11] B. Ya. Levin, *Lectures on Entire Functions*, American Mathematical Society, 1996.
- [12] L. S. Maergoiz, *Asymptotic Characteristics of Entire Functions and Their Applications in Mathematics and Biophysics*, Kluwer Academic Publishers, New York, 2003.
- [13] M. J. Miles, An extremum property of convex functions, *American Mathematical Monthly*, Vol. 76, 1969, pp. 921-922. doi:10.2307/2317948
- [14] M. M. Peixoto, On the existence of derivatives of generalized convex functions, *Summa Brazilian Mathematics*, Vol. 2, No. 3, 1948, pp. 35-42.
- [15] M. M. Peixoto, Generalized convex functions and second order differential inequalities, *Bulletin of the American Mathematical Society*, Vol. 55, No. 6, 1949, pp. 563- 572. doi:10.1090/S0002-9904-1949-09246-7
- [16] A. W. Roberts and D. E. Varberg, *Convex Functions*, Academic Press, New York, 1973.

¹MATHEMATICS, COLLEGE OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO BOX 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

E-mail address: sever.dragomir@vu.edu.au

URL: <http://rgmia.org/dragomir>

²DST-NRF CENTRE OF EXCELLENCE, IN THE MATHEMATICAL AND STATISTICAL SCIENCES, SCHOOL OF COMPUTER SCIENCE & APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, JOHANNESBURG 2050, SOUTH AFRICA