

Hermite based poly-Bernoulli polynomials with a q -parameter

Ugur Duran¹, Mehmet Acikgoz² and Serkan Araci^{3,*}

¹Department of Mathematics, Faculty of Arts and Science,
University of Gaziantep, TR-27310 Gaziantep, Turkey
E-Mail: mtdrnugur@gmail.com

²Department of Mathematics, Faculty of Arts and Science,
University of Gaziantep, TR-27310 Gaziantep, Turkey
E-Mail: acikgoz@gantep.edu.tr

³Department of Economics, Faculty of Economics,
Administrative and Social Science,
Hasan Kalyoncu University, TR-27410 Gaziantep, Turkey
E-Mail: mtsrkn@hotmail.com

***Corresponding Author**

Abstract: We introduce the Hermite based poly-Bernoulli polynomials with a q parameter and give some of their basic properties including not only addition property, but also derivative properties and integral representations. We also define the Hermite based λ -Stirling polynomials of the second kind, and then provide some relations. Moreover, we derive several correlations and identities including the Hermite-Kampé de Fériet (or Gould-Hopper) family of polynomials, the Hermite based poly-Bernoulli polynomials with a q parameter and the Hermite based λ -Stirling polynomials of the second kind.

2010 Mathematics Subject Classification: Primary 33C45; Secondary 11B68, 11B73.

Keywords and Phrases: Hermite-Kampé de Fériet polynomials; Hermite Bernoulli polynomials; poly Bernoulli polynomials; Stirling numbers of second kind; Polylogarithm functions.

1. INTRODUCTION

Special polynomials and numbers possess a lot of importances in many fields of mathematics, physics, engineering and other related disciplines including the topics such as differential equations, mathematical analysis, functional analysis, mathematical physics, quantum mechanics and so on. One of the most considerable polynomials in the theory of special polynomials is the Hermite-Kampé de Fériet (or Gould-Hopper) polynomials (see [1]) and one other is Bernoulli polynomials (see [10], [16]). Nowadays, these type polynomials and their several generalizations have been studied and used by many mathematicians and physicsics, see [1-17] and references therein. Araci et al. [2] introduced a new concept of the Apostol Hermite-Genocchi polynomials by using the modified Milne-Thomson's polynomials and also derived several implicit summation formulae and general symmetric identities arising from different analytical means and generating functions method. Bretti et al. [4] defined multidimensional extensions of the Bernoulli and Appell polynomials by using the Hermite-Kampé de Fériet polynomials and gave the differential equations, satisfying by the corresponding $2D$ polynomials, derived from exploiting the factorization method. Bayad et al. [3] considered poly-Bernoulli polynomials and numbers and proved a collection of extremely important and fundamental identities satisfied by the poly-Bernoulli polynomials and numbers. Cenkci et al. [5] considered poly-Bernoulli numbers and polynomials with a q parameter and developed some aritmetical and number theoretical properties. Dattoli et al. [6] applied the method of generating function to introduce new forms of Bernoulli numbers and polynomials, which were exploited to derive further classes of partial sums involving generalized many index many variable polynomials. Khan et al. [7] introduce the Hermite poly-Bernoulli polynomials and numbers of the second kind and examined some of their applications in combinatorics, number theory and other fields of mathematics. Kurt et al. [8] studied on the Hermite-Kampé de Fériet based second kind Genocchi polynomials and presented some relationships of them. Ozarslan [11] introduced an unified family of Hermite-based Apostol-Bernoulli, Euler and Genocchi polynomials and then, acquired some symmetry identities between these polynomials and the generalized sum of integer powers. Ozarslan also gave explicit closed-form formulae for this unified family and proved a finite series relation between this unification and $3d$ -Hermite polynomials. Pathan [12] defined a new class of generalized Hermite-Bernoulli

polynomials and derived several implicit summation formulae and symmetric identities by using different analytical means applying generating functions. Pathan et al. [13] introduced a new class of generalized polynomials associated with the modified Milne-Thomson's polynomials $\Phi_n^{(\alpha)}(x, v)$ of degree n and order α and provided some of their properties.

The usual notations \mathbb{C} , \mathbb{R} , \mathbb{Z} , \mathbb{N} and \mathbb{N}_0 are referred to the set of all complex numbers, the set of all real numbers, the set of all integers, the set of all natural numbers and the set of all nonnegative integers, respectively, in the content of this paper.

An outline of this paper is as follows. Section 2 contains the definitions of the Hermite based poly-Bernoulli polynomials ${}_H\mathcal{B}_{n,q}^{(k,j)}(x, y)$ with a q parameter and the Hermite based λ -Stirling polynomials $S_2^{(\lambda,j)}(n, m; x, y)$ of the second kind, and then provides some properties and relationships for ${}_H\mathcal{B}_{n,q}^{(k,j)}(x, y)$ and $S_2^{(\lambda,j)}(n, m; x, y)$. Section 3 examines several correlations including the Hermite-Kampé de Fériet polynomials, the Hermite based poly-Bernoulli polynomials with a q parameter and the Hermite based λ -Stirling polynomials of the second kind.

2. PRELIMINARY RESULTS

The exponential generating function for the Hermite-Kampé de Fériet (or Gould-Hopper) family of polynomials is

$$\sum_{n=0}^{\infty} H_n^{(j)}(x, y) \frac{t^n}{n!} = e^{xt+yt^j} \quad (\text{see [4]}), \quad (2.1)$$

where $j \in \mathbb{N}$ with $j \geq 2$. In the case $j = 1$, the corresponding $2D$ polynomials are simply expressed by the Newton binomial formula. Upon setting $j = 2$ in (2.1) gives the two variable Hermite polynomials $H_n^{(2)}(x, y)$ and the mentioned polynomials have been used to define $2D$ extensions of some special polynomials, such as Bernoulli and Euler polynomials (see [6]).

Recalling that the polynomials $H_n^{(j)}(x, y)$ are the native solution of the generalized heat equation:

$$\frac{\partial}{\partial t} F(x, y) = \frac{\partial^j}{\partial x^j} F(x, y) \quad \text{with } F(x, 0) = x^n$$

and satisfies the following formula

$$H_n^{(j)}(x, y) = n! \sum_{r=0}^{\lfloor \frac{n}{j} \rfloor} \frac{x^{n-jr} y^r}{r! (n-jr)!},$$

where $\lfloor \cdot \rfloor$ is Gauss' notation, and represents the maximum integer which does not exceed a number in the square brackets.

For $k \in \mathbb{Z}$ with $k > 1$, the k -th polylogarithm function is defined by

$$Li_k(t) = \sum_{m=1}^{\infty} \frac{t^m}{m^k} \quad (t \in \mathbb{C} \text{ with } |t| < 1). \quad (2.2)$$

We always assume $|t| < 1$ along this paper. When $k = 1$, $Li_1(t) = -\log(1-t)$. In the case $k \leq 0$, $Li_k(t)$ are the rational functions:

$$Li_0(t) = \frac{t}{1-t}, \quad Li_{-1}(t) = \frac{t}{(1-t)^2}, \quad Li_{-2}(t) = \frac{t^2+t}{(1-t)^3}, \quad Li_{-3}(t) = \frac{t^3+4t^2+t}{(1-t)^4}, \dots$$

2.1. The Hermite based poly-Bernoulli polynomials with a q parameter. Let $n, k \in \mathbb{Z}$ in conjunction with $n \geq 0$ and $k > 0$ and let $q \in \mathbb{R} - \{0\}$. We introduce the Hermite based poly-Bernoulli polynomials with a q parameter via the following exponential generating function to be

$$\sum_{n=0}^{\infty} \mathcal{B}_{n,q}^{(k,j)}(x, y) \frac{t^n}{n!} = \frac{q Li_k\left(\frac{1-e^{-qt}}{q}\right)}{1-e^{-qt}} e^{xt+yt^j}. \quad (2.3)$$

Upon setting $x = y = 0$, we then get ${}_H\mathcal{B}_{n,q}^{(k,j)}(0,0) =: {}_H\mathcal{B}_{n,q}^{(k,j)}$ which is called the poly-Bernoulli numbers with a q parameter (see [5]).

Some special cases of ${}_H\mathcal{B}_{n,q}^{(k,j)}(x,y)$ are listed by the following consecutive remarks.

Remark 1. Letting $y = 0$, we have ${}_H\mathcal{B}_{n,q}^{(k,j)}(x,0) := {}_H\mathcal{B}_{n,q}^{(k)}(x)$ called poly-Bernoulli polynomials with a q parameter (see [5]).

Remark 2. In the case $q = 1$, ${}_H\mathcal{B}_{n,q}^{(k,j)}(x,y)$ reduces to the the Hermite based poly-Bernoulli polynomials ${}_H\mathcal{B}_n^{(k,j)}(x,y)$ (see [11]).

Remark 3. When $q = 1$ and $y = 0$, we obtain the poly-Bernoulli polynomials $B_n(x)$ (see [10] [15], [16]).

Remark 4. When $q = k = 1$ and $y = 0$, we obtain the usual Bernoulli polynomials $B_n(x)$ (see [10] [15], [16]).

The addition formula for ${}_H\mathcal{B}_{n,q}^{(k,j)}(x,y)$ is provided in the following proposition.

Proposition 1. (Addition Property) We have

$${}_H\mathcal{B}_{n,q}^{(k,j)}(x_1 + x_2, y_1 + y_2) = \sum_{m=0}^n \binom{n}{m}_H \mathcal{B}_{n-m,q}^{(k,j)}(x_1, y_1) H_m^{(j)}(x_2, y_2). \quad (2.4)$$

Proof. In view of (2.3), with the series manipulation procedure, we see

$$\begin{aligned} \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x_1 + x_2, y_1 + y_2) \frac{t^n}{n!} &= \frac{qLi_k\left(\frac{1-e^{-qt}}{q}\right)}{1-e^{-qt}} e^{(x_1+x_2)t+(y_1+y_2)t^j} \\ &= \frac{qLi_k\left(\frac{1-e^{-qt}}{q}\right)}{1-e^{-qt}} e^{x_1t+y_1t^j} e^{x_2t+y_2t^j} \\ &= \left(\sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \frac{t^n}{n!}\right) \left(\sum_{n=0}^{\infty} H_n^{(j)}(x,y) \frac{t^n}{n!}\right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{m=0}^n \binom{n}{m}_H \mathcal{B}_{n-m,q}^{(k,j)}(x_1, y_1) H_m^{(j)}(x_2, y_2)\right) \frac{t^n}{n!}, \end{aligned}$$

which implies the asserted result (2.4). \square

An immediate output of Proposition 1 is stated in the following corollary.

Corollary 1. The following holds true:

$${}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) = \sum_{m=0}^n \binom{n}{m}_H \mathcal{B}_{n-m,q}^{(k,j)} H_m^{(j)}(x,y). \quad (2.5)$$

The derivative properties of ${}_H\mathcal{B}_{n,q}^{(k,j)}(x,y)$ are stated in the following proposition.

Proposition 2. (Derivative Properties) Each of the following formulas holds true:

$$\frac{\partial {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y)}{\partial x} = n {}_H\mathcal{B}_{n-1,q}^{(k,j)}(x,y) \quad \text{and} \quad \frac{\partial {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y)}{\partial y} = (n)_j {}_H\mathcal{B}_{n-j,q}^{(k,j)}(x,y),$$

where $(n)_j = n(n-1)(n-2)\cdots(n-j+1)$ called the falling factorial function (see [16]).

Proof. The proof follows from (2.3). So we omit them. \square

The integral representations of ${}_H\mathcal{B}_{n,q}^{(k,j)}(x,y)$ are stated in the following proposition.

4

Proposition 3. (Integral Representations) The following equalities hold true:

$$\int_v^\mu {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) dx = \frac{{}_H\mathcal{B}_{n+1,q}^{(k,j)}(\mu,y) - {}_H\mathcal{B}_{n+1,q}^{(k,j)}(v,y)}{n+1}$$

and

$$\int_\gamma^\zeta {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) dy = \frac{{}_H\mathcal{B}_{n+j,q}^{(k,j)}(x,\zeta) - {}_H\mathcal{B}_{n+j,q}^{(k,j)}(x,\gamma)}{(n+1)^{(j)},}$$

where $(n)^{(j)} = n(n+1)(n+2)\cdots(n+j-1)$ called the rising factorial function (see [16]).

Proof. Using the derivative properties of ${}_H\mathcal{B}_{n,q}^{(k,j)}(x,y)$ given in Proposition 2, we easily get the asserted results. So, we omit them. \square

We have the following proposition.

Proposition 4. The following formula is valid:

$${}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) = \sum_{m=0}^{\lfloor \frac{n}{j} \rfloor} {}_H\mathcal{B}_{n-jm,q}^{(k)}(x) \frac{(n)_m y^m}{(n-jm)!}. \quad (2.6)$$

Proof. By (2.3), we have

$$\begin{aligned} \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \frac{t^n}{n!} &= \frac{qLi_k\left(\frac{1-e^{-qt}}{q}\right)}{1-e^{-qt}} e^{xt} e^{yt^j} \\ &= \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k)}(x) \frac{t^n}{n!} \sum_{n=0}^{\infty} y^n \frac{t^{jn}}{n!} \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^{\lfloor \frac{n}{j} \rfloor} {}_H\mathcal{B}_{n-jm,q}^{(k)}(x) \frac{t^{n-jm}}{(n-jm)!} y^m \frac{t^m}{m!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{m=0}^{\lfloor \frac{n}{j} \rfloor} {}_H\mathcal{B}_{n-jm,q}^{(k)}(x) \frac{(n)_m y^m}{(n-jm)!} \right) \frac{t^n}{n!}, \end{aligned}$$

which implies the desired result (2.6). \square

2.2. The Hermite based λ -Stirling polynomials of the second kind. We introduce the Hermite based λ -Stirling polynomials of the second kind is defined by

$$\sum_{n=0}^{\infty} S_2^{(\lambda,j)}(n,m;x,y) \frac{t^n}{n!} = \frac{(\lambda e^t - 1)^m}{m!} e^{xt+yt^j}. \quad (2.7)$$

Several specific circumstances of ${}_H\mathcal{B}_{n,q}^{(k,j)}(x,y)$ are listed via the following consecutive remarks.

Remark 5. Upon setting $\lambda = 1$, we have $S_2^{(1,j)}(n,m;x,y) := S_2^{(j)}(n,m;x,y)$ called the Hermite based Stirling polynomials of the second kind.

Remark 6. Letting $y = 0$, $S_2^{(\lambda,j)}(n,m;x,y)$ reduces to the $S_2^\lambda(n,m;x)$ called the weighted λ -Stirling numbers of the second kind (see [5]).

Remark 7. When $\lambda = 1$ and $y = 0$, $S_2^{(j)}(n,m;x,y)$ reduces to the $S_2(n,m;x)$ called the weighted Stirling numbers of the second kind (see [5]).

Remark 8. Setting $x = y = 0$, we have $S_2^{(\lambda,j)}(n,m;0,0) := S_2^\lambda(n,m)$ called the familiar λ -Stirling numbers of the second kind (see [5]).

Remark 9. In the case $\lambda = 1$ and $x = y = 0$, we have $S_2^{(1,j)}(n, m; 0, 0) := S_2(n, m)$ called the familiar Stirling numbers of the second kind (see [7], [16]).

We give some relations and properties belonging to the Hermite based λ -Stirling polynomials of the second kind by the following consecutive propositions.

Proposition 5. We have

$$\begin{aligned} S_2^{(\lambda,j)}(n, m; x, y) &= \sum_{l=0}^n \binom{n}{l} S_2^\lambda(l, m) H_{n-l}^{(j)}(x, y) \\ S_2^{(\lambda,j)}(n, m; x, y) &= \sum_{l=0}^n \binom{n}{l} S_2^\lambda(l, m; 0, y) x^{n-l} \\ S_2^{(\lambda,j)}(n, m; x, y) &= \sum_{l=0}^{\lfloor \frac{n}{j} \rfloor} S_2^\lambda(l, m; x, 0) \frac{(n)_l y^l}{(n-jl)!}, \end{aligned}$$

where $(n)_l$ equals to $n(n-1)(n-2)\cdots(n-l+1)$ called the falling factorial function.

Proposition 6. We have

$$S_2^{(q^{-1},j)}(n+1, m+1; x, y) = \frac{1}{q} \sum_{l=0}^n \binom{n}{l} S_2^{q^{-1}}(l, m; x, y).$$

Proposition 7. (Derivative Properties) Each of the following formulas holds true:

$$\frac{\partial}{\partial x} S_2^{(\lambda,j)}(n, m; x, y) = n S_2^{(\lambda,j)}(n-1, m; x, y) \quad \text{and} \quad \frac{\partial}{\partial y} S_2^{(\lambda,j)}(n, m; x, y) = (n)_j S_2^{(\lambda,j)}(n-j, m; x, y),$$

where $(n)_j$ equals to $n(n-1)(n-2)\cdots(n-j+1)$ called the falling factorial function.

Proposition 8. (Integral Representations) The following equalities hold true:

$$\int_v^\mu S_2^{(\lambda,j)}(n, m; x, y) dx = \frac{S_2^{(\lambda,j)}(n+1, m; \mu, y) - S_2^{(\lambda,j)}(n+1, m; v, y)}{n+1}$$

and

$$\int_\gamma^\zeta S_2^{(\lambda,j)}(n, m; x, y) dy = \frac{S_2^{(\lambda,j)}(n+j, m; x, \zeta) - S_2^{(\lambda,j)}(n+j, m; x, \gamma)}{(n+1)^{(j)},}$$

where $(n)^{(j)} = n(n+1)(n+2)\cdots(n+j-1)$ called the rising factorial function.

3. MAIN RESULTS

This part includes our main results.

A correlation including ${}_H\mathcal{B}_{n,q}^{(k,j)}(x, y)$ and $H_n^{(j)}(x, y)$ is given by the following theorem.

Theorem 1. The polynomials ${}_H\mathcal{B}_{n,q}^{(k,j)}(x, y)$ and $H_n^{(j)}(x, y)$ satisfies the following relation:

$$\begin{aligned} &\sum_{m=0}^n \binom{n}{m} {}_H\mathcal{B}_{n-m,q}^{(k,j)}(x, y) q^m - {}_H\mathcal{B}_{n,q}^{(k,j)}(x, y) \\ &= \sum_{m=0}^{\infty} \frac{q^{-m}}{(m+1)^k} \sum_{r=0}^{m+1} \binom{m+1}{r} (-1)^r H_n^{(j)}(x+q-rq, y). \end{aligned} \quad (3.1)$$

Proof. By (2.3), we write

$$\sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x, y) \frac{t^n}{n!} = \frac{q Li_k\left(\frac{1-e^{-qt}}{q}\right)}{e^{qt}-1} e^{(x+q)t+yt^j}.$$

6

Then, we consider that

$$(e^{qt} - 1) \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x, y) \frac{t^n}{n!} = qLi_k \left(\frac{1 - e^{-qt}}{q} \right) e^{(x+q)t+yt^j}. \quad (3.2)$$

LHS of (3.2):

$$\begin{aligned} LHS &= \sum_{n=0}^{\infty} q^n \frac{t^n}{n!} \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x, y) \frac{t^n}{n!} - \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x, y) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{m=0}^n \binom{n}{m} {}_H\mathcal{B}_{n-m,q}^{(k,j)}(x, y) q^m - {}_H\mathcal{B}_{n,q}^{(k,j)}(x, y) \right) \frac{t^n}{n!}. \end{aligned} \quad (3.3)$$

By (2.1), RHS of (3.2):

$$\begin{aligned} RHS &= q \sum_{m=1}^{\infty} \frac{\left(\frac{1-e^{-qt}}{q} \right)^m}{m^k} e^{(x+q)t+yt^j} = q \sum_{m=0}^{\infty} \frac{\left(\frac{1-e^{-qt}}{q} \right)^{m+1}}{(m+1)^k} e^{(x+q)t+yt^j} \\ &= \sum_{m=0}^{\infty} \frac{q^{-m}}{(m+1)^k} \sum_{r=0}^{m+1} \binom{m+1}{r} (-1)^r e^{(x+q-rq)t+yt^j} \\ &= \sum_{m=0}^{\infty} \frac{q^{-m}}{(m+1)^k} \sum_{r=0}^{m+1} \binom{m+1}{r} (-1)^r \sum_{n=0}^{\infty} H_n^{(j)}(x+q-rq, y) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{m=0}^{\infty} \frac{q^{-m}}{(m+1)^k} \sum_{r=0}^{m+1} \binom{m+1}{r} (-1)^r H_n^{(j)}(x+q-rq, y) \right) \frac{t^n}{n!}. \end{aligned}$$

LHS and RHS yield to the desired result (3.1). \square

A correlation between ${}_H\mathcal{B}_{n,q}^{(k,j)}(x, y)$ and $S_2^{(j)}(n, m; x, y)$ is stated in the following theorem.

Theorem 2. *The following correlation holds true:*

$$\begin{aligned} &\sum_{m=0}^n \binom{n}{m} {}_H\mathcal{B}_{n-m,q}^{(k,j)}(x, y) q^m - {}_H\mathcal{B}_{n,q}^{(k,j)}(x, y) \\ &= \sum_{m=0}^{\infty} \frac{(-1)^{n+m+1}}{(m+1)^k} q^{n-m} (m+1)! S_2^{(j)} \left(n, m; -\frac{x}{q} - 1, (-1)^j \frac{y}{q^j} \right). \end{aligned} \quad (3.4)$$

Proof. By (2.3), we write

$$(e^{qt} - 1) \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x, y) \frac{t^n}{n!} = qLi_k \left(\frac{1 - e^{-qt}}{q} \right) e^{(x+q)t+yt^j}.$$

Using the series manipulation procedure, LHS of (2.7):

$$LHS = \sum_{n=0}^{\infty} \left(\sum_{m=0}^n \binom{n}{m} {}_H\mathcal{B}_{n-m,q}^{(k,j)}(x, y) q^m - {}_H\mathcal{B}_{n,q}^{(k,j)}(x, y) \right) \frac{t^n}{n!}.$$

By (3.2), RHS of (2.7):

$$\begin{aligned}
 RHS &= q \sum_{m=1}^{\infty} \frac{\left(\frac{1-e^{-qt}}{q}\right)^m}{m^k} e^{(x+q)t+yt^j} = q \sum_{m=0}^{\infty} \frac{\left(\frac{1-e^{-qt}}{q}\right)^{m+1}}{(m+1)^k} e^{(x+q)t+yt^j} \\
 &= \sum_{m=0}^{\infty} \frac{(-1)^{m+1}}{(m+1)^k} q^{-m} (e^{-qt} - 1)^{m+1} e^{(x+q)t+yt^j} \\
 &= \sum_{m=0}^{\infty} \frac{(-1)^{m+1}}{(m+1)^k} q^{-m} (m+1)! \sum_{n=0}^{\infty} S_2^{(j)} \left(n, m; -\frac{x}{q} - 1, (-1)^j \frac{y}{q^j} \right) \frac{(-qt)^n}{n!} \\
 &= \sum_{n=0}^{\infty} \left(\sum_{m=0}^{\infty} \frac{(-1)^{n+m+1}}{(m+1)^k} q^{-m} (m+1)! S_2^{(j)} \left(n, m; -\frac{x}{q} - 1, (-1)^j \frac{y}{q^j} \right) \right) \frac{t^n}{n!}.
 \end{aligned}$$

LHS and RHS gives the desired result (3.3). \square

From (2.2), we readily derive that

$$\frac{d}{dt} Li_k(f(t)) = \frac{f'(t)}{f(t)} Li_{k-1}(f(t)). \quad (3.5)$$

An relation for the Hermite based poly-Bernoulli polynomials with a q parameter is given by the following theorem.

Theorem 3. *We have*

$$\begin{aligned}
 &\sum_{m=0}^n \binom{n}{m} {}_H\mathcal{B}_{n+1-m,q}^{(k,j)}(x,y) q^m - {}_H\mathcal{B}_{n+1,q}^{(k,j)}(x,y) \\
 &= q {}_H\mathcal{B}_{n,q}^{(k-1,j)}(x,y) + x \sum_{m=0}^n \binom{n}{m} {}_H\mathcal{B}_{n-m,q}^{(k,j)}(x,y) q^m - (x+q) {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \\
 &\quad + yj \frac{(n-j+1)!}{n!} \left(\sum_{m=0}^{n-j+1} \binom{n-j+1}{m} {}_H\mathcal{B}_{n-j+1-m,q}^{(k,j)}(x,y) q^m - {}_H\mathcal{B}_{n+j-1,q}^{(k,j)}(x,y) \right). \quad (3.6)
 \end{aligned}$$

Proof. In the light of (3.5), we get

$$\frac{d}{dt} Li_k \left(\frac{1-e^{-qt}}{q} \right) = \frac{q}{e^{qt}-1} Li_{k-1} \left(\frac{1-e^{-qt}}{q} \right).$$

Differentiate both sides of (3.2) with respect to t , we derive

$$\begin{aligned}
 \frac{d}{dt} \left(e^{qt} \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \frac{t^n}{n!} - \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \frac{t^n}{n!} \right) &= q e^{qt} \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \frac{t^n}{n!} \\
 &\quad + e^{qt} \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n+1,q}^{(k,j)}(x,y) \frac{t^n}{n!} - \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n+1,q}^{(k,j)}(x,y) \frac{t^n}{n!}
 \end{aligned}$$

8

and

$$\begin{aligned}
& \frac{d}{dt} \left(qLi_k \left(\frac{1 - e^{-qt}}{q} \right) e^{(x+q)t+yt^j} \right) \\
&= \frac{q^2}{e^{qt} - 1} Li_{k-1} \left(\frac{1 - e^{-qt}}{q} \right) e^{(x+q)t+yt^j} \\
&\quad + qLi_k \left(\frac{1 - e^{-qt}}{q} \right) \left[(x+q) e^{(x+q)t+yt^j} + yjt^{j-1} e^{(x+q)t+yt^j} \right] \\
&= \frac{q^2}{e^{qt} - 1} Li_{k-1} \left(\frac{1 - e^{-qt}}{q} \right) e^{(x+q)t+yt^j} \\
&\quad + e^{qt} (x+q) \frac{qLi_k \left(\frac{1 - e^{-qt}}{q} \right)}{e^{qt} - 1} e^{(x+q)t+yt^j} - (x+q) \frac{qLi_k \left(\frac{1 - e^{-qt}}{q} \right)}{e^{qt} - 1} e^{(x+q)t+yt^j} \\
&\quad + e^{qt} yjt^{j-1} \frac{qLi_k \left(\frac{1 - e^{-qt}}{q} \right)}{e^{qt} - 1} e^{(x+q)t+yt^j} - yjt^{j-1} \frac{qLi_k \left(\frac{1 - e^{-qt}}{q} \right)}{e^{qt} - 1} e^{(x+q)t+yt^j} \\
&= q \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k-1,j)}(x,y) \frac{t^n}{n!} \\
&\quad + (x+q) \sum_{n=0}^{\infty} q^n \frac{t^n}{n!} \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \frac{t^n}{n!} - (x+q) \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \frac{t^n}{n!} \\
&\quad + yjt^{j-1} \sum_{n=0}^{\infty} q^n \frac{t^n}{n!} \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \frac{t^n}{n!} - yjt^{j-1} \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \frac{t^n}{n!} \\
&= q \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k-1,j)}(x,y) \frac{t^n}{n!} + (x+q) \sum_{n=0}^{\infty} \left(\sum_{m=0}^n \binom{n}{m} {}_H\mathcal{B}_{n-m,q}^{(k,j)}(x,y) q^m - {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \right) \frac{t^n}{n!} \\
&\quad + yj \sum_{n=0}^{\infty} \left(\sum_{m=0}^n \binom{n}{m} {}_H\mathcal{B}_{n-m,q}^{(k,j)}(x,y) q^m - {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \right) \frac{t^{n+j-1}}{n!},
\end{aligned}$$

which gives the desired result (3.6). □

Here, we give the following theorem.

Theorem 4. Let $n \in \mathbb{N}_0$ and $k \in \mathbb{N}$. We then have

$${}_H\mathcal{B}_{n,q}^{(-k,j)}(x,y) = \sum_{m=0}^{\min(n,k)} (m!)^2 S_2^{(j)} \left(n, m; \frac{x}{q} + 1, \frac{y}{q^j} \right) q^n S_2^{q^{-1}}(k, m; 1).$$

Proof. By inspiring the proof given by Cenkci and Komatsu [5], by (2.3), we write

$$\sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(-k,j)}(x,y) \frac{t^n}{n!} = \frac{q}{1 - e^{-qt}} \sum_{m=0}^{\infty} (m+1)^k \left(\frac{1 - e^{-qt}}{q} \right)^{m+1} e^{xt+yt^j}.$$

Then, using (2.7), for $|z| < 1$, we obtain

$$\begin{aligned}
& \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(-k,j)}(x,y) \frac{t^n}{n!} \frac{z^k}{k!} \\
&= \frac{q}{1-e^{-qt}} \sum_{k=0}^{\infty} \sum_{m=0}^{\infty} (m+1)^k \left(\frac{1-e^{-qt}}{q} \right)^{m+1} \frac{z^k}{k!} e^{xt+yt^j} \\
&= \frac{q}{1-e^{-qt}} \sum_{m=0}^{\infty} \left(\frac{1-e^{-qt}}{q} \right)^{m+1} e^{(m+1)z} e^{xt+yt^j} \\
&= \frac{q}{1-e^{-qt}} \sum_{m=0}^{\infty} \left(\frac{e^z(1-e^{-qt})}{q} \right)^{m+1} e^{xt+yt^j} \\
&= \frac{qe^{z+qt}}{q-(e^{qt}-1)(e^z-q)} e^{xt+yt^j} \\
&= \sum_{m=0}^{\infty} e^{(x+q)t+yt^j} (e^{qt}-1)^m e^z (q^{-1}e^z-1)^m \\
&= \sum_{m=0}^{\infty} \left[\sum_{n=0}^{\infty} m! S_2^{(j)} \left(n, m; \frac{x}{q} + 1, \frac{y}{q^j} \right) q^n \frac{t^n}{n!} \right] \cdot \left[m! \sum_{k=0}^{\infty} S_2^{q^{-1}}(k, m; 1) \frac{z^k}{k!} \right] \\
&= \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} \left(\sum_{m=0}^{\infty} (m!)^2 S_2^{(j)} \left(n, m; \frac{x}{q} + 1, \frac{y}{q^j} \right) q^n S_2^{q^{-1}}(k, m; 1) \right) \frac{t^n}{n!} \frac{z^k}{k!},
\end{aligned}$$

which finalize this theorem. \square

A correlation including $S_2^{(j)}(n, m; x, y)$ and ${}_H\mathcal{B}_{n,q}^{(k,j)}(x, y)$ is given by the following theorem.

Theorem 5. *The following holds true:*

$${}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) = q \sum_{u=0}^n \binom{n}{u} \sum_{s=0}^{\infty} s^u \sum_{m=0}^{\infty} \frac{(m+1)!}{(m+1)^k} S_2^{(j)} \left(n-u, m+1; -\frac{x}{q}, (-1)^j \frac{y}{q^j} \right) (-q)^{n-m-1}. \quad (3.7)$$

Proof. From (2.3) and utilizing (2.7), with the series manipulation procedure, we acquire

$$\begin{aligned}
\sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \frac{t^n}{n!} &= \frac{q}{1-e^{-qt}} \sum_{m=0}^{\infty} \frac{(-q)^{-m-1}}{(m+1)^k} (e^{-qt}-1)^{m+1} e^{xt+yt^j} \\
&= q \sum_{s=0}^{\infty} \sum_{u=0}^{\infty} \frac{(-sqt)^u}{u!} \sum_{m=0}^{\infty} \frac{(-q)^{-m-1}}{(m+1)^k} (e^{-qt}-1)^{m+1} e^{xt+yt^j} \\
&= q \sum_{s=0}^{\infty} \sum_{u=0}^{\infty} (-sq)^u \frac{t^u}{u!} \sum_{m=0}^{\infty} \frac{(-q)^{-m-1} (m+1)!}{(m+1)^k} \\
&\quad \cdot \sum_{n=0}^{\infty} S_2^{(j)} \left(n, m+1; -\frac{x}{q}, (-1)^j \frac{y}{q^j} \right) (-q)^n \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \left(q \sum_{s=0}^{\infty} (-sq)^n \sum_{m=0}^{\infty} \frac{(-q)^{-m-1} (m+1)!}{(m+1)^k} \right) \frac{t^n}{n!} \\
&\quad \cdot \sum_{n=0}^{\infty} S_2^{(j)} \left(n, m+1; -\frac{x}{q}, (-1)^j \frac{y}{q^j} \right) (-q)^n \frac{t^n}{n!},
\end{aligned}$$

which implies the desired result (3.7). \square

We give the following theorem.

Theorem 6. *The following relation is valid:*

$${}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) = \sum_{s=0}^{\infty} \sum_{m=0}^{\infty} \frac{q^{-m}}{(m+1)^k} \sum_{l=0}^{m+1} \binom{m+1}{l} (-1)^l H_n^{(j)}(x-lq-sq,y). \quad (3.8)$$

Proof. By means of (2.1), (2.3) and (2.7), we derive

$$\begin{aligned} \sum_{n=0}^{\infty} {}_H\mathcal{B}_{n,q}^{(k,j)}(x,y) \frac{t^n}{n!} &= \frac{q}{1-e^{-qt}} \sum_{m=0}^{\infty} \frac{(-q)^{-m-1}}{(m+1)^k} (e^{-qt}-1)^{m+1} e^{xt+yt^j} \\ &= \sum_{s=0}^{\infty} \sum_{m=0}^{\infty} \frac{q^{-m}}{(m+1)^k} \sum_{l=0}^{m+1} \binom{m+1}{l} (-1)^l e^{(x-lq-sq)t+yt^j} \\ &= \sum_{n=0}^{\infty} \left(\sum_{s=0}^{\infty} \sum_{m=0}^{\infty} \frac{q^{-m}}{(m+1)^k} \sum_{l=0}^{m+1} \binom{m+1}{l} (-1)^l H_n^{(j)}(x-lq-sq,y) \right) \frac{t^n}{n!}. \end{aligned}$$

Thus, we arrive at the desired identity (3.8). \square

References

- [1] Appell, P., Kampé de Fériet, J., Fonctions hypergéométriques et hypersphériques. Polyômes d'Hermite, *Gauthier-Villars*, Paris, 1926.
- [2] Araci, S., Khan, W.A., Acikgoz, M., Özel, C., Kumam, P. A new generalization of Apostol type Hermite-Genocchi polynomials and its applications, *SpringerPlus*, 2016, doi. 10.1186/s40064-016-2357-4.
- [3] Bayad, A., Hamahata, Y. Polylogarithms and poly-Bernoulli polynomials, *Kyushu J. Math.*, 65 (2011), 15-24.
- [4] Bretti, G., Ricci, P., E. Multidimensional extensions of the Bernoulli and Appell polynomials, *Taiwanese J. Math.*, 8(3) (2004), 415-428.
- [5] Cenkci, M., Komatsu, T. Poly-Bernoulli numbers and polynomials with a q parameter, *J. Number Theory*, 152 (2015), 38-54.
- [6] Dattoli, G., Lorenzutta S., Cesarano, C. Finite sums and generalized forms of Bernoulli polynomials, *Rend. Math. Appl.*, 19 (1999), 385-391.
- [7] Khan, W., A., Khan, N., U., Zia, S. A note on Hermite poly-Bernoulli numbers and polynomials of the second kind, *Turkish J. Anal. Number Theory*, 3(5) (2015), 120-125.
- [8] Kurt, B., Simsek, Y. On the Hermite based Genocchi polynomials, *Adv. Stud. Contemp. Math.*, 23 (2013), 13-17.
- [9] Kurt, B. Identities and Relation on the Poly-Genocchi Polynomials with a q -Parameter, *J. Inequal. Spec. Funct.*, (to appear).
- [10] Mahmudov, N.I. On a class of q -Benoulli and q -Euler polynomials, *Adv. Difference Equ.*, 2013, doi: 10.1186/1687-1847-2013-108.
- [11] Ozarslan, M.A. Hermite-based unified Apostol-Bernoulli, Euler and Genocchi polynomials, *Adv. Difference Equ.*, 2013, doi: 10.1186/1687-1847-2013-116.
- [12] Pathan, M., A. A new class of generalized Hermite-Bernoulli polynomials, *Georgian Math. J.*, 19 (2012), 559-573.
- [13] Pathan, M., A., Khan, W., A. Some implicit summation formulas and symmetric identities for the generalized Hermite-Bernoulli polynomials. *Mediterr. J. Math.*, 12 (2015) 679-695.
- [14] Pathan, M., A., Khan, W., A. A new class of generalized polynomials associated with Hermite and Bernoulli polynomials, *Le Matematiche*, LXX (2015), 53-70
- [15] Srivastava, H.M. Some generalizations and basic (or q -) extensions of the Bernoulli, Euler and Genocchi polynomials, *Appl. Math. Inform. Sci.*, 5 (2011), 390-444.
- [16] Srivastava, H.M., Choi, J. Zeta and q -Zeta functions and associated series and integrals; *Elsevier Science Publishers*: Amsterdam, The Netherlands, 2012, 674 p.
- [17] Widder, D., V. The Heat Equation, *Academic Press*, New York, 1975.